

A Cross-sectional, Multicenter, Observational Study to Assess the Prophylaxis of Venous Thromboembolism in Lebanese and Jordanian Hospitals

Imad Hajj (✉ Ahmed.elgebaly@ray-cro.com)

Saint George's Hospital <https://orcid.org/0000-0002-8324-0960>

Mahmoud AL -Masri (✉ malmasri@khcc.jo)

King Hussein Cancer Center

Kaldoun Bashaireh

Jordan University of Science and Technology Faculty of Medicine

Mohammed Bani Hani

Jordan University of Science and Technology Faculty of Medicine

Shadi Hamouri

Jordan University of Science and Technology Faculty of Medicine

Joe Khouzami

Sanofi

Nisrine Sabra

Sanofi

Chahine Fadel

Sanofi

Research

Keywords: Venous thromboembolism, VTE Prophylaxis, ACCP guideline

Posted Date: July 20th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42401/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on February 10th, 2021. See the published version at <https://doi.org/10.1186/s12959-021-00261-2>.

Abstract

Background

There is a growing body of evidence showing substantial underuse of appropriate venous thromboembolism (VTE) prophylaxis in patients at risk. In the present study, our goal was to assess the current practices in the use rate of VTE prophylaxis among hospitalized patients in Jordan and Lebanon.

Methods

A cross-sectional, multicenter, observational study was conducted on 40 centers across Lebanon and Jordan. We included patients who were admitted to the participating hospitals for the treatment of a serious medical or surgical illness. The patients' records were screened for the fulfillment of inclusion/exclusion criteria during a single assessment visit. The proportion of medical and surgical patients who were at risk of VTE and the thrombo-prophylactic measures employed by physicians for these patients were assessed according to the American College of Chest Physicians (ACCP 2016) guidelines.

Results

The present study included 704 patients (400 from Jordan and 304 from Lebanon) with a mean age of 54.9 ± 17.5 years. Almost 59% of the patients received prophylaxis treatment in form of pharmacological anticoagulant prophylaxis and/or mechanical prophylaxis. Low molecular weight heparin was the most commonly used anticoagulant for VTE prophylaxis in 366 out of the total 704 (51.9%) patients in the analysis cohort. Two hundred and sixteen patients (52%, 95% confidence interval [47.1% – 56.9%]) received appropriate prophylactic agents out of 415 patients who were eligible for prophylaxis according to the ACCP 2016 guidelines. On the other hand, 199 (72.1%, 95% confidence interval [66.4% – 77.3%]) patients received prophylaxis out of 276 ineligible patients.. The rate of compliance to guidelines showed wide variations according to the type of hospital, specialty, and the patients' age. The multivariate logistic regression analysis showed that only age was a significant predictor of appropriate VTE prophylaxis (odds ratio [OR] 1.05, $P < 0.001$).

Conclusion

The rates of the appropriate use of VTE prophylaxis are low in Lebanon and Jordan. There is a lack of compliance to guidelines for VTE prophylaxis use for hospitalized patients in both countries.

1. Introduction

Venous thromboembolism (VTE) is a life-threatening disorder and a major cause of morbidities and mortality among hospitalized patients. The condition is characterized by the development of thrombosis of deep veins of the leg or pelvis (DVT) that propagate to the pulmonary circulation leading to pulmonary embolism (PE)(1). According to previous epidemiological studies, VTE is the s the third most common cardiovascular disease with reported incidence rates of 130 and 100 per 100,000 persons every year for men and women, respectively(2). Old age, obesity, prolonged immobility, acute heart failure, malignancy, hyperestrogenemia, and genetic susceptibility are major risk factors for VTE development(3). Hospitalized patients are at increased risks of VTE due to the presence of multiple risk factors that are usually cumulative. Despite that VTE has been previously thought as a complication of major surgery, recently published evidence shows that hospitalized patients with the medical illness have comparable risks of VTE to that of the patients undergoing major surgery(4). Previous reports demonstrated that up to 20% of hospitalized medical patients are expected to develop VTE during hospital stay(5). VTE is a major cause of mortality in hospitalized patients as well, up to 10% of fatality cases during hospitalization were attributed to VTE in autopsy-based studies(6). Therefore, effective VTE prophylaxis among hospitalized at-risk patients, either surgical or medical, is critical in improving patients' outcomes and survival.

Anticoagulants represent the cornerstone for the VRE prophylaxis among hospitalized patients. Historically, unfractionated heparin (UFH) was the anticoagulant of choice for VTE prophylaxis; however, the current body of evidence shows that low molecular weight heparin (LMWH) has comparable efficacy profile, with more favorable safety data than UFH(7). Therefore, LMWH and fondaparinux are currently the first-line prophylaxis options in hospitalized patients(8). Other VTE prophylaxis strategies include graduated compression stockings (GCSs) and intermittent pneumatic compression (IPC), with equivocal data regarding their efficacy(9).

Although multiple international guidelines recommend the use of VTE prophylaxis for at-risk hospitalized patients, the use rates of prophylaxis strategies are substantially low in many regions. The ENDORSE study reported that less than 40% of hospitalized medical patients receiving appropriate prophylaxis, with great variations in prophylaxis use between countries, regions, and hospitals.

Therefore, in this study, we aimed to assess the current practices of VTE prophylaxis among hospitalized patients in Jordan and Lebanon.

2. Patient And Methods

We followed the STROBE (Strengthening the Reporting of Observational Studies in Epidemiology Statement) guidelines (Supplementary file no.1) during the preparation of this manuscript(10). The present study run in concordance with the principles of the declaration of Helsinki and applicable local regulatory laws. Written informed consent was obtained from every eligible patient, or their relatives, prior to the study's enrollment.

2.1. Study design and Setting

We conducted a multicenter, observational, cross-sectional study across Lebanon and Jordan hospitals. Participating physicians from 40 centers across both counties were randomly selected and asked to recruit eligible patients consecutively. The selection process of the participating physicians was stratified according to the type of hospital, specialties, and geographical area.

2.2. Participants

Adults' patients (≥ 40 years old for medical patients and ≥ 18 years old for surgical patients) of both sexes, who were admitted to any of the participating centers for serious medical illness or surgical indication, were included. We excluded patients with current or recent (one month before the study) deep venous thrombosis or pulmonary embolism, history of intake of anticoagulant for another co-morbidity in the last month before study's enrollment, weight below 40 kg or above 100 kg, impaired kidney or liver functions, concomitant participation to a clinical, or missing hospital chart. Pregnant or lactating women were excluded as well.

2.3. Sample size calculation and sampling methods:

Under the assumption that almost 40% of the eligible patients for VTE prophylaxis are actually receiving appropriate prophylaxis treatment globally, the calculated sample size for the present study was 700 patients within a margin of error of about 3.6% and 95% confidence interval. This assumes that half ($N = 350$) of those 700 recruited patients for the study would be eligible for VTE prophylaxis, a conservative assumption based on the results of AVAIL ME study¹ and its extension(11, 12).

The present study utilized a non-probability, consecutive, sampling technique to recruit eligible patients. Investigators were asked to recruit consecutive eligible patients and each selected physician included about 10 to 20 patients who met the inclusion criteria per consulting session.

2.4. Data collection and study's outcomes

The following data were collected from every eligible patient: demographic characteristics, anthropometric measures (weight, height, and body mass index [BMI]), cause of hospital admission, length of hospital stay, type and duration surgery for surgical patients, risk factors for VTE, type and frequency of VTE prophylaxis, and eligibility for VTE prophylaxis according to American College of Chest Physicians (ACCP 2016) guidelines(13).

The primary outcome of the present study was the percentage of patients receiving appropriate VTE prophylaxis among the patients eligible for prophylaxis. The secondary outcomes were the rate of patients receiving prophylaxis treatment without being eligible for such treatment as per the guideline, the profile of the medically ill and surgical hospitalized patients and their VTE risk, the compliance rate of VTE prophylaxis in the different specialties/type of surgery, VTE prophylaxis differences between different regions and geographies, and the predictive factors or barriers for appropriate prophylaxis.

2.5. Statistical methods

All variables recorded during the study were summarized. Frequencies and percentages (with 95% confidence interval [CI] for the primary endpoint) were provided for categorical variables. Mean and standard deviation were provided for continuous variables. The analysis was stratified by type of patient (surgical vs. medical), type of surgery, country, and specialty of the doctors. Multivariate logistic regression models were conducted using the baseline factors to assess their effect as predictors of the appropriate use of VTE prophylaxis. A p-value of less than 5% was considered statistically significant. All statistical tests were performed using SPSS program version 25 (IBM, Armonk, NY, USA).

Results

The present study screened 705 patients from 40 sites across Lebanon and Jordan. Of them, 704 patients (400 from Jordan and 304 from Lebanon) (99.9%) were eligible for final analysis and one patient (0.1%) was excluded as the weight was above 100 kg. Regarding the demographic characteristics of the included patients, the patients' age ranged from 18 to 93 years with a mean age of 54.9 ± 17.5 years. Almost 48% of the patients were males. The mean weight and height of the included patients were 75.7 ± 13.4 kg and 165.9 ± 9.3 cm, respectively. The BMI of the included patients ranged from 16.1 to 44.2 kg/m^2 with a mean BMI of $27.5 \pm 4.8 \text{ kg/m}^2$. In terms of vital signs, the mean SBP and DBP of the included patients were 124.1 ± 15.4 and 73.8 ± 9.7 mmHg, respectively. Sixty percent of the patients had one or more current medical conditions. The most commonly encountered medical condition was hypertension (34.8%), followed by diabetes mellitus (22.6%) and coronary artery disease (10.1%). Two-hundred and forty-six (34.9%) patients were admitted for medical causes only, 449 (63.8%) patients were admitted for surgical causes only, and 9 (1.3%) patients were admitted for both medical and surgical causes. The average hospital stay of the included patients was 5.8 ± 8.4 days. The Demographic and clinical characteristics of the included patients were summarized in Table 1.

Table 1
Demographic and clinical characteristics of the participants at baseline

Variables	Patients (N = 704)
Age in years , mean (SD)	54.96 (17.5)
Male , No. (%)	340 (48.3%)
Weight in kg , mean (SD)	75.72 (13.4)
Height in cm , mean (SD)	165.96 (9.3)
BMI in kg/m² , mean (SD)	27.47 (4.8)
Systolic blood pressure in mmHg , mean (SD)	124.1 (15.4)
Diastolic blood pressure in mmHg , mean (SD)	73.75 (9.7)
Pulse rate in beat/min , mean (SD)	81.38 (12.5)
Temperature in C , mean (SD)	36.87 (0.46)
Hypertension , No. (%)	245 (34.8)
Diabetes , No. (%)	159 (22.6)
History of other chronic illness , No. (%) *	503 (71.4)
Cause of admission , No. (%)	246 (34.94%)
- Medical Causes	449 (63.78%)
- Surgical Causes	9 (1.28%)
- Medical and Surgical Causes	
Type of Surgery , No. (%)	327 (46.45%)
- Open	131 (18.61%)
- Laparoscopic	
Hospital Stay in days , mean (SD)	5.83 (8.4)
Risk factors for VTE , No. (%)	391 (55.5%)
- Surgical risk	221 (31.4%)
- Medical risk	4 (0.6%)
- Medical and surgical risk	

Variables	Patients (N = 704)
Surgical risk factors for VTE, No. (%)	237 (33.7)
- Obesity	117 (16.7%)
- Age > 60 years old	59 (8.38)
- Laparoscopic Surgery (> 45 min)	523 (74.3)
- Other factors*	
Degree of surgical risk factors for VTE, No. (%)	162 (23.1)
- Low	109 (15.48)
- Moderate	86 (12.22)
- High	38 (5.4)
- Highest	
Medical risk factors for VTE, No. (%)	94 (13.35)
- Elderly Age (\geq 70 Years)	93 (13.2)
- Reduced Mobility (At Least 72 h)	86 (12.22)
- Obesity	181 (25.7%)
- Other factors	
Risk factors associated with increased bleeding, No. (%)	60 (8.5)
Risk factors associated with Mechanical Prophylaxis	25 (3.6%)
<i>*Patient may have more than one chronic condition</i>	
<i>VTE; venous thromboembolism, SD; standard deviation</i>	

Six hundred and sixteen patients (87.5%) had one or more risk factors for VTE which were either surgical (55.5%), medical (31.4%), or surgical and medical risk factors (0.6%). Sixty patients (8.5%) had risk factors associated with increased bleeding such as active bleeding (1.1%) and low platelet count. In addition, 25 (3.5%) patients had risk factors associated with mechanical prophylaxis which were severe peripheral arterial disease (0.4%), congestive heart failure (2.8%), and acute superficial/deep vein thrombosis (0.3%).

Among the 704 patients who were eligible for the final analysis, 415 (58.9%) patients received prophylaxis treatment in form of pharmacological anticoagulant prophylaxis (n = 371, 52.7%), mechanical prophylaxis (n = 13, 1.8%), and pharmacological plus mechanical prophylaxis (n = 31, 4.4%). LMWH was the most commonly used anticoagulant for VTE prophylaxis (n = 366); however, the unfractionated heparin was administrated in 56 patients only as seen in Table 2. In Lebanon as well as

Jordan, LMWH was the most commonly used anticoagulant for VTE prophylaxis (N = 192, 48%) and (N = 174, 57.2%), respectively (Table 3).

Table 2
VTE prophylaxis/treatment of the included patients

Variables	Patients (N = 704)
No. of patients received prophylaxis, No. (%)	415 (58.9)
- Pharmacological anticoagulant prophylaxis	371 (52.7)
- Mechanical prophylaxis	13 (1.8)
- Pharmacological and Mechanical prophylaxis	31 (4.4)
Types of anticoagulant, No. (%)	366 (51.9)
- LMWH	56 (7.9)
- Unfractionated Heparin	
Timing of anticoagulants for surgical patients, No. (%)	138 (59.23)
- Preoperatively	94 (40.34)
- Postoperatively	1 (0.43)
- Preoperatively/ Postoperatively	
No. of patients continued prophylaxis after discharge, No. (%)	128 (30.84)
Duration of LMWH after discharge, mean (SD)	17.58 (80.9)
Duration of Warfarin after discharge, mean (SD)	28 (16.2)
Concomitant treatment No. (%)	460 (65.34)
<i>LMWH; low molecular weight heparin, SD; standard deviation</i>	

Table 3
VTE prophylaxis differences according to the included countries

Variables, No (%)	Jordan (N = 400)	Lebanon (N = 304)
LMWH	192 (48.0%)	174 (57.2%)
Unfractionated Heparin	30 (7.5%)	28 (9.2%)
Other Anticoagulants	3 (0.8%)	1 (0.3%)
Mechanical Prophylaxis	12 (3.0%)	32 (10.5%)
<i>LMWH; low molecular weight heparin</i>		

Among surgical patients who received anticoagulants (N = 233), 59.2% of them received the drug preoperatively and 40.3% received it postoperatively. Only 6.3% of the patients received mechanical prophylaxis in the form of graduated compression stockings or intermittent pneumatic compression. Almost 31% of the patients continued anticoagulants treatment after discharge in the form of LMWH (85.9%), aspirin (10.2%), warfarin (3.1%), and Fondaparinux (0.8%).

Among the total 704 eligible patients, 415 (58.9%) patients received VTE prophylaxis, while 289 (41.1%) did not received prophylaxis. For those who received VTE prophylactic treatment, 216 (52%, 95% CI [47.1% – 56.9%]) received appropriate prophylactic agents according to ACCP guideline. For those who were not treated with prophylactic agents, 212 (73.4%, 95% CI [67.9% – 78.4%]) were actually eligible for VTE prophylaxis according to ACCP guideline as presented in Table 4.

Table 4
Appropriate VTE prophylaxis according to ACCP guidelines

Variables, No (%)	Eligible for prophylaxis*	Not Eligible for prophylaxis*	Total
Received prophylaxis	216 (30.68%)	199 (28.27%)	415 (58.95%)
Not received prophylaxis	212 (30.11%)	77 (10.94%)	289 (41.05%)
Total	428 (60.79%)	276 (39.21%)	704 (100%)
<i>* According to ACCP guidelines</i>			

About 60.9% (95% CI 51.9–69.4%) of the patients (n = 78) received VTE prophylaxis out of the medical patients who were eligible for prophylaxis according to ACCP 2016 guideline (n = 128). While in surgical patients, only 45.1% (95% CI 39.3% – 51%) of the patients (n = 133) received VTE prophylaxis out of those who were eligible for prophylaxis (n = 295), Fig. 1. All patients with combined medical and surgical conditions received appropriate prophylaxis.

Regarding the orthopedic surgery, most of the patients received appropriate VTE prophylaxis according to ACCP 2016 guideline. While in non-orthopedic surgery, the number of patients who received appropriate VTE prophylaxis (N = 170) was lower than the number of patients who were eligible for prophylaxis (N = 227) as seen in Fig. 2.

The supplementary file no.2 shows the distribution of appropriate VTE prophylaxis according to doctors' specialty and the type of surgery.

Overall, the rate of compliance to ACCP guidelines was higher in private hospitals than in public hospitals (85% versus 57.8%, respectively). The rate of compliance to ACCP guidelines was higher among oncologists (73.3%) and general family specialists (72%) than other specialized doctors. The rate of compliance was higher among cases of orthopedic surgeries (100%) and oncological surgeries (79%) than other types of surgery as presented in Table 5.

Table 5
Rates of Compliance to ACCP guidelines

Variables, No (%)	Compliant to ACCP guidelines	Not Compliant to ACCP guidelines	Total
Type of hospital,	125 (85.03)	22 (14.97)	147
- Private Hospital	168 (57.8)	22 (11.58)	190
- Public Hospital			
Risk Assessment Method implementation	170 (37.86)	279 (62.14)	449
Age of patients	262 (47.4)	291 (52.6)	553
- ≥ 40 years old	31 (20.5%)	120 (79.5%)	151
- < 40 years old			
Physician Speciality	33 (47.14)	37 (52.86)	70
- Pneumology	32 (40.51)	47 (59.49)	79
- Cardiology	37 (39.78)	56 (60.22)	93
- Internal Medicine	11 (73.33)	4 (26.67)	15
- Oncology	6 (26.09)	17 (73.91)	23
- Infectious Disease	18 (72.0)	7 (28.0)	25
- General/Family Medicine	53 (42.74)	71 (57.26)	124
- Orthopaedics Surgery	24 (60.0)	16 (40.0)	40
- Vascular Surgery	84 (38.53)	134 (61.47)	218
- General Surgery	26 (34.67)	49 (65.33)	75
- Other Surgery			

Variables, No (%)	Compliant to ACCP guidelines	Not Compliant to ACCP guidelines	Total
Type of surgery	16 (100.00%)	0	16
- Hip replacement	19 (100.00%)	0	19
- Knee replacement	21 (100.00%)	0	21
- Hip fracture	0 (0.00%)	6 (100)	6
- Curative arthroscopy	1 (3.03%)	32 (96.97)	33
- Other Ortho trauma	17 (51.52%)	16 (48.48)	33
- Colon /small bowel	7 (63.64%)	4 (36.36)	11
- Rectosigmoid	6 (31.58%)	13 (68.42)	19
- Gastric	30 (45.45%)	36 (54.55)	66
- Hepatobiliary	0 (0.00%)	1 (100)	1
- Vascular	4 (28.57%)	10 (71.43)	14
- Thoracic	15 (78.95%)	4 (21.05)	19
- Oncologic	60 (28.57%)	150 (71.43)	210
- Others			
<i>ACCP; American College of Chest Physicians</i>			

The multivariate logistic regression analysis showed that only age was a significant predictor of appropriate VTE prophylaxis in the present study (OR 1.05, 95% CI [1.04–1.07], P < 0.001) (Table 6).

Table 6
Multivariate logistic regression analysis to identify the significant Predictor Variables for appropriate prophylaxis.

	OR	95% C.I. for OR		P value
		Lower	Upper	
Age	1.05	1.042	1.071	< 0.001*
Gender	1.11	0.764	1.627	0.573
BMI	0.99	0.954	1.033	0.719
SBP	0.98	0.974	1.004	0.145
DBP	1.00	0.983	1.031	0.567
HR	1.01	0.994	1.027	0.226
Temperature	1.13	0.733	1.768	0.564
Hospital Stay	1.02	0.996	1.047	0.094
Included hospitals	0.97	0.918	1.041	0.480
Type of hospital	1.48	0.872	2.505	0.147
Site of hospitals	1.23	0.986	1.558	0.066
Medical Reasons	2.95	0.536	16.281	0.213
Surgical Reasons	0.68	0.125	3.771	0.664
<i>*Patients' age is the only significant predictor for appropriate prophylaxis</i>				
<i>OR; odds ratio, CI; confidence interval</i>				

Discussion

There is a growing body of evidence that shows substantial underuse of VTE prophylaxis in patients at risk(5). However, the extent of VTE prophylaxis underuse in the Middle East is unclear. In the present observational, multinational, study, we found that the rates of the appropriate use of VTE prophylaxis were 60.93% and 45.08% of the medical and surgical hospitalized patients, who were eligible for VTE prophylaxis, respectively. Notably, there were wide variations in the rates of the appropriate use of VTE prophylaxis according to the type of hospitals, geographical areas, and the specialty of treating physicians. Finally, in the present study, we observed that the rate of an appropriate VTE prophylaxis according to the ACCP guidelines was increasing with the age of the patients: patients with old age (≥ 40 years) were associated with a higher rate of compliance to ACCP guidelines than young patients (< 40 years) [47.4% versus 20.5%]. In addition, the multivariate logistic regression analysis showed that only

age was a significant predictor of appropriate VTE prophylaxis in the present study (OR 1.05, 95% CI [1.042–1.07], $P < 0.001$).

In 2017, Levine and colleagues showed that increasing age (OR 0.97, $P < 0.001$) of patients and a primary cardiovascular diagnosis (OR 0.18, $P < 0.001$) (chest pain, congestive heart failure, syncope/near-syncope, chronic ischemic heart disease, sinus tachycardia) decreased the likelihood of VTE prophylaxis(14).

Factors related to underuse included primary cardiovascular diagnosis (OR = 0.18, $P < .0001$) and age (OR = 0.97, $P < .0001$)

The risk of VTE is a 10-fold higher in patients who are hospitalized after trauma, surgery, immobilizing medical illness, or pregnant and puerperal women than the general population. Accordingly, recent clinical guidelines strongly recommend the provision of pharmacological VTE prophylaxis in acutely or critically ill inpatients at risk(13, 15). However, the present study showed that there is substantial underuse of VTE prophylaxis among hospitalized patients in Lebanon and Jordan; less than two-thirds of eligible medical patients received appropriate prophylaxis, while this rate was even much lower in surgical patients. The rate of appropriate VTE prophylaxis apparently varied across different hospitals, geographical areas, and specialties; while the rate of compliance to ACCP guidelines was as low as 26% in some hospitals. These findings reflect the lack of standardized protocols for VTE prophylaxis use for hospitalized patients in both countries. In accordance with our findings, a multicenter study from Lebanon in hospitalized patients reported that the rate of appropriate VTE prophylaxis was 65% of patients at low risk, 30% of patients at moderate risk, and 61% of patients at high risk(16). A more recent report from Jordan reported that the rate of appropriate VTE prophylaxis was 67% in patients admitted to Jordan University Hospital; the study also demonstrated low compliance with the institutional guideline(17). Other reports showed low rates of appropriate VTE prophylaxis use and guidelines compliance in Saudi Arabia(18) and Iran(19). The AVAIL ME study reported that the overall rate of VTE prophylaxis use in the Middle East among eligible patients was 47.8% for medical patients and 60% for surgical patients; while the rate of compliance to ACCP guidelines was only 36%(11). Globally, the multinational multicenter IMPROVE study reported that the appropriate use of VTE prophylaxis in eligible medical patients was 60%(20).

Although the underutilization of VTE prophylaxis in at-risk patients represents a major cause of in-hospital mortality and morbidity, inappropriate use of VTE prophylaxis in low-risk patients, when not medically indicated, can have a negative impact on patients' outcomes leading to bleeding and drugs interaction(21). In the present study, we found that 77.97% of the medical patients, who were not eligible for VTE prophylaxis, received prophylactic agents, compared to 66.88% of the surgical patients who were not eligible for VTE prophylaxis. Similar to our findings, a multicenter study from the United States (US) reported that 77.9% of hospitalized patients received excessive VTE prophylaxis without appropriate indication(22). Another retrospective cohort study reported that pharmacological VTE prophylaxis was present in 74% of low-risk patients, who were not eligible for prophylaxis(21).

In conclusion, the rates of the appropriate use of VTE prophylaxis among hospitalized patients and guideline's compliance are low in Lebanon and Jordan. There are wide variations in the rates of the appropriate use of VTE prophylaxis according to the type of hospitals, geographical areas, and the specialty of treating physicians; therefore, awareness campaigns about the appropriate VTE prophylaxis should be performed.

Lis Of Abbreviations

ACCP: American College of Chest Physicians

BMI: Body mass index

CI: Confidence interval

DVT: Deep venous thrombosis

GCSs: graduated compression stockings

IPC: Intermittent pneumatic compression

LMWH: Low molecular weight heparin

VTE: Venous thromboembolism

UFH: unfractionated heparin

US: United States

Declarations

- **Ethics approval and consent to participate:** The study was approved by responsible ethics committee of health authorities in Jordan and Lebanon. Written informed consent was obtained from every eligible patient, or their relatives, prior to the study's enrollment.
- **Consent for publication:** Not applicable
- **Availability of data and materials:** Not applicable
- **Competing interests:** The authors declare that they have no competing interests
- **Funding:** This study was sponsored and funded by Sanofi. Editorial support was provided by Ahmed Elgebaly MD, of RAY-CRO, and paid for by Sanofi
- **Authors' contributions:** IH developed the study design, shared in data collection, interpreted the data, and revised the manuscript; MA developed the study design, shared in data collection, interpreted the data, and revised the manuscript; KB shared in data collection, analyzed and interpreted the data, and revised the manuscript; MB shared in data collection, analyzed and interpreted the data, and revised the manuscript; SH shared in data collection, analyzed and interpreted the data, and revised

the manuscript; JK shared in data collection and manuscript writing; NS shared in data collection and manuscript writing; CF shared in data collection and manuscript writing

- **Acknowledgements:** The authors thank the study participants, trial staff, and investigators for their participation.

References

1. Martinelli I. Risk factors in venous thromboembolism. In: *Thrombosis and Haemostasis*. 2001. p. 395–403.
2. Tagalakis V, Patenaude V, Kahn SR, Suissa S. Incidence of and mortality from venous thromboembolism in a real-world population: The Q-VTE study cohort. *Am J Med*. 2013;126(9):832.e13–21.
3. Scheres LJJ, Lijfering WM, Cannegieter SC. Current and future burden of venous thrombosis: Not simply predictable. *Res Pract Thromb Haemost*. 2018;2(2):199–208.
4. Guidelines ECP, Geerts WH, Bergqvist D, Pineo GF, Heit J, Samama CM, et al. Prevention of venous thromboembolism: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines (8th Edition). *Chest* [Internet]. 2008;133(6 Suppl):381S – 453S. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18574271>.
5. Cohen AT, Bergmann JF, Lloret-Linares C, Rami A, Garay RP, Kakkar AK, et al. Venous thromboembolism risk and prophylaxis in the acute hospital care setting (ENDORSE study): a multinational cross-sectional study. *Lancet*. 2008;37:387–94.
6. Heriot GS, Pitman AG, Gonzales M, McKelvie P. The four horsemen: Clinicopathological correlation in 407 hospital autopsies. *Intern Med J*. 2010;40(9):626–32.
7. Mismetti P, Laporte-Simitsidis S, Tardy B, Cucherat M, Buchmüller A, Juillard-Delsart D, et al. Prevention of venous thromboembolism in internal medicine with unfractionated or low-molecular-weight heparins: A meta-analysis of randomised clinical trials. *Thromb Haemost*. 2000;83(1):14–9.
8. Lloyd NS, Douketis JD, Moinuddin I, Lim W, Crowther MA. Anticoagulant prophylaxis to prevent asymptomatic deep vein thrombosis in hospitalized medical patients: A systematic review and meta-analysis. *J Thromb Haemost*. 2008;Vol. 6:405–14.
9. Dobromirski M, Cohen AT. How I manage venous thromboembolism risk in hospitalized medical patients. *Blood*. 2012;Vol. 120:1562–9.
10. Elm E, Von, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP, et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for reporting observational studies *. *Int J Surg*. 2014;12(12):1495–9.
11. Taher AT, Aoun J, Salameh P. The AVAIL ME study: A multinational survey of VTE risk and prophylaxis. *J Thromb Thrombolysis*. 2011;31(1):47–56.
12. Mokhtari M, Salameh P, Kouчек M, Kashani BS, Taher A, Waked M. The AVAIL ME Extension: A multinational Middle Eastern survey of venous thromboembolism risk and prophylaxis. *J Thromb*

Haemost. 2011;9(7):1340–9.

13. Kearon C, Akl EA, Ornelas J, Blaivas A, Jimenez D, Bounameaux H, et al. Antithrombotic Therapy for VTE Disease: CHEST Guideline and Expert Panel Report. *Chest* [Internet]. 2016;149(2):315–52. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26867832>.
14. Levine RL, Hergenroeder GW, Miller CC, Davies A. Venous thromboembolism prophylaxis in emergency department admissions. *J Hosp Med*. 2007 Mar;2(2):79–85.
15. Schünemann HJ, Cushman M, Burnett AE, Kahn SR, Beyer-Westendorf J, Spencer FA, et al. American Society of Hematology 2018 guidelines for management of venous thromboembolism: prophylaxis for hospitalized and nonhospitalized medical patients. Vol. 2, *Blood advances*. 2018. p. 3198–225.
16. Zeitoun AA, Dimassi HI, El Kary DY, Akel MG. An evaluation of practice pattern for venous thromboembolism prevention in Lebanese hospitals. *J Thromb Thrombolysis*. 2009;28(2):192–9.
17. Gharaibeh L, Albsoul-Younes A, Younes N. Evaluation of venous thromboembolism prophylaxis after the introduction of an institutional guideline: Extent of application and implementation of its recommendations. *J Vasc Nurs*. 2015;33(2):72–8.
18. Abba AA, Al Ghonaim MA, Rufai AM. Physicians' practice for prevention of venous thromboembolism in medical patients. *J Coll Physicians Surg Pakistan*. 2004;14(4):211–4.
19. Mokhtari M, Attarian H, Norouzi M, Kouchek M, Kashani BS, Sirati F, et al. Venous thromboembolism risk assessment, prophylaxis practices and interventions for its improvement (AVAIL-ME Extension Project, Iran). *Thromb Res*. 2014;133(4):567–73.
20. Tapson VF, Decousus H, Pini M, Chong BH, Froehlich JB, Monreal M, et al. Venous thromboembolism prophylaxis in acutely ill hospitalized medical patients: Findings from the international medical prevention registry on venous thromboembolism. *Chest*. 2007;132(3):936–45.
21. Pavon JM, Sloane RJ, Pieper CF, Colón-Emeric CS, Cohen HJ, Gallagher D, et al. Poor adherence to risk stratification guidelines results in overuse of venous thromboembolism prophylaxis in hospitalized older adults. *J Hosp Med*. 2018.
22. Grant PJ, Conlon A, Chopra V, Flanders SA. Use of Venous Thromboembolism Prophylaxis in Hospitalized Patients. *JAMA Intern Med*. 2018;178(8):1122.

Supplementary Information Note

The two Supplementary Files were omitted in this version

Figures




Figure 1

VTE prophylaxis out of the patients eligible for prophylaxis according to ACCP guidelines. ACCP; American College of Chest Physicians, VTE; venous thromboembolism




Figure 2

VTE prophylaxis in surgical patients out of the patients eligible for prophylaxis according to ACCP guidelines. ACCP; American College of Chest Physicians, VTE; venous thromboembolism