Cross-Cultural Adaptation, Translation, and Moroccan Validation of CareGiver Oncology Quality of Life (CarGoQoL) questionnaire

AHLAM AITOUA (✉ ahlamorim2005@gmail.com)

national cancer institute

Mrabti Hind

Universite Mohammed V de Rabat Faculte de Medecine et de Pharmacie Rabat

Haddou Rahou Bouchra

Universite Ibn Tofail Kenitra Faculte des Sciences

Razine Rachid

Universite Mohammed V de Rabat Faculte de Medecine et de Pharmacie Rabat

Errihani Hassan

Universite Mohammed V de Rabat Faculte de Medecine et de Pharmacie Rabat

Research note

Keywords: Caregivers, palliative cancer ,CarGoQoL ,Reliability, Validity

Posted Date: August 3rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42616/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Objective

The aim of this study is to validate the Moroccan version of The Caregivers Oncology Quality of Life (CarGoQoL) questionnaire to measure the caregiver's quality of life of palliative cancer patients in Morocco.

Results

The psychometric properties of CarGoQoL were generally good. A Cronbach's coefficient of 0.82 was obtained indicating a good internal consistency except for the private life dimension ($\alpha = 0.14$). A good reliability of the test re-test was also observed, with an item internal consistency of 0.9. Missing data and ceiling effect were low; some ceiling effects were observed for Relationship with healthcare (38.9%); correlations were confirmed between the different dimensions that corresponding to the same domain; significant association was identified for caregiver and patient characteristics.

Introduction

Due to the continuing advances in cancer treatment, cancer is likely to increasingly become a chronic disease [1]. Cancer is a catastrophic event for the family and can impose serious stress on patients and their family members [2]. The caregiver accompanies his or her loved one at the time of diagnosis, during treatment and follows the news about the disease's progression. He is afraid of the future and death which are stressful events [3]. Informal caregivers of patients with a life-threatening disease such as cancer are often deeply involved in the patient's disease and provide extraordinary and uncompensated care. The caregivers may take on a range of disease related task and provide emotional support [4–6].

It is recognized that caregiving has negative effects on the caregiver's quality of life (QoL) [7, 8]. The QoL of informal caregivers is an important issue both in the context of active/curative and palliative/supportive treatment [9].

In order to measure the quality of life of caregiver several scientific studies have created and validated different scales.

Recently; a French team has developed the CarGoQoL questionnaire [10] which has already been validated by a study on caregivers of glioma patients [7] and by a validation study of an American version [11]. It has the advantage of a multidimensional approach and is based exclusively on the caregiver's perspective.

In Morocco; no studies have been conducted on the measurement of health-related quality of life among caregivers of palliative cancer patients.[12] The availability of a valid and reliable Moroccan version of quality of life measurement instruments is essential to collect such measures. There are several advantages to culturally adapting an existing instrument rather than composing a new one[13]

Most adaptation processes involve two main steps: the assessment of equivalence between the source and adapted instrument and the assessment of the measurement properties of the adapted instrument [12]. The intercultural adaptation of a quality of life instrument is a step-by-step process [13–16].

In this context; the objective of this study was to carry out a transcultural adaptation and validity of the CarGoQoL questionnaire to evaluate the psychometric properties of the translated version.

Methods

Caregiver characteristics:

For caregiver characteristics; we collected information on: gender, age, marital status and nature of the relationship with the patient and duration of care. For patient we collected information on: sex, age, cancer location, disease duration, stage and Karnovsky Index.

Characteristics of the CarGoQoL original version:

The CarGOQoL is a specific questionnaire that includes 29 questions describing 10 dimensions: psychological well-being, burden, relationship to health care, administration and finance, adaptation, physical well-being, self-esteem, leisure, social support and private life. The CarGOQoL questionnaire gave scores on a scale from 0 to 100, where 0 represents the value of the lowest quality of life and 100 represents the highest quality of life.

Translation and cultural adaptation of the CarGoQoL

The process of cross-cultural adaptation began with a user Agreement reference number # 102016 from the mapi research trust .The CarGoQoL translation guide produced by mapi was used. Reports of the translation process have been prepared and copies are sent to MAPI RESEARCH TRUST.

The methodological process included forward and backward translation. The original version of the CarGoQoL has been translated by two independent translators from French to the Moroccan Arabic dialect. Two other translators unfamiliar with the original version carried out a back-translation of the synthesis of the two translated

versions. The translated and back-translated versions were reviewed by a committee of experts. The final Moroccan Arabic dialect version was drafted after a pre-test in about 20 caregivers of palliative cancer patients admitted to the Rabat National Institute of Oncology and declared palliative.

Subjects recruitment

The study was conducted with caregivers of patients with advanced palliative cancer between December 2017 and July 2018 at the national institute of oncology in RABAT.

It is the first reference center for the diagnosis and treatment of cancer in Morocco.

The caregivers included were the carers of cancer patients declared palliative (out of all therapeutic resources) admitted to the national oncology institute of RABAT

The recommendations of the Streiner guide were followed to estimate the sample size. The number of subjects required which indicates that for a value of intra-class correlation coefficient around 0.70 and an accuracy of 0.10. The number of subjects should be at least 100[16].

The number of caregivers of advanced cancer patients reported palliative (out of all therapeutic resources) who reported during the data collection period was approximately 120

Instruments and procedures

The Moroccan version of CarGoQoL was administered to caregivers of palliative diagnosed cancer patients .The same questionnaire was administered again from the third to the seventh day later to assess reproducibility. Participants provided socio-demographic and clinical data.

Statistical analysis

The psychometric tests were carried out according to the following procedure: Evaluate missing data and score distribution (mean, interval, floor and ceiling effects). The reliability of internal consistency was assessed by the Cronbach coefficient α .

A value of 0.70 or more was considered adequate [17]. Inter-observer reproducibility was tested using the intra-class correlation coefficient (ICC). The acceptability of the CarGoQoL was assessed with the response rate, percentage of missing data and time required to complete the questionnaire. p value of < 0.05 was considered statistically significant.

Multi-treatment scale analysis was used to examine the following elements:

Convergent validity and discriminate validity by determining the extent to which items are in correlation with the dimension assumed to be represented. Items must have a significantly higher correlation with their own dimension than with other dimensions. The correlations of the items to their own dimensions must be > 0.40 .

The discriminate validity of CarGOQoL was assessed using the clinical and sociodemographic characteristics. All statistical analyses were performed using SPSS 13.0.

Results

120 caregivers were included in this study. Participants were aged 17 to 67 years.

86.7% were women and 45% had a secondary education. The most common primary cancer diagnoses were respectively 43% gynecological and breast, 37.5% digestive, 22% lung. The median care giving duration was 13.06 months. Other socio-demographic and clinical characteristics are presented in (Table 1)

Table 1
Caregiver's sociodemographic and clinical characteristics

Caregivers		N = 120
Age(Years)	Mean $\pm$ SD	45.47 $\pm$ 11.7
Gender	Women	105(86.7)
	Men	16(13.3)
Residence	Urban	98(81.7)
	Rural	19(15.8)
	NP	3(2.5)
Relationship with the patient	Partner	35(29.2)
	Parent	40(33.4)
	Children	18(15)
	Brother/Sister	25(20.9)
	Others	2(1.6)
Education	illiterate	25(20.8)
	Primary	41(34.2)
	secondary	45(37.5)
	university	9(7.5)
Employmentstatus	Unemployed	70(58.3)
	Employed	47(39.1)
	NP	3(2.5)
Primary cancer diagnostis	Lung	22(18.3)
	Digestive	45(37.5)
	Gynecological and Breast	43(35.8)
	urological	5(4.2)
	Others	5(4.2)
Karnofsky index	≤ 30	31(25.8)
	[40–50]	79(65.9)
	≥ 60	19(8.4)
Duration of patient assistance (Months)	Median	13.06 [2–72]
Time to transition to palliative care (Months)	Median	22.89[1- 204]
SD standard deviation friend(1) other family member Cavum(2) thyroid(1) unknown primary(2)		

The average time to complete the questionnaire was 9 minutes with missing data of 1.7% for the Social support dimension and 29.2% for the private life dimension. The floor effect ranged from 0.8–2.5% and the ceiling effect from 16.7% to 38.9%.(Table 2)

Table 2
Dimensions' characteristics of CarGoQoL

Dimension/ Item	N	M ± SD	Flooreffect (%)	Ceilingeffect (%)	Cronbach'sα	IIC Min- Max	IDV Min- Max	Reliability ICC Test-retest
PsWB (4)	120	± 68.43 31.50	0.8	23.3	0.96	0.89– 0.94	0.41– 0.94	0.83
B(4)	120	± 66.47 29.10	0.8	16.7	0.91	0.82– 0.94	0.30– 0.65	0.77
RHC(3)	120	± 67.98 18.55	0.8	38.9	0.94	0.91– 0.95	0.42– 0.77	0.95
AF(3)	120	± 58.75 18.26	2.5	29.2	0.87	0.69– 0.92	0.42– 0.76	0.93
COP(3)	120	± 80.06 17.22	0.8	36.7	0.66	0.66– 0.87	0.48– 0.76	0.98
PhWB (4)	120	± 70.64 21.87	0.8	18.3	0.96	0.90– 0.91	0.42– 0.69	0.98
SE(2)	120	± 24.47 19.13	0.8	32.2	0.68	0.89	0.76	0.94
LEI (2)	120	± 74.68 12.49	0.8	20.8	0.65	0.87	0.76	0.86
SS (2)	118	± 71.61 21.22	0.8	38.3	0.82	0.93	0.76	0.92

± mean ± standard deviation. ⬇percentage of the lowest modality. ⬆percentage of the highest modality, PsWB: psychological well being, B:burden;RHC:relationship with healthcare; AF: administration and finances; COP :coping, PhWB: physical well being; SE: self-esteem; LEI: leisure time; SS: social support; PL: private life, IIC: item-internal consistency, IDV: item-discriminant validity ,ICC: intraclass correlation coefficients.

Dimension/ Item	N	M ± SD	Flooreffect (%)	Ceilingeffect (%)	Cronbach's α	IIC Min- Max	IDV Min- Max	Reliability ICC Test-retest
PL(2)	85	± 65.73 12.22	0.8	28.8	0.14	0.65	0.62	0.76
$\bar{x}$ mean $\pm$ standard deviation. $\bar{L}$ percentage of the lowest modality. $\bar{H}$ percentage of the highest modality, PsWB: psychological well being, B:burden;RHC:relationship with healthcare; AF: administration and finances; COP :coping, PhWB: physical well being; SE: self-esteem; LEI: leisure time; SS: social support; PL: private life, IIC: item-internal consistency, IDV: item-discriminant validity ,ICC: intraclass correlation coefficients.								

Internal consistency for the overall index was $\alpha = 0.82$. Reliability was explored using test-retest data with 20 caregivers, including CarGoQoL administered twice in a 3–7 day interval. (CCI) was from 0.76 to 0.98. The correlation between each item and its own dimension was higher than the correlation with the other dimensions. (Table 2)

Interdimensional correlations were significant ($r \geq 0.4$) with the exception of administration and finance (AF), self-esteem (SE) and leisure time (LEI).

A high correlation was noted between some dimensions; the Well Being (PsWB) and Physical Well Being (PhWB), Coping (COP) and (PsWB) ,(PhWB)and (COP) with a range from 0.55 to 0.77.(Table S1)

The discriminate validity of CarGoQoL was assessed using socio-demographic characteristics (Table 3). Women reported statistically significant lower scores for PhWB and AF dimensions than men. A physical burden (PhWB) was reported when the caregiver is a parent of the patient. An activity limit (LEI) was reported in the children of patients. A privacy disruption was reported when the caregiver is a sibling of the patient. The least senior caregivers reported a poor quality of life for the dimensions PsWB, Burden, COP and PhWB. Overall, no difference was found with respect to the patient's gender. Caregivers reported a worse quality of life in terms of emotional dimensions : PsWB,COP and physical dimension (PhWB) when the patient is younger. When the patient is older; caregivers reported lower quality of life scores in dimensions B,SE,LEI and SS. Caregivers reported a lower quality of life for the AF and SE dimensions when the duration of assistance increases and for the LEI and PL dimensions when the duration of the illness is long.

Table 3

:Comparisons (mean $\pm$ standard deviation) and correlations of CarGOQoL scores with respect to caregivers' and patients' characteristics

	PsWB	B	RHC	AF	COP	PhWB	SE	LEI	SS	PL
Caregiver's gender										
women	68.7 $\pm$ 31.7	68.4 $\pm$ 28,2	67.1 $\pm$ 18.6	57.2 $\pm$ 17.1	80.3 $\pm$ 16.9	70.1 $\pm$ 21.7	25.0 $\pm$ 19.2	75.2 $\pm$ 12.5	71.7 $\pm$ 20.8	66.1 $\pm$ 12.1
men	66.5 $\pm$ 31.02	53.7 $\pm$ 32,5	73.4 $\pm$ 17.8	68.7 $\pm$ 22.2	78.1 $\pm$ 19.6	73.8 $\pm$ 22.8	21.0 $\pm$ 18.6	71.0 $\pm$ 11.8	70.5 $\pm$ 24.8	63.4 $\pm$ 12.9
P value	0.8	0.04	0.2	0.01	0.6	0.03	0.4	0.2	0.8	0.4
Relationship status										
Partner	58.8 $\pm$ 36.7	64.4 $\pm$ 30.1	64.7 $\pm$ 17.1	60.2 $\pm$ 20.9	76.4 $\pm$ 17.2	57.3 $\pm$ 26.3	17.5 $\pm$ 15.2	81.4 $\pm$ 11.8	72.1 $\pm$ 19.4	59.6 $\pm$ 10.0
Parent	72.7 $\pm$ 30.3	71.2 $\pm$ 29.2	64.1 $\pm$ 19.9	56.2 $\pm$ 17.0	81.0 $\pm$ 18.3	77.6 $\pm$ 17.2	27.5 $\pm$ 20.8	73.1 $\pm$ 10.4	70.3 $\pm$ 22.5	70.8 $\pm$ 10.2
Child	65.7 $\pm$ 29.7	69.2 $\pm$ 30.5	73.6 $\pm$ 21.0	66.2 $\pm$ 16.2	76.8 $\pm$ 19.0	66.6 $\pm$ 18.6	30.5 $\pm$ 24.7	69.4 $\pm$ 17.7	66.6 $\pm$ 26.4	68.7 $\pm$ 13.5
Brother/sister	75.2 $\pm$ 24.7	58.6 $\pm$ 27.1	75.0 $\pm$ 14.0	55.3 $\pm$ 17.3	85.0 $\pm$ 13.3	80.0 $\pm$ 14.3	24.0 $\pm$ 14.3	71.5 $\pm$ 8.4	76.6 $\pm$ 17.7	67.1 $\pm$ 14.3
P value	0.1	0.4	0.08	0.3	0.2	0.00	0.07	0.003	0.6	0.009
Caregiver's age class										
≤ 40	51.9 $\pm$ 30.1	48.1 $\pm$ 27.4	74.5 $\pm$ 20.5	55.1 $\pm$ 16.4	74.2 $\pm$ 18.3	61.8 $\pm$ 22.1	20.9 $\pm$ 18.1	76.1 $\pm$ 13.8	78.7 $\pm$ 17.2	62.5 $\pm$ 9.4
40–60	72.4 $\pm$ 30.3	70.4 $\pm$ 25.4	67.3 $\pm$ 16.8	61.0 $\pm$ 18.7	81.2 $\pm$ 17.0	71.6 $\pm$ 20.5	27.4 $\pm$ 19.4	11.9 $\pm$ 1.4	71.2 $\pm$ 21.6	66.4 $\pm$ 13.1
≥ 60	86.7 $\pm$ 24.1	89.1 $\pm$ 27.0	56.1 $\pm$ 16.2	56.1 $\pm$ 19.2	87.7 $\pm$ 10.8	85.8 $\pm$ 18.6	18.3 $\pm$ 18.2	75.0 $\pm$ 12.5	57.5 $\pm$ 21.0	66.3 $\pm$ 10.6
P value	0.00	0.00	0.005	0.2	0.02	0.001	0.1	0.7	0.005	0.5
Patient's gender										
Women	69.7 $\pm$ 29.5	65.0 $\pm$ 28.0	71.0 $\pm$ 18.8	58.0 $\pm$ 16.4	80.0 $\pm$ 17.8	74.2 $\pm$ 18.2	24.2 $\pm$ 20.1	73.3 $\pm$ 10.7	74.2 $\pm$ 21.8	68.3 $\pm$ 11.3
Men	67.4 $\pm$ 34.3	68.0 $\pm$ 30.8	63.5 $\pm$ 17.7	60.1 $\pm$ 20.7	80.6 $\pm$ 15.8	65.5 $\pm$ 25.2	25.0 $\pm$ 18.2	76.5 $\pm$ 14.6	67.8 $\pm$ 19.7	62.8 $\pm$ 12.8
P value	0.3	0.7	0.08	0.5	0.08	0.06	0.9	0.4	0.1	0.1
Patient' age class(years)										
≤ 30	54.5 $\pm$ 39.2	78.6 $\pm$ 35.4	70.2 $\pm$ 23.0	52.3 $\pm$ 12.4	60.7 $\pm$ 26.2	60.7 $\pm$ 27.4	42.8 $\pm$ 32.1	83.9 $\pm$ 9.4	64.2 $\pm$ 31.8	70.0 $\pm$ 6.8
30–70	60.7 $\pm$ 33.0	58.0 $\pm$ 28.5	69.8 $\pm$ 17.1	57.5 $\pm$ 17.7	77.5 $\pm$ 17.1	65.4 $\pm$ 22.2	22.6 $\pm$ 17.2	75.5 $\pm$ 12.9	76,1 $\pm$ 16.4	64.0 $\pm$ 12.6

	PsWB	B	RHC	AF	COP	PhWB	SE	LEI	SS	PL
70	86.1 ± 16.7	80.9 ± 22.2	63.8 ± 20.2	62.2 ± 19.9	88.5 ± 9.7	82.7 ± 14.1	24.6 ± 18.4	71.3 ± 11.2	63.8 ± 25.1	67.4 ± 12.0
P value	0.00	0.00	0.2	0.2	0.00	0.00	0.02	0.03	0.009	0.3
Caregiving duration (months)										
6	75.5 ± 29.3	76.3 ± 26.3	67.0 ± 18.2	60.8 ± 21.2	82.9 ± 17.3	77.4 ± 17.5	35.8 ± 21.7	73.9 ± 11.2	65.9 ± 27.3	71.8 ± 11.6
6–24	65.3 ± 32.2	64.8 ± 28.4	68.3 ± 19.2	56.8 ± 16.7	78.5 ± 17.8	68.2 ± 22.9	22.5 ± 17.5	74.8 ± 13.3	72.3 ± 19.3	63.9 ± 11.6
24	77.0 ± 29.2	57.1 ± 38.1	65.0 ± 12.9	71.6 ± 20.1	85.0 ± 10.2	73.7 ± 19.0	12.5 ± 13.1	73.7 ± 7.0	77.5 ± 21.0	67.1 ± 14.8
p value	0.2	0.1	0.8	0.04	0.3	0.1	0.001	0.9	0.2	0.06
Patient's disease duration (months)										
12	68.6 ± 32.6	71.2 ± 28.1	68.0 ± 19.9	60.7 ± 18.7	80.3 ± 18.1	70.6 ± 21.5	27.5 ± 21.3	74.5 ± 11.5	69.1 ± 22.3	68.7 ± 11.3
12–48	65.3 ± 32.4	62.5 ± 31.7	65.3 ± 16.9	57.5 ± 17.8	78.8 ± 17.5	67.3 ± 24.3	20.9 ± 16.7	77.9 ± 11.8	72.0 ± 21.2	61.4 ± 11.8
48	72.9 ± 27.0	55.9 ± 24.0	75.5 ± 17.9	51.6 ± 16.1	80.5 ± 14.3	78.7 ± 15.4	20.0 ± 13.1	68.3 ± 14.8	79.4 ± 14.3	67.8 ± 15.9
P value	0.7	0.1	0.1	0.2	0.9	0.2	0.14	0.03	0.2	0.03

Discussion

In this article we present the results of a transcultural validation of the Moroccan version of CarGoQoL. This questionnaire is well designed and validated to measure and evaluate the quality of life of caregivers of cancer patients. It was developed by a French research team and has an American version conducted in the United States [10, 11].

For the development of our Moroccan Arabic version, we followed a rigorous scientific methodology recommended in the field [13, 16, 17].

The CarGoQoL questionnaire is exclusive in identifying dimensions specific to carers. The cross-cultural adaptation process was carried out in close collaboration with the Mapi Research Trust team that granted the conduct of this study.

The validation was conducted with a diverse sample including spouses, parents and children siblings to identify the different links with the palliative cancer patient and covers all aspects of caregivers' quality of life.

The psychometric properties of the Moroccan Arabic version of CarGoQoL are considered satisfactory. The average time required to complete the questionnaire was similar to that of the two French and American validation studies [10, 11].

Only one dimension of privacy that is unsatisfactory in terms of its Internal consistency ($\alpha = 0.14$) and missing data rate 29.2%, this is almost similar to the American study [11].

The test-retest time was three to seven days, this meets the recommended guidelines suggesting that the retest interval of 2 to 10 days is usually used[17].

In our study, there were more women than male caregivers. This is similar in the French and American study [10, 11]. In Morocco, it is not surprising that it is women who assume the role of primary caregiver because in our culture they are often expected to be the caregiver.

In our study, caregivers experienced a ripple effect and physical burden when they are the patients' parents .This was also in other contexts [18, 21].

All items were strongly correlated with their own dimensions than with others. Discriminate validity was determined between certain caregiver characteristics and quality of life scores. Previous studies demonstrated that the characteristics of patients impose a significant impact on the need for family care and the level of emotional distress of the family caregivers [19–21].

Our study confirmed that women had lower quality of life scores than men. This was strongly studied and confirmed in the literature [22, 23].

A strong correlation has been demonstrated in our study between the dimensions psychological well-being and physical well-being. This could be justified by the fact that psychological problems have an impact on the caregiver's physical state leading to significant physiological changes.

Also, meanings have been noted in the administrative and financial dimension among female carers and when the duration of the assistance exceeds 24 months. It is explained by the material and moral commitment towards the patient. This has also been demonstrated by a study carried out in Morocco; in most cases it is the family that assists, materially and morally helps the patient. [24]

Conclusion

The Moroccan Arabic version of CarGOQoL is a useful, reliable and valid tool to measure the quality of life of carers of palliative cancer patients in Morocco. Having a Moroccan version of CarGoQoL will make it easier for researchers to choose valid instruments to conduct studies that focus on carers.

Limitations

Some limitations should be taken into account: the failure to carry out external validation on a similar scale given its unavailability. The CarGoQoL is a self-administered questionnaire but the problem of the low literacy in Morocco yielded to a self-administration or administration by the same investigator following the level of education of the patients .

Our version is in Moroccan Arabic who is the language spoken by the majority of Moroccans but the cultural diversity in Morocco emerges from regional languages such as "Tarifit", "Tamazight" and "Tachelhit" which must be taken into account in subsequent studies.

Abbreviations

CarGoQoL: CareGiver Oncology Quality of Life; PsWB:psychological well being, B:burden;RHC:relationship with healthcare;AF:administration and finances;COP:coping, PhWB:physical well being;SE:self-esteem;LEI:leisure time;SS:social support;PL:private life ;ICC:item internal consistency;IDV:item discriminant validity; α :Alpha Cronbach's alpha

Declarations

Availability of data and materials

All data generated or analysed during this study are included in this manuscript

Ethics approval and consent to participate

All participants signed a written consent. The study has been approved by the ethics committee for biomedical research of the MOHAMED V Faculty of Medicine and Pharmacy in RABAT (N/R: folder number 15/18)

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests for this study.

Funding

There was no funding for this study

Authors' contributions

AA designed the study and analyzed the data and wrote the manuscript. **B.HR** was involved in the translation and critical revision of the manuscript. **RR** contributed to the design, analysis, and interpretation of the data and was involved in the writing of the manuscript. **EH** and **MH** were critically involved in analysis, interpretation of data, the revision of the manuscript and gave final approval of the version to be published. All author's have read and approved the manuscript.

Acknowledgment

The authors thank the team of Mapi research for permitting the use of CarQoQoL. We also thank the staff at the National Oncology Institute in Rabat, Morocco for this study. We thank Ms FOUZIA HAMMOUCH,, Dr LAILA LAHLOU and professor SABER BOUTAYEB for their collaboration all along the completion of this study. We thank the patients and families for their cooperation.

References

1. Bergwelt-Baildon MV, Hallek MJ, Shimabukuro-Vornhagen AA, Kochanek M. CCC meets ICU. Redefining the role of critical care of cancer patients. *BMC Cancer*. 2010;10:612.
2. Stenberg U, Ruland CM, Miaskowski C. Review of the literature on the effects of caring for a patient with cancer. *Psycho-Oncology*. 2009;19:1013–25.
3. Lengacher CA, Kip KE, Barta M, Post-White J, Jacobsen PB, Groer M, Lehman B, Moscoso MS, Kadel R, Le N, Loftus L, Stevens CA, Malafa MP. et M. M. Shelton. A pilot study evaluating the effect of mindfulness-based stress reduction on psychological status, physical status, salivary cortisol, and interleukin-6 among advanced-stage cancer patients and their caregivers. *J Holist Nurs Sep*. 2012;30(13):170–85.
4. Given BA, Given CW, Kozachik S. Family support in advanced cancer. *CA Cancer J Clin*. 2001;51:213–31.
5. Molassiotis A, Wilson B, Blair S, Howe T, Cavet J. Living with multiple myeloma: experiences of patients and their informal caregivers. *Support Care Cancer*. 2011;19:101–11.
6. Ellis J. The impact of lung cancer on patients and carers. *Chronic Respir Dis*. 2012;95(1):39–47.
7. Minaya Flores P, Berbis J, Chinot O, Auquier P. Assessing the quality of life among caregivers of patients with glioma. *Neurooncol Pract*. 2014 ;1 :191–7.
8. El –Jawahri AR, Traeger LN, Kuzmuk K, Eusebio JR, Vandusen HB, Shin JA, Keenan T, Gallagher ER, Greer JA, Pirl WF, and al. Quality of life and mood of patients and family caregivers during hospitalization for hematopoietic stem. cell transplantation *Cancer*. 2015;121:951–9.
9. Siegel K, Reveis VH, Houts P, Mor V. Caregiver burden and unmet patient needs. *Cancer*. 1991;68:1131–40.
10. Minaya P, Baumstarck K, Berbis J, Concalves A, Barlesi F, Michel J, and al. The caregiver oncology quality of life questionnaire (CarGoQoL): Development and validation of an instrument to measure the quality of life of the caregivers patients with cancer. *Eur J Cancer*. 2012 Apr;48 :904 – 11.
11. Kaveney S, Baumstarck K. Minaya P, Shannon T, Symes P, Loundou and al. Validation of the American version of the Caregiver Oncology Quality of life (CarGoQoL) questionnaire. *Health Qual Life Outcomes*. 2016;14:82.
12. *Nejjari C, El Fakir S, Bendahhou K, El Rhazi K* and al. Translation and validation of European organization for research and treatment of cancer quality of life Questionnaire -C30 into Moroccan version for cancer patients in Morocco. *BMC Research Note*. 2014; 7:228
13. Guillemin F, Bombardier C, Beaton D. Cross-cultural adaptation of health-related quality of life measures: literature review and proposed guidelines. *J Clin Epidemiol*. 1993;46(12):1417–32.
14. Lohr KN, Aaronson NK, Alomso J, Burnam MA, Patrick DL, Perrin EB, Roberts JS. Evaluating quality-of-life and health-status instrument: development of scientific review criteria. *Clin Ther*. 1996;18:979–92.
15. Acquadro C, Conway K, Giroudet C, Mearl. Linguistic validation manual for health outcome assessments. *Mapi* 2012.
16. Streiner DL, Geoffrey RN. Health measurement scales. A practical guide to their development and use 2005. 3d edition. ISBN 978-0-19-968521-9.
17. Cronbach LJ. Coefficient alpha and the internal structure of tests. *Psychometrika*. 1951;16:297–334.
18. Lund. et al. **BMC Cancer. Cancer caregiving tasks and consequences and their associations with caregiver status and the caregiver's relationship to the patient: a survey.** 2014; 14:541.
19. Northouse L, Williams AL, Given B, McCorkle R. Psychosocial care for family caregivers of patients with cancer. *J Clin Oncol*. 2012;30:1227–34.

20. Yu H, Li L, Liu C, Huang W, Zhou J, Fu W, Ma Y, Li S, Chang Y, Liu G, Wu Q. Factors associated with the quality of life of family caregivers for leukemia patients in China. *Health Qual Life Outcomes*. 2017;15:55.
21. W.Awadalla et al. Factors associated with quality of life of outpatients with breast cancer and gynecologic cancers and their family caregivers: a controlled study.*BMC Cancer*.2007; 7:102.
22. Janda M, Steginga S, Langbecker D, et al. Quality of life among patients with a brain tumor and their carers. *J Psychosom Res*. 2007;63(6):617–23.
23. Parvataneni R, Polley MY, Freeman T, et al. Identifying the needs of brain tumor patients and their caregivers. *J Neurooncol*. 2011;104(3):737–44.
24. H.Errihani et al.*Rev Francoph Psycho-Oncologie* .2005(2):80–85.DOI 10.1007/s10332-005-0060-8.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1AHLAMresearchnote.docx](#)