

When to Resume Bariatric Surgery After COVID-19 Pandemic? – Results of Patients’ and Surgeons’ Survey.

Alicja Dudek

Uniwersytet Jagiellonski w Krakowie

Michał Wysocki

Uniwersytet Jagiellonski w Krakowie Collegium Medicum

Maciej Walędziak

Uniwersytet Jagiellonski w Krakowie Collegium Medicum

Jacek Szeliga

Uniwersytet Jagiellonski w Krakowie Collegium Medicum

Monika Proczko-Stepaniak

Uniwersytet Jagiellonski w Krakowie Collegium Medicum

Michał Pedziwiatr

Uniwersytet Jagiellonski w Krakowie Collegium Medicum

Piotr Major (✉ majorpiotr@gmail.com)

Jagiellonian University Medical College <https://orcid.org/0000-0001-6552-7979>

Research article

Keywords: bariatric care after the pandemic, bariatric surgery, SARS-CoV-2 pandemic

Posted Date: August 12th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-42874/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on March 15th, 2021. See the published version at <https://doi.org/10.1186/s12893-021-01145-y>.

Abstract

Background

Coronavirus Disease 2019 is affecting most countries around the world, including Poland. In response, all elective surgeries have been postponed. We asked patients and surgeons when they want bariatric surgery to resume after pandemic. The main aim of the study was to determine patients' and surgeons' expectations about when to resume bariatric surgery regarding COVID-19 pandemic state.

Methods

The study was conducted in two groups: Group 1 – bariatric patients; Group 2 – bariatric surgeons. Two online surveys were distributed.

Results

A total of 895 patients, 299 before, 596 after surgery and 32 surgeons took part in survey. All patients and surgeons declared willingness to resume bariatric surgeries after pandemic and responded that they should be resumed immediately the World Health Organization announces end of pandemic (42%). The majority of patients before surgery answered that bariatric procedures should be resumed immediately the number of daily incidents begins to decrease (53%). In the patient group, current body mass index ($p < 0.001$) and contact with COVID+/quarantined persons ($p < 0.001$) had impact on the response to resumption of bariatric procedures.

Conclusions

Patients opted to wait for bariatric surgery until the oncological queue has become shorter. Surgeons presented a readiness to resume both procedures in parallel.

Trial registration: 1072.6120.103.2020

Background

The COVID-19 is affecting most countries around the world, including Poland[1]. The coronavirus pandemic has had a radical impact on the functioning of healthcare systems worldwide. The health crisis has also brought new challenges to surgical care, including bariatric surgery. In response to the COVID-19 pandemic, a strategy of postponing elective surgery has been adopted by most surgical societies[2,3]. Bariatric surgery is one of the first disciplines that has largely implemented this strategy. According to *The International Federation for the Surgery of Obesity and Metabolic Disorders* (IFSO) recommendations, all elective surgical and endoscopic cases for metabolic and bariatric surgery should be postponed in the interim. Furthermore, clinic and hospital visits are not recommended[4], as other surgical and non-urgent oncological procedures have also been recommended to be cancelled. However, promises to resume procedures at the earliest available date once the pandemic

subsidies have been offered by the healthcare authorities. The extended waiting lists at surgeries will be another important issue after the pandemic.

The number of bariatric procedures performed before the pandemic was insufficient in relation to the rising prevalence of obesity and the needs of bariatric patients worldwide[5,6]. In addition, the waiting list for bariatric procedures is growing rapidly as a result of the pandemic. The increasing negative effects of a pandemic outbreak affect bariatric surgeons, surgical centers, patients and health care systems. Current efforts are mainly focused on reorganizing the functioning of hospitals and wards to defeat the coronavirus. There is a lack of data on plans for post-pandemic bariatric care organizations.

To effectively prepare bariatric teams, bariatric care organizations should implement their strategies immediately after the pandemic has ended. An appropriate timing for the resumption of bariatric procedures should be determined first.

Methods

The aim of the study was to define an appropriate time to resume bariatric surgery after the end of the SARS-COV-2 pandemic.

Study design

The study was conducted in two groups: Group 1 – bariatric patients, Group 2 – bariatric surgeons. Two anonymous online surveys were designed. Surgeons from four bariatric centers were responsible for distribution of survey and gathering of material. The first survey was prepared for patients from the Polish *Association of Bariatric Patients* (CHLO) and distributed online among patients. The second survey was sent to all bariatric surgeons associated with the *Metabolic and Bariatric Chapter at the Association of Polish Surgeons* (SCMiB) who were invited to take part in the study by recruiting surgeons via e-mail through the SCMiB mailing list. An invitation to the project was also published on the official SCMiB website. The surveys were conducted between 8 and 22 April. The invitation to the project and the introduction to the questionnaire give details of the study design, assumptions of the study and types of questions. At the beginning of the survey, detailed technical instructions were issued on how to complete the form. Participation in both surveys was voluntary. Participants were required to give their consent to participate in the study as their response to the first survey question, which also supplied information on how to withdraw consent.

Inclusion and exclusion criteria

Group 1 included patients who were members of CHLO and who answered all the questions in the survey. The group contained patients before and after surgery. Cancer patients or patients with complications after bariatric interventions whose surgery could not be postponed were advised not to complete the

questionnaire. Group 2 included surgeons and general surgery residents who are members of SCMiB working in bariatric centers. Both certified and non-certified bariatric surgeons were included in the study. Retired surgeons, surgeons who were not performing bariatric procedures and medical doctors who are not practicing medicine during the pandemic (because of sick leave or vacation) were excluded.

Survey

Questionnaires for both study groups were accessible online. There was no time limitation on completion. Both surveys were divided into two parts.

The first part consisted of questions covering consent to participate and basic demographic data characterizing the groups of patients and surgeons. The group of patients was asked to state their age, sex, current body mass index (BMI), obesity comorbidities, whether they had undergone bariatric procedures in the past, contact with COVID+/quarantined persons and COVID status of the hospital taking care of them. Questions to the surgeon group concerned age, sex, stage of surgical training, type of employing hospital, COVID status of employing hospital and number of bariatric procedures performed in 2019 and 2020 until the start of the pandemic.

The second part included the same questions in both surveys. The patient group and surgeon group were first asked when bariatric surgery should be resumed. In view of the lengthening waiting list for other elective procedures, especially oncological operations, both surveys then asked for the respondent's opinion on how to reconcile the waiting lists for bariatric procedures with those for other elective surgery, including oncological procedures (Fig. 1).

Statistical analysis

Results were presented as mean plus standard deviation or median with interquartile range. To assess factors that could potentially influence the responses provided, univariate and multivariate regression models were run on Statistica 13.5 PL (Statsoft, USA).

Results

Overall, 927 participants took part in the survey (895 patients/32 surgeons). The study group included 792 (85%) females and 135 (15%) males. The median age was 39 (33–45). Respondents were associated with both COVID-dedicated hospitals (22%) and non-COVID-dedicated hospital (77%).

Group 1 characteristics (patients)

The median age of the patient group was 39 (33–45). The study group included 786 females (88%) and 109 males (12%). The median current BMI was 34.77 (29.3–41.18). 74% of patients presented with

obesity comorbidities. The patients who participated in the study were from both COVID-dedicated hospitals (198; 22.1%) and non-COVID-dedicated hospitals (697; 77.9%). 56 of the patients (6.26%) had had contact with COVID+ or a person who was quarantined. The patient study group was divided into two subgroups: patients before surgery and patients and after surgery (299/596). Basic characteristics of both subgroups are presented in Table 1.

Table 1
Basic characteristics of group 1

	All patients	Patients before surgery	Patients after surgery
n (%)	895 (100%)	299 (33%)	596 (67%)
Median age, years (IQR)	39 (33-45)	38 (32-44)	39 (33-45)
Males/Females, n (%)	109/786 (12%/88%)	39/260 (13%/87%)	69/527 (12%/88%)
Median BMI, kg/m ² (IQR)	34.77 (29.3-41.18)	42.61 (38.99-47.63)	31.21 (27.54-47.63)
Obesity comorbidities, n (%)	661 (74%)	239 (80%)	422 (71%)
Contact with COVID+/ quarantined person	56 (6.26%)	10 (3.34%)	46 (7.72%)
Hospital of patient care			
COVID-dedicated hospital	198 (22.12%)	79 (26.42%)	119 (19.97%)
Non-COVID dedicated hospital	697 (77.88%)	220 (73.58%)	477 (80.03%)

Group 2 characteristics (surgeons)

The mean age of surgeons who participated the survey was 42 ± 10. The group contained 26 males (81%) and 6 females (19%). The group contained 26 specialist surgeons (81%) and 6 residents (19%). 53% of the respondents worked in university hospitals, 38% of them in district hospitals and 9% in state departments. 12 surgeons were working in the COVID-dedicated hospitals (38%), while 20 in non-COVID-dedicated hospitals (63%). Median number of bariatric procedures performed overall in the hospital in 2019 was 150 (82–200). Median number of bariatric procedures performed overall in 2020 until the start of the pandemic was 29 (20–50) (Table 2).

Table 2
Basic characteristics of group 2

	Bariatric surgeons
n (%)	32 (100%)
Mean age, years±SD	42±10
Males/Females, n (%)	26/6 (81%/19%)
Stage of surgical training:	
Specialist surgeons, n (%)	26 (81%)
Residents, n (%)	6 (19%)
Type of employing hospital	
University hospital, n (%)	17 (53%)
State department, n (%)	3 (9%)
District hospital/city hospital, n (%)	12 (38%)
COVID status	
COVID-dedicated hospital	12 (38%)
Non-COVID dedicated hospital	20 (63%)
Numbers of bariatric procedures	
Median number of bariatric procedures in 2019, n (IQR)	150 (82-200)
Median number of bariatric procedures in 2020 until pandemic, n (IQR)	29 (20-50)

When bariatric procedures should be resumed

All patients and surgeons declared a high willingness to resume bariatric surgery after the SARS-CoV-2 pandemic, with a median score of 10 (10–10). There was no statistically significant difference between patients and surgeons in their answers to questions on the timing for resuming bariatric procedures. The majority in both groups indicated **immediately World Health Organization (WHO) announces the end of the pandemic**: 377 (42.4%), 14 (43.8%). The distribution of the remaining answers was as follows: immediately the number of daily incidents in the country begins to decrease: 251 (28.1%), 7 (21.9%); only after the last patient with COVID-19 has been discharged from hospital: 121 (13.5%), 9 (28.1%); only after the introduction of SARS- CoV-2 vaccinations: 95 (10.6%), 1 (3.1%); no opinion: 51 (5.4%), 1 (3.1%) (Fig. 2).

Answers given by the two subgroups of patients (before and after surgery) presented two statistically-significant different approaches to the question asked. The majority of patients in the before surgery subgroup answered that bariatric procedures should be resumed **immediately the number of daily incidents in the country begins to decrease**: 157 (52.5%), while the most popular answer in the after surgery subgroup was **immediately WHO announces the end of the pandemic**: 291 (48.8%).

Being associated with a COVID-dedicated hospital might have had an impact on the opinions of both groups on restoring bariatric surgery: after WHO announces the end of the pandemic ($p = 0.02$) and only after the last patient with COVID-19 has been discharged ($p < 0.001$).

Among the patient group, current BMI ($p < 0.001$) and contact with a COVID+/quarantined person ($p < 0.001$) influenced opinions on resumption of bariatric procedures after daily incidents in the country start to decrease. Answering that resumption should take place immediately WHO announces the end of the pandemic might be related to the status of surgery for that patient ($p < 0.001$). The statistical analysis indicated that factors like stage of surgical training, type of employing hospital, COVID status of employing hospital and number of bariatric procedures in 2019 or in 2020 until the start of the pandemic did not influenced the opinions given by surgeons ($p > 0.050$).

Resumption of bariatric procedures in relation to oncological surgery

The answers of the patient and surgeon groups to the question when bariatric procedures should be resumed in relation to oncological surgery differed significantly. The majority of patients opted for resumption of bariatric procedures only after managing the oncological queue: 396 (46%). However, most surgeons stated that bariatric procedures should be resumed in parallel with oncological procedures: 15 (48.4%). Analyzing the two subgroups, both in patients before and after surgery, the majority pointed out that shortening the queue for oncological procedures should be the priority: 90 (30.1%), 306 (54.5%). However, there was a significant difference in opinions on the parallel resumption of oncological and bariatric procedures between patients before and after surgery: (74 (25%) vs. 67 (11.4%)) (Fig. 3).

Discussion

The study presented here is the result of a national survey, the aim of which was to begin discussion with both surgeons and patients on the most appropriate time to resume bariatric surgery. The study showed a high willingness by all participants to resume bariatric surgery immediately after the SARS-CoV-2 pandemic ends. The survey affirmed high patient expectations for bariatric operations to be resumed as soon as WHO announces the end of the pandemic.

According to the latest data, obesity may present a higher risk for severe illness from COVID-19 and it can significantly worsen the prognosis of pneumonia[7,8]. Sockalingam et al. point out the tremendous psychological impact of the pandemic; an increase in eating psychopathology and binge eating

syndrome can have a strong negative effect on obesity management and gaining weight[9]. The group of patients who experienced delays in the preoperative period showed a greater desire to resume surgery as soon as the pandemic curve flattens compared to patients after surgery. The dominant opinion among patients who had completed their treatment was to wait until WHO announces the end of the pandemic. Current BMI and contact with a COVID+/ quarantined person should be taken into consideration as the most likely factors to influence patient opinions. Those findings may be related to patients who fear that they are in the high-risk group feeling a desperate need for surgery. On the other hand, many scientists are warning of the likelihood of a second wave of the epidemic in the fall[10]. The prospect of a rebound of the epidemic may also be increasing patients' desperation for surgery. The surgeon group showed a high readiness to provide bariatric surgery treatment as soon as possible after the end of the pandemic. The COVID status of the employing hospital does not seem to have had an influence on the answers given.

In the context of oncological treatment, patients opted to wait for bariatric surgery until the oncological queue is shorter. However, surgeons showed a readiness to resume both procedures in parallel. A prioritization strategy should be established that is appropriate to the readiness of surgeons and the needs of patients.

This study is associated with several limitations. Firstly, possible limitations of our study include recall bias and the subjectivity of patients and surgeons in stating their opinions. Secondly, free text comments provided by participants were not collected in the survey. It might be worthwhile extending the study to include a qualitative analysis. Additionally, the survey was limited to patients who are members of the Polish *Association of Bariatric Patients* (CHLO) and may not reflect the entire bariatric population; however, it is worth noting that CHLO members include a significant proportion of bariatric patients from all over the country. The second survey was dedicated only to bariatric surgeons. The group of surgeons specializing in bariatrics in Poland is limited. Asking for the opinion of all surgeons could be valuable. As a final point, it should be noted that we concentrated only on Polish bariatric departments and Polish patients so it may be difficult to extrapolate our results to other countries; however, we anticipate that most bariatric surgeons and patients have similar problems.

Conclusions

A timing for the resumption of bariatric procedures should be established in line with the needs of patients and the readiness of surgeons. The overall opinion of both patients and surgeons was to resume bariatric procedures immediately WHO announces the end of the pandemic. Patients before surgery preferred to resume bariatric procedures immediately the number of daily incidents in the country begins to decrease. Patients preferred to wait for bariatric surgery until the oncological queue has become shorter. Surgeons demonstrated a willingness to resume bariatric procedures in parallel with oncological surgery immediately after the pandemic ends.

List Of Abbreviations

WHO - World Health Organization

IFSO - *The International Federation for the Surgery of Obesity and Metabolic Disorders*

CHLO - *Association of Bariatric Patients*

SCMiB - *Metabolic and Bariatric Chapter at the Association of Polish Surgeons*

BMI - body mass index

Declarations

Ethics approval and consent to participate

The study was anonymous and performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki and its later amendments (Fortaleza). Participants were informed about the aim of the study and written informed consent was obtained electronically prior to the beginning of the survey. The study was approved by the Bioethics Committee of the Jagiellonian University (1072.6120.103.2020).

Consent for publication

Not applicable

Availability of data and materials

The datasets generated during and/or analysed during the current study are available in the Jagiellonian University Medical College repository. Data contained on the server supported to the Jagiellonian University Medical College. Backup policy of the Jagiellonian University Medical College. Copies automatically use 24 hours. Data copies are used on two separate media, geographically dislocated, in accordance with best practices. Access to the server where a production instance of the system is available and to the server where copies are available are physically, technically environmentally secured.

Competing interests

Authors declare no conflict of interest.

Funding

Not applicable

Authors' contributions

AD has made a substantial contribution to the manuscript conception and design of the work.

MW has made a substantial contribution to the manuscript conception and design of the work.

MW has made a substantial contribution to the data acquisition and analysis.

JS has made a substantial contribution to the data acquisition and analysis.

MPS has made a substantial contribution to the data acquisition and analysis.

MP has made a substantial contribution to study conception and manuscript revision.

PM has made a substantial contribution to study conception and manuscript revision.

All authors read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. WHO. WHO characterizes COVID-19 as a pandemic. <https://www.who.int/emergencies/diseases/novel-coronavirus-2019/events-as-they-happen> n.d.
2. American College of Surgeons releases recommendations for surgical management of elective operations during COVID-19 pandemic. <https://www.facs.org/about-acs/covid-19/information-for-surgeons> n.d.
3. Rymarowicz J, Pędziwiatr M, Major P, Nowakowski M. How we prepared our operating theatre for patients with SARS-CoV-2 virus. *Videosurgery Other Miniinvasive Tech i Inne Tech Małoinwazyjne* 2020. doi:10.5114/wiitm.2020.95090.
4. Yang W, Wang C, Shikora S, Kow L. Recommendations for Metabolic and Bariatric Surgery During the COVID-19 Pandemic from IFSO. *Obes Surg* 2020. doi:10.1007/s11695-020-04578-1.
5. Chooi YC, Ding C, Magkos F. The epidemiology of obesity. *Metabolism* 2019. doi:10.1016/j.metabol.2018.09.005.
6. Arroyo-Johnson C, Mincey KD. Obesity Epidemiology Worldwide. *Gastroenterol Clin North Am* 2016;45:571-9. doi:10.1016/j.gtc.2016.07.012.
7. Wang D, Hu B, Hu C, Zhu F, Liu X, Zhang J, et al. Clinical Characteristics of 138 Hospitalized Patients with 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. *JAMA - J Am Med Assoc* 2020. doi:10.1001/jama.2020.1585.

8. Gupta R, Ghosh A, Singh AK, Misra A. Clinical considerations for patients with diabetes in times of COVID–19 epidemic. *Diabetes Metab Syndr Clin Res Rev* 2020. doi:10.1016/j.dsx.2020.03.002.
9. 10.1002/oby.22836
Sockalingam S, Leung SE, Cassin SE. The Impact of COVID–19 on Bariatric Surgery: Re-Defining Psychosocial Care. *Obesity* 2020. doi:10.1002/oby.22836.
10. Wu ZY. Asymptomatic and pre-symptomatic cases of COVID–19 contribution to spreading the epidemic and need for targeted control strategies. *Zhonghua Liu Xing Bing Xue Za Zhi* 2020. doi:10.3760/cma.j.cn112338-20200406-00517.

Figures

First part of questionnaires	
Patients survey	Surgeons survey
<i>Please enter Your age:</i>	<i>Please enter Your age:</i>
<i>Please select Your sex: F/M</i>	<i>Please select Your sex: F/M</i>
<i>Please enter Your current BMI:</i>	<i>Please choose Your stage of surgical training:</i> <input type="checkbox"/> specialist surgeons <input type="checkbox"/> residents
<i>Please choose if You have any of listed obesity comorbidities:</i> <input type="checkbox"/> no comorbidities <input type="checkbox"/> insulin resistance <input type="checkbox"/> hypertension <input type="checkbox"/> obstructive sleep apnea (OSA) <input type="checkbox"/> lipid disorders <input type="checkbox"/> arthritis <input type="checkbox"/> others	<i>Please choose type of hospital in which You are employed:</i> <input type="checkbox"/> university hospital <input type="checkbox"/> district hospital <input type="checkbox"/> state department
<i>What is Your stage of bariatric treatment?</i> <input type="checkbox"/> I am before bariatric procedure. <input type="checkbox"/> I have undergone bariatric procedure.	<i>What is the COVID status of the hospital where You are employed:</i> <input type="checkbox"/> COVID-dedicated hospital <input type="checkbox"/> non-COVID
<i>Have You had contact with COVID+ or quarantinated person?</i> <input type="checkbox"/> Yes <input type="checkbox"/> No	<i>Please enter the numbers of bariatric procedures performed in 2019:</i>
<i>What is the COVID status of the hospital which take care of You?</i> <input type="checkbox"/> COVID-dedicated hospital <input type="checkbox"/> non-COVID dedicated hospital	<i>Please enter the numbers of bariatric procedures performed in 2020 until pandemic:</i>
Second part of both questionnaires	
What is Your opinion on when to resume bariatric operations?	
<input type="checkbox"/> Only after the introduction of SARS-CoV-2 vaccination.	
<input type="checkbox"/> Immediately when the number of daily incidents in the country begins to decrease.	
<input type="checkbox"/> Only after the last patient with COVID-19 will be discharged from the hospital.	
<input type="checkbox"/> Immediately when WHO announces the end of a pandemic.	
<input type="checkbox"/> No opinion.	
What is Your opinion on when to resume bariatric procedures in relation to oncological surgery?	
<input type="checkbox"/> Only when the queue for oncological procedures will be shortened.	
<input type="checkbox"/> At the same time as oncological procedures.	
<input type="checkbox"/> Due to the low risk of complications in bariatrics, it can be resumed before oncological procedures.	
<input type="checkbox"/> No opinion.	

Figure 1

Questions asked in surveys - patients and surgeons group

Time when bariatric procedures should be resumed




Figure 2

Time when bariatric procedures should be resumed.

Resumption of bariatric procedures in relation to oncological surgery




Figure 3

Resumption of bariatric procedures in relation to oncological surgery.