

# Intratumoral Administration of Astatine-211-labeled Gold Nanoparticle for Alpha Therapy

Hiroki Kato (✉ [kato-h@umin.org](mailto:kato-h@umin.org))

Osaka University Graduate School of Medicine <https://orcid.org/0000-0003-0838-6772>

Xuhao Huang

Osaka University Graduate School of Science

Yuichiro Kadonaga

Institute for Radiation Sciences, Osaka University

Daisuke Katayama

Osaka University Graduate School of Medicine

Kazuhiro Ooe

Osaka University Graduate School of Medicine

Atsushi Shimoyama

Osaka University Graduate School of Science

Kazuya Kabayama

Osaka University Graduate School of Science

Atsushi Toyoshima

Institute for Radiation Sciences, Osaka University

Atsushi Shinohara

Osaka University Graduate School of Science

Jun Hatazawa

Research Center for Nuclear Physics of Osaka University

Koichi Fukase

Osaka University Graduate School of Science

---

## Research Article

**Keywords:** astatine-211, alpha emitters, gold nanoparticles, radiolabeling, cancer therapy

**Posted Date:** April 21st, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-430857/v1>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

**Version of Record:** A version of this preprint was published at Journal of Nanobiotechnology on July 28th, 2021. See the published version at <https://doi.org/10.1186/s12951-021-00963-9>.

# Abstract

## Background:

$^{211}\text{At}$  is a high-energy  $\alpha$ -ray emitter with a relatively short half-life and a high cytotoxicity for cancer cells. Its dispersion can be imaged using clinical scanners, and it can be produced in cyclotrons without the use of nuclear fuel material. This study investigated the biodistribution and the antitumor effect of  $^{211}\text{At}$ -labeled gold nanoparticles ( $^{211}\text{At}$ -AuNP) administered intratumorally.

## Results:

AuNP with a diameter of 5, 13, 30, or 120 nm that had been modified with poly (ethylene glycol) methyl ether (mPEG) thiol and labeled with  $^{211}\text{At}$  ( $^{211}\text{At}$ -AuNP-S-mPEG) were incubated with tumor cells, or intratumorally administered to C6 glioma or PANC-1 pancreatic cancers subcutaneously transplanted into rodent models. Systemic and intratumoral distributions of the particles in the rodents were then evaluated using scintigraphy and autoradiography, and the changes in tumor volumes were followed for about 40 days.  $^{211}\text{At}$ -AuNP-S-mPEG was cytotoxic when it was internalized by the tumor cells. After intratumoral administration,  $^{211}\text{At}$ -AuNP-S-mPEG became localized in the tumor and did not spread to systemic organs during a time period equivalent to 6 half-lives of  $^{211}\text{At}$ . Tumor growth was strongly suppressed for both C6 and PANC-1 by  $^{211}\text{At}$ -AuNP-S-mPEG. In the C6 glioma model, the strongest antitumor effect was observed in the group treated with  $^{211}\text{At}$ -AuNP-S-mPEG with a diameter of 5 nm.

## Conclusions:

The intratumoral single administration of a simple nanoparticle,  $^{211}\text{At}$ -AuNP-S-mPEG, was shown to suppress the growth of tumor tissue strongly in a particle size-dependent manner without radiation exposure to other organs caused by systemic spread of the radionuclide.

## Background

As a local radiation therapy for cancer and in addition to external irradiation with  $\gamma$ -rays or X-rays, the effectiveness of brachytherapy using a sealed X-ray source for prostate cancer,[1] breast cancer,[2, 3] uterine cancer,[4, 5] head and neck cancer,[6, 7] and brain tumors,[8] is well known. However, the invasiveness of the procedure, extra-lesion displacement of the sealed radiation source, or adverse effects caused by the leakage of X-rays from radionuclides with a long half-life into adjacent organs are problematic issues. Adverse effects on other organs and psychological adverse effects of brachytherapy with  $\gamma$ -rays and X-rays have also been reported.[9] Brachytherapy for breast cancer is prone to causing skin ulcers, infections, rib fractures, and local pain, and its therapeutic effect is equivalent[10] or inferior[11] to external beam radiation therapy (EBRT). As for brachytherapy for brain tumors,[12–14] although the therapeutic effect has been better than ERBT in some cases[14] or about the same,[13] adverse effects such as vomiting, headaches and cerebral edema are relatively common.[8]

$\beta$ -Ray-emitting or  $\alpha$ -ray-emitting nuclides have a high linear energy transfer (LET) and relative biological effectiveness and are particularly toxic to proliferating cells. Because of the short range of  $\beta$ -ray or  $\alpha$ -ray emitters, normal tissues are minimally exposed if the radiation sources are appropriately distributed. This property can overcome the adverse effects of conventional EBRT and brachytherapy caused by  $\gamma$ -rays or X-rays. Thus, brachytherapy using high-LET radiation is feasible using locally distributed small seeds labeled with  $\beta$ - or  $\alpha$ -ray emitting nuclides. Yook et al. showed that the  $\beta$ -ray-emitting nuclide  $^{177}\text{Lu}$  could be used to label gold nanoparticles (AuNP) that could then be locally administered to breast cancer xenografts to suppress tumor growth.[15] Zhu et al. clarified that poly(cyclotriphosphazene-copolyethylenimine) nanospheres labeled with radionuclide  $^{131}\text{I}$  had anticancer capabilities when injected intratumorally [16]. Salvanou et al. demonstrated that  $\alpha$ -ray-emitting nuclide  $^{225}\text{Ac}$ -labeled AuNP could be used to inhibit tumor growth when administered intratumorally to glioma xenografts [17]. However, these studies also showed that radionuclides could spread systemically to other organs after intratumoral administration at time points that were much earlier than the half-lives of the radionuclides. Systemic exposures to long half-life nuclides, such as  $^{177}\text{Lu}$ ,  $^{131}\text{I}$  or  $^{225}\text{Ac}$ , are detrimental and should be minimized whenever radionuclides are administered intratumorally. Assuming that the radionuclides are not scattered externally from the tumor, extremely high doses of radioactivity could be administered repeatedly without exposing normal organs and tissues to radiation. On the other hand, since their ranges are extremely short,  $\alpha$ -ray emitters release energy only within their short distribution ranges. Thus, controlling the tissue distribution after administration is pivotal, and might be possible by adjusting the seed size and performing additional seed modifications.

$^{211}\text{At}$ , which can be produced in cyclotrons without the use of nuclear fuel material, emits high-energy  $\alpha$ -rays with a relatively short half-life and can be clearly imaged by clinical scanners or autoradiography. In the present study, we investigated the systemic and intratumoral distributions and verified the antitumor effect of  $^{211}\text{At}$ -labeled AuNP administered intratumorally.

## Methods

### Animal models

Male nude rats (F344/NJcl-rnu/rnu; 7 weeks old) were purchased from CLEA Japan, Inc. (Tokyo, Japan), and male nude mice (BALB/cSlc-nu/nu) were purchased from Japan SLC Inc. (Tokyo, Japan).

C6 glioma cells, which are a rat cell line derived from a glial cell tumor induced by N-nitrosomethylurea, were used as tumor cells. C6 glioma cells are suitable for assessing growth inhibition and the tissue retention of nanoparticles against wash-out in blood flow because they generally have a high growth rate and a high level of vascularization. C6 glioma cells obtained from RIKEN BRC were cultured using modified Eagle medium (MEM) (Sigma-Aldrich Japan, Tokyo, Japan) supplemented with 10% fetal bovine serum in a humidified incubator at 37°C under 5%  $\text{CO}_2$ .

Human pancreatic cancer cells (PANC-1) were obtained from the American Type Culture Collection. The cells were cultured in RPMI1640 medium with L-glutamine and phenol red (Fujifilm Wako Pure Chemical, Tokyo, Japan) with 10% heat-inactivated fetal bovine serum and 1% penicillin-streptomycin.

C6 glioma cells ( $0.9 \times 10^7$  cells/50  $\mu$ L MEM) supplemented with 50  $\mu$ L matrigel (Corning, New York, USA) were subcutaneously transplanted into both shoulders of the rats under anesthesia with 2.0% isoflurane in oxygen. PANC-1 cells ( $1.0 \times 10^7$  cells/50  $\mu$ L RPMI1640) supplemented with 50  $\mu$ L matrigel were transplanted into the left shoulders of the mice by subcutaneous injection.

## In vitro cell toxicity study

The C6 rat glioma cell line and the PANC-1 cell line were cultured in Dulbecco's MEM medium (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan) containing 10% fetal bovine serum (FBS) (Gibco™, Life Technologies, Carlsbad, CA USA) and 1% penicillin-Streptomycin Solution (FUJIFILM Wako Pure Chemical Corporation, Osaka, Japan). The C6 rat glioma cells ( $2 \times 10^4$  cells/well in 100  $\mu$ L medium) and PANC-1 cells ( $1 \times 10^4$  cells/well in 100  $\mu$ L medium) were then seeded onto 96-well plates and cultured for 1 day. Different sizes of AuNP modified with poly (ethylene glycol) methyl ether (mPEG) thiol (AuNP-S-mPEG) and  $^{211}\text{At}$ -AuNP-S-mPEG were synthesized (see the supplementary document) [18] and serially diluted; samples were then added to each well (25  $\mu$ L/well). The mass concentrations of AuNP-S-mPEG are shown in Supplementary Table S1. After 24 hours of incubation, we measured cell viability using a Cell Counting Kit 8 (CCK8) (Dojindo Laboratories, Kumamoto, Japan) and a microplate reader at 450 nm.

## Evaluation of intracellular uptake of AuNP-S-mPEG

C6 rat glioma cells were seeded onto 35 mm glass bottom dishes ( $4 \times 10^5$  cells/dish in 2 mL medium) and cultured for 1 day. Different kinds of AuNP-S-mPEG nanoparticles were added to the dish, and the cultures were incubated for 24 hours. After incubation, Hoechst solution was added to the dish, and the cultures were incubated for 10 minutes while being protected from light. The cells were washed 3 times with PBS, and 4% paraformaldehyde phosphate buffer solution was added to fix the samples for 30 minutes. After fixation, the samples were washed 3 times with PBS.

The cellular uptake of AuNP-S-mPEG was investigated using reflectance imaging from the cells.[19] Images were taken using a Nikon A1R + inverted confocal microscope (Nikon Corp., Tokyo, Japan) and a Plan Apo VC water-immersion objective lens (60 $\times$ , NA = 1.20; Nikon, Tokyo, Japan).

## Administration of AuNP-S-mPEG

Nanoparticles were intratumorally administered to 12 rats (24 tumors) 13 days after C6 glioma cell transplantation and to 12 mice (12 tumors) 14 days after PANC-1 cell inoculation under anesthesia with 2.0% isoflurane in oxygen. Solutions of saline (for 3 rats, 6 tumors), AuNP-S-mPEG + saline (for 3 rats, 6 tumors), 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG + saline (for 4 rats, 8 tumors), 30 nm  $^{211}\text{At}$ -AuNP-S-mPEG + saline (for 4 rats, 8 tumors), 13 nm  $^{211}\text{At}$ -AuNP-S-mPEG + saline (for 3 rats, 6 tumors, and for 6 mice, 6 tumors), and 5 nm  $^{211}\text{At}$ -AuNP-S-mPEG + saline (for 3 rats, 6 tumors) were prepared in a manner such that the

solution volume was equal to the corresponding tumor volume. Solutions of saline (for 3 rats, 6 tumors), unlabeled 30 nm AuNP-S-mPEG + saline (for 3 rats, 6 tumors), and unlabeled 13 nm AuNP-S-mPEG + saline (for 6 mice, 6 tumors) were also prepared in the same way for use in the control subjects.

Each of the solutions was slowly administered into the tumor over a time period of about 1 minute. For administration, the linear probe of an ultrasonic device (ProSound  $\alpha 6$ ; Hitachi-Aloka Medical, Ltd., Tokyo, Japan) was used so that the needle tip of the syringe (Myjector 29G; Terumo Co. Ltd., Tokyo, Japan) was placed in the center of the tumors. The radioactivity administered per tumor was  $1.4 \pm 0.4$  MBq for rats and  $1.2 \pm 0.1$  MBq for mice (Table S2 in the supplementary document).

## Follow-up and evaluation after administration

After intratumoral administration, the rats were monitored for 38 or 39 days; body weight and tumor size were measured under anesthesia using 2.0% isoflurane in oxygen. Forty days after administration, the tumor was excised and the tumor mass was measured. After the experiments, the animals were euthanized with an excessive dose of isoflurane.

## Scintigraphy

Whole-body scintigraphy was performed at 4 and 19 hours after the administration of  $^{211}\text{At}$ -AuNP-S-mPEG to rats; scintigraphy was performed using a gamma camera (E.cam; Siemens Healthcare, Erlangen, Germany) equipped with a low-energy, high-resolution and parallel-hole collimator. The detectors were set at a position 5 cm from the subject. Each animal was immobilized in a prone position under anesthesia with 2.0% isoflurane in oxygen, and imaging was performed for 10 minutes at 4 and 19 hours after the administration of  $^{211}\text{At}$  to obtain planar images of the front and rear surfaces with a  $256 \times 256$  matrix size (pixel size: 1.2 mm). For the 5 nm and 13 nm  $^{211}\text{At}$ -AuNP-S-mPEG groups, image acquisitions were also performed at 42 hours after the administration of  $^{211}\text{At}$ -AuNP-S-mPEG.

## Autoradiography

All the tumors were excised from the rats on the day after the administration of  $^{211}\text{At}$ -AuNP-S-mPEG with a diameter of 120 nm ( $n = 2$ ) or 30 nm ( $n = 2$ ). The tumor specimens were then immediately frozen at  $-80^\circ\text{C}$  and sliced to a thickness of about 30  $\mu\text{m}$  using a cryostat (CryoStar NX70; Thermo Scientific Inc., MA, USA), then attached to slide glasses. The frozen sections were then immediately dried with a dryer and placed in contact with an imaging plate for about 1 hour. Imaging was performed using a nonconfocal variable mode laser scanner (Typhoon FLA 7000; GE Healthcare Life Sciences, Buckinghamshire, England). After the tumors were resected, the rats were euthanized by the administration of an excessive amount of isoflurane.

## Statistics

Statistical calculations were performed using SPSS 17.0. To compare tumor masses, a one-way ANOVA and the Levene test were performed, followed by the Tukey honestly significant difference (HSD) post-hoc test.

## Results

### In vitro cytotoxicity study

C6 glioma cells were cultured with unlabeled 5 nm, 13 nm, 30 nm, or 120 nm AuNP-S-mPEG for 24 hours, but no obvious effect on viability was observed even at a high concentration. AuNP-S-mPEG was thus determined to be non-cytotoxic regardless of the concentration or particle size (Fig. 1A-D).

A significant decrease in viability was observed only in C6 glioma cells and PANC-1 cells cultured with 1 MBq/mL of 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG (Fig. 1E-L).

When C6 glioma cells were cultured in the presence of AuNP-S-mPEG for 24 hours, the 5 nm, 13 nm, 30 nm, and 120 nm AuNP-S-mPEG were internalized into the cells in a concentration-dependent manner (Fig. 2B, D, F, G). For the internalization of the 5 nm, 13 nm and 30 nm AuNP-S-mPEG, however, a concentration higher than that required by the 120 nm AuNP-S-mPEG was necessary (Fig. 2A, C, E). Precipitation, which was thought to have increased the concentration at the bottom of the culture significantly, was only observed in the 120 nm AuNP-S-mPEG group (Fig. 2H).

### Scintigraphy

No RI distribution was detected in any region other than the C6 tumor for all sizes of  $^{211}\text{At}$ -AuNP-S-mPEG at 4 and 19 hours after administration (Fig. 3A-D). In the 5 nm and 13 nm  $^{211}\text{At}$ -AuNP-S-mPEG groups, no systemic RI distribution was detected even at 42 hours after administration (Fig. 3C, D). The 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG were found to accumulate in spots within the tumor tissue, whereas the 5 nm, 13 nm, and 30 nm  $^{211}\text{At}$ -AuNP-S-mPEG diffused over a wider range than that observed for the 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG. For PANC-1 pancreatic cancer as well as C6 glioma, RI accumulated exclusively within the tumor after the administration of 13-nm  $^{211}\text{At}$ -AuNP-S-mPEG (Fig. 3E).

### Autoradiography of tumors

The RI was observed in tumors excised from rats injected with 120 nm or 30 nm  $^{211}\text{At}$ -AuNP-S-mPEG. The distribution of the 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG was confirmed to be remarkably uneven (Fig. 4A), whereas the 30 nm  $^{211}\text{At}$ -AuNP-S-mPEG were distributed almost throughout the entire tumor (Fig. 4B).

### Study of treatment efficacy

All the animals survived for 38 days after administration. No abnormal changes in body weight were seen in any of the rats, and no significant differences in body weight were seen among the groups (Fig. 5A). On the other hand, the body weight in the control mice treated with unlabeled AuNP-S-mPEG was significantly lower than that in the mice treated with  $^{211}\text{At}$ -AuNP-S-mPEG. The C6 gliomas treated with the 5 nm  $^{211}\text{At}$ -AuNP-S-mPEG had the lowest growth rate, based on tumor size (Fig. 5B). Regarding the mass

of the excised tumors, the group treated with the 5 nm  $^{211}\text{At}$ -AuNP-S-mPEG had the smallest tumor masses, which were significantly smaller than those of the control group. The mass of the excised tumor was proportional to the size of the  $^{211}\text{At}$ -AuNP-S-mPEG that had been used (Fig. 5C). Unlabeled AuNP-S-mPEG had no effect on tumor size.

Significant treatment effects were also confirmed in the PANC-1 xenograft models (Fig. 5D-F).

## Discussion

Consistent with the finding that the binding between AuNP and  $^{211}\text{At}$  is tight and stable,[20] the labeling rate of  $^{211}\text{At}$ -AuNP-S-mPEG and the size of the particles did not appear to change in the present study even after a time period equivalent to 6 half-lives of  $^{211}\text{At}$ . An in vitro experiment evaluating cytotoxicity against tumor cells showed that unlabeled AuNP-S-mPEG at nanoparticle sizes of 5 to 120 nm did not affect cell viability even after the cellular internalization of the nanoparticles had been confirmed. Thus, the toxicity of AuNP-S-mPEG itself was suggested to be negligible.

In C6 glioma cells, significant cytotoxicity was observed after in vitro treatment with  $^{211}\text{At}$ -AuNP-S-mPEG with a diameter of 120 nm labeled with 1 MBq/mL of radioactivity. In the PANC-1 cells, the cytotoxicity of  $^{211}\text{At}$ -AuNP-S-mPEG was similar to that seen in the C6 glioma cells. Thus, the cell toxicity of  $^{211}\text{At}$ -AuNP-S-mPEG as a non-targeted agent may not depend on the cell type.

The degree of cellular internalization of the 120 nm AuNP-S-mPEG was shown to be very high (Fig. 2G). For smaller AuNP-S-mPEG, however, higher concentrations were required for cellular internalization (Fig. 2B, D, F). Although the mechanism responsible for cellular internalization remains unknown, the 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG might have precipitated to the bottom of the well used in the presently reported in vitro system, creating an extremely elevated concentration around cells that had adhered to the bottom of the culture. As a result, frequent cellular contact caused by close proximity with a high concentration of AuNP-S-mPEG might be involved in cellular internalization. Cytotoxicity was only confirmed (Fig. 1H) under conditions where internalization was observed (Fig. 2G). These facts suggest that the cytotoxicity of  $^{211}\text{At}$ -AuNP-S-mPEG is caused by  $\alpha$  irradiation, which is enhanced by the cellular internalization of the nanoparticles. When injected into cancer tissue, the  $^{211}\text{At}$ -AuNP-S-mPEG are thought to be first distributed in the intercellular fluid.[21] Intercellular fluid accounts for about 15% of the whole tissue, and the diffusion capability of nanoparticles is lower than that of a membrane-permeable solvent. Therefore, at the time of injection, the AuNP is probably spatially concentrated around the tumor cells. The prolonged contact of cells in tissues with a high concentration of  $^{211}\text{At}$ -AuNP-S-mPEG may cause the nanoparticles to be internalized, thereby suppressing tumor cell growth.

Observations of the intratumoral distribution using autoradiography revealed that the 30 nm  $^{211}\text{At}$ -AuNP-S-mPEG had a high diffusivity in the tumor tissue, while the 120 nm  $^{211}\text{At}$ -AuNP-S-mPEG had a relatively low diffusivity. These observations suggest that the smaller the nanoparticle size, the greater the

diffusivity in the tumor tissue. The tumor growth suppression effect of the  $\alpha$ -rays from  $^{211}\text{At-AuNP-S-mPEG}$  was proportional to the size of the particles under the conditions of this study. Thus, for particles that are a minimum of 5 nm in diameter or larger, smaller  $^{211}\text{At-AuNP-S-mPEG}$  are considered to have greater diffusivity in tumor tissues and hence a greater tumor growth suppression effect. Unlabeled AuNP-S-mPEG did not show obvious cytotoxicity in either in vivo or in vitro studies. Thus, the antitumor property of the nanoparticles is clearly due to the cytotoxic effect of  $\alpha$ -rays.  $^{211}\text{At-AuNP-S-mPEG}$  with diameters ranging between 5 and 120 nm did not accumulate systemically in any organs for at least 42 hours, i.e. 6 half-lives of  $^{211}\text{At}$  after intratumoral administration. These results show that  $^{211}\text{At-AuNP-S-mPEG}$  does not undergo back-diffusion from the tumor tissue into blood vessels. On the other hand, AuNP with a diameter of 10 nm or more reportedly exhibit vascular permeability and can be dispersed to various organs in a size-dependent manner after intravenous or gastrointestinal administration.[22–24] These inconsistencies with previous studies indicate that the vascular permeability of AuNP may depend on particle modification.

The effectiveness of targeting a specific protein for the purpose of nanoseed brachytherapy is controversial.[15, 25] In this study, even non-targeted nanoseeds were shown to have a tumor growth inhibitory effect and to exert cytotoxicity in two different types of tumor cells both in vitro and in vivo. In principle, this method is likely to be independent of the histological type of the malignant tumor and could therefore be easily used as a brachytherapy for many tumor types. The labeling of  $^{211}\text{At-AuNP-S-mPEG}$  is completed by a simple operation involving the stirring of AuNP-S-mPEG and  $^{211}\text{At}$  together. Therefore, the nanoparticles can be generated immediately before administration, allowing them to be labeled with a high specific radioactivity. In the present study,  $^{211}\text{At-AuNP-S-mPEG}$  showed a high and exclusive retention at the injected site, where  $^{211}\text{At}$  radioactivity was detected. These findings and the properties of  $\alpha$ -rays suggest that very high-dose treatments with simple nanoparticles  $^{211}\text{At-AuNP-S-mPEG}$  might be feasible on an outpatient basis. In addition, Au and PEG are already clinically used as medicines, and their low toxicity has already been recognized.[26, 27]

The present study had the following limitations. First, the mass concentration, particle concentration, and osmotic pressure of the administered solutions were difficult to control because the administration of a constant radioactivity concentration was prioritized. Second, since the radioactivity could not be further increased due to restrictions on the supply of  $^{211}\text{At}$ , the radioactivity level administered to the tumor was set at around 1.4 MBq. In a study examining the intratumoral administration of Lu-177-labeled AuNP-S-mPEG, however, a much stronger tumor growth suppression was observed at an administration level of 4.5 MBq, compared with 3.0 MBq administration.[25] Further study is thus necessary to optimize the dose in experiments using higher doses.

## Conclusions

The intratumoral single administration of  $^{211}\text{At-AuNP-S-mPEG}$  was capable of strongly suppressing the tumor growth of C6 glioma and PANC-1 cells.  $^{211}\text{At-AuNP-S-mPEG}$  ranging in diameter from 5 nm to 30

nm were shown to diffuse well locally after intratumoral administration without spreading to other organs throughout the body for at least 6 half-lives of  $^{211}\text{At}$ . A smaller particle size was thought to be more effective for suppressing tumor growth. Locally administered nanoseed brachytherapy using  $^{211}\text{At}$ -AuNP-S-mPEG was considered to be an effective and safe non-targeted anticancer treatment.

## Abbreviations

EBRT  
External beam radiation therapy, LET:Linear energy transfer, AuNP:Gold nanoparticles, MEM:Modified Eagle medium, FBS:Fetal bovine serum, mPEG:Methoxypolyethylene glycol, PEG:Poly ethylene glycol, CCK8:Cell Counting Kit 8.

## Declarations

### Acknowledgements

In this study, we gratefully acknowledge Dr. Akihiro Ito at Analytical Instrument Facility, Graduate School of Science, Osaka University for support of TEM measurement, Mr. Shigeru Tamiya at Center for Scientific Instrument Renovation and Manufacturing Support, Osaka University for support of ICP-OES measurement, and Mr. Takanori Kobayashi for his efforts in performing the experiments.

### Authors' contributions

Conceptualization, H.K.; methodology, H.K., Y.K., A.S., and K.K.; formal analysis, H.K.; investigation, H.K., X.H., Y.K., D.K., and K.O.; resources, X.H.; data curation, H.K. and Y.K.; writing—original draft preparation, H.K. and Y.K.; writing—review and editing, H.K. and Y.K.; visualization, H.K. and Y.K.; supervision, A.S., J.H., and K.F.; project administration, H.K.; funding acquisition, H.K. and K.F. All authors have read and agreed to the published version of the manuscript.

### Funding

This research was funded by JSPS KAKENHI, grant number JP18K07674, JP20H05675, JP20H00404, and QiSS program of the OPERA grant number JPMJOP1721.

$^{211}\text{At}$  was supplied from RIKEN through the material transfer agreement and was given through Supply Platform of Short-lived Radioisotopes supported by JSPS Grant-in-Aid for Scientific Research on Innovative Areas, Grant Number JP16H06278.

### Availability of data and materials

All data generated or analyzed during this study are included in this published article.

### Ethics approval and consent to participate

The study was conducted according to the guidelines of the ARRIVE (Animal Research: Reporting In Vivo Experiments) and the Osaka University Animal Experiment Regulations, and approved by the Osaka University Animal Experiment Committee (pro-protocol code 30-103-006 and date of approval 2019/02/20).

### Consent for publication

All authors agree to be published.

### Competing interests

The author(s) have no potential competing of interest to declare with respect to the conduct of the research, the authorship, and/or the publication of this article.

### Author details

1. Department of Nuclear Medicine and Tracer Kinetics, Osaka University Graduate School of Medicine
2. Department of Chemistry, Graduate School of Science, Osaka University
3. Division of Science, Institute for Radiation Sciences, Osaka University
4. Research Center for Nuclear Physics, Osaka University


## References

1. Smith GD, Pickles T, Crook J, Martin AG, Vigneault E, Cury FL, Morris J, Catton C, Lukka H, Warner A, et al: **Brachytherapy improves biochemical failure-free survival in low- and intermediate-risk prostate cancer compared with conventionally fractionated external beam radiation therapy: a propensity score matched analysis.***Int J Radiat Oncol Biol Phys* 2015, **91**:505-516.
2. Skowronek J, Chichel A: **Brachytherapy in breast cancer: an effective alternative.***Prz Menopauzalny* 2014, **13**:48-55.
3. Roddiger SJ, Kolotas C, Filipowicz I, Kurek R, Kuner RP, Martin T, Baltas D, Rogge B, Kontova M, Hoffmann G, et al: **Neoadjuvant interstitial high-dose-rate (HDR) brachytherapy combined with systemic chemotherapy in patients with breast cancer.***Strahlenther Onkol* 2006, **182**:22-29.
4. Lin AJ, Hassanzadeh C, Markovina S, Schwarz J, Grigsby P: **Brachytherapy and survival in small cell cancer of the cervix and uterus.***Brachytherapy* 2019, **18**:163-170.
5. Robin TP, Amini A, Schefter TE, Behbakht K, Fisher CM: **Brachytherapy should not be omitted when treating locally advanced neuroendocrine cervical cancer with definitive chemoradiation therapy.***Brachytherapy* 2016, **15**:845-850.
6. Inoue T, Yoshida K, Yoshioka Y, Shimamoto S, Tanaka E, Yamazaki H, Shimizutani K, Teshima T, Furukawa S: **Phase III trial of high- vs. low-dose-rate interstitial radiotherapy for early mobile tongue cancer.***Int J Radiat Oncol Biol Phys* 2001, **51**:171-175.
7. Kovacs G: **Modern head and neck brachytherapy: from radium towards intensity modulated interventional brachytherapy.***J Contemp Brachytherapy* 2015, **6**:404-416.

8. Barbarite E, Sick JT, Berchmans E, Bregy A, Shah AH, Elsayyad N, Komotar RJ: **The role of brachytherapy in the treatment of glioblastoma multiforme.***Neurosurg Rev* 2017, **40**:195-211.
9. Ferenc S, Rzymiski P, Skowronek J, Karczewski J: **Physical and psychosocial side-effects of brachytherapy: a questionnaire survey.***J Contemp Brachytherapy* 2015, **7**:381-386.
10. Polgar C, Ott OJ, Hildebrandt G, Kauer-Dorner D, Knauerhase H, Major T, Lyczek J, Guinot JL, Dunst J, Miguelez CG, et al: **Late side-effects and cosmetic results of accelerated partial breast irradiation with interstitial brachytherapy versus whole-breast irradiation after breast-conserving surgery for low-risk invasive and in-situ carcinoma of the female breast: 5-year results of a randomised, controlled, phase 3 trial.***Lancet Oncol* 2017, **18**:259-268.
11. Smith GL, Xu Y, Buchholz TA, Giordano SH, Jiang J, Shih YC, Smith BD: **Association between treatment with brachytherapy vs whole-breast irradiation and subsequent mastectomy, complications, and survival among older women with invasive breast cancer.***JAMA* 2012, **307**:1827-1837.
12. Teixeira MJ, Lepski G, Correia C, Aguiar PH: **Interstitial irradiation for CNS lesions.***Stereotact Funct Neurosurg* 2003, **81**:24-29.
13. Waters JD, Rose B, Gonda DD, Scanderbeg DJ, Russell M, Alksne JF, Murphy K, Carter BS, Lawson J, Chen CC: **Immediate post-operative brachytherapy prior to irradiation and temozolomide for newly diagnosed glioblastoma.***J Neurooncol* 2013, **113**:467-477.
14. Chan TA, Weingart JD, Parisi M, Hughes MA, Olivi A, Borzillary S, Alahakone D, Detorie NA, Wharam MD, Kleinberg L: **Treatment of recurrent glioblastoma multiforme with GliaSite brachytherapy.***Int J Radiat Oncol Biol Phys* 2005, **62**:1133-1139.
15. Yook S, Cai Z, Lu Y, Winnik MA, Pignol JP, Reilly RM: **Intratumorally Injected <sup>177</sup>Lu-Labeled Gold Nanoparticles: Gold Nanoseed Brachytherapy with Application for Neoadjuvant Treatment of Locally Advanced Breast Cancer.***J Nucl Med* 2016, **57**:936-942.
16. Zhu W, Zhao L, Fan Y, Zhao J, Shi X, Shen M: **(<sup>131</sup>I)-Labeled Multifunctional Polyphosphazene Nanospheres for SPECT Imaging-Guided Radiotherapy of Tumors.***Adv Healthc Mater* 2019, **8**:e1901299.
17. Salvanou EA, Stellas D, Tsoukalas C, Mavroidi B, Paravatou-Petsotas M, Kalogeropoulos N, Xanthopoulos S, Denat F, Laurent G, Bazzi R, et al: **A Proof-of-Concept Study on the Therapeutic Potential of Au Nanoparticles Radiolabeled with the Alpha-Emitter Actinium-225.***Pharmaceutics* 2020, **12**.
18. Ziegler C, Eychmüller A: **Seeded Growth Synthesis of Uniform Gold Nanoparticles with Diameters of 15–300 nm.***The Journal of Physical Chemistry C* 2011, **115**:4502-4506.
19. Guggenheim EJ, Khan A, Pike J, Chang L, Lynch I, Rappoport JZ: **Comparison of Confocal and Super-Resolution Reflectance Imaging of Metal Oxide Nanoparticles.***PLoS One* 2016, **11**:e0159980.
20. Dziawer L, Kozminski P, Meczynska-Wielgosz S, Pruszynski M, Lyczko M, Was B, Celichowski G, Grobelny J, Jastrzebski J, Bilewicz A: **Gold nanoparticle bioconjugates labelled with At-211 for targeted alpha therapy.***Rsc Advances* 2017, **7**:41024-41032.


21. Stine CA, Munson JM: **Convection-Enhanced Delivery: Connection to and Impact of Interstitial Fluid Flow.***Front Oncol* 2019, **9**:966.
22. De Jong WH, Hagens WI, Krystek P, Burger MC, Sips AJ, Geertsma RE: **Particle size-dependent organ distribution of gold nanoparticles after intravenous administration.***Biomaterials* 2008, **29**:1912-1919.
23. Hillyer JF, Albrecht RM: **Gastrointestinal persorption and tissue distribution of differently sized colloidal gold nanoparticles.***J Pharm Sci* 2001, **90**:1927-1936.
24. Sonavane G, Tomoda K, Makino K: **Biodistribution of colloidal gold nanoparticles after intravenous administration: effect of particle size.***Colloids Surf B Biointerfaces* 2008, **66**:274-280.
25. Cai Z, Yook S, Lu Y, Bergstrom D, Winnik MA, Pignol JP, Reilly RM: **Local Radiation Treatment of HER2-Positive Breast Cancer Using Trastuzumab-Modified Gold Nanoparticles Labeled with (177)Lu.***Pharm Res* 2017, **34**:579-590.
26. Boisselier E, Astruc D: **Gold nanoparticles in nanomedicine: preparations, imaging, diagnostics, therapies and toxicity.***Chem Soc Rev* 2009, **38**:1759-1782.
27. Webster R, Didier E, Harris P, Siegel N, Stadler J, Tilbury L, Smith D: **PEGylated proteins: evaluation of their safety in the absence of definitive metabolism studies.***Drug Metab Dispos* 2007, **35**:9-16.

## Figures



**Figure 1**


please see the manuscript file for the full caption



**Figure 2**


C6 glioma cells were seeded onto 35-mm glass bottom dishes and the cells were incubated with 5 nm, 13 nm, 30 nm or 120 nm AuNP-S-mPEG for 24 hours. The intracellular uptake of these nanoparticles was examined using reflectance imaging. In each image, the leftmost section shows the bright field image of the cells, the second section shows the reflectance image of the particles taken up by the cells, the third section shows the fluorescent Hoechst staining for DNA, and the rightmost section shows the fused

image. The C6 glioma cells were incubated with AuNP-S-mPEG with the following sizes and mass concentrations: (A) 5 nm and 12.7 mg/L, (B) 5 nm and 138.4 mg/L, (C) 13 nm and 6.9 mg/L, (D) 13 nm and 68.6 mg/L, (E) 30 nm and 8.7 mg/L, (F) 30 nm and 51.6 mg/L, and (G) 120 nm and 33.6 mg/L. (H) AuNP-S-mPEG aqueous solutions containing each nanoparticle size (5 nm, 13 nm, 30 nm and 120 nm) were allowed to stand and were then observed 24 hours later. The 120 nm AuNP-S-mPEG had precipitated after 24 hours, and the concentration at the bottom of the container had increased.




**Figure 3**

Scintigraphy of C6 glioma-bearing rats was performed at 4 and 19 hours after the administration of 120 nm (A), 30 nm (B), 13 nm (C), or 5 nm (D)  $^{211}\text{At}$ -AuNP-S-mPEG. For the 13 nm (C) and 5 nm (D)  $^{211}\text{At}$ -AuNP-S-mPEG, imaging was also performed at 42 hours after administration. Scintigraphy of PANC-1 bearing mice was also performed in a same way as rats at 4, 19, and 42 hours after the administration of 13nm  $^{211}\text{At}$ -AuNP-S-mPEG (E). The radioactivity distributions (upper rows) are also shown using surface plots (lower rows) that display a square field of view of the dorsal detector with side length 30 cm for rats (A-D), or 15 cm for mice (E), which covers the whole body of an animal. No systemic accumulation of radioactivity was observed in any of the organs. S: Superior, I: inferior, R: right, L: left.



**Figure 4**

Twenty-four hours after the administration of 120 nm (A) and 30 nm (B)  $^{211}\text{At}$ -AuNP-S-mPEG, the tumors were excised, sliced and attached to a slide glass (right). Autoradiography (left) was performed after placing both tumor sections in contact with the same imaging plate for an hour.



**Figure 5**

Changes in the body weights of the rats (A) and mice (D) and changes in the tumor volumes of the C6 glial cells (B) and PANC-1 cells (E) after intratumoral administration ( $1.4 \pm 0.4$  MBq/tumor for rats and  $1.2 \pm 0.1$  MBq/tumor for mice). The C6 glial (C) and PANC-1 (F) tumors were excised 40 days after administration and their masses were measured. The error bars indicate the standard deviation. The tumor mass after the administration of  $^{211}\text{At}$  was significantly smaller than that seen in the controls. The body weights of the mice in the control group were significantly lower than those in the group treated with  $^{211}\text{At}$  ( $*0.01 \leq p < 0.05$ ,  $**0.001 \leq p < 0.01$ ,  $***p < 0.001$ ).

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Graphicalabstract20210409.pptx](#)
- [Supplement20210409.docx](#)