

Microvirga Amygdalina sp. nov., and *Microvirga Alba* sp. nov., Two New Species Isolated from Soil in South Korea

Hyejin Oh

Seoul Women's University

Myungkyum Kim

Seoul Women's University

Sathiyaraj Srinivasan (✉ drsini@swu.ac.kr)

Seoul Women's University <https://orcid.org/0000-0001-6019-7421>

Research Article

Keywords: Methylobacteriaceae, *Microvirga*, Radiation, Taxonomy

Posted Date: June 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-431927/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Two novel Gram-stain-negative, aerobic, rod-shaped, circular, convex, light-pink and white-coloured bacterial strains BT291^T and BT350^T were isolated from soil collected in Uijeongbu city (37° 44' 55" N, 127° 2' 20" E) and Jeju island (33° 22' 48" N, 126° 31' 48" E), respectively, South Korea. Phylogenetic analysis based on 16S rRNA gene sequences revealed that strains BT291^T and BT350^T belong to a distinct lineage within the genus *Microvirga* (family *Methylobacteriaceae*, order *Rhizobiales*, class *Alpha Proteobacteria*, phylum *Proteobacteria*, kingdom *Bacteria*). The 16S rRNA gene sequence similarity between the two strains BT291^T and BT350^T was 97.4 %. The two strains were found to have the same quinone system, with Q-10 as the major respiratory quinone. The major polar lipids of strains BT291^T and BT350^T were phosphatidylethanolamine (PE), diphosphatidylglycerol (DPG), phosphatidylcholine (PC) and phosphatidylglycerol (PG). The major cellular fatty acids of strain BT291^T were C_{18:1} ω7c (58.2 %) and cyclo-C_{19:0} ω8c (25.7 %). The major cellular fatty acids of strain BT350^T were C_{18:1} ω7c (38.5 %) and cyclo-C_{19:0} ω8c (27.7 %). Based on the polyphasic analysis (phylogenetic, chemotaxonomic and biochemical), strains BT291^T and BT350^T can be suggested as two novel bacterial species within the genus *Microvirga* and the proposed names are *Microvirga amygdalina* and *Microvirga alba*, respectively. The type strain of *Microvirga amygdalina* is BT291^T (= KCTC 72368^T = NBRC 114845^T) and the type strain of *Microvirga alba* is BT350^T (= KCTC 72385^T = NBRC 114848^T).

Introduction

The genus *Microvirga* was first described by Kanso and Patel (2003) allocated to the family *Methylobacteriaceae* order *Rhizobiales*. At the time of writing (January 2021), the genus comprises 23 published and not validly published species (<http://www.bacterio.net/Microvirga.html>). *Microvirga* species have been retrieved from various polar environments in the last years, e.g. from regoliths from Tibet hot spring sediments (Liu et al. 2020), roots of rapeseed plants (Jimenez-Gomez et al. 2019), root nodule (Wang et al. 2019), forest soil (Zhang et al. 2019), rhizospheric soil (Li et al. 2020) and root nodule (Msaddak et al. 2020). In genus *Microvirga*, cells are Gram-stain-negative, C_{18:1} ω7c and cyclo-C_{19:0} ω8c was contained in the major fatty acids, and the genome size and DNA G + C content were 3.53–9.63 Mb and 61.1–65.1 %, respectively (Zhang et al. 2019).

In this study, both strains BT291^T and BT350^T were newly isolated from a soil sample Uijeongbu city (37° 44' 55" N, 127° 2' 20" E) and Jeju island (33° 22' 48" N, 126° 31' 48" E), respectively, South Korea. In the present study, we conducted a phylogenetic analysis based on the 16S rRNA gene sequences and phenotypic, genotypic and chemotaxonomic characteristics to determine the taxonomic position of strains BT291^T and BT350^T. The results suggested that strains BT291^T and BT350^T represent two novel species of the genus *Microvirga*, for which the name *Microvirga amygdalina alba* sp. nov. and *Microvirga alba* sp. nov. are proposed.

Materials And Methods

Isolation and cultural conditions

Strain BT291^T was isolated from Uijeongbu city (37° 44' 55" N, 127° 2' 20" E) and strain BT350^T was isolated from Jeju island (33° 22' 48" N, 126° 31' 48" E) located in South Korea. Colonies were isolated using Reasoner's 2A (R2A) agar medium (Difco) after incubation at 25°C for 9 days. After that, the strains were routinely cultured on R2A agar at 25°C, maintained at 4°C and stored in 10 % (w/v) glycerol suspension at - 80°C. The 16S rRNA gene sequences of the purified colonies were identified using the EzBioCloud server (<https://www.ezbiocloud.net/>).

Morphology, physiology and biochemical analysis

The cell morphology was examined using transmission electron microscopy (JEOL, JEM1010) with negative staining. The Gram-staining reaction was performed using a kit, following the manufacturer's instruction (bioMérieux). Catalase activity was examined with 3 % (w/v) H₂O₂ solution and oxidase activity was examined by the addition of 1 % (w/v) tetramethyl-p-phenylenediamine (Cappuccino and Sherman 2002). The growth of strains BT291^T and BT350^T was tested on Reasoner's 2A (R2A) agar, Luria-Bertani (LB) agar, Tryptic Soy Agar (TSA), Nutrient Agar (NA) and on MacConkey (MAC) agar. Growth was tested at different temperatures (10, 15, 25 and 30°C) under various pH conditions (5 to 9, 1 pH intervals) and different NaCl concentrations (1 % to 5 % [w/v %], 1 % intervals). API 20NE and API ZYM tests were performed according to the manufacturer's instruction (bioMérieux).

Phylogenetic analysis

The 16S rRNA genes of strains BT291^T (1,428 bp) and BT350^T (1,435 bp) were amplified and sequenced using two universal bacterial primers 27F and 1492R (Weisburg et al. 1991) using the BT291^T and BT350^T genomic DNA as a template. The sequencing was then done using the 337F, 518R, 785F, and 926R universal primers (Macrogen). To determine the taxonomic positions of strains BT291^T and BT350^T, 16S rRNA sequences of similarity searches were obtained from EzBioCloud (Yoon et al. 2017) and compared with of strains BT291^T and BT350^T using EzEditor2 server. Phylogenetic trees were reconstructed using the MEGAX program (Kumar et al. 2018) with the neighbor-joining (Saitou and Nei 1987), maximum-likelihood (Felsenstein 1981) and maximum-parsimony algorithms (Fitch 1971). The stability of the tree topologies was evaluated by bootstrap analysis based on 1,000 resampling method (Felsenstein 1985). Evolutionary distances were calculated according to the Kimura two-parameter model (Kimura 1983).

Genome sequencing

The genomic DNA was extracted using a genomic DNA extraction kit according to the manufacturer's instruction (Solgent). The sequencing libraries were prepared using the Nextera DNA Flex Library Prep Kit (Illumina), and whole-genome sequencing was performed by iSeq 100. The genome sequences were assembled using SPAdes 3.10.1 (Algorithmic Biology Lab, St. Petersburg Academic University of the

Russian Academy of Sciences). The whole-genome sequences of strains BT291^T and BT350^T were deposited in GenBank (www.ncbi.nlm.nih.gov/) database. The genome sequences of strains BT291^T and BT350^T were annotated by the National Center for Biotechnology Information Prokaryotic Genome Annotation Pipeline (PGAP) (Tatusova et al. 2016). The average nucleotide identity (ANI) was calculated using the EzBioCloud (<https://www.ezbiocloud.net>) and the digital DNA-DNA hybridization (dDDH) was calculated using the Genome-to Genome Distance Calculator (GGDC) with the recommended formula 2 (Table 1) (Meier-Kolthoff et al. 2013).

Chemotaxonomic characteristics

For analysis of cellular fatty acid, polar lipid and quinone strains BT291^T and BT350^T were grown on R2A agar at 25°C for three days and cells were freeze-dried. Polar lipids of strains BT291^T and BT350^T were extracted as described previously. The total lipids, glycolipids, phosphatidylcholine and amino groups were separated using two-dimensional thin-layer chromatography (TLC). The polar lipid spots were detected by spraying the proper detection reagents (Komagata and Suzuki 1987; Minnikin et al. 1984). The fatty acids were purified by saponification, methylation and extraction procedures (Sasser 1990). The quinones were extracted using the Sep-Pak Vac cartridges (Waters) and analyzed by high-performance lipid chromatography (HPLC) based on the previous methods (Hiraishi et al. 1996). The fatty acid methyl esters (FAME) were identified using the Sherlock Microbial Identification System V6.01 (MIS, database TSBA6, MIDI Inc).

Results And Discussion

Morphology, Physiology and Biochemical analysis

Strains BT291^T and BT350^T were Gram-staining-negative bacteria and they showed rod-shaped morphology (Fig. S1). Colonies of strains BT291^T and BT350^T were circular, convex and smooth after incubation for three days at 25°C. Colonies color of strains BT291^T and BT350^T were light-pink and white, respectively. Cells of strain BT291^T could survive at 10 to 30°C (optimum 25°C) and pH 6.0–9.0 (optimum 8.0) in R2A medium and the cells of strain BT350^T could survive at 10 to 30°C (optimum 25°C) and pH 5.0–9.0 (optimum 8.0) in R2A medium. Differential features between the new strains and reference strains were provided in Table 1. In addition, the negative reaction of strains BT291^T and BT350^T on API kits were given as supplementary tables (Table S2 and S3).

Phylogenetic and Genome sequence analysis and analysis

Based on the 16S rRNA gene sequence similarities, strains BT291^T and BT350^T were affiliated with the family *Methylobacteriaceae* and showed high sequence similarities with the genus *Microvirga*. The strain BT291^T was closely related to *Microvirga aerophila* 5420S-12^T (97.5 % 16S rRNA gene similarity) and *Microvirga subterranean* DSM 14364^T (97.2 %). The strain BT350^T was closely related to *Microvirga aerophila* 5420S-12^T (97.6 %) and *Microvirga brassicacearum* CDVBN77^T (96.8 %). The 16S rRNA gene

sequence similarities of strains BT291^T and BT350^T with the closely related type strains were less than 97.5 % and 97.6 %, respectively and with other *Microvirga* species were less than 96.9 %. These values were around or below the 98.7 % 16S rRNA gene sequence similarity recently used as the threshold for differentiating among bacterial species (Chun et al. 2018). The remaining *Microvirga* species exhibited sequence similarities lower than 97.0 %. After the reconstruction of neighbor-joining, maximum-likelihood (Fig. S2) and maximum-parsimony (Fig. S3) trees strain BT350^T clustered with *M. brassicacearum* CDVBN77^T and *M. flavescens* c27j1^T and strain BT291^T clustered independently (Fig. 1). The phylogenetic analysis results clearly showed that strains BT291^T and BT350^T are two new species within the genus *Microvirga*.

The draft genome of strain BT291^T was 4.77 Mb (51.2×) long and consisted of 4,473 protein-coding genes, 57 RNA genes (6 rRNA genes, 50 tRNA genes) and 8 pseudogenes. The draft genome of strain BT350^T was 4,42 Mb (29.9×) long and consisted of 4,014 protein-coding genes, 51 RNA genes (4 rRNA genes, 47 tRNA genes) and 66 pseudogenes. The genome sequence of the strains BT291^T and BT350^T have been deposited in GenBank under the accession numbers NZ_JAFEMB000000000 and NZ_JADQDO010000000, respectively. The DNA G + C contents of strains BT291^T and BT350^T were 64.7 mol% and 61.9 mol%, respectively. These values were within the range of the G + C contents for the genus *Microvirga* as previously reported (63.5–64.3 mol%). The digital DNA-DNA hybridization values between strains BT291^T and BT350^T and other related type strains of genus *Microvirga* were less than 23.1 %, respectively (Table S1), which are below the cutoff (70 %) point (Meier-Kolthoff et al. 2013). Average nucleotide identity (ANI) values between strains BT291^T and BT350^T and other related type strains of genus *Microvirga* were less than 79.1 %, respectively (Table S1). These values are below the ANI species threshold (95–96 % ANI value) as described by Ritcher and Rossello-Mora (2009).

Chemotaxonomic characterization

The fatty acid profiles of strains BT291^T and BT350^T and three reference strains of genus *Microvirga* were presented in Table 2. The major fatty acids of strain BT291^T C_{18:1} ω7c (58.2 %) and cyclo-C_{19:0} ω8c (25.7 %). Strain BT291^T has smaller amounts of C_{16:0} (5.8 %), summed feature 2 (iso-C_{16:1} I / C_{14:0} 3OH) (2.2 %) and C_{18:0} (1.5 %), whereas other closely related *Microvirga* species (*M. aerophila* 5420S-12^T, *M. subterranean* DSM 14364^T and *M. subterranea* Fail4^T) have larger amounts of corresponding fatty acids. Strain BT291^T contained C_{19:0} 10-methyl (1.1 %) and C_{14:0} (> 1 %), but other closely related *Microvirga* species (*M. aerophila* 5420S-12^T, *M. subterranean* DSM 14364^T and *M. subterranea* Fail4^T) did not contain those fatty acids.

The major fatty acid profiles of strain BT350^T were C_{18:1} ω7c (38.5 %) and cyclo-C_{19:0} ω8c (27.7 %). Strain BT350^T has larger amount of C_{16:0} (10.4 %), cyclo-C_{17:0} (6.4 %) and C_{18:0} 3OH (1.8 %), whereas

other closely related *Microvirga* species (*M. aerophila* 5420S-12^T, *M. subterranean* DSM 14364^T and *M. subterranea* Fail4^T) have smaller amounts of corresponding fatty acids. Strain BT350^T did not contain C_{17:1} ω8c, C_{17:1} ω6c and C_{17:1} ω7c 11-methyl, but other closely related *Microvirga* species (*M. aerophila* 5420S-12^T, *M. subterranean* DSM 14364^T and *M. subterranea* Fail4^T) contained those fatty acids.

The polar lipids of strain BT291^T consisted of a phosphatidylethanolamine (PE), an unknown diphosphatidylglycerol (DPG), unknown phosphatidylcholine (PC), an unknown phosphatidylglycerol (PG), an unknown aminolipid (AL), an unknown aminophospholipid (APL), an unknown phospholipid (PL), an unknown glycolipid (GL) and two unknown lipids (L) (Fig. S4). In contrast, strain BT350^T consisted of a phosphatidylethanolamine (PE), an unknown diphosphatidylglycerol (DPG), an unknown phosphatidylglycerol (PG), unknown phosphatidylcholine (PC), an unknown amino lipid (AL) and an unknown lipid (L) (Fig. S5). The dominant respiratory quinone of strains BT291^T and BT350^T was Q-10. These results supported that chemotaxonomic characteristic of strains BT291^T and BT350^T are similar to those of the other species in the genus *Microvirga*. Based on phenotypic, phylogenetic and biochemical features, it is concluded that strains BT291^T and BT350^T represent two novel species of the genus *Microvirga*, for which the name *Microvirga amygdalina* and *Microvirga alba* are proposed.

Description of *Microvirga amygdalina* sp. nov.

Microvirga amygdalina (a.myg.da.li'na L. fem. adj. *amygdalina* of almonds).

Cells are Gram-stain-negative, aerobic, rod-shaped, 0.6–1.3 μm in diameter and about 1.6–2.7 μm in length, non spore forming and non-motile. Colonies are irregular, convex and light-pink-colored on Reasoner's 2A (R2A) agar plates after growth for three days at 25°C. Growth is observed at temperatures ranging from 10 to 30°C (optimum 25°C). The pH range for growth is 6.0–9.0 (optimum pH 8.0) on R2A agar. Normal cell growth occurs at 10–30°C (optimum 25°C) and pH 6.0–9.0 (optimum 8.0). Cells grow on Reasoner's 2A agar (R2A), Luria-Bertani agar (LB), Tryptic Soy Agar (TSA), Nutrient Agar (NA) and Macconkey (MAC) agar (weakly). Cells are positive for oxidase and catalase activity. The major respiratory quinone is Q-10. The dominant cellular fatty acids are C_{18:1} ω7c (58.2 %) and cyclo-C_{19:0} ω8c (25.7 %). The major polar lipids are phosphatidylethanolamine (PE), diphosphatidylglycerol (DPG), phosphatidylcholine (PC), phosphatidylglycerol (PG). Positive for nitrate reduction, arginine dihydrolase and urease (API 20NE). Positive for esterase (C4) and acid phosphatase (API ZYM). The whole-genome sequence of strain BT291^T has been deposited in GenBank under the accession number NZ_JAFEMB000000000 (4.77 Mb). The genome-based G + C content is 64.7 mol%. The GenBank accession number for the 16S rRNA gene sequence of strain BT291^T is MT795755 (1,422 bp). The type strain BT291^T (= KCTC 72368^T = NBRC 114845^T) was isolated from a soil sample collected in Uijeongbu city (37° 44' 55" N, 127° 2' 20" E), South Korea.

Description of *Microvirga alba* sp. nov.

Microvirga alba (al'ba. L. fem. adj. *alba* white).

Cells are Gram-stain-negative, aerobic, rod-shaped, 0.4–0.9 µm in diameter and about 0.5–1.2 µm in length, non spore forming and non-motile. Colonies are irregular, convex and white colored on Reasoner's 2A (R2A) agar plates after growth for three days at 25°C. Growth is observed at temperatures ranging from 10 to 30°C (optimum 25°C). The pH range for growth is 5.0–9.0 (optimum pH 8.0) on R2A agar. Cells grow on Reasoner's 2A agar (R2A), Tryptic Soy Agar (TSA) and Nutrient Agar (NA) but not on Luria-Bertani agar (LB) and Macconkey (MAC) agar. Cells are positive for oxidase and catalase activity. The major respiratory quinone is Q-10. The dominant cellular fatty acids are C_{18:1} ω7c (38.5 %) and cyclo-C_{19:0} ω8c (27.7 %). The major polar lipids are phosphatidylethanolamine (PE), diphosphatidylglycerol (DPG), phosphatidylcholine (PC) and phosphatidylglycerol (PG). Weakly positive for trisodium citrate (API 20NE). Positive for alkaline phosphatase and esterase (C4) (API ZYM).

The whole-genome sequence of strain BT350^T has been deposited in GenBank under the accession number NZ_JADQD001000000 (4,42 Mb). The genome-based G + C content is 61.9 mol%. The GenBank accession number for the 16S rRNA gene sequence of strain BT350^T is MT795757 (1,416 bp). The type strain BT350^T (= KCTC 72385^T = NBRC 114848^T) was isolated from a soil sample collected in Jeju island (33° 22' 48" N, 126° 31' 48" E), South Korea.

Declarations

Acknowledgements: This work was supported by a research grant from Seoul Women's University (2021) and by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE) of the Republic of Korea (NIBR202002108). In addition, we are grateful to Dr. Aharon Oren (The Hebrew University of Jerusalem, Israel) for helping with the etymology.

Conflicts of interest: The authors declare that there are no conflicts of interest.

Ethical Approval: This article does not contain any studies with human participants or animals.

References

1. Cappuccino JG, Sherman N (2002) Microbiology- A laboratory manual, 6th edn. Pearson Education, Inc. Benjamin Cummings, California
2. Chun J, Oren A, Ventosa A, Christensen H, Arahal DR et al (2018) Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 68:461-466. <https://doi.org/10.1099/ijsem.0.002516>
3. Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368-376. <https://doi.org/10.1007/BF01734359>
4. Felsenstein J (1985) Confidence limit on phylogenies: an approach using the bootstrap. *Evolution* 39:783-791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>

5. Fitch WM (1971) Toward defining the course of evolution: minimum change for a specific tree topology. *Syst Zool* 20:406-416. <https://doi.org/10.2307/2412116>
6. Hiraishi A, Ueda Y, Ishihara J, Mori T (1996) Comparative lipoquinone analysis of influent sewage and activated sludge by high performance liquid chromatography and photodiode array detection. *J Gen Appl Microbiol* 42:457-469. <https://doi.org/10.2323/jgam.42.457>
7. Jimenez-Gomez A, Saati-Santamaria Z, Igual JM, Rivas R, Mateos PF, Garcia-Fraile P (2019) Genome Insights into the Novel Species *Microvirga brassicacearum*, a Rapeseed Endophyte with Biotechnological Potential. *Microorganisms* 7:0
8. Kanso S, Patel BK (2003) *Microvirga subterranea* gen. nov., sp. nov., a moderate thermophile from a deep subsurface Australian thermal aquifer. *Int J Syst Evol Microbiol* 53:401-406. <https://doi.org/10.1099/ijs.0.02348-0>
9. Kimura M (1983) *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press
10. Komagata K, Suzuki K (1987) 4 Lipid and cell-wall analysis in bacterial systematics. *Method Microbiol* 19:161-207. [https://doi.org/10.1016/S0580-9517\(08\)70410-0](https://doi.org/10.1016/S0580-9517(08)70410-0)
11. Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018) MEGA X: Molecular Evolutionary Genetics Analysis across Computing Platforms *Mol Biol Evol* 35(6):1547-1549. <https://doi.org/10.1093/molbev/msy096>
12. Li J, Gao R, Chen Y, Xue D, Han J, Wang J, Dai Q, Lin M, Ke X, Zhang W (2020) Isolation and Identification of *Microvirga thermotolerans* HR1, a Novel Thermo-Tolerant Bacterium, and Comparative Genomics among *Microvirga* Species. *Microorganisms* 8:0
13. Li J, Gao R, Chen Y, Xue D, Han J, Wang J, Dai Q, Lin M, Ke X, Zhang W (2020) Isolation and Identification of *Microvirga thermotolerans* HR1, a Novel Thermo-Tolerant Bacterium, and Comparative Genomics among *Microvirga* Species. *Microorganisms* 8:0
14. Liu ZT, Xian WD, Li MM, Liu L, Ming YZ, Jiao JY, Fang BZ, Xiao M, Li WJ (2020) *Microvirga arsenatis* sp. nov., an arsenate reduction bacterium isolated from Tibet hot spring sediments. *Antonie Van Leeuwenhoek* 113:1147-1153. <https://doi.org/10.1007/s10482-020-01421-6>
15. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M (2013) Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC bioinformatics* 14:60
16. Minnikin DE, O'Donnell AG, Goodfellow M, Alderson G, Athalye M, Schaal A, Parlett JH (1984) An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Microbiol Meth* 2:233-241. [https://doi.org/10.1016/0167-7012\(84\)90018-6](https://doi.org/10.1016/0167-7012(84)90018-6)
17. Msaddak A, Rejili M, Duran D, Mars M, Palacios JM, Ruiz-Argueso T, Rey L, Imperial J (2019) *Microvirga tunisiensis* sp. nov., a root nodule symbiotic bacterium isolated from *Lupinus micranthus* and *L. luteus* grown in Northern Tunisia. *Syst Appl Microbiol* 42:126015. <https://doi.org/10.1016/j.syapm.2019.126015>
18. Richter M, Rossello-Mora R (2009) Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* 106:19126-19131

19. Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Bio Evol* 4:406-425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>
20. Sasser M (1990) Identification of Bacteria by Gas Chromatography of Cellular Fatty Acids. MIDI Technical Note 101. Newark, DE: MIDI Inc
21. Tapase SR, Mawlankar RB, Sundharam SS, Krishnamurthi S, Dastager SG, Kodam KM (2017) *Microvirga indica* sp. nov., an arsenite-oxidizing Alphaproteobacterium, isolated from metal industry waste soil. *Int J Syst Evol Microbiol* 67:3525-3531. <https://doi.org/10.1099/ijsem.0.002157>
22. Tatusova T, DiCuccio M, Badretdin A et al (2016) NCBI prokaryotic genome annotation pipeline. *Nucleic Acids Res* 44:6614-6624. <https://doi.org/10.1093/nar/gkw569>
23. Wang F, Yang L, Deng J, Liu X, Lu Y, Chen W, Wu J (2019) *Microvirga calopogonii* sp. nov., a novel alphaproteobacterium isolated from a root nodule of *Calopogonium mucunoides* in Southwest China. *Antonie Van Leeuwenhoek* 112:1593-1602. <https://doi.org/10.1007/s10482-019-01285-5>
24. Weisburg WG, Barns SM, Pellerier DA, Lane DJ (1991) 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* 173:697-703. <https://doi.org/10.1128/jb.173.2.697-703.1991>
25. Weon HY, Kwon SW, Son JA, Jo EH, Kim SJ, Kim YS, Kim BY, Ka JO (2010) Description of *Microvirga aerophila* sp. nov. and *Microvirga aerilata* sp. nov., isolated from air, reclassification of *Balneimonas flocculans* Takeda et al. 2004 as *Microvirga flocculans* comb. nov. and emended description of the genus *Microvirga*. *Int J Syst Evol Microbiol* 60:2596-2600. <https://doi.org/10.1099/ijms.0.018770-0>
26. Yoon S, Ha S, Kwon S, Lim J, Kim Y et al (2017) Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613-16. <https://doi.org/10.1007/s10482-017-0844-4>
27. Zhang XJ, Zhang J, Yao Q, Feng GD, Zhu HH (2019) *Microvirga flavescens* sp. nov., a novel bacterium isolated from forest soil and emended description of the genus *Microvirga*. *Int J Syst Evol Microbiol* 69:667-671. <https://doi.org/10.1099/ijsem.0.003189>

Tables

Table 1. Different characteristics of *Microvirga amygdalina* and *Microvirga alba* and closely related species of genus *Microvirga*.

Taxa: 1, strain BT291^T (data was obtained in this study); 2, strain BT350^T (data was obtained in this study); 3, *M. aerophila* 5420S-12^T (data was taken Tapase et al. 2017); 4, *M. subterranea* FaiI4^T (data was taken Tapase et al. 2017); 4, *M. brassicacearum* CDVBN77^T (data was taken Jimenez-Gomez et al. 2019).

All strains were negative for production of indole, *N*-acetyl- β -glucosaminidase, α -fucosidase, β -galactosidase (PNPG), α -glucosidase (starch hydrolysis), β -glucosidase, β -glucuronidase, α -mannosidase and valine arylamidase.

+, positive; -, negative; w, weak positive

Characteristic	1	2	3	4	5
Size (µm long)	1.6-2.7	0.5-1.2	1.-3.3	4	1.2-1.5
Size (µm wide)	0.6-1.3	0.4-0.9	1.2-1.5	1-1.5	0.7-0.9
Colony color	light pink	white	light pink	light pink	white
Nitrate reduction					
Nitrate reduction to NO ₂	-	-	-	+	+
Nitrate reduction to N ₂	+	-	-	+	+
Production of acid from glucose	-	-	-	w	+
Enzyme activity					
acid phosphatase	+	-	-	+	+
alkaline phosphatase	-	+	-	w	+
arginine dihydrolase	+	-	-	-	-
esterase (C4)	+	+	-	+	+
esterase (C8)	-	-	-	+	+
α-galactosidase	-	-	+	-	-
β-galactosidase (ONPG)	-	-	+	-	-
β-glucosidase (Esculin hydrolysis)	-	-	-	+	+
leucine arylamidase	-	-	-	+	+
lipase (C14)	-	-	-	+	-
naphtol-AS-BI-phosphohydrolase	-	-	+	+	+
protease (gelatin hydrolysis)	-	-	-	+	-
urease	+	-	-	-	-
Assimilation					
citrate	-	w	+	+	+
L-arabinose	-	-	+	-	-
D-glucose	-	-	-	+	-
D-maltose	-	-	-	+	+
D-mannose	-	-	+	+	-
D-mannitol	-	-	+	-	+
N-acetyl-D-glucosamine	-	-	-	-	+
G+C content	64.7 %	61.9 %	62.2 %	63.5 %	62.3 %

Table 2. Cellular fatty acid profiles of *Microvirga amygdalina* sp. nov., and *Microvirga alba* sp. nov., and closely related species of genus *Microvirga*.

Taxa: 1, strain BT291^T (data was obtained in this study); 2, strain BT350^T (data was obtained in this study); 3, *M. aerophila* 5420S-12^T (data was taken Weon et al. 2010); 4, *M. subterranean* DSM 14364^T (data was taken Weon et al. 2010); 4, *M. subterranea* FaiI4^T (data was taken Jimenez-Gomez et al. 2019). All strains were grown on R2A agar at 25-28 °C. For

unsaturated fatty acids, the location of the double bond was presented by counting the number from the methyl (ω) end of the carbon chain.

-, not detected; tr, trace amount (<1 %).

Fatty acids	1	2	3	4	5
Saturated					
14:0	tr	tr	1.1	-	-
16:0	5.8	10.4	7.6	5.5	9.2
17:0	-	2.7	-	6.7	-
17:0 cyclo	1.4	6.4	1.1	-	3.8
18:0	1.5	2.2	1.0	3.8	6.1
19:0 10-methyl	1.1	-	-	-	-
19:0 cyclo $\omega 8c$	25.7	27.7	3.6	2.4	24.3
Unsaturated					
17:1 $\omega 8c$	-	-	-	2.0	-
17:1 $\omega 6c$	-	-	-	1.4	-
18:1 $\omega 7c$ 11-methyl	-	-	-	-	4.2
18:1 $\omega 7c$	58.2	38.5	69.6	73.4	39.3
18:0 3OH	1.2	1.8	tr	-	1.7
Summed Feature 2 (16:1 iso I / 14:0 3OH)	2.2	3.5	4.3	3.3	4.5
Summed Feature 3 (16:1 $\omega 6c$ / 16:1 $\omega 7c$)	2.1	6.1	11.0	1.5	4.5

Figures




Figure 1

Neighbor-joining phylogenetic tree reconstructed from a comparative analysis of 16S rRNA gene sequences showing the relationships of strains BT291T and BT350T with closely related validly published species. Bootstrap values (based on 1,000 replications) greater than 70 % based on neighbor-joining method is shown at the branch nodes. Circles indicate that the corresponding nodes were also recovered in the maximum-parsimony tree. Triangles indicate that the corresponding nodes were recovered in the maximum-likelihood tree. *Phyllobacterium loti* S658T was used as an outgroup. Bar, 0.01 substitutions per nucleotide position. The compact triangles represent species of the genus *Methylobacterium* and *Chelatococcus*.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarydata.docx](#)