Predictors of adherence to prescribed exercise programs for older adults with non-musculoskeletal indications for exercise: a systematic review

Julia F Shaw

University of Ottawa Faculty of Medicine

Sophie Pilon

Ottawa Hospital

Matthieu Vierula

Ottawa Hospital

Daniel I McIsaac (✉ dmcisaac@toh.ca)

University of Ottawa Faculty of Medicine <https://orcid.org/0000-0002-8543-1859>

Research article

Keywords: exercise, systematic review, prognosis, evidence-based medicine

Posted Date: July 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-43573/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

BACKGROUND AND OBJECTIVES:

Prescribed exercise to treat medical conditions and to prepare for surgery is a promising intervention to prevent adverse health outcomes for older adults; however, adherence to exercise programs may be low. Our objective was to identify and grade the quality of predictors of adherence to prescribed exercise in older adults.

METHODS

After registration (CRD42018108242), prospective experimental studies were identified using a peer-reviewed search strategy applied to MEDLINE, EMBASE, Cochrane and CINAHL from inception until April 23, 2019. Following independent and duplicate review of titles, abstracts and full texts, we included prospective studies with an average population age ≥ 65 years, where exercise was formally prescribed for a medical or surgical condition. We excluded studies where exercise was prescribed for a chronic musculoskeletal condition. Risk of bias was assessed using the Quality in Prognostic studies tool or Cochrane risk of bias tool, as appropriate. Predictors of adherence were identified, pooled, and graded for quality using an adaptation of the Grading of Recommendations Assessment, Development and Evaluation (GRADE) framework for predictor studies.

RESULTS

We included 19 observational studies and 4 randomized controlled trials ($n = 5785$). Indications for exercise included cardiac ($n = 6$), pulmonary rehabilitation ($n = 7$), or other ($n = 10$; surgical, medical, and neurologic). Overall adherence rate was reported in 20 studies (range 21%-93%; mean 68%, standard deviation 23%). Moderate-quality evidence suggested that positive predictors of adherence were self-efficacy and good self-rated mental health; negative predictors were depression (high quality) and distance from the exercise facility. Moderate-quality evidence suggested that comorbidity and age were not predictive of adherence.

CONCLUSIONS

These findings can inform design of future exercise programs as well as identification of individuals who may require extra support to benefit from prescribed exercise.

Background

Western populations are aging at a rapid rate; it is estimated that by 2050, seniors could account for up to 30% of our population.(1) The declining physical function that accompanies older age is associated with increased disability, institutionalisation, and mortality.(2) Additionally, frailty, a multidimensional syndrome related to age- and disease-related deficits, increases in prevalence with age and results in vulnerability to stressors and adverse health outcomes.(3, 4) Therefore, a large proportion of older individuals facing physiologic stressors, such as surgery or chronic medical conditions, are at risk of suffering worse outcomes compared to those who are more physically fit.

Older individuals preparing for major interventions or who have medical problems may benefit from interventions that target increasing their physical reserve to improve outcomes. Exercise has been identified as a promising perioperative intervention to improve postoperative outcomes in vulnerable older adults having surgery,(5) and has been shown to reduce mortality after cardiac events.(6) While exercise shows encouraging results for the treatment and prevention of adverse health outcomes in older adults, participants must adhere to the prescribed program in order to benefit from the exercise intervention. However, it is well-documented that older adults' adherence to prescribed exercise programs is low, especially in those with complex health conditions.(7) To support successful implementation of exercise programs for older adults, we must first identify what factors influence adherence to these programs to ensure that participants are willing and able to comply. To our knowledge, no studies have synthesized and graded the strength of evidence for patient- and program-level factors that predict exercise adherence.

To address this gap in the literature, our objective was to identify and grade the quality of predictors of adherence to prescribed exercise in older adults with either a medical or surgical indication. This systematic review will provide knowledge to inform current care and future research regarding the implementation and design of exercise programs for older adults with medical and surgical indications for physical activity.

Methods

Design

This was a systematic review that followed best practice recommendations from the Cochrane Collaboration(8) and for systematic reviews of observational and prognostic studies.(9, 10) We pre-registered our protocol (PROSPERO 2018 CRD42018108242) and reported findings using the Preferred Reporting Items for Systematic reviews and Meta Analyses guidelines.(11) All stages of the review were conducted using Distiller SR (Evidence Partners, Ottawa, Canada), a cloud-based systematic review platform.

Search strategy

A search strategy was developed in consultation with an information specialist (Supplementary Table S1) and peer-reviewed.(12) Citations in English or French were extracted from MEDLINE, Embase, Cochrane and CINAHL from inception until March 2018.

Eligibility criteria

Studies were eligible for inclusion if the following criteria were met: (1) average age of participants ≥ 65 years; (2) participants had a medical or surgical condition as an indication for exercise; and (3) participants were prescribed or recommended a formal exercise program. Prior to beginning our review, we recognized that exercise programs for chronic musculoskeletal conditions (e.g., low back pain or chronic joint pain or arthritis) *versus* other indications would be a primary source of heterogeneity. We also identified several syntheses of adherence in chronic musculoskeletal conditions that were already available, (13–17) therefore, we excluded studies where chronic musculoskeletal conditions were the indication for exercise. Study designs were limited to prospective experimental studies (to minimize the effects of misclassification bias and measurement error) and effect estimates predictive of adherence were limited to those that underwent multivariable adjustment (to minimize confounding bias), as recommended by best practice guidelines.(10) This meant that we included 1) adjusted associations between participant or program characteristics and adherence reported from prospective cohort studies or the experimental arm of randomized trials of prescribed exercise, 2) or the effect estimate from a randomized trial if it compared the effect of two different program features on adherence.

Study selection and data extraction

Title and abstract screening was performed in duplicate; any studies reviewed as 'yes' or 'unsure' by either reviewer were advanced to full-text review (agreement between both reviewers was required to exclude a study). Full-text articles were also assessed in duplicate and reasons for exclusion at this stage were recorded and categorized (wrong age group, no exercise program, no predictors of adherence, no medical or surgical condition, wrong study design, and other). Disagreements between reviewers during full-text review were resolved by consensus after discussion with the senior author (DIM).

A unique data extraction form was created for this study. The form was piloted in a sample of 8 studies by two extractors, which were then reviewed with the senior author. Following piloting, data was extracted by one reviewer and independently reviewed and checked for accuracy by a second reviewer. Extracted data included publication details (author, year), study design, sample size, average age, medical/surgical condition indicating exercise, and whether frailty status was assessed. We also extracted characteristics at the exercise program level, including inpatient/outpatient, supervised/unsupervised, and type of program (i.e., cardiac rehabilitation, pulmonary rehabilitation or other). Our primary outcome, program adherence, was recorded, including the definition used to quantify adherence and the overall adherence rate reported.

Risk of bias assessment

Two reviewers independently evaluated risk of bias, and disagreements were resolved through discussion with a senior author. Randomized controlled trials were assessed using the Cochrane risk of bias tool(8) while observational studies were assessed using the Quality in Prognostic Studies (QUIPS) tool.(18)

Synthesis of results and analysis

Our primary analysis was structured to support the Grading of Recommendations Assessment, Development and Evaluation (GRADE) adaptation for prognostic factor research framework.(19)

First, we categorized studies based on the indication for exercise (cardiac rehabilitation, pulmonary rehabilitation, and other). Next, prognostic factors were identified and categorized within themes (based on a consensus meeting within the investigative team). Where a prognostic factor was reported by two or more studies, the strength and quality of the association of the predictive factor with adherence was assigned using the GRADE framework. This process applies 8 criteria that can upgrade or downgrade the quality of evidence supporting a prognostic factor, and allows for evidence of a review of prognostic factors to be efficiently summarized for end-users.(19)

We also calculated descriptive statistics for the overall collection of included studies, as well as by indication for exercise. Overall adherence rates were calculated and averaged across all studies, as well as by exercise indication category. Adherence measures were separated based on measurement on a continuous scale (i.e., proportion of prescribed exercise completed) or as binary measurements (i.e., adherent vs not adherent).

Results

Study selection

The search strategy identified 982 records; 970 remained after duplicates were removed. Following title and abstract screening, 247 full text articles were assessed for eligibility and 23 were included. Study selection and reasons for exclusion are presented in Fig. 1.

Study characteristics

Study characteristics are presented in Table 1. Nineteen observational studies and 4 randomized controlled trials were included. A total of 5785 individuals were prescribed exercise across all studies (sample sizes ranged from 23-1218 participants) and average age ranged from 66–79 years. Indications for exercise included cardiac rehabilitation (n = 6), pulmonary rehabilitation (n = 7), and other (n = 10; including surgical, medical and neurologic indications). Most (20/23 (87%)) exercise programs were supervised.

Table 1
Study Characteristics

Author	Year	Design	N	Average age	Medical indication	Exercise program	Adherence	Adherence definition
Ades et al.(33)	1992	OBS	226	70	MI or CABG	CR ^a	21% ^c	Entry into the CR
Aherne et al.(46)	2017	OBS	98	69	PVD	Other ^a	N/A ^b	Number of sessions attended
Brown et al.(34)	2016	OBS	440	66	COPD	PR ^a	52% ^c	Number of sessions attended
Casey et al.(20)	2008	OBS	600	66	CVD	CR ^a	78% ^c	Staff judgement
Covey et al.(47)	2014	RCT	113	68	COPD	PR ^a	93% ^b	Percent of exercise completed
Cox et al.(35)	2013	OBS	85	68	Cognitive impairment	Other	78% ^b	Self-reported
Craike et al.(21)	2016	OBS	52	67	Prostate cancer	Other ^a	80% ^b	Number of sessions attended
Fan et al.(22)	2008	OBS	1218	67	COPD	PR ^a	79% ^b	Number of sessions attended
Gallagher et al.(23)	2003	OBS	196	67	CVD	CR ^a	32% ^c	Number of sessions attended
Hogg et al.(24)	2012	OBS	812	> 65	COPD	PR ^a	54% ^c	Number of sessions attended
Jensen et al.(25)	2016	OBS	50	69	Bladder cancer	Other	66% ^c	Self-reported
Mangione et al.(48)	2005	RCT	23	79	Hip fracture	Other ^a	98% ^b	Number of sessions attended
Messer et al.(36)	2007	OBS	164	66	Incontinence	Other ^a	70% ^c	Self-reported
Mudge et al.(26)	2013	OBS	140	> 65	CVD, pulmonary disease	Other ^a	42% ^b	Number of sessions attended
Pakzad et al.(27)	2013	OBS	30	66	CVD	CR ^a	N/A ^b	Number of sessions attended
Pandey et al.(37)	2017	RCT	40	67	Diabetes	Other ^a	70% ^b	Self-reported
Pickering et al.(28)	2013	OBS	70	73	Parkinson's disease	Other ^a	79% ^b	Percent of exercise completed
Rizk et al.(38)	2015	RCT	35	67	COPD	PR ^a	75% ^b	Percent of exercise completed
Selzler et al.(29)	2016	OBS	64	69	COPD	PR ^a	81% ^b	Number of sessions attended
Selzler et al.(30)	2012	OBS	814	68	COPD	PR ^a	83% ^b	Number of sessions attended
Tiedemann et al.(39)	2012	OBS	76	67	Stroke	Other ^a	60% ^b	Number of sessions attended
Tooth et al.(31)	1993	OBS	30	66	MI	CR	93% ^{bf} , 87% ^{bg}	Percent of exercise completed
van Montfort et al. (32)	2016	OBS	409	66	PCI	CR ^a	N/A ^b	Number of sessions attended

CABG = coronary artery bypass graft; COPD = chronic obstructive pulmonary disease; CR = cardiac rehabilitation; CVD = cardiovascular disease; FEV1 = forced expiratory volume in 1 second; HADS = Hospital Anxiety and Depression Scale; HDL = high density lipoprotein; IMD = Index of Multiple Deprivation; MI = myocardial infarction; MRC = Medical Research Council; OBS = observational; PCI = primary coronary intervention; PR = pulmonary rehabilitation; PVD = peripheral vascular disease; RCT = randomized controlled trial; a = supervised exercise program; b = adherence as a continuous measure, c = adherence as a categorical threshold; d = % participation; e = % completion; f = % duration; g = % frequency

Adherence to prescribed exercise rates

Exercise adherence was measured as a continuous variable in 16 studies and as a categorical outcome with a specified cut-off (demarking adherent vs not) in the remaining 7 studies. Overall adherence rate was reported in 20 studies and ranged from 21–93% (mean 68%, standard deviation (SD) 23%). Adherence was highest for pulmonary rehabilitation (71%, SD 15%), and other indications (74%, SD 13%); cardiac rehabilitation had lower rates (55%, SD 33%). However, lack of variance measures around adherence estimates limited our ability to perform formal comparative meta-analysis or meta-regression.

Predictors of exercise adherence

Predictors of exercise adherence were grouped into the following clusters: demographic, psychological, program-related, medical condition severity, comorbidities, and other (see Supplementary Tables S1, S2, S3; Additional Files 1, 2, 3). Demographic factors were evaluated by 13 studies(20, 21, 30–32, 22–29) (Table 2), psychological factors by 14 studies(20, 21, 32–36, 22–24, 27–31) (Table 3), program-related factors by 2 studies,(37, 38) medical condition severity by 11 studies,(21, 22, 39, 23, 24, 28–32, 34) comorbidities by 8 studies(20, 25, 27, 29–31, 34, 35) and other predictors by 5 studies.(24, 26, 31, 33, 39)

Table 2
Demographic Predictors of Exercise Adherence

Study	Predictors	Direction	Theme
Casey et al. (2008)(20)	Age (years)	+	Age
	Employed (vs not employed/retired)	0	Employment
	Gender (male vs female)	0	Sex
Craike et al. (2016)(21)	Highest level of education (less than university degree vs university degree or higher)	0	Education
Fan et al. (2008)(49)	Age (per 1 year change)	0	Age
	Female gender	0	Sex
	Education reference: < high school		Education
	High school	+	
	Some college	+	
	> College	+	
Gallagher et al. (2003)(23)	Unemployed or retired (vs employed)	-	Employment
	Age > 70 (vs 55–70)	-	Age
Hogg et al. (2012)(24)	Deprivation quintile (IMD score) reference: IMD 6.86–28.1		Social status
	IMD 28.11–35.02	0	
	IMD 35.03–39.57	0	
	IMD 39.58–43.85	-	
	IMD 43.86–60.41	-	
Jensen et al. (2016)(25)	Gender (women vs men)	0	Sex
	Age (< 70 vs ≥ 70)	0	Age
Mudge et al. (2013)(26)	Retired from workforce (vs "working" and "not working")	+	Employment
	Age < 65 vs 65+	0	Age
	Sex (male vs female)	0	Sex
	Living alone vs living with family/others	0	Living status
Pakzad et al. (2013)(27)	Identity	0	
Pickering et al. (2013)(28)	Gender (male vs female)	0	Sex
	Living status (alone vs partner vs family/friends vs other)	0	Living status
	Age multiplicative decrease per 10 years	-	Age
Selzler et al. (2016)(29)	Age (years)	0	Age
Selzler et al. (2012)(30)	Age (years)	+	Age
Tooth et al. (1992)(31)	Scale of Status and Prestige (high score = lower social standing)	-	Social status
	Age (years)	0	Age
	Education (years)	0	Education
van Montfort et al. (2016)(32)	Female sex (vs male)	0	Sex
	Age (years)	0	Age

IMD = Index of Multiple Deprivation (0, the least deprived, to 86, the most deprived); Scale of Status and Prestige (1 to 7, where 1 represents occupations of the highest social standing); + = significant positive effect; 0 = no significant effect; - = significant negative effect

Table 3
Psychological Predictors of Exercise Adherence

Study	Predictors	Direction	Theme
Ades et al. (1992)(33)	Presence of depression before hospitalization	-	Depression
Brown et al. (2016)(34)	Beck Depression Index	0	Depression
Casey et al. (2008)(20)	Beck Depression Index (high scores, more depressed)	-	Depression
Cox et al. (2013)(35)	Baseline self-efficacy (higher)	+	Self-efficacy
Craike et al. (2016)(21)	Role functioning (higher)	+	
	Sexual activity	0	
Fan et al. (2008)(22)	State-Trait Anxiety Index ≥ 36	-	Anxiety
	Beck Depression Index ≥ 5	-	Depression
Gallagher et al. (2003)(23)	Perceived control	0	Control
	Personal stressful event	-	
Hogg et al. (2012)(24)	Hospital Anxiety and Depression Score "Not depressed" 0–7	reference	Depression
	"Risk of depression" 8–10	0	
	"Depressed" 11	-	
Messer et al. (2007)(36)	Task self-efficacy summary scores (higher)	+	Self-efficacy
	Regulatory self-efficacy summary scores (higher)	+	
	Knowledge self-efficacy	0	
van Montfort et al. (2016)(32)	Positive affect	0	
Pakzad et al. (2013)(27)	State-Trait Anxiety Index (higher)	+	Anxiety
	Consequences	0	
	Chronology (acute/chronic)	0	
	Treatment control	0	
	Personal control	0	
Pickering et al. (2013)(28)	EQ-5D state of health thermometer	+	
	EQ-5D No pain/discomfort	reference	
	EQ-5D Moderate pain/discomfort	0	
	EQ-5D Extreme pain/discomfort	-	
	EQ-5D Not anxious/depressed	reference	Anxiety, Depression
	EQ-5D Moderate anxious/depressed	-	
	EQ-5D Extreme anxious/depressed	-	
	Mental health problem (self-reported)	-	Mental health
Selzler et al. (2012)(30)	Social functioning (36-Item Short Form Survey)	+	
	Mental health (36-Item Short Form Survey)	+	Mental health
	Role emotional (36-Item Short Form Survey)	+	
Selzler et al. (2016)(29)	Task self-efficacy	+	Self-efficacy
	Coping self-efficacy	0	
	Scheduling self-efficacy	0	
Tooth et al. (1992)(31)	Expectations (higher)	+	
	Psychological status (profile of mood states score)	0	
+ = significant positive effect; 0 = no significant effect; - = significant negative effect			

GRADE recommendations

Prognostic factors, categorized by themes, reported by at least 2 observational studies were assessed using the GRADE framework (Table 4).

Table 4
Grading of Recommendations Assessment, Development and Evaluation

Predictors	Participants	Studies	+	0	-	Phase	Limitations	Inconsistency	Indirectness	Imprecision	Publication bias	↑ effect size	Do not
Demographic													
Age (older)	3591	10	2	6	2	2	✓	X	✓	✓	✓	∅	∅
Sex (male)	2487	6		6		2	✓	✓	✓	✓	✓	∅	∅
Employed	936	3	1	1	1	2	✓	X	✓	✓	X	∅	∅
More education	1300	3	1	2		2	✓	X	✓	X	X	∅	∅
Living alone	210	2		2		2	✓	X	✓	✓	X	∅	∅
Lower SES	842	2			2	2	✓	✓	✓	X	X	∅	∅
Psychological													
Anxiety	1318	3	1		2	2	✓	X	✓	✓	X	∅	∅
Depression	3366	6		1	5	2	✓	✓	✓	✓	✓	∅	D
Higher self-efficacy	313	3	3			2	✓	✓	✓	X	X	∅	D
Higher control	226	2		2		2	✓	✓	✓	X	X	∅	∅
Good mental health	884	2	2			2	✓	✓	✓	✓	X	∅	∅
Comorbidities													
High BMI	1848	3		3		2	✓	✓	✓	✓	X	∅	∅
Smoker	1446	5	1	2	2	2	✓	X	✓	✓	X	∅	∅
High cholesterol	158	3	1	2		2	✓	X	✓	X	X	∅	∅
Hypertension	128	2		2		2	✓	✓	✓	X	X	∅	∅
higher CCI	1268	2		2		2	✓	✓	✓	✓	X	∅	∅
Condition severity													
Better respiratory function	878	2	1	1		2	✓	X	✓	✓	X	∅	∅
Higer FEV1	1658	2	1	1		2	✓	X	✓	✓	X	∅	∅
Program													
Farther distance	1444	2			2	2	✓	✓	✓	✓	X	∅	∅
Continuous exercise (vs intermittent)	75	2*	1		1	2	✓	X	✓	✓	X	∅	∅
Other													
Exercise history	160	2	1	1		2	✓	X	✓	X	X	∅	∅
+ = number of studies with a significant positive effect; 0 = number of studies with no significant effect; - = number of studies with a significant negative effect; Limitations: X = serious limitations; D = present; AE = not present; * randomized controlled trials													

Demographics

Demographic predictors included age, sex or gender, employment, education, social status and living situation. There was low-quality evidence that lower socioeconomic status predicted lower adherence. High-quality evidence suggested that sex was not predictive of adherence and moderate-quality evidence suggests that age does not predict adherence. Low, low and very low-quality evidence, respectively, suggested a lack of prediction of adherence for employment status, living status and education.

Psychological factors

Psychological predictors included anxiety, depression, self-efficacy, control, and self-rated mental health. High-quality evidence supported a negative association between the presence of depression and adherence. Individuals who had good self-rated mental health and who had good self-efficacy were more likely to be adherent (moderate-quality evidence). Low-quality evidence suggested that anxiety and perception of control did not predict adherence.

Comorbidities

Identified comorbidities reported as predictors of exercise adherence were Body Mass Index (BMI), smoking status, hypercholesterolemia, hypertension, and Charleston Comorbidity Index (CCI). None of these were predictive of exercise adherence, which was supported by moderate-quality evidence for BMI and CCI, low-quality evidence for smoking status and hypertension, and very low-quality evidence for hypercholesterolemia. Frailty was not assessed or reported in any of the studies.

Medical condition severity

Measures of respiratory disease severity were not found to be predictive of adherence, but this was only supported by low-quality evidence.

Program factors

The type of exercise program (continuous vs interval exercise) was evaluated by two randomized controlled trials. Although randomized trials are considered to provide high-quality evidence, we downgraded the evidence of no association to moderate quality, given that trial findings were contradictory (one trial reported better adherence to interval exercise, one reported better adherence with continuous exercise). Moderate-quality evidence suggests that living a further distance from the exercise facility decreased adherence.

Other

Low-quality evidence suggests that a history of exercise participation is not predictive of exercise adherence.

Risk of bias within studies

Nine observational studies were deemed to be at low risk of bias and 10 were at moderate risk of bias; no studies were at high risk of bias (Supplementary Table S5). Importantly, prognostic factor measurement and study confounding components of the tool scored low risk of bias across all studies. All four randomized trials were assessed as high risk of bias due to lack of blinding, however, this is recognizably difficult in exercise interventions (Supplementary Table S6). All other domains were low or unclear risk of bias.

Discussion

In this systematic review of predictors of exercise adherence in older adults with non-musculoskeletal indications for prescribed exercise, we found that positive predictors of adherence, supported by moderate-quality evidence, were higher self-efficacy and good self-rated mental health. Negative predictors included depression (high-quality) and distance from the exercise facility (moderate quality). Comorbidity status, sex and age did not appear to be predictive of adherence, supported by moderate- to high-quality evidence. As prescribed exercise programs are less likely to be effective without high levels of adherence, these findings provide important insights into current practice and future research. In current practice, identification of negative predictors, with a particular focus on mental health, could allow for increased personalization and targeting of support. The small number of identified predictors with at least moderate-quality evidence and sparse data available for many predictors suggest that future research is needed to better understand and predict poor exercise adherence in older adults.

Numerous studies have estimated exercise adherence rates in a variety of populations, typically reporting similar or slightly higher adherence rates than those identified in our study. For example, Bullard et al.(40) reported a pooled adherence rate of 77% (95% CI 68%, 84%) across 30 studies of adults with cancer, cardiovascular disease or diabetes. However, few studies have evaluated what patient- and program-factors predict adherence, and to our knowledge, none have evaluated the strength of this evidence using a standard framework such as GRADE. Most available data currently focuses on program-related factors. Similar to our findings, Morgan et al.(41) identified program location as a barrier to participation and adherence, while Sheill et al.(42) found that difficulties travelling to exercise locations were a substantial barrier for individuals with advanced cancer. We found no evidence that the type of exercise program (i.e., interval vs continuous exercise) was predictive of adherence, which is consistent with recommendations that the act of engaging in exercise is likely of greater importance than the specific type of exercise performed.(40, 42)

Some authors have advocated the identification of participant-level 'red flags' to adherence as a way to personalize exercise program design and support.(40) However, this approach requires a thorough understanding of what participant characteristics may act as red flags. At the participant level, consistent findings from our study and from others suggest that aspects of mental health are likely key predictors of adherence. Self-efficacy has previously been reported as a predictor of adherence in a systematic review of home-based physiotherapy,(43) which is consistent with our findings and aligns with other systematic reviews that have found one's intentions to engage in health-changing behaviors to be strongly predictive of adherence.(44) We also found that the presence of depression was a strong predictor of poor adherence and the only predictor supported by high-quality evidence. The related concept of good self-rated mental health (to some degree the inverse of depression) had moderate quality evidence supporting its role as a positive predictor of adherence. Whether anxiety predicts adherence in older people remains to be determined; we found no clear evidence of an association, as the strength and quality of evidence was low and reflected findings from only 3 studies. Interestingly, we did not find evidence that comorbidities, sex, or age were important predictors of adherence, as none suggested a directional association. Obesity and multimorbidity were also the only comorbidities with at least moderate quality evidence. Many comorbidities were not assessed and the impact of frailty was not reported in any studies, suggesting a need for future research. Finally, absent from the literature and related reviews is the consideration that program factors may interact with participant factors when predicting adherence. Although we were unable to identify any evidence of this phenomenon in our review, future evaluation is likely warranted to understand how, for example, participant-level red

flags such as poor mental health may potentially be modified by specifically targeted aspects of program design. Such efforts could lead to better personalization and potentially higher adherence in individuals at risk of poor participation.

Strengths and limitations

Our study's findings should be considered in the context of its strengths and limitations. First, we conducted our review according to best-practice methodologies, which included protocol pre-registration, peer-review of our search strategy, review of multiple databases, a focus on adjusted estimates and contextualisation of our findings within the GRADE strength of evidence framework. Furthermore, our results are based on identified studies that were generally at low or moderate risk of bias (apart from blinding issues in randomized trials, which is typical of exercise studies). However, despite pre-specifying a defined population of interest, included studies represented a somewhat heterogeneous group of participants who engaged in exercise for cardiovascular, pulmonary and other indications. We were also unable to identify adequately homogenous data to support quantitative meta-analyses. This may, in part, reflect the number of largely unvalidated measures used to define exercise adherence in clinical research.⁽⁴⁵⁾ Accordingly, we classified our studies based on whether adherence was measured using a continuous or binary definition; however, this may not have completely captured the heterogeneity in underlying adherence measures.

Conclusions

Design of prescribed exercise programs for older adults requires an understanding of how program and participant characteristics impact exercise adherence. Based on the GRADE Framework for prognostic research, mental health factors appear to be the most important patient-level predictors, while a longer distance from the exercise facility was the only clear program-related factor predicting adherence. These findings can help to inform the design of current programs and personalization of support for participants. Future research is needed to evaluate the impact of other patient- and program-level predictors.

Abbreviations

QUIPS: Quality in Prognostic Studies; GRADE: Grading of Recommendations Assessment, Development and Evaluation; BMI: Body Mass Index; CCI: Charleston Comorbidity Index

Declarations

Ethics Approval and Consent to Participate:

Not applicable

Consent for publication:

Not applicable

Availability of Data and Materials:

Not applicable

Competing Interests:

Use of Distiller SR was supported by the Department of Anesthesiology & Pain Medicine at The Ottawa Hospital. All authors declare that they have no competing interests.

Funding:

This research was supported by the Canadian Frailty Catalyst Grant. DIM acknowledges salary support from The Ottawa Hospital Anesthesiology Alternate Funds Association and the Canadian Anesthesiologists' Society Career Scientist Award. This research was conducted with financial support from the Canadian Frailty Network (CFN), which is supported by the Government of Canada through the Networks of Centres of Excellence (NCE) program. CFN had no role in the study design, data collection, analysis or interpretation of the results, or preparation of the manuscript.

Authors' Contributions:

All authors made substantial contributions to conception and design, or acquisition of data, or analysis and interpretation of data; drafting the manuscript or revising it critically for important intellectual content; and final approval of the version to be published.

Acknowledgements:

Not applicable.

References

1. Statistics Canada. Population Projections for Canada (2018 to 2068), Provinces and Territories (2018 to 2043). 2019.
2. Guralnik JM, Ferrucci L, Simonsick EM, Salive ME, Wallace RB. Lower-extremity function in persons over the age of 70 years as a predictor of subsequent disability. *N Engl J Med*. 1995;332(9):556–62.
3. Fried LP, Tangen CM, Walston J, Newman AB, Hirsch C, Gottdiener J, et al. Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci*. 2001;56(3).
4. Rockwood K, Song X, MacKnight C, Bergman D, Hogan DB, McDowell I, et al. A global clinical measure of fitness and frailty in elderly people. *C Can Med Assoc J = J l'Association médicale Can*. 2005;173(5):489–95.
5. McIsaac DI, Jen T, Mookerji N, Patel A, Lalu MM. Interventions to improve the outcomes of frail people having surgery: A systematic review. Quinn TJ, editor. *PLoS One*. 2017 Dec;12(12):e0190071.
6. Taylor RS, Brown A, Ebrahim S, Jolliffe J, Noorani H, Rees K, et al. Exercise-based rehabilitation for patients with coronary heart disease: systematic review and meta-analysis of randomized controlled trials. *Am J Med*. 2004;116(10):682–92.
7. Picorelli AMA, Pereira LSM, Pereira DS, Felício D, Sherrington C. Adherence to exercise programs for older people is influenced by program characteristics and personal factors: A systematic review. *J Physiother*. 2014;60(3):151–6.
8. Higgins J, Green S, editors. *Cochrane Handbook for Systematic Reviews of Interventions*. 5.1.0. The Cochrane Collaboration; 2011.
9. Stroup DF, Berlin JA, Morton SC, Olkin I, Williamson GD, Rennie D, et al. Meta-analysis of observational studies in epidemiology: a proposal for reporting. Meta-analysis Of Observational Studies in Epidemiology (MOOSE) group. *JAMA*. 2000 Apr;283(15):2008–12.
10. Riley RD, Moons KGM, Snell KIE, Ensor J, Hooft L, Altman DG, et al. A guide to systematic review and meta-analysis of prognostic factor studies. *BMJ*. 2019;364.
11. Moher D, Liberati A, Tetzlaff J, Altman DG. The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *PLoS Med*. 2009;6(7).
12. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol*. 2016;75:40–6.
13. Jordan JL, Holden MA, Mason EE, Foster NE. Interventions to improve adherence to exercise for chronic musculoskeletal pain in adults. *Cochrane database Syst Rev*. 2010 Jan;(1):CD005956.
14. Crandall S, Howlett S, Keysor JJ. Exercise Adherence Interventions for Adults With Chronic Musculoskeletal Pain. *Phys Ther*. 2013 Jan;93(1):17–21.
15. Jordan JL, Foster NE, Holden MA, Mason EE. Interventions to improve adherence to exercise for chronic musculoskeletal pain in adults. In: Jordan JL, editor. *Cochrane Database of Systematic Reviews*. Chichester: John Wiley & Sons, Ltd; 2006.
16. Geneen LJ, Moore RA, Clarke C, Martin D, Colvin LA, Smith BH. Physical activity and exercise for chronic pain in adults: an overview of Cochrane Reviews. *Cochrane Database Syst Rev*. 2017 Apr.
17. Nicolson PJA, Bennell KL, Dobson FL, Van Ginckel A, Holden MA, Hinman RS. Interventions to increase adherence to therapeutic exercise in older adults with low back pain and/or hip/knee osteoarthritis: a systematic review and meta-analysis. *Br J Sports Med*. 2017 May;51(10):791–9.
18. Hayden JA, van der Windt DA, Cartwright JL, Co P. Assessing Bias in Studies of Prognostic Factors. *Ann Intern Med*. 2013;158(4):280–6.
19. Huguët A, Hayden JA, Stinson J, McGrath PJ, Chambers CT, Tougas ME, et al. Judging the quality of evidence in reviews of prognostic factor research: adapting the GRADE framework. *Syst Rev*. 2013 Sep;2:71.
20. Casey E, Hughes JW, Waechter D, Josephson R, Rosneck J. Depression predicts failure to complete phase-II cardiac rehabilitation. *J Behav Med*. 2008;31(5):421–31.
21. Craike M, Gaskin CJ, Courneya KS, Fraser SF, Salmon J, Owen PJ, et al. Predictors of adherence to a 12-week exercise program among men treated for prostate cancer: ENGAGE study. *Cancer Med*. 2016;5(5):787–94.
22. Fan VS, Giardino ND, Blough DK, Kaplan RM, Ramsey SD, Fishman AP, et al. Costs of pulmonary rehabilitation and predictors of adherence in the National Emphysema Treatment Trial. *COPD J Chronic Obstr Pulm Dis*. 2008;5(2):105–16.
23. Gallagher R, McKinley S, Dracup K. Predictors of women's attendance at cardiac rehabilitation programs. *Prog Cardiovasc Nurs*. 2003;18(3):121–6.
24. Hogg L, Garrod R, Thornton H, McDonnell L, Bellas H, White P. Effectiveness, attendance, and completion of an integrated, system-wide pulmonary rehabilitation service for COPD: Prospective observational study. *COPD J Chronic Obstr Pulm Dis*. 2012;9(5):546–54.
25. Jensen BT, Laustsen S, Jensen JB, Borre M, Petersen AK. Exercise-based pre-habilitation is feasible and effective in radical cystectomy pathways—secondary results from a randomized controlled trial. *Support Care Cancer*. 2016;24(8):3325–31.
26. Mudge A, Adsett J. Factors Predicting Successful Transition to Community-based Maintenance Exercise Programs Following Exercise Rehabilitation. *Cardiopulm Phys Ther J*. 2013;24(4):18–24.
27. Pakzad S, Charette Y, Bourque P. Liens entre la perception de la maladie cardiovasculaire et la participation aux programmes de réadaptation cardiaque: une étude exploratoire. *Can J Cardiovasc Nurs*. 2013;23(4):19–25.
28. Pickering RM, Fitton C, Ballinger C, Fazakarley L, Ashburn A. Self reported adherence to a home-based exercise programme among people with Parkinson's disease. *Park Relat Disord*. 2013;19(1):66–71.
29. Selzler AM, Rodgers WM, Berry TR, Stickland MK. The importance of exercise self-efficacy for clinical outcomes in pulmonary rehabilitation. *Rehabil Psychol*. 2016;61(4):380–8.

30. Selzler AM, Simmonds L, Rodgers WM, Wong EYL, Stickland MK. Pulmonary rehabilitation in chronic obstructive pulmonary disease: Predictors of program completion and success. *COPD J Chronic Obstr Pulm Dis*. 2012;9(5):538–45.
31. Tooth L, McKenna K, Colquhoun D. Prediction of Compliance with a Post-Myocardial Infarction Home-Based Walking Program. *Aust Occup Ther J*. 1993;40(1):17–22.
32. van Montfort E, Denollet J, Widdershoven J, Kupper N. Interrelation and independence of positive and negative psychological constructs in predicting general treatment adherence in coronary artery patients – Results from the THORESCI study. *J Psychosom Res*. 2016;88:1–7.
33. Ades PA, Waldmann ML, Mccann WJ, Weaver SO. Predictors of Cardiac Rehabilitation Participation in Older Coronary Patients. *Arch Intern Med*. 1992;152(5):1033–5.
34. Brown AT, Hitchcock J, Schumann C, Wells JM, Dransfield MT, Bhatt SP. Determinants of successful completion of pulmonary rehabilitation in COPD. *Int J COPD*. 2016;11(1):391–7.
35. Cox KL, Flicker L, Almeida OP, Xiao J, Greenop KR, Hendriks J, et al. The FABS trial: A randomised control trial of the effects of a 6-month physical activity intervention on adherence and long-term physical activity and self-efficacy in older adults with memory complaints. *Prev Med (Baltim)*. 2013;57(6):824–30.
36. Messer KL, Hines SH, Raghunathan TE, Seng JS, Diokno AC, Sampsel CM. Self-efficacy as a predictor to PFMT adherence in a prevention of urinary incontinence clinical trial. *Heal Educ Behav*. 2007;34(6):942–52.
37. Pandey A, Suskin N, Poirier P. The Impact of Burst Exercise on Cardiometabolic Status of Patients Newly Diagnosed With Type 2 Diabetes. *Can J Cardiol*. 2017;33(12):1645–51.
38. Rizk AK, Wardini R, Chan-Thim E, Bacon SL, Lavoie KL, Pepin V. Acute responses to exercise training and relationship with exercise adherence in moderate chronic obstructive pulmonary disease. *Chron Respir Dis*. 2015 Nov;12(4):329–39.
39. Tiedemann A, Sherrington C, Dean CM, Rissel C, Lord SR, Kirkham C, et al. Predictors of adherence to a structured exercise program and physical activity participation in community dwellers after stroke. *Stroke Res Treat*. 2012;2012.
40. Bullard T, Ji M, An R, Trinh L, MacKenzie M, Mullen SP. A systematic review and meta-analysis of adherence to physical activity interventions among three chronic conditions: Cancer, cardiovascular disease, and diabetes. *BMC Public Health*. 2019;19(1):1–11.
41. Morgan F, Battersby A, Weightman AL, Searchfield L, Turley R, Morgan H, et al. Adherence to exercise referral schemes by participants – what do providers and commissioners need to know? A systematic review of barriers and facilitators. *BMC Public Health*. 2016;16(1):227.
42. Sheill G, Guinan E, Brady L, Hevey D, Hussey J. Exercise interventions for patients with advanced cancer: A systematic review of recruitment, attrition, and exercise adherence rates. *Palliat Support Care*. 2019 Dec;17(6):686–96.
43. Essery R, Geraghty AWA, Kirby S, Yardley L. Predictors of adherence to home-based physical therapies: a systematic review. *Disabil Rehabil*. 2017 Mar;39(6):519–34.
44. Husebø AML, Dyrstad SM, Søreide JA, Bru E. Predicting exercise adherence in cancer patients and survivors: a systematic review and meta-analysis of motivational and behavioural factors. *J Clin Nurs*. 2013 Jan;22(1–2):4–21.
45. McLean S, Holden MA, Potia T, Gee M, Mallett R, Bhanbhro S, et al. Quality and acceptability of measures of exercise adherence in musculoskeletal settings: A systematic review. *Rheumatol (United Kingdom)*. 2017;56(3):426–38.
46. Aherne TM, Kheirleiseid EAH, Boland M, Carr S, Al-Zabi T, Bashar K, et al. Supervised exercise therapy in the management of peripheral arterial disease – An assessment of compliance. *Vasa - Eur J Vasc Med*. 2017;46(3):219–22.
47. Covey MK, Collins EG, Reynertson SI, Dilling DF. Resistance training as a preconditioning strategy for enhancing aerobic exercise training outcomes in COPD. *Respir Med*. 2014;108(8):1141–52.
48. Mangione KK, Craik RL, Tomlinson SS, Palombaro KM. Can Elderly Patients Who Have Had a Hip Fracture Perform Moderate- to High-Intensity Exercise at Home? *Phys Ther*. 2005;85(8):727–39.
49. Ugolini G, Pasini F, Ghignone F, Zattoni D, Bacchi Reggiani ML, Parlanti D, et al. How to select elderly colorectal cancer patients for surgery: a pilot study in an Italian academic medical center. *Cancer Biol Med*. 2015 Dec;12(4):302–7.

Figures

Figure 1. PRISMA Flow Diagram for Study Selection and Inclusion

Figure 1

PRISMA Flow Diagram for Study Selection and Inclusion

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile6.docx](#)
- [AdditionalFile5.docx](#)
- [AdditionalFile4.docx](#)
- [AdditionalFile3.docx](#)
- [AdditionalFile2.docx](#)
- [AdditionalFile1.docx](#)