

Methicillin Resistant Staphylococcus aureus nasal carriage and associated factors in a rural tertiary hospital in Eastern Uganda: A prospective cross-sectional study

Nobert Thembo (✉ nobertt50@gmail.com)

Mbale Regional Referral Hospital

George Masifa

Mbale Regional Referral Hospital

Gerald Kamugisha

Mbarara University of Science and Technology

Robinah Nabitaka

Mbarara University of Science and Technology

Benjamin Akais

Mbarara University of Science and Technology Faculty of Medicine

Peter Olupot-Olupot

Mbale Regional Referral Hospital

Edward Kiyonga

Mbarara University of Science and Technology Faculty of Medicine

Rita Muhindo

Mbale Regional Referral Hospital

Lucas Ampaire

Mbarara University of Science and Technology

Research article

Keywords: Nasal carriage, Staphylococcus aureus, Methicillin Resistance

Posted Date: August 28th, 2019

DOI: <https://doi.org/10.21203/rs.2.13651/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Asymptomatic carriage of Methicillin-Resistant *Staphylococcus aureus* (MRSA) can predispose the host to a wide array of infections that would pose a challenge in the management of the cases in a current era encompassed with antibiotic resistance. To inform public health strategies, the study sought to describe MRSA nasal carriage frequencies and the associated factors concerning nasal carriage among patients attending Mbale Regional Referral Hospital (MRRH).

Methods: Two hundred eighteen consented participants presenting to the hospital for medical care between January and April 2018 were recruited to participate in this study. Sampling was done on both anterior nares using a pre-moistened swab and then transported to the laboratory at Room temperature for the detection of MRSA. Culture was performed on blood agar and plates incubated at 37°C for 24 hours. Identification of *Staphylococcus aureus* was done using conventional biochemical tests. MRSA was detected phenotypically using cefoxitin (30µg) as a surrogate test as per the Clinical Laboratory Standard Institute guidelines 2017 version. Patient demographic characteristics and the MRSA nasal carriage factors were collected using a pre-tested questionnaire. The collected raw data was entered into excel and later imported into STATA software for analysis.

Results: Overall, Majority of the participants were in-patients (138, 63.3%) with the proportions of both females and males among the participants being 154/218 (70.6%) and 64/218 (29.3%) respectively. Mean age for both female and male participants was 40.16 (SD± 17.04) years respectively. *Staphylococcus aureus* nasal carriage rate among the participants was 22.9% (50/218), with 72% (36/50) of the harboured strains phenotypically expressing methicillin resistance. Colonization with Methicillin Resistant *Staphylococcus aureus* did not show any significant relationship with all the studied factors.

Conclusion: There was a moderate *Staphylococcus aureus* nasal carriage among the participants in Mbale Regional Referral Hospital. We also observed a highly displayed phenotypic expression of methicillin resistance among the isolated *Staphylococcus aureus* strains. The studied factors indicated an independent influence on the rate of nasal carriage. For surveillance purposes to combat future outbreaks, there is a need to do a larger study with better power to better draw generalizable conclusions of carriage in the population.

Background

Staphylococcus aureus is among the most significant human bacterial pathogens worldwide (1), with Methicillin-Resistant *Staphylococcus aureus* (MRSA) currently being the most common antibiotic-resistant strain in most of the hospital and community settings (2)(3). *Staphylococcus aureus* is also the second leading cause of nosocomial bacteremia (4) and has been highlighted as a priority organism of interest by the WHO (5). Asymptomatic nasal carriage of Methicillin-Resistant *Staphylococcus aureus*

(MRSA) can predispose the host to a wide array of infections that would pose a challenge in the management of the cases in the current era of antibiotic resistance (6)(7).

The magnitude of MRSA remains greatly unknown in developing countries yet surveillance systems to guide interventions require expertise and resources, which are inadequate(8). Studies conducted elsewhere by Conceição *et al.*, and Aiken *et al.*, documented MRSA prevalence rates of 26.9% and 7.0% respectively (9)(10). However, in resource-limited settings, not much has been done to explore on MRSA nasal carriage as reported by Bebell *et al.*, 2016 and yet MRSA is currently recognized as the leading cause of hospital-acquired infections (11). Empirical and inappropriate use of antibiotics, multiple-pathology and use of invasive devices have attributed to a high rate of MRSA hence extended hospital duration and high treatment costs among patients (12).

Identifying the source, reservoirs and vectors for the spread of antibiotic-resistant bacteria poses significant challenges in tracking antimicrobial resistance. The patient's carried endogenous microflora may all play a vital role in the cause of nosocomial infections (13)(14). In a study conducted on surgical wound infections in the hospital indicated that 65.9% of the cases were caused by MRSA strains (15). There are currently no documented reports on nasal colonization and carriage rates of MRSA among patients and associated carriage factors in Mbale Regional Referral Hospital. With most patients coming from different settings, our study aimed to determine the prevalence of *Staphylococcus aureus* and MRSA nasal colonization, and further identify the risk factors associated with asymptomatic carriage of MRSA.

Methods

Study design

This was a descriptive cross-sectional study carried out between January and April 2018 at Mbale Regional Referral Hospital on both in and Outpatients.

Study area

The study was conducted at both the Outpatient and In-patient departments (wards) of Mbale Regional Referral Hospital located in Mbale municipality. The hospital serves as both a regional referral and a tertiary teaching hospital for the medical school of Busitema University and School of clinical officers. It has a bed capacity of 400 and serves a population of approximately 4 million people.

Target population

The study targeted all patients seeking medical care at Mbale RRH who are asymptomatic carriers of MRSA. These included those from the inpatient (medical, surgical and maternity wards) and outpatient departments.

Inclusion Criteria

We included Patients aged 15 years and above, seeking medical care at either the In-patient or the Outpatient department at MRRH and had consented to take part in the study. Written informed consent for children aged 16 and 15 years was sought from their parents or guardians.

Recruitment of study participants

Consent was sought prior to recruiting of the participants. A cluster random sampling technique was used whereby participants were selected randomly with respect to the two broad clusters i.e. OPD and in-patients to ensure equal representative sampling. A questionnaire was administered to capture both demographic factors and predictor factors for MRSA nasal carriage.

Sample collection and transportation

From each participant, specimens for *Staphylococcus aureus* culture were collected from both the anterior nares using sterile broth moistened dry swabs. The swabs were immediately placed into a tube with Cary-Blair transport media (BBL™, BD bioscience), labelled with the participants' study number, initials, date and time of sample collection. Samples were immediately delivered to the Clinical Microbiology Laboratory, Mbale RRH for processing.

Isolation and identification of Staphylococcus aureus

The swabs were first inoculated on Blood Agar and incubated at 37°C aerobically for 24 hours after which plate reading was done. Gram staining was performed on discrete colonies displaying the culture characteristics of *Staphylococcus aureus* to confirm for Gram-positive cocci in clusters. A series of conventional biochemical tests such as catalase test using 3% hydrogen peroxide, bound and free coagulase using reconstituted rabbit lyophilized plasma (Remel, Europe, Ltd. Dartford, Kent DA2 6PT, UK) were performed to identify the *Staphylococcus aureus*. Further confirmation was done by sub-culturing of the isolated colonies of *Staphylococcus aureus* onto Mannitol Salt Agar (MSA, Oxoid, CM0085. Thermo-scientific, US) and incubated at 37°C for 24 hours aerobically. After 24 hours, the plates were examined for growth with interest in mannitol fermenting colonies that appeared as yellow colonies, measuring 1–2 mm in diameter and slightly raised.

Preservation of the Isolates

By use of a sterile pre-flamed wire loop, pure growing colonies were scraped off the Blood agar purity plates and suspended in 1ml of 15% glycerol broth the frozen at –80°C till needed for MRSA phenotypic testing.

Phenotypic detection for MRSA

Agar disk diffusion (Kirby Bauer) technique on 4% sodium chloride Mueller Hinton Agar (BBL™, BD) was employed. The inoculum was prepared by picking distinct colonies from a fresh pure loon culture on Blood agar and suspended into 5mls of a 0.85% saline water to make a bacterial suspension. The suspension was vortexed for 15 seconds and the turbidity adjusted visually by adding sufficient saline water to achieve a 0.5 McFarland standard. A sterile cotton swab was dipped into the suspension, rotated several times and the excess fluid removed by pressing on the sidewalls of the tube. The dried surface of the agar was evenly streaked with the sterile swab aseptically rotating the plate at 60° to ensure uniform distribution. A ceftioxin disc (30 µg, HIMedia, Mumbai, India) was placed on the surface of the inoculated plate using sterile forceps. The setup was incubated at 37°C for 24 hours aerobically. The plates were read after 24 hours of incubation and the zone diameters of inhibition were measured using a Vernier calliper. The measured zones of inhibition were recorded and interpretations were done following the interpretive cut-offs (MRSA_ ≤ 21 mm and MSSA _ ≥ 22 mm) as per the Clinical Laboratory Standard Institute (CLSI) guidelines (31).

Data Analysis

Raw data were entered into Microsoft Excel and later exported to Stata Corp. version 13 for analysis. Descriptive statistics including proportions, means were used to describe the participants and to determine prevalence. Associations were generated using Odds ratio at 95% CI and a *P-value* of ≤0.05. We performed a logistic regression to determine any association between the factors collected as predictors and MRSA nasal carriage as the outcome. We considered a univariate analysis at a level of significance of 0.2 to describe the categorical variables by sex and obtain baseline characteristics.

Quality Control

Standard Operating procedures were followed, aseptic transfer techniques were performed, reagents were kept at 2–8°C, and the questionnaire was pretested before the commencement of the study.

Staphylococcus aureus ATCC 25923 was used as the positive control strain for the identification of biochemical tests and susceptibility tests on 4% sodium chloride Muller Hinton Agar.

Results

Demographic characteristics of patients

The study recruited 218 participants aged 15 years and above presenting to Mbale Regional Referral Hospital for medical care. Majority of the participants were in-patients 138/218 (63.3%) with the proportions of both females and males among the participants being 154/218 (70.6%) and 64/218 (29.3%) respectively. Mean age for both female and male participants was 40.16 (SD± 17.04) years respectively as shown in table 1 below.

The proportion of patients with Staphylococcus aureus and MRSA nasal carriage attending Mbale Regional Referral Hospital

The prevalence rate for *Staphylococcus aureus* nasal carriage among the participants was 22.9% (50/218) as indicated in table 2 below. Proportions for MRSA nasal carriage among the participants were found to be 16.5% (36/218) and 6.4% (14/218) of whom were colonized with MSSA (table 3).

Distribution of Methicillin resistance among Staphylococcus aureus Isolates.

Of the 50 studied *Staphylococcus aureus* isolates, 72% (36/50) of the isolates were found to be phenotypically expressing methicillin resistance while 28% (14/50) were methicillin sensitive as seen in figure 1.

Multivariate analysis of studied factors for association with Methicillin Resistant Staphylococcus aureus nasal colonization.

Colonization with Methicillin Resistant *Staphylococcus aureus* was independently associated with age (OR = 0.9825889; 95% CI = 0.9612351–1.004417, *P*-value = 0.117), history of hospitalization (OR = 1.156944; 95% CI = 0.5689258–2.352712; *P*-value = 0.687), contact with animals (OR = 1.141084; 95% CI = 0.8069104–1.613653; *P*-value = 0.455), antibiotic use (OR = 0.7516896; 95% CI = 0.3804775–1.485974; *P*-value = 0.411) and HIV status (OR = 0.7215959; 95% CI = 0.2871594–2.107922; *P*-value = 0.551).

Discussion

Prevalence of *Staphylococcus aureus* nasal carriage

Results from this study indicated that the nasal carriage rates for *Staphylococcus aureus* among the participants was 22.9%, a rate significantly higher than that reported from studies from Thika Hospital, Kenya (8.9%), Nigeria (18.3%), and China (2.4%) (16)(17)(18). Studies conducted elsewhere by Bebell *et al*, Ouedraogo *et al*, Moniri *et al* and Ateba *et al* revealed slightly higher rates of 28.5%, 29.1%, 32.9%, 38% and 29% respectively compared to the finding in our study (11)(19)(20)(21). However, our finding was similar to reports from North Germany by Mehraj *et al.*, Northwest Ethiopia and Italy which documented *Staphylococcus aureus* nasal carriage rates of 21.9% and 23% respectively (22)(23). The difference in the various findings would have been attributed to the sample size used giving the different studies a better power to detect the *Staphylococcus aureus* nasal carriage. In addition, the studies were conducted in different geographical regions and the regions might have differing antibiotic usage practices hence differences in the carriage bio-films.

MRSA carriage Prevalence

In our study, we determined the prevalence of MRSA and MSSA among randomly selected participants who presented to the hospital for medical care. Our study documented an overall prevalence rate of 16.5% (36/218) for MRSA among the participants who were included in the study. However, we noted a very high 72% (36/50) methicillin resistance expression among the carried *Staphylococcus aureus* strains. The finding for the overall carriage rate (16.5%) among the participants was lower than the reported prevalence of 46% from Mulago hospital, Kampala, Uganda (24). However, our finding was higher than a prevalence rate of 2.8% reported by Bebell *et al* from Mbarara Hospital, Uganda (11). Our observed high (72%) methicillin resistance expression among the isolated *Staphylococcus aureus* strains was contrary to reports from three regional hospitals in Tanzania that is Muhimbili National Hospital (10.5%), Mwananyamala and Amana regional hospitals (24.7%) (25)(26). This could have been attributed to the high numbers of in-patients who were included in the study and duration of hospital stay could have had an effect of acquisition. In addition, the choice of the participants included could have affected the prevalence since our study did not consider the children age group.

5.4 Factors associated with Methicillin Resistant *Staphylococcus aureus* nasal carriage

In our study, colonization by Methicillin Resistant *Staphylococcus aureus* did not indicate any significant statistical association with all the tested variables. Nonetheless, we observed different levels of consistency with the findings of previous studies. Our study indicated that antibiotic usage did not increase the likelihood of colonization by Methicillin Resistant *Staphylococci aureus* (OR = 0.7516896; 95% CI = 0.3804775–1.485974; *P*-value = 0.411), while HIV status was found to have no effect on the nasal carriage rate even with low immunity (OR = 0.7215959; 95% CI = 0.2871594–2.107922; *P*-value = 0.551). The statistical association findings by Bhattacharya *et al* that reported no association with immune status or recent antibiotic use (27), were similar to the findings in our study. In contrary, studies by Soltan *et al*, Moniri *et al*, and reported findings indicating that previous antibiotic usage was significantly associated with MRSA carriage (28)(20). In addition, Studies by Amorim *et al*, Portugal and Rasamiravaka *et al*, Madagascar revealed an association of MRSA nasal carriage to the history of hospitalization and previous antibiotic use which was contrary to our findings that found no association of the studied factors to MRSA nasal carriage (29)(30). The no association with all the variable factors might have been attributed to by the difference in the characteristics and varied geographical location for the participants.

Conclusions

Our study revealed that there was a moderate *Staphylococcus aureus* nasal carriage among the participants but with high methicillin resistance expressed (72%) among the isolated strains. Assessment of the factors with the nasal carriage rates revealed that none of the factors significantly influenced carriage. This study has been the first in Mbale, Eastern Uganda to provide baseline data on *Staphylococcus aureus* and MRSA nasal carriage. Therefore, there is need to do extensional surveillance to understand the drivers and transmission patterns for *Staphylococcus aureus* and MRSA within the hospital and the community.

Abbreviations

MCRI: Mbale Clinical Research Institute

MRSA: Methicillin-Resistant *Staphylococcus aureus*

MRRH: Mbale Regional Referral Hospital

WHO: World Health Organization

OPD: Out Patient Department

CI: Confidence Interval

CLSI: Clinical Laboratory Standard Institute

ATCC: America Type Culture Control

SD: Standard Deviation

HIV: Human Immune Virus

MSSA: Methicillin Susceptible *Staphylococcus aureus*

OR: Odds ratios

MRRH-REC: Mbale Regional Referral Hospital Research Ethics Committee

CSV: Comma Separated Version Files

CoNS: Coagulase Negative *Staphylococcus*

N/A: Not Applicable.

Declarations

Acknowledgements

Ethics approval and consent to participate

Ethical approval was obtained from the Faculty of Medicine Research Committee at Mbarara University of Science and Technology (Ref: DMS/6) and Mbale Regional Referral Hospital Ethics Review Committee (Ref: MRRH-REC-IN-COM 110/2017). Written informed consent was obtained from the patients who had satisfied the study eligibility criteria for inclusion. The consenting process was done in a private room and all the participants were allocated identification numbers to ensure the confidentiality of the participants.

Consent for publication

Not applicable

Availability of data and materials

The data sets in comma separated version files (CSV) are available and have been attached as supplementary information files.

Competing interests.

The authors have no competing interests

Funding

The authors pooled funds to support this project from design, data collection, and data analysis through manuscript writing.

Author contributions

This work was conducted in collaboration between all authors. Authors NT, GK, RN, BA and EK designed the study, wrote the protocol, participated in the data and statistical analysis process, literature searches and wrote the first draft of the manuscript. Author GM did the original conceptualization, supervised the fieldwork laboratory analysis and participated in the write-up of the manuscript. Authors POO and RM participated in the manuscript writing and proof reading. Author LA did the overall supervision of the work right from the development of the protocol, statistical analysis and manuscript conceptualization. All authors read and approved the final manuscript.

We thank the administration and theatre staff of Mbale regional referral hospital for their cooperation and technical guidance. We also thank the academic staff of Mbarara University of Science and Technology, Faculty of Medicine for their tremendous mentorship.

References

1. Oguzkaya-Artan, M., Baykan, Z., Artan, C., & Avsarogullari L. Prevalence and risk factors for Methicillin Resistant Staphylococcus aureus carriage among emergency department workers and bacterial contamination on touch surfaces in Erciyes University Hospital, Kayseri, Turkey. *African Heal Sci.* 2015;15(4):1289–1294.
2. Shibabaw, A., Abebe, T., & Mihret A. Nasal carriage rate of Methicillin Resistant Staphylococcus aureus among Dessie Referral Hospital health care workers; Dessie, Northeast Ethiopia. *Antimicrob Resist Infect Control.* 2013;2(1):25.
3. El Aila, N. A., Al Laham, N. A., & Ayes BM. Nasal carriage of Methicillin Resistant Staphylococcus aureus among health care workers at Al Shifa hospital in Gaza Strip. *BMC Infect Dis.* 2017;17(1):28.

4. Bazira, J., Boum II, Y., Sempa, J., Iramiot, J., Nanjebe, D., Sewankambo, N., & Nakanjako D. Trends in antimicrobial resistance of *Staphylococcus aureus* Isolated from clinical samples at Mbarara Regional Referral Hospital in Rural Uganda. *Br Microbiol Res J*. 2014;4(10):1084.
5. Who. Global priority list of antibiotic-resistant bacteria to guide research, discovery, and development of new antibiotics [Internet]. 2017. Available from: https://www.who.int/.../WHO-PPL-Short_Summary_25Feb-ET_NM_WHO.pdf%0A
6. Daini, O. A., & Akano SA. Plasmid-mediated antibiotic resistance in *Staphylococcus aureus* from patients and non-patients. *Sci Res Essays*. 2009;4(4):346–350.
7. Albrich WC1 HS. Health-care workers: source, vector, or victim of MRSA? *Lancet Infect Dis*. 2008;8(5):289–301.
8. Bagheri Nejad S., B. Allegranzi, S. B. Syed BE and DP. Health-care-associated infections in Africa: a systemic review. *World Heal Organ*. 2011;89:757–76.
9. Conceição, T., Santos Silva, I., de Lencastre, H., & Aires-de-Sousa M. *Staphylococcus aureus* nasal carriage among patients and health care workers in São Tomé and Príncipe. *Microb Drug Resist*. 2014;20(1):57–66.
10. Aiken, A.M., Mutuku, I. M., Sabat, A. J., Akkerboom, V., Mwangi, J., Scott, J. A. G., ... & Grundmann H. Carriage of *Staphylococcus aureus* in Thika Level 5 Hospital, Kenya: a cross-sectional study. *Antimicrob Resist Infect Control*. 2014;3(1):22.
11. Bebell, L. M., Ayebare, A., Boum, Y., Siedner, M. J., Bazira, J., Schiff, S. J., ... & Firth PG. Prevalence and correlates of MRSA and MSSA nasal carriage at a Ugandan regional referral hospital. *J Antimicrob Chemother*. 2016;72(3):888–892.
12. Klein, E., Smith, D. L., & Laxminarayan R. Hospitalizations and deaths caused by Methicillin-Resistant *Staphylococcus aureus*, United States, 1999–2005. *Emerg Infect Dis*. 2007;13(12):1840.
13. Vonberg, R. P., Stamm-Balderjahn, S., Hansen, S., Zuschneid, I., Rüden, H., Behnke, M., & Gastmeier P. How often do asymptomatic healthcare workers cause methicillin-resistant *Staphylococcus aureus* outbreaks? A systematic evaluation. *Infect Control Hosp Epidemiol*. 2006;27(10):1123–7.
14. Dalstrom DJ, Venkatarayappa I, Manternach AL, Palcic MS, Heyse BA PM. Time-dependent contamination of opened sterile operating-room trays. *J Bone Jt Surg Am*. 2008;90(5):1022–5. pmid:18451394.
15. George, M., Iramiot, J. S., Muhindo, R., Olupot-Olupot, P., & Nanteza A. Bacterial Aetiology and Antibiotic Susceptibility Profile of Post-Operative Sepsis among Surgical Patients in a Tertiary Hospital in Rural Eastern Uganda. *Microbiol Res J Int*. 2018;24(2).
16. Aiken, A.M., Mutuku, I. M., Sabat, A. J., Akkerboom, V., Mwangi, J., Scott, J. A. G., ... & Grundmann H. Carriage of *Staphylococcus aureus* in Thika Level 5 Hospital, Kenya: a cross-sectional study. *Antimicrob Resist Infect Control*. 2014;3(1):22.
17. M. Okwu, S. Bamgbala and WA. Prevalence of nasal carriage of community-associated Methicillin-Resistant *Staphylococcus aureus* (CA-MRSA) among healthy primary school children in Okada, Nigeria. *Prevalence*. 2012;2(4):2224–3186.

18. J. Deng, G. Xiao, Y. Zhu, W. Zhou and CW. Staphylococcus aureus nasal carriage and its antibiotic resistance profiles in Tibetan school children in Southwest China. *Hong Kong J Pediatr.* 2014;19:75–78.
19. Ouedraogo, A. S., Dunyach-Remy, C., Kissou, A., Sanou, S., Poda, A., Kyelem, C. G., Solassol, J., Bañuls, A. L., Van De Perre, P. Ouédraogo, R. and Jean-Pierre, H. High nasal carriage rate of Staphylococcus aureus containing panton-valentine leukocidin-and EDIN-encoding genes in community and hospital settings in Burkina Faso. *Front Microbiol.* 2016;7:p.1406.
20. Moniri, R., Musav, G. A. and Fadavi N. The prevalence of nasal carriage Methicillin-Resistant Staphylococcus aureus in hospitalized patients. *Pak J Med Sci.* 25(4):pp.656–659.
21. Ngoa UA, Schaumburg F, Adegnika AA, Kösters K, Möller T, Gaus E, Fernandes JF, Alabi A, Issifou S, Becker K, Grobusch MP, Kreamsner PG LB. Epidemiology and population structure of Staphylococcus aureus in various population groups from a rural and semi urban area in Gabon, Central Africa. *Acta Trop.* 2012;124(1):42–7.
22. Mehraj J, Akmatov MK, Strömpl J, Gatzemeier A, Layer F, Werner G, Pieper DH, Medina E, Witte W, Pessler F KG. Methicillin-Sensitive and Methicillin-Resistant Staphylococcus aureus Nasal Carriage in a Random Sample of Non-Hospitalized Adult Population in Northern Germany. *PLoS One.* 2014;9(9):e107937.
23. Tigabu, A., Tiruneh, M., & Mekonnen F. Nasal Carriage Rate, Antimicrobial Susceptibility Pattern, and Associated Factors of Staphylococcus aureus with Special Emphasis on MRSA among Urban and Rural Elementary School Children in Gondar, Northwest Ethiopia: A Comparative Cross-Sectional Study. *Adv Prev Med.* 2018;
24. Kateete DP, Namazzi S, Okee M, Okeng A, Baluku H, Musisi NL, Katabazi FA, Joloba ML, Ssentongo R NF. High prevalence of methicillin resistant Staphylococcus aureus in the surgical units of Mulago hospital in Kampala, Uganda. *BMC Res notes.* 2011;4(1):326.
25. Moyo SJ, Nkinda L, Majigo M, Mmbaga E, Mbembati N, Aboud S LE. Prevalence of Methicillin-Resistant Staphylococcus aureus carriage on admission among patients attending regional hospitals in Dar es Salaam, Tanzania. *BMC Res Notes.* 2017;10(1):417.
26. oyo SJ, Aboud S, Blomberg B, Mkopi N, Kasubi M, Manji K, Lyamuya EF, Maselle SY LN. High nasal carriage of Methicillin-Resistant Staphylococcus aureus among healthy Tanzanian under–5 children. *Microb drug Resist.* 2014;1;20(1):82–8.
27. Alsan M, Schoemaker L, Eggleston K, Kammili N, Kolli P BJ. Out-of-pocket health expenditures and antimicrobial resistance in low-income and middle-income countries: an economic analysis. *Lancet Infect Dis.* 2012;15(10):1203–1210.
28. Soltani B, Ardakani AT, Moravveji A, Erami M, Rezaei MH, Moniri R NM. Risk factors for Methicillin Resistant Staphylococcus aureus nasal colonization of healthy children. *Jundishapur J Microbiol.* 7(9).
29. Amorim, M. L., Vasconcelos, C., Oliveira, D.C., Azevedo, A., Calado, E., Faria, N. A., ... & Cabeda JM. Epidemiology of Methicillin-Resistant Staphylococcus aureus (MRSA) nasal colonization among

patients and healthcare workers in a Portuguese hospital: a pre-intervention study toward the control of MRSA. *Microb Drug Resist.* 2009;15(1):19–26.

30. Rasamiravaka T, Rasoanandrasana S, Zafindraibe NJ, Alson AO RA. Evaluation of Methicillin-Resistant *Staphylococcus aureus* nasal carriage in Malagasy patients. *J Infect Dev Ctries.* 17;7(4):318–22.
31. Clinical and Laboratory Standards Institute. M100 Performance Standards for Antimicrobial Susceptibility Testing, 27th. 2017. 224 p.

Tables

Tables

Table 1: Demographic characteristics of patients attending Mbale RRH.

Demographic factors	Characteristics	Males (n=64)	Females (n=154)
Age	Mean age (40.16; SD± 17.04) in years	45	38
Animal contact	Yes	42	87
	No	22	67
Residence	Resident in town	18	62
	Resident in village	46	92
History of hospitalization	Has ever been hospitalized	21	52
	Never been hospitalized	42	102
Antibiotic usage	Use antibiotics	34	81
	Does not use antibiotics	30	73
HIV sero-status	Negative	45	107
	Positive	5	21
	Unknown	14	26
Isolated organism	CoNS	44	72
	No growth	9	43
	<i>Staphylococcus aureus</i>	11	39
Methicillin sensitivity	MRSA	11	25
	MSSA	0	14
	N/A	53	114

Key: MRSA- Methicillin Resistant *Staphylococcus aureus*, MSSA- Methicillin Susceptible *Staphylococcus aureus*, CoNS- Coagulase Negative *Staphylococcus*, n- proportions per gender

Table 2: Proportions of patients with *Staphylococcus aureus* nasal carriage.

Isolated Organism	Frequency (n)	Percentage (%)
<i>Staphylococcus aureus</i>	50	22.9
CoNS	116	53.2
Not Applicable	52	23.8
Total	218	100

Key: n-number of participants, %-prevalence rate

Methicillin Sensitivity	Frequency	Percentage	Cumulative frequency
MRSA	36	16.51	16.51
MSSA	14	6.42	22.94
Not Applicable	168	77.06	100.00
Total	218	100.00	

Table 3: Proportions of MRSA and MSSA among patients at Mbale RRH.

Key: MRSA-Methicillin Resistant *Staphylococcus aureus*, MSSA-Methicillin Susceptible *Staphylococcus aureus*

Methicillin Sensitivity	OR	P-value	95% [Confidence Interval]	
Age	0.9825889	0.117	0.9612351	1.004417
Residence	0.7119681	0.347	0.3505714	1.445921
History of Hospitalization	1.156944	0.687	0.5689258	2.352712
Contact with animals	1.141084	0.455	0.8069104	1.613653
Antibiotic use	0.7516896	0.411	0.3804775	1.485974
HIV status (Positive)	0.7215959	0.551	0.2470208	2.107922
(Unknown)	0.71886	0.481	0.2871594	1.799557

Table 4: The multivariate analysis of studied factors for association with Methicillin Resistant *Staphylococcus aureus* nasal colonization.

Key: OR-odds ratios

Figures




Figure 1

Distribution of Methicillin Sensitivity among carriers of Staphylococcus aureus Key: MSSA-Methicillin Susceptible Staphylococcus aureus, MRSA-Methicillin Resistant Staphylococcus aureus