

# Can indices of myocardial damage predict carbon monoxide poisoning outcomes?

**Hitoshi Koga**

St. Mary's Hospital

**Hideki Tashiro** (✉ [mdhidekit@mac.com](mailto:mdhidekit@mac.com))

St Mary's Hospital <https://orcid.org/0000-0002-8275-6375>

**Kouta Mukasa**

St. Mary's Hospital

**Tomohiro Inoue**

St Mary's Hospital

**Aya Okamoto**

St. Mary's Hospital

**Shougo Urabe**

St. Mary's Hospital

**Shuuichirou Sagara**

St Mary's Hospital

**Kazumi Yano**

St. Mary's Hospital

**Kouhei Onitsuka**

St Mary's Hospital

**Hisashi Yamashita**

St. Mary's Hospital

---

## Original Research

**Keywords:** Carbon monoxide poisoning, carboxyhemoglobin, QT interval, QT dispersion, troponin I

**Posted Date:** July 27th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-43628/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

## Background

Carbon monoxide causes electrical, functional, and morphological changes in the heart. It is not clear, however, whether the indices of myocardial damage can predict the patient's prognosis after carbon monoxide poisoning. This retrospective study aimed to investigate the relation between the carboxyhemoglobin level and electrocardiographic (ECG) changes and whether the ECG changes and troponin I levels are related to the patient's prognosis after carbon monoxide poisoning.

## Results

Carboxyhemoglobin, troponin I, and ECG parameters were measured in 70 patients with carbon monoxide poisoning. The QT and RR intervals were measured for each ECG lead in all patients, and the corrected QT interval and corrected QT dispersion were calculated. The correlation between the maximum corrected QT interval and the carboxyhemoglobin level was significant ( $P = 0.0072$ ,  $R^2 = 0.1017$ ), as were the relationships between QT dispersion and carboxyhemoglobin ( $P < 0.001$ ,  $R^2 = 0.2358$ ) and the corrected QT dispersion and carboxyhemoglobin ( $P < 0.001$ ,  $R^2 = 0.2613$ ). The multivariate logistic analysis showed that the significant predictors of sequential disability were hyperbaric oxygen therapy ( $P = 0.0182$ ), corrected QT dispersion ( $P = 0.0062$ ), and troponin I level ( $P = 0.0002$ ).

## Conclusions

Patients' prognosis following carbon monoxide poisoning can be predicted based on corrected QT dispersion and the troponin I level. Patients with myocardial damage should be monitored not only for their cardiovascular outcome but also for their neurological outcome and their prognosis.

## Background

Carbon monoxide is a colorless, odorless, non-irritating gas. It is produced endogenously in small amounts as a byproduct of the catabolism of heme molecules. It can also be inhaled when hydrocarbon-containing fuels are not completely burned<sup>1,2</sup>. Survivors of carbon monoxide poisoning may suffer neurological and psychiatric sequelae. Carbon monoxide causes electrical, functional, and morphological changes in the heart. The QT interval has long been known to vary significantly among the individual leads of a surface 12-lead electrocardiogram<sup>3</sup>.

A potential clinical application of this inter-lead difference was proposed in 1990 by Day et al., who suggested that the inter-lead difference in the QT interval might provide a measure of repolarization inhomogeneity, which they called "QT dispersion"<sup>4</sup>. Although it has been established that carbon monoxide induces electrocardiographic (ECG) changes and alterations of cardiac biomarkers<sup>3,5</sup>, it is not

clear whether the indices of myocardial damage can predict the patient's prognosis after carbon monoxide toxicity. Hence, this study aimed to investigate the relation between carboxyhemoglobin and ECG changes and whether the ECG changes and troponin I levels are related to the prognosis of patients with carbon monoxide poisoning.

## Methods

The study group of this retrospective study included 70 consecutive patients with carbon monoxide poisoning (42 men, 28 women; age  $52 \pm 18$  years) who had been admitted to St. Mary's Hospital, Kurume, Japan for treatment between June 2013 and September 2019. Clear electrocardiograms were available for each patient. Patients' carbon monoxide poisoning had been confirmed by arterial blood analysis. Carboxyhemoglobin measurements were performed in the ambulance and/or at admission. The highest values were adopted.

### QT interval measures and cardiac enzymes

All 12-lead electrocardiograms were obtained at a paper speed of 25mm/sec with standard lead positions. QT and RR intervals were measured on each electrocardiogram in all patients. The QT interval was measured from the beginning of the QRS complex to the end of the T wave. The QT intervals for each lead were measured and corrected for heart rate (QTc) using Bazett's formula ( $QT/\sqrt{RR}$ )<sup>6</sup>. The QTc dispersion was the difference between the leads with the shortest and longest QTc intervals<sup>4</sup>. QT intervals were measured upon admission to the emergency department. Additionally, blood samples were obtained and the troponin I level was determined.

### Statistical analysis

Retrospective statistical analyses were performed using JMP and SAS university edition software (SAS Institute Inc., Cary, NC, USA). Results are presented as means  $\pm$  SDs. Spearman correlation analysis and logistic analysis were used to examine the relationships between carboxyhemoglobin levels and clinical variables. A value of  $P < 0.05$  was considered to indicate statistical significance.

## Results

### Patients

Altogether, 70 patients were included in the study. Clinical characteristics of the patients are presented in Table 1. Two patients died after admission. In all, 12 patients were diagnosed with encephalopathy at or after discharge, whereas no patients exhibited cardiomyopathy at or after discharge.

## Correlation between carboxyhemoglobin and QT intervals

The correlation between maximum QT intervals and the carboxyhemoglobin levels was not significant. Conversely, the correlation between the maximum QTc interval and the carboxyhemoglobin level was significant ( $P=0.0072$ ,  $R^2=0.1017$ ; Fig. 1), as were the relationships between QT dispersion and carboxyhemoglobin ( $P<0.001$ ,  $R^2=0.2358$ ; Fig. 1) and the QTc dispersion and carboxyhemoglobin ( $P<0.001$ ,  $R^2=0.2613$ ; Fig. 1).

## Logistic analyses

Death, carbon monoxide encephalopathy, and cardiomyopathy at or after discharge were defined as sequential disabilities. However, none of the patients in this study had cardiomyopathy. The reference values for the maximum QTc interval, QTc dispersion, carboxyhemoglobin level, and troponin I level were established based on the Classification and Regression Tree. Univariate logistic analysis showed that the significant predictors of sequential disability were smoking ( $P=0.0361$ ), consciousness disorder on admission ( $P=0.0072$ ), maximum QTc interval time ( $>484$  ms;  $P=0.0009$ ), QTc dispersion ( $>46$  ms;  $P=0.0003$ ), high concentration of carboxyhemoglobin ( $>44\%$ ;  $P=0.0044$ ), and high troponin I level ( $>0.36$  ng/ml;  $P<0.001$ ; Table 2). The multivariate logistic analysis indicated that the significant predictors of sequential disability were hyperbaric oxygen therapy ( $P=0.0182$ ), QTc dispersion ( $P=0.0062$ ), and troponin I level ( $P=0.0002$ ; Table 2).

## Discussion

Carbon monoxide poisoning, a serious health problem, is associated with a high incidence of severe morbidity and mortality. It causes myocardial toxicity and life-threatening arrhythmias. Carbon monoxide reduces the oxygen-carrying capacity of blood and binds with cardiac myoglobin, causing a rapid decrease in myocardial oxygen reserves<sup>6</sup>. Several studies have shown that carbon monoxide intoxication causes increased QT intervals and QT dispersion<sup>6-8</sup>. This study showed that QTc dispersion and carboxyhemoglobin are significantly related. Furthermore, QTc dispersion and troponin I are predictors of sequential disability.

### Correlation between carboxyhemoglobin and QT intervals

The *QT interval* is an indicator of ventricular repolarization on the electrocardiogram. A *prolonged QT interval* reflects impaired myocardial refractoriness. *QT dispersion* reflects the physiological variability of regional ventricular repolarization. *Increased QT dispersion* is related to the heterogeneity of regional ventricular repolarization and is accepted as a marker for arrhythmia and sudden death<sup>9</sup>. In this study, the QTc interval and QTc dispersion correlated with the carboxyhemoglobin level. Hanci et al. also reported that the QTc interval and QTc dispersion show good correlations with carboxyhemoglobin<sup>10</sup>.

Increased QTc dispersion and intervals in carbon monoxide toxicity might be caused by carbon monoxide on the myocardium, which causes homogeneous impulse formation in the ventricles.

Relation between the prognosis of patients with carbon monoxide toxicity and cardiac markers

The univariate logistic analysis indicated that the predictors of sequential disability in patients with carbon monoxide toxicity were smoking, consciousness disorder, maximum QTc interval, QTc dispersion, carboxyhemoglobin, and troponin I. The multivariate analysis revealed that the significant predictors of sequential disability were hyperbaric oxygen therapy, QTc dispersion, and troponin I. Note that the multivariate analysis did not find that carboxyhemoglobin was related to a poor outcome. The prognosis of patients with carbon monoxide toxicity could be predicted by the duration of carbon monoxide exposure and its concentration. Conversely, carboxyhemoglobin was not necessarily related to the exposure time or the concentration of the carbon monoxide. Because patients with carbon monoxide toxicity usually receive treatment immediately after rescue, carboxyhemoglobin was not measured at its peak concentration.

Hampson et al. found that carboxyhemoglobin measurement was a poor predictor of clinical status in patients with carbon monoxide poisoning <sup>11</sup>. Moreover, mortality was associated with the absolute difference in carboxyhemoglobin <sup>11</sup>. Satran et al., however, reported that moderate-to-severe carbon monoxide poisoning causes myocardial injury when assessed by electrocardiography or biomarkers <sup>5</sup>. Henry et al. reported that patients with myocardial injury (cardiac troponin I > 0.7 ng/ml, the creatine kinase-MB level, and/or diagnostic ECG changes) had increased mortality <sup>12</sup>.

Carbon monoxide-mediated toxicity results from several factors. Carbon monoxide binds to hemoglobin with an affinity 200–250 times that of oxygen. Thus, exposure to carbon monoxide, even in low concentrations, results in competitive binding to hemoglobin, reduced oxygen delivery, and profound tissue hypoxia <sup>13</sup>. Carbon monoxide also binds to cytochrome-c oxidase, directly interfering with cellular respiration <sup>14</sup>. These mechanisms are believed to cause neurological injury and likely contribute to myocardial injury as well <sup>12</sup>. Although there were no patients with cardiomyopathy at/after discharge in this study, the indices of myocardial injury could lead to neurological injury or even mortality. The mechanisms described influence this result.

Limitations

The primary limitations of this study were the small sample population, single-center design, and retrospective nature. Despite these limitations, the study may act as a good basis for further study of the topic. The findings must be confirmed in prospective, multicenter studies with larger populations.

## Conclusion

The prognosis for patients with carbon monoxide poisoning may be predicted based on the QTc dispersion value and the troponin I level. Patients with myocardial damage should be monitored not only for their cardiovascular outcome but also for their neurological outcome and their prognosis.

# Abbreviations

ECG

electrocardiogram

QTc

corrected QT interval

# Declarations

## Availability of data and materials

The datasets used in the current study are available from the corresponding author on reasonable request.

## DISCLOSURES

The protocol for this research project was approved by a suitably constituted ethics committee of our institution (St. Mary's Hospital; Approval No. 19-0710). The study conforms to the provisions of the Declaration of Helsinki.

## Consent for publication

Not applicable

## Competing interests

There are no competing interests to declare in this study.

## ACKNOWLEDGMENT

We thank Nancy Schatken, BS, MT(ASCP), from Edanz Group (<https://en-author-services.edanzgroup.com/>), for editing a draft of this manuscript.

## Funding

none

## Author information

### Affiliations

Emergency Department, St. Mary's Hospital, 422 Tsubuku-honmachi, Kurume 830-8543, Japan

Hitoshi Koga, Kouta Mukasa, Tomohiro Inoue, Aya Okamoto, Shougo Urabe, Shuuichirou Sagara, Kazumi Yano, Kouhei Onitsuka, Hisashi Yamashita

Division of Cardiology and Emergency Department, St. Mary's Hospital, 422 Tsubuku-honmachi, Kurume 830-8543, Japan

Hideki Tashiro

### Contributions

HT, KY and HK conducted the study design. KY, HK, KM, TI, AO, SU, SS and KO collected the data. HT performed the statistical analysis and prepared this manuscript. HT, HK and HY finalized the manuscript. The authors read and approved the final manuscript.

### Corresponding author

Correspondence to Hideki Tashiro

## References

1. Roth D, Hubmann N, Habel C, Herkner H, Schreiber W, Laggner A. Victim of carbon monoxide poisoning identified by carbon monoxide oximetry. *J Emerg Med.* 2011;40(6):640–2.
2. Koyuncu S, Bol O, Ertan T, Gunay N, Akdogan HI. The detection of occult CO poisoning through noninvasive measurement of carboxyhemoglobin; A cross-sectional study. *Am J Emerg Med* 2019.158383 doi.org/10.1016/j.ajem. 2019.158383.
3. Yelken B, Tanriverdi B, Centinbas F, Memis D, Sut N. The assessment of QT intervals in acute carbon monoxide poisoning. *Anadolu Kardiyol Derg.* 2009;9:397–400.
4. Day CP, McComb JM, Campbell RWF. QT dispersion: an indication of arrhythmia risk in patients with long QT intervals. *Br Heart J.* 1990;63:342–4.

5. Satran D, Henry CR, Adkinson C, Nicholson CI, Bracha Y, Henry TD. Cardiovascular manifestations of moderate to severe carbon monoxide poisoning. *J Am Coll Cardiol*. 2005;45:1513–6.
6. Atescelik M, Bozdemir MN, Yildiz M, Gurbuz S, Ayranci M. QT dispersion in carbon monoxide poisoning. *Eur Rev Med Pharmacol Sci*. 2012;16(Suppl 1):25–9.
7. Erglu M, Omer U, Zafer I, Yalcin M, Yildirim AO, Kardesglu E. Carbon monoxide poisoning increase Tpeak-Tend dispersion and QTc dispersion. *Cardiovasc J Afr*. 2014;25:106–9.
8. Gürkan Y, Canatay H, Toprak A, Urai E, Toker K. Carbon monoxide poisoning- a cause of increased QT dispersion. *Acta Anaesthesiol Scand*. 2002;46(2):180–3.
9. Shimizu H, Ohnishi Y, Inoue T, Yokoyama M. QT/QTc dispersion in patients with monomorphic or polymorphic ventricular tachycardia/ventricular fibrillation. *J Electrocardiol*. 2001;34:119–25.
10. Hanci V, Ayoğu H, Yurtlu S, Yıldırım N, Okyay D, Erdoğan G. et. al. Effects of acute monoxide poisoning on the P-wave and QT interval dispersions. *Anadolu Kardiyol Derg*. 2011;1:48–52.
11. Hampson NB, Niels M, Hauff BE. Carboxyhemoglobin levels in carbon monoxide poisoning: do they correlate with the clinical picture? *Am J Emerg Med*. 2008;26(6):665–9.
12. Henry CR, Satran D, Lindgren B, Adkinson C, Nicholson CI, Henry TD. Myocardial injury and long-term mortality following moderate to severe carbon monoxide poisoning. *JAMA* 2006;295:398–402.
13. Ernst A, Zibrak JD. Carbon monoxide poisoning. *N Engl J Med*. 1998;339(22):1603–8.
14. Tibbles PM, Edelsberg JS. Hyperbaric-Oxygen therapy. *N Engl J Med*. 1996;334:1642–8.

## Tables


**Table 1.** Patients' characteristics

Male sex (%)	42 patients (60%)
Age (mean $\pm$ SD)	51.6 $\pm$ 18.2 years
Smoker	26 patients (37.1%)
Hypertension	10 patients (14.3%)
Diabetes mellitus	4 patients (5.7%)
Asymptomatic patients (%)	23 patients (33%)
Consciousness disorder on admission (%)	37 patients (53%)
Cardiopulmonary arrest (%)	3 patients (4%)
Hyperbaric oxygen therapy (%)	49 patients (70%)
Carboxyhemoglobin level (%) (normal range 0.5-1.5)	21.8 $\pm$ 14.8
Troponin I (ng/ml) (normal range 0.00-0.09)	0.85 $\pm$ 3.36
Mean emergency department arrival time (min)	165 $\pm$ 148

**Table 2.** Findings of the univariate and multivariate analyses

	Univariate analysis		Multivariate analysis	
	Odds ratio (95% CI)	p	Odds ratio (95% CI)	p
Sex (M)	1.875(0.528-7.498)	0.3291		
Age	0.568 (0.236-6.610)	0.6481		
Age (>30)	1.56 (0.236-30.803)	0.6903		
Smoke	0.222 (0.0326-0.915)	0.0361	1.270 (0.131 – 11.347)	0.8261
Consciousness disorder	5.667((1.565-27.183)	0.0072	0.760(0.081 – 6.484)	0.7598
Hyperbaric oxygen therapy	0.488 (0.145-1.696)	0.2516	0.0880 (0.007 – 0.674)	0.0182
Corrected maximum QT interval	10.2 (2.607- 44.108)	0.0009		
Corrected QT dispersion	10.227 (2.865-43.383)	0.0003	19.273(2.187 – 336.036)	0.0062
Carbon monoxide Hemoglobin (>44)	9.814 (2.063-52.237)	0.0044	3.781 (0.204 – 92.689)	0.3703
Troponin I (>0.36)	17.333 (4.25-83.958)	<0.001	66.792 (6.090 – 1850.641)	0.0002

## Figures



**Figure 1**

Top left. Relation between the maximum QT interval and the carboxyhemoglobin level. Top right. Relation between the maximum corrected QT interval and the carboxyhemoglobin level. Bottom left. Relation between QT dispersion and the carboxyhemoglobin level. Bottom right. Relation between corrected QT dispersion and the carboxyhemoglobin level. The relationships between the maximum QT interval and carboxyhemoglobin is not significant, whereas the other relationships are all significant.