

Development And Validation Of Models To Predict Cesarean Delivery Among Low-Risk Nulliparous Women At Term: A Retrospective Study In China

Fangcan Sun

First Affiliated Hospital of Soochow University

Bing Han (✉ hanbing@suda.edu.cn)

First Affiliated Hospital of Soochow University <https://orcid.org/0000-0001-7496-6850>

Fangfang Wu

Suzhou Ninth people's hospital

Qianqian Shen

Sihong county people's hospital

Minhong Shen

First Affiliated Hospital of Soochow University

Youguo Chen

First Affiliated Hospital of Soochow University

Research article

Keywords: Cesarean delivery after failure of trial of labor, cesarean, induction of labor, prediction, nulliparous, nomogram

Posted Date: July 27th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-44296/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Cesarean delivery after failure of trial of labor is associated with adverse maternal and perinatal outcomes. A prediction algorithm to identify women with high risk of an emergency cesarean could help reduce morbidity and mortality associated with labor. The objective of the present study was to derive and validate a simple model to predict cesarean delivery for low-risk nulliparous women in Chinese population.

Methods: This retrospective study analyzed the low-risk nulliparous women with singleton cephalic full-term fetus delivered in two medical centers. After the clinical data of the women who delivered at the tertiary referral center (n=6 551) was collected and was used univariate and multivariable logistic regression analysis, the prediction model was fitted. We performed external validation using data from nulliparous who delivered from another hospital(secondary referral center, n=7 657). A new nomogram was established based on the development cohort to predict the cesarean. The ROC curve, calibration plot and decision curve analysis were used to assess the predictive performance.

Results: The cesarean delivery rates in the development cohort and the external validation cohort were 8.79% (576/6 551) and 7.82% (599/7 657). Multivariable logistic regression analysis showed that maternal age, height, BMI, weight gained during pregnancy, gestational age, induction method, meconium-stained amniotic fluid and neonatal sex were independent factors affecting cesarean outcome. Because sex of the fetuses were unknown until they born(China's Fertility Policy), we established two prediction models according to fetal sex was involved or not. The AUC was 0.782 and 0.774, respectively. The Hosmer-Lemeshow goodness-of-fit test showed that these two models fitted well. Decision curve analysis demonstrated that the models were clinically useful. And internal validation using Bootstrap method showed that these prediction models perform well. On the external validation set, the AUC were 0.775 and 0.775, respectively. The calibration plots for the probability of cesarean showed a good correlation. The online web server was constructed based on the nomogram for convenient clinical use.

Conclusions: Both two models established by these factors have good prediction efficiency and high accuracy, which can provide the reference for clinicians to guide pregnant women to choose an appropriate delivery mode.

Background

Cesarean is a life-saving surgical procedure within the obstetric domain. However, it may have risks for subsequent pregnancies as well as long-term effects that are still undertaking research[1–3]. With the development of urbanization, the relaxation of the one-child policy and the introduction of the universal two-child policy in China, a growing number of pregnant women are using the Internet to obtain for medical information about pregnancy and childbirth. It lets them realize the disadvantages and risks of cesarean section, they are more willing to deliver vaginally[4, 5]. Sometimes spontaneous vaginal delivery

doesn't go so well, and pregnancies have to undergo the emergency operation. And yet, maternal and perinatal complications are higher when a failed trial of labor results in an emergency cesarean delivery. Arduous birth experience could impose life-long negative effects and have an unpleasant procedure on the life of the mother, child, family, clinic, and society[6, 7]. In the era of foreseeable medicine, a prediction algorithm to determine women at risk of an emergency cesarean could help reduced labor associated morbidity and mortality.

Currently, several published models studied on prediction of cesarean delivery were mostly focused on the white populations[8–12], whereas the data of the majority of Han ethnic backgrounds of the Chinese population are insufficient. To date, China lacks the relevant researches, and there is limited empirical evidence and clinical experience reported.

Therefore, the objective of our study was to derive and validate a clinical prediction model of cesarean delivery for nulliparous women in low-risk at term. The use of calculators may give physicians an evidence-based tool to assist with patient counseling and to provide individualized guidance of delivery mode.

Methods

Participants

This retrospective study was conducted in the Department of Obstetrics and Gynecology of The First Affiliated Hospital of Soochow University (tertiary referral center, hospital 1) and (secondary referral center, hospital 2). For the development cohort, we used data from hospital 1 between January 1, 2011 and August 31, 2017. External validation sampled from hospital 2 between January 1, 2013 to December 31, 2019. Institutional Review Board approval by these two hospitals was obtained for the study waiving informed consent for this retrospective study. Methods and reporting guidelines were followed by the TRIPOD (Transparent Reporting of a multivariable prediction model for Individual Prognosis Or Diagnosis) statement[13] (Additional files 1).

Inclusion and exclusion criteria

Low-risk pregnancy nulliparous women undergoing the labor with singleton, term(37 0/7 weeks of gestation or greater) and cephalic pregnancies were recruited. Exclusion criteria were as follows:

- (1) Women had complications during pregnancy(e.g., cardiac failure, severe liver and kidney diseases, hypertensive disorders of pregnancy, diabetes, oligohydramnios, placenta previa, vasa previa and fetal growth restriction);
- (2) Women had a scarred uterus(e.g., myomectomy);
- (3) Women had contraindications to vaginal delivery;

(4) Women with the cesarean delivery on maternal request.

The pregnant women who meet the criteria were divided into the emergency cesarean delivery group and the vaginal delivery group according to their delivery modes.

Data collection

Data on maternal characteristics and perinatal parameters were collected from the institution's obstetrics database, which was obtained by the patient's medical record review.

Characteristics

The outcome of interest was defined as cesarean delivery. A cesarean delivery was performed if there was fetal distress, arrested active phase, prolonged latent phase, prolonged second stage, arrested descent, fever indicating chorioamnionitis, and other medical indications, such as threatened uterine rupture. It is worth noting that we only collected the major indications. The candidate predictor variables had to be easily accessible through characteristics' data. To identify predictor variables, a systematic review of the literature was conducted[8-12, 14-16]. The following variables were recorded: maternal age, height, weight, baseline body mass index (BMI), weight change during pregnancy, gestational age at delivery, premature rupture of membranes (PROM), epidural analgesia, meconium-stained amniotic fluid, induction methods(oxytocin, amniotomy, disposable cervical dilator balloon, prostaglandin(Propess or Misoprostol)), neonatal sex, and neonatal birth weight.

Operational definitions

The relevant guidelines[17, 18] were used to determine cesarean delivery indications such as arrest of descent and a prolonged second stage of labor. BMI was calculated as $\text{weight (kg)}/[\text{height (m)}]^2$. Baseline BMI was defined as pre-pregnancy BMI. Gestational age was calculated by the date of the last menstrual period and confirmed by ultrasound examination during first-trimester (by measuring the crown-rump length) or second-trimester(by measuring biparietal diameter, abdominal circumference and femur length). We used the following modes for grouping for induction methods.

(1) Oxytocin Induction group: initial cervical dilation < 6 cm with only oxytocin induction. This was defined as Induction method 1;

(2) Amniotomy group: initial cervical dilation < 6 cm with artificial rupture of membranes with or without oxytocin induction(amniotomy after Prostaglandin E2 and Disposable cervical dilator balloon were not included). This was defined as Induction method 2;

(3) Disposable Cervical Dilator Balloon: with Disposable cervical dilator balloon induction. This was defined as Induction method 3;

(4) Prostaglandin E2 group: with Prostaglandin E2(Propess) induction. This was defined as Induction method 4;

(5) Cervical Dilation \geq 6cm group: initial cervical dilation \geq 6cm experience augmentation of labor(experience amniotomy and/or oxytocin);

(6) Unreceived Labor Intervention group: women entered labor naturally and the labor did not be intervened.

Induction group was defined as group (1) plus (2) plus (3) plus (4), augmentation group was defined as group (5). Considering that the women in the augmentation group naturally entered the active stage of labor, the augmentation group and the unreceived labor intervention group were regarded as a group, which serves as a reference group for the induction method (1)-(4). For convenience, spontaneous labor group was defined as group (5) plus group (6).

Statistical analysis

Data analysis was conducted by using the statistical software package SPSS (24.0) and R (3.6.2).

Univariate analysis was performed for all clinical data. For continuous variables, the tests of normality were performed first. The Student's t-test was used to compare the continuous variables with a normal distribution. The Mann-Whitney test was used to compare discrete or continuous variables without a normal distribution. The chi-square test and Fisher exact test were used, as appropriate, for the categorical variables. Standard descriptive statistics (mean \pm standard deviations or median and interquartile range) were used to summarize continuous variables. Percentage and frequency were used for categorical variables.

Baseline variables that were considered clinically relevant or that showed a univariate relationship with outcome(candidate variables with a p value <0.05 on univariate analysis) were entered into the multivariable logistic regression model. Variables for inclusion were carefully chosen, given the number of events available, to ensure parsimony of the final model. The results of logistic regression models were presented as odds ratio (OR) with their 95% confidence intervals (CIs).

The discrimination and calibration of the prediction model were evaluated. Discrimination is the extent to which patients with cesarean delivery is identified likely to have this positive outcome. Calibration refers to the extent to which the calculated risks reflect the actual percentage of women with the outcome in each group. It is the agreement between observed outcomes and predictions. The area under the receiver operating characteristic curve(AUC ROC) was calculated to assess the discrimination ability. AUC ROC was interpreted using following categories: non-informative (AUC ROC = 0.5), poor accuracy (0.5 < AUC ROC <0.7), moderate accuracy (0.7 < AUC ROC <0.9), high accuracy (0.9 < AUC ROC < 1); and perfect accuracy (AUC ROC = 1)[19]. The calibration of the prediction model was assessed using the Hosmer-Lemeshow goodness-of-fit test($P>0.05$ was taken to indicate good fitting) and/or calibration plot. Decision curve analysis (DCA) was used to assess the clinical value of the model, which was a method

for evaluating the net benefit. We assessed internal validity with a bootstrapping technique (resample 1000 times) to show the performance of model. We conducted an external verification of the final model in another hospital. We reported the predictive performance in the validation cohort also using the measures of discrimination and calibration. A nomogram was established using the predictive model by R. Established a dynamic nomogram by using the rms DynNom, and built a web online applications through shinyapps. All P values were two-tailed, and a significance level of 5% was used.

Results

Study population and outcomes

Between 1 January 2011 and 31 August 2017, 18,228 deliveries were managed at our center. After exclusion of ineligible patients, 6,551 women were enrolled (Additional files 2). Among the recruited subjects, 576(8.79%) women gave birth by cesarean after failure of trial of labor. Most cesarean deliveries were performed for concerns with fetal heart rate patterns in labor (48.44% = 279/576) and arrest of dilation in the active phase of the first stage of labor (20.83% = 120/576).

Characteristics of pregnancies

Participants were divided into vaginal delivery and cesarean delivery groups, and the comparisons of demographics and clinical characteristics are displayed in Table 1.

Table 1
 Characteristics of maternal and neonatal by mode of delivery

Variable	Cesarean Delivery (n = 576)	Vaginal Delivery (n = 5975)	P*
Maternal characteristics			
Maternal age (y)	27.50 ± 3.12	26.84 ± 3.03	< 0.001
Baseline height (cm)	159.54 ± 4.24	161.7 ± 4.61	< 0.001
Baseline BMI (kg/m ²)	21.49 ± 2.60	20.75 ± 2.49	< 0.001
Weight change during pregnancy (kg)	14.77 ± 4.21	14.14 ± 4.22	0.001
Gestational age (w)	40.23 ± 1.00	39.76 ± 1.00	< 0.001
Labour and others characteristics			
PROM	152 (26.39%)	1238(20.72%)	0.001
Induction methods			< 0.001
Spontaneous Labor group	114(19.79%)	2596(43.45%)	
Oxytocin Induction group	152(26.39%)	976(16.33%)	
Amniotomy group	268(46.53%)	2189(36.64%)	
Disposable Cervical Dilator Balloon group	15(2.60%)	65(1.09%)	
Prostaglandin E2 group	27(4.69%)	149(2.49%)	
Epidural analgesia	23(3.99%)	161(2.69%)	0.072
Meconium-stained amniotic fluid			< 0.001
0	434(75.35%)	5511(92.23%)	
1	30(5.21%)	209(3.50%)	
2	33(5.73%)	150(2.51%)	

BMI, body mass index; PROM, premature rupture of membranes;

Data are mean ± standard deviation, n/(N %)

* Two-sided P based on the χ^2 for categorical variables, and the t test for continuous variables.

Variable	Cesarean Delivery (n = 576)	Vaginal Delivery (n = 5975)	P*
⊠/ bloody	79(13.72%)	105(1.76%)	
Neonatal characteristics			
Neonatal sex(male)	334(57.99%)	2934(49.10%)	< 0.001
Neonatal birth weight(g)	3528.18 ± 389.13	3343.93 ± 367.50	< 0.001
BMI, body mass index; PROM, premature rupture of membranes;			
Data are mean ± standard deviation, n/(N %)			
* Two-sided P based on the χ^2 for categorical variables, and the t test for continuous variables.			

It can be found that maternal age, BMI, weight gain during pregnancy, gestational age at delivery, level of meconium-stained amniotic fluid, rate of PROM and male infants who underwent cesarean were higher than the vaginal delivery group. Also, their height was significantly lower than those in the vaginal group. The composition ratios of methods of induction were statistically significant between the two groups. There were no significant differences in the rate of delivering with labor epidural analgesia between the two groups. Women who aged more than 26.5 years, height less than 160.5 cm, gestational age more than 279 days, had pre-pregnancy BMI more than 21.4 kg/m² and gained pregnancy weight more than 13.3 kg were more likely to undergo cesarean.

Model development

Factors that were statistically significantly associated with cesarean delivery in univariate analysis or that were considered clinically relevant were entered into the multivariable logistic regression. The results of the analysis showed in Figure1.

Derivation of a model

The results of multivariable logistic regression analysis showed that in the model, maternal age, height, gestational age, pre-pregnancy BMI, Weight change during pregnancy, PROM, methods of induction, level of meconium-stained amniotic fluid and neonatal sex were predictors of cesarean delivery. Developed with data from 6,551 women with complete data, the final equation 1 was:

$$\text{logit}(P_1) = -7.351 + 0.075 * \text{Maternal age (y)} - 0.113 * \text{height(cm)} + 0.072 * \text{pre-pregnancy BMI (kg/m}^2\text{)} + 0.053 * \text{Weight change during pregnancy (kg)} + 0.062 * \text{Gestational age (day)} + 0.711 * (\text{indicator for occurrence of PROM}) + 0.929 * (\text{indicator for occurrence of Induction method 1}) + 0.938 * (\text{indicator for occurrence of Induction method 2}) + 1.459 * (\text{indicator for occurrence of Induction method 3}) + 0.902 * (\text{indicator for occurrence of Induction method 4}) + 0.700 * (\text{level of Meconium-stained amniotic fluid}) + 0.482 * (\text{indicator for occurrence of male fetus}) \quad (\text{Model 1})$$

At the same time, due to China's relevant policies that the sex of the fetus was unknown until they had born, we also established a prediction model that did not include fetal sex. The equation 2 was:

$\text{logit}(P_2) = -6.472 + 0.073 * \text{Maternal age (y)} - 0.112 * \text{height(cm)} + 0.072 * \text{pre-pregnancy BMI (kg/m}^2\text{)} + 0.050 * \text{Weight change during pregnancy (kg)} + 0.060 * \text{Gestational age (day)} + 0.710 * (\text{indicator for occurrence of PROM}) + 0.942 * (\text{indicator for occurrence of Induction method 1}) + 0.956 * (\text{indicator for occurrence of Induction method 2}) + 1.444 * (\text{indicator for occurrence of Induction method 3}) + 0.938 * (\text{indicator for occurrence of Induction method 4}) + 0.689 * (\text{level of Meconium-stained amniotic fluid})$ (Model 2)

Discrimination, calibration and decision curve analysis

The areas under the receiver operating characteristic curve, calibration plot and decision curve analysis were presented in Figure 2. For model 1, the AUC curves were 0.782(95%CI: 0.771-0.791). According to the ROC curve, the ideal cutoff point for predicted probability was obtained, that was when the predicted probability was greater than 7.45%, the AUC was the largest. At this time, there were 452 cases of cesarean delivery, the incidence rate was 17.6%. The sensitivity of the prediction model 1 was 78.47% and the specificity was 64.55%. The calibration plot for the probability of cesarean delivery showed a good correlation between the predicted and actual probabilities(Hosmer-Lemeshow test: P=0.263, Figure 2B). The AUC curves for model 2 were 0.774(95%CI: 0.763-0.784). Using a cut-off of predicted probability of 8.7% to define a positive test classified women with good accuracy (sensitivity 70.66%; specificity 70.59%).The Hosmer-Lemeshow test for goodness of fit showed a good fit of the model (P=0.817). The calibration plot of the Model 2 showed that the observed outcomes were in good agreement with the predicted outcomes(Figure 2C).

DCA is a novel method that can examine diagnostic and prognostic strategies, can be used to evaluate and compare different predictive models, and identify the net benefit of a prediction model[20]. Thus, we used DCA for models to predict the correct diagnosis of cesarean delivery. The results (Figure 2:D-F) indicated that two models were useful for threshold probabilities of 4% to 60%. There was no significant difference in the net benefit between them. The clinical impact curves also demonstrated the clinical utility of the models.

Internal and External validation and predictive performance

Results of the internal validation using Bootstrap method indicated that the cesarean delivery rates predicted by both models were consistent with the real data(Figure 3 A-B).

We validated the two formulas by using a separate data set which was derived from another hospital(hospital 2). External validation of the logistic regression equations using a cohort that included 7 657 low-risk nulliparous women with the same inclusion criteria as the original data set used to create the model(Additional files 3). It is worth noting that cervical dilator balloon and Propess were not routinely used as induction methods in the secondary-grade A class hospital in this study. In this hospital, 25 micrograms of misoprostol was used prostaglandin for induction of cervical ripening. Considering that misoprostol is also a prostaglandin drug induction method, therefore, when we verified externally, it replaced Propess in induction method 4. That is, if misoprostol was used to induce labor, the value of induction method 4 in the prediction formula was 1. When these two formulas were applied to this cohort, it achieved AUC of 0.775 (95% CI 0.755–0.796) and 0.775 (95% CI 0.754–0.796)(Figure 3 C-D). The calibration plots presented an acceptable agreement in the validation cohort between the prediction and actual observation for the Model1 and Model 2 (Figure 3 E-F).

Nomogram

The formula used a nomogram to calculate the probability of cesarean (Figure 4). For a given woman, each characteristic was aligned with the corresponding number of points on the points axis, and a total summated point was derived. The sum of all points lined with predicted probability of cesarean delivery. We also developed a friendly software-based calculator that can give the percentage likelihood of cesarean delivery. It can be found at: <https://fangcan.shinyapps.io/CSsexDynNomapp/> , <https://fangcan.shinyapps.io/CSDynNomapp/>. We used two examples to illustrate the dynamic nomogram on web (Additional files 4).

Discussion

Cesarean delivery after failure of trial of labor has been concerned by many researchers. This study was aiming to identify the predictors for cesarean experience among low-risk women and develop a clinical prediction model. The model allows physicians to assess the individual and systematic risks of pregnant women before childbirth. It can be used during counseling to increase acceptance of vaginal delivery for those with a high chance of success while minimizing procedures performed on women with low chances for a successful vaginal delivery. Screening for high-risk women and applying adequate interventions may reduce the risk of adverse outcomes. To achieve the best outcomes for mothers and babies, medical staff need to provide the pregnant women with the proper suggestion of delivery.

The models highlight the importance of variables such as maternal age, maternal height, pre-pregnancy BMI, gestational age, weight gain during pregnancy, degree of meconium-stained amniotic fluid, indicator for occurrence of premature rupture of membranes and fetal sex in determining the cesarean delivery when women failed the trial of labor. These results are consistent with the existing literature on the risk of cesarean delivery [16, 21–23]. Past studies also found gestational age of induced labor and induction methods were related to the incidence of cesarean [24, 25]. The ethnic disparity was observed in cesarean delivery rates, Stark et al. showed that the frequency of cesarean was lowest in non-Hispanic white women and highest in non-Hispanic black women [26]. In addition, delivery time, clinicians' personal beliefs and pregnant women themselves' views of decision-making also have an impact on delivery mode [27–29].

Currently, given the maternal and fetal implications of failed trial of labor, a number of prediction models aimed at determining the likelihood of cesarean delivery had been developed. A study by Levine et al. [9] used a nomogram to develop and validate a predictive model for women undergoing an induction of labor with an unfavorable cervix. Nulliparity, BMI, gestation age ≥ 40 weeks, modified Bishop score, and height were significantly associated with cesarean. This model with an AUC of 0.79 in the development cohort and 0.73 in the validation cohort. Jochum et al. [10] had developed a scoring system for predicting cesarean delivery after labor induction with cervical ripening based on a secondary data analysis. Height, BMI, gestational age, parity, dilation, effacement, fetal head station, medical indication, suspicion of macrosomia, PROM and concerning fetal status were found to be strongly associated with cesarean delivery. The AUC ROC in the derivation set and internal validation set were 0.76 and 0.74, respectively. Rossi et al. [12] developed and validated a predictive risk calculator for cesarean among women undergoing induction of labor. Seven independent risk factors had been associated with an increased risk

of cesarean included prior vaginal delivery, maternal weight at delivery, height, age, prior cesarean delivery, gestational age at induction, and maternal race. The model had an area under the curve for the receiver operating characteristic curve of 0.787 (95% CI 0.786–0.788), and it performed well on external validation (0.783, 95% CI 0.764–0.802).

However, it is uncertain whether these models are applicable to Chinese women. At the same time, China is a yellow race and people are relatively thin and have a low BMI. To date, China lacks the studies in relevant researches. Therefore, based on the analysis of clinically relevant factors of existing prediction models and the Fertility Policy of China (fetal sex identification is prohibited), this study established risk prediction models suitable for low-risk pregnant women for cesarean delivery through the multivariable Logistic regression analysis. These prediction models discriminated well (Model 1: AUC ROC was 0.782(95%CI: 0.771–0.791); Model 2: AUC ROC was 0.774(95%CI: 0.763–0.784)). Moreover, an external validation of women from another secondary medical center was conducted, which demonstrated a consistent measure of discrimination with the AUC of 0.775 (95% CI 0.755–0.796) and 0.775 (95% CI 0.754–0.796), respectively. The results of external validation showed that our prediction model can be further extended to the data set of another research center, indicating that the application value of the model has been greatly expanded. The online web server was constructed based on the nomogram to facilitate clinical practice. Clinicians can carry out a risk assessment and provide appropriate suggestion on patients at any time through mobile phones.

To our knowledge, the model developed in this study using information on maternal factors and using robust modeling methods is the first model applicable to Chinese low-risk women. These models still show good prediction ability in both internal and external validation populations. Further investigation of model validity and impact before generalizing is important and should be undertaken.

The limitation of this study may include the following aspects. Firstly, this is a retrospective study in nature. Some data is inevitably missing and inherently biased. For example, the success of induced labor mainly depends on the cervical ripening, and the Bishop's score has most often been used to describe cervical ripeness. This was not well documented in retrospective cases, so we did not include this factor in the final prediction model. The literature showed that the Bishop's score was originally designed to predict the likelihood of multiparous women at term to enter spontaneous labor, this could result in making it less predictive of outcome after labor induction in nulliparas. A systematic review also concluded that the Bishop score was a poor predictor for the outcome of induced labor at term[30]. In comparison with other prediction models, there are variations in induction rates and variables most likely to influence successful vaginal delivery. The sample size of the study was relatively small, and the included subjects were only the single-centered population in the region. The selection of the population also targeted only low-risk nulliparous. Given the diversity of geography, economy, medical level and environment throughout China, our findings may not representative the Chinese population in many jurisdictions. The neonatal birth weight had been associated with an increased risk of cesarean, but it was not included in the final prediction model because it could not be accurately known before delivery. Nowadays, the ultrasonography is the main method to estimate fetal weight, but it is not accurate

enough. Ethnicity, the time interval between ultrasonic examination and delivery, fetal sex, fetal position, maternal BMI and seniority of sonographers are influencing factors on sonographic estimated fetal weight. Compared with the Jochum[10] model, which was established in the scoring system, the prediction model was developed by Logistic regression analysis in this study had a more complex process. But it was accurate in calculation and had high sensitivity and specificity. The nomogram was established, which can be convenient for clinicians in the clinic. At the same time, we developed a software-based calculator that gives the percentage likelihood of cesarean delivery.

In conclusion, multivariable analysis showed that maternal age, height, BMI, weight gain during pregnancy, gestational age, mode of labor induction, meconium-stained amniotic fluid, presence of PROM and neonatal sex were an independent risk factor for cesarean delivery in this study. With the adjustment in China's family planning strategy and release of the second-child policy, more pregnant women are faced with the choice of delivery mode. The prediction model established by these factors has a better prediction performance. Obstetricians and midwives can use the tool to predict which women will need surgery. Those who as high-risk women could be offered elective cesarean, which could avoid additional adverse effects. Those at low risk should be comforted and encouraged to adopt vaginal delivery. However, the benefits of using models should be demonstrated before routine introduction into clinical practice. Further study is warranted to optimize these models by conducting multicenter researches studies with large samples. It is worth noting that these models should be combined with the clinical practice of patients rather than applied in isolation. It provides evidence-based knowledge to support their delivery mode choices and to improve maternal and perinatal outcomes and optimize the allocation of resources.

Declarations

Ethics approval and consent to participate

Institutional Review Board approval by The First Affiliated Hospital of Soochow University and Sihong county People's Hospital were obtained for the study waiving informed consent for this retrospective study. A total of 14 208 study participants were recruited in two ways. First, there was retrospective recruitment of consecutive 6551 pregnant women in the final analysis delivered from January 1, 2011 to August 31, 2017 (2018019). Second, 7 657 women came from Sihong county People's Hospital between January 1, 2013 and December 31, 2018 were recruited retrospectively into a validation cohort.

Consent for publication

Not applicable.

Availability of data and materials

The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests

Funding

No funding

Authors' contributions

HB had the original idea of the study. SFC, SQQ, WFF, SMH and CYG contributed to data collection. SFC analyzed the data. SFC wrote the first draft of the paper. HB and SFC provided critical review and interpretation of the results. HB provided professional language. All authors have read the manuscript and approved to publish.

Acknowledgements

Not applicable

References

1. Betrán AP, Ye J, Moller AB, Zhang J, Gülmezoglu AM, Torloni MR: **The Increasing Trend in Caesarean Section Rates: Global, Regional and National Estimates: 1990-2014.** *PloS one* 2016, **11**(2):e0148343.
2. Chu S, Zhang Y, Jiang Y, Sun W, Zhu Q, Wang B, Jiang F, Zhang J: **Cesarean section without medical indication and risks of childhood allergic disorder, attenuated by breastfeeding.** *Sci Rep* 2017, **7**(1):9762.
3. Zhang T, Sidorchuk A, Sevilla-Cermeno L, Vilaplana-Perez A, Chang Z, Larsson H, Mataix-Cols D, Fernandez de la Cruz L: **Association of Cesarean Delivery With Risk of Neurodevelopmental and Psychiatric Disorders in the Offspring: A Systematic Review and Meta-analysis.** *JAMA network open* 2019, **2**(8):e1910236.
4. Liang J, Mu Y, Li X, Tang W, Wang Y, Liu Z, Huang X, Scherpbier RW, Guo S, Li M *et al*: **Relaxation of the one child policy and trends in caesarean section rates and birth outcomes in China between 2012 and 2016: observational study of nearly seven million health facility births.** *BMJ (Clinical research ed)* 2018, **360**:k817.
5. Li HT, Luo S, Trasande L, Hellerstein S, Kang C, Li JX, Zhang Y, Liu JM, Blustein J: **Geographic Variations and Temporal Trends in Cesarean Delivery Rates in China, 2008-2014.** *Jama* 2017, **317**(1):69-76.
6. Hosseini Tabaghdehi M, Kollahdozan S, Keramat A, Shahhossein Z, Moosazadeh M, Motaghi Z: **Prevalence and factors affecting the negative childbirth experiences: a systematic review.** *The journal of maternal-fetal & neonatal medicine : the official journal of the European Association of Perinatal Medicine, the Federation of Asia and Oceania Perinatal Societies, the International Society of Perinatal Obstet* 2019:1-8.

7. Størksen HT, Garthus-Niegel S, Adams SS, Vangen S, Eberhard-Gran M: **Fear of childbirth and elective caesarean section: a population-based study.** *Bmc Pregnancy & Childbirth* 2015, **15**(1):221.
8. Tolcher MC, Holbert MR, Weaver AL, McGree ME, Olson JE, El-Nashar SA, Famuyide AO, Brost BC: **Predicting Cesarean Delivery After Induction of Labor Among Nulliparous Women at Term.** *Obstetrics and gynecology* 2015, **126**(5):1059-1068.
9. Levine LD, Downes KL, Parry S, Elovitz MA, Sammel MD, Srinivas SK: **A validated calculator to estimate risk of cesarean after an induction of labor with an unfavorable cervix.** *American journal of obstetrics and gynecology* 2018, **218**(2):254.e251-254.e257.
10. Jochum F, Le Ray C, Blanc-Petitjean P, Langer B, Meyer N, Severac F, Sananes N: **Externally Validated Score to Predict Cesarean Delivery After Labor Induction With Cervi Ripening.** *Obstetrics and gynecology* 2019, **134**(3):502-510.
11. Alavifard S, Meier K, Shulman Y, Tomlinson G, D'Souza R: **Derivation and validation of a model predicting the likelihood of vaginal birth following labour induction.** *BMC pregnancy and childbirth* 2019, **19**(1):130.
12. Rossi RM, Requarth E, Warshak CR, Dufendach KR, Hall ES, DeFranco EA: **Risk Calculator to Predict Cesarean Delivery Among Women Undergoing Induction of Labor.** *Obstetrics and gynecology* 2020, **135**(3):559-568.
13. Moons KG, Altman DG, Reitsma JB, Ioannidis JP, Macaskill P, Steyerberg EW, Vickers AJ, Ransohoff DF, Collins GS: **Transparent Reporting of a multivariable prediction model for Individual Prognosis or Diagnosis (TRIPOD): explanation and elaboration.** *Annals of internal medicine* 2015, **162**(1):W1-73.
14. Burke N, Burke G, Breathnach F, McAuliffe F, Morrison JJ, Turner M, Dornan S, Higgins JR, Cotter A, Geary M *et al*: **Prediction of cesarean delivery in the term nulliparous woman: results from the prospective, multicenter Genesis study.** *American journal of obstetrics and gynecology* 2017, **216**(6):598.e591-598.e511.
15. Levine LD, Downes KL, Parry S, Elovitz MA, Srinivas SK: **468: Predicting cesarean delivery for women undergoing an induction of labor with an unfavorable cervix.** *American Journal of Obstetrics & Gynecology* 2017, **216**(1):S277.
16. Kawakita T, Reddy UM, Huang CC, Auguste TC, Bauer D, Overcash RT: **Predicting Vaginal Delivery in Nulliparous Women Undergoing Induction of Labor at Term.** *Am J Perinatol* 2018, **35**(7):660-668.
17. **ACOG Practice Bulletin Number 49, December 2003: Dystocia and augmentation of labor.** *Obstetrics and gynecology* 2003, **102**(6):1445-1454.
18. Caughey AB, Cahill AG, Guise JM, Rouse DJ: **Safe prevention of the primary cesarean delivery.** *American journal of obstetrics and gynecology* 2014, **210**(3):179-193.
19. Hanley JA, McNeil BJ: **The meaning and use of the area under a receiver operating characteristic (ROC) curve.** *Radiology* 1982, **143**(1):29-36.
20. Vickers AJ, Elkin EB: **Decision curve analysis: a novel method for evaluating prediction models.** *Medical decision making : an international journal of the Society for Medical Decision Making* 2006, **26**(6):565-574.

21. Bergholt T, Egil Skjeldestad F, Pyykonen A, Rasmussen SC, Tapper AM, Bjarnadottir RI, Smarason A, Masdottir BB, Klungsoyr K, Albrechtsen S *et al*: **Maternal age and risk of cesarean section in women with induced labor at term - a Nordic register-based study**. *Acta obstetrica et gynecologica Scandinavica* 2019.
22. Hikita N, Haruna M, Matsuzaki M, Sasagawa E, Murata M, Yura A, Oidovsuren O: **Is High Maternal Body Mass Index Associated with Cesarean Section Delivery in Mongolia? A Prospective Observational Study**. *Asian/Pacific Island nursing journal* 2019, **4**(3):128-134.
23. Lipschuetz M, Cohen SM, Israel A, Baron J, Porat S, Valsky DV, Yagel O, Amsalem H, Kabiri D, Gilboa Y *et al*: **Sonographic large fetal head circumference and risk of cesarean delivery**. *American journal of obstetrics and gynecology* 2018, **218**(3):339.e331-339.e337.
24. Grobman WA, Rice MM, Reddy UM, Tita ATN, Silver RM, Mallett G, Hill K, Thom EA, El-Sayed YY, Perez-Delboy A *et al*: **Labor Induction versus Expectant Management in Low-Risk Nulliparous Women**. *The New England journal of medicine* 2018, **379**(6):513-523.
25. Nwabuobi C, Gowda N, Schmitz J, Wood N, Pargas A, Bagiardi L, Odibo L, Camisasca-Lopina H, Kuznicki M, Sinkey R *et al*: **Risk factors for Cesarean delivery in pregnancies with a small-for-gestational-age fetus undergoing induction of labor**. *Ultrasound in obstetrics & gynecology : the official journal of the International Society of Ultrasound in Obstetrics and Gynecology* 2019.
26. Stark EL, Grobman WA, Miller ES: **The Association between Maternal Race and Ethnicity and Risk Factors for Primary Cesarean Delivery in Nulliparous Women**. *Am J Perinatol* 2019.
27. Del Carmen GA, Stapleton S, Qadan M, Del Carmen MG, Chang D: **Does the Day of the Week Predict a Cesarean Section? A Statewide Analysis**. *The Journal of surgical research* 2019, **245**:288-294.
28. Panda S, Begley C, Daly D: **Clinicians' views of factors influencing decision-making for caesarean section: A systematic review and metanalysis of qualitative, quantitative and mixed methods studies**. *PloS one* 2018, **13**(7):e0200941.
29. Loke AY, Davies L, Mak YW: **Is it the decision of women to choose a cesarean section as the mode of birth? A review of literature on the views of stakeholders**. *BMC pregnancy and childbirth* 2019, **19**(1):286.
30. Kolkman DG, Verhoeven CJ, Brinkhorst SJ, van der Post JA, Pajkrt E, Opmeer BC, Mol BW: **The Bishop score as a predictor of labor induction success: a systematic review**. *Am J Perinatol* 2013, **30**(8):625-630.

Figures




Figure 1

Univariate and multivariable logistic regression analysis of risk factors for Cesarean Delivery




Figure 2

The discrimination, calibration and decision curves with prediction models A: ROC curves of two models. B-C: Calibration plot for the the logistic regression prediction model 1 and model 2. D: DCA for two

models. E-F: Use Model 1 and Model 2 to predict risk stratification for 1 000 people. The red curve or green curve indicated the number of people classified as positive by the Model 1 or Model 2 at each threshold probability, the blue curve was the number of true.




Figure 3

Internal and external validation and predictive performance A-B: Calibration curves of the two models using 1000 bootstrap re-samples. C-D: The red ROC curve represents data from the Model 1 and green ROC curve represents data from the Model 2. E-F: Calibration curves of the two models in the external validation set.




Figure 4

The nomograms of the two models

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfiles1.docx](#)
- [Additionalfiles4.tif](#)
- [Additionalfiles3.docx](#)
- [Additionalfiles2.tif](#)