

# Monitoring priority substances in biota under the Water Framework Directive: How effective is a tiered approach based on caged invertebrates? A proof of concept study targeting PFOS in French rivers.

Marc Babut (✉ [marc.babut@inrae.fr](mailto:marc.babut@inrae.fr))

INRAE <https://orcid.org/0000-0002-6919-1314>

Benoit J.D. Ferrari

Swiss Centre for Applied Ecotoxicology

Patrick Jame

ISA UMR5280 CNRS Université de Lyon

Azziz Assoumani

Institut National de l'Environnement Industriel et des Risques

François Lestremau

Institut National de l'Environnement Industriel et des Risques

Nicolas Hette-Tronquart

Office français de la biodiversité

Cécile Miege

INRAE

Olivier Perceval

Office français de la biodiversité

---

## Research

**Keywords:** Water Framework Directive, chemical status, biota, monitoring, caging, *Gammarus fossarum*, PFOS, compliance checking, trophic magnification factor

**Posted Date:** September 22nd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-44363/v2>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published on October 7th, 2020. See the published version at <https://doi.org/10.1186/s12302-020-00416-4>.

# Monitoring priority substances in biota under the Water Framework Directive: How effective is a tiered approach based on caged invertebrates? A proof of concept study targeting PFOS in French rivers.

Authors: Marc BABUT<sup>1,\*</sup>, Benoit J.D. FERRARI<sup>2</sup>, Patrick JAME<sup>3</sup>, Azziz ASSOUMANI<sup>4</sup>, François LESTREMAU<sup>4</sup>, Nicolas HETTE-TRONQUART<sup>5</sup>, Cécile MIEGE<sup>1</sup> and Olivier PERCEVAL<sup>5</sup>

Affiliations

<sup>1</sup>INRAE, RIVERLY, F-69625, Villeurbanne, France

<sup>2</sup> Swiss Centre for Applied Ecotoxicology (Ecotox Center), EPFL ENAC IIE-GE, Station 2, CH-1015 Lausanne, Switzerland

<sup>3</sup>ISA, Université de Lyon, CNRS, Université Claude Bernard Lyon 1, Institut des Sciences Analytiques, UMR 5280, 5 rue de la Doua, F-69100 Villeurbanne, France

<sup>4</sup>INERIS, Unité Méthodes et développements en analyses pour l'environnement, F-60550, Verneuil-en-Halatte, France

<sup>5</sup>OFB (Office français de la biodiversité, Direction de la recherche et de l'appui scientifique), F-94300 Vincennes, France

\* Corresponding author: [marc.babut@inrae.fr](mailto:marc.babut@inrae.fr)

## Abstract

**Background:** This study aims to describe and test a tiered approach for assessing compliance to Environmental Quality standards (EQSs) for priority substances in biota in line with the European Water Framework Directive. This approach is based on caged gammarids and trophic magnification factors (TMFs) at the first tier, with fish analysed at the second tier at sites predicted to exceed the EQS at the first tier. A dataset was implemented by monitoring perfluorooctane sulfonate (PFOS) in caged gammarids exposed at 15 sites in French rivers, and in fish muscle and rest-of-body from the same sites. Isotopic ratios ( $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$ ) were also measured in gammarids and fish. Two scenarios were developed to compare measured PFOS concentrations in fish against predicted concentrations based on measures in caged gammarids and TMFs. Scenario (1) compared measured PFOS concentrations in fish fillets with predicted PFOS concentrations based on measured concentrations in caged gammarids and  $\delta^{15}\text{N}$ . Scenario (2) tested whether or not EQS exceedance was correctly predicted based on measured concentrations in caged gammarids and trophic levels (TLs) from wild fish and gammarid populations.

**Results:**  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  variations showed that caged gammarids used local food resources during exposure in the field. PFOS concentrations in gammarids were fairly variable through time at each site. In fish, concentrations ranged from  $<1 \text{ ng g}^{-1}$  to  $250 \text{ ng g}^{-1}$  (wet weight). After adjustment to the TL at which the EQS is set, 12 sites were above the EQS for PFOS. In scenario (1), predicted concentrations were almost correct at 7 sites out of 15. Most

incorrect predictions were overestimations that were slightly improved by applying a lower (neutral) TMF. In scenario (2) we tested several variants for parameters involved in the predictions. The most efficient combination yielded two wrong predictions out of 15. This result was obtained with a higher (more conservative) TMF value, mean concentrations in gammarids from several field exposures during a year, and a TL for gammarids at the median of the distribution in French rivers.

**Conclusion:** The proposed tiered approach was thus efficient. However, the number of sites was relatively limited, and the dataset was biased towards EQS exceedance. The tiered approach warrants further validation.

**Keywords**

Water Framework Directive – chemical status – biota – monitoring – caging – *Gammarus fossarum* - PFOS – compliance checking – trophic magnification factor

## 1 Background

The Water Framework Directive (WFD) originally laid down a number of requirements including a periodic assessment of the 'chemical status' of European water bodies comparing the concentrations of a set of priority substances (PSs) measured in water, sediment, or biota against their respective environmental quality standards (EQSs) (E.P. and E.C. 2000). EQSs are thresholds defining a good chemical status based on risks to aquatic ecosystems and human health. Nevertheless, the primary EQS implementation focused on water only, as there were no EQS available at the time for the targeted PS in sediment, nor in biota.

A first set of EQSs targeting biota was introduced in 2008 in the daughter directive updating the WFD for mercury, hexachlorobenzene and hexachlorobutadiene (E.C. 2008). An extended list of PS was adopted in 2013 along with the corresponding EQSs, with biota targeted for 11 PSs out of 45 (E.P. and E.C. 2013). Most of these chemicals are hydrophobic and prone to (or suspected of) biomagnification, presumably putting predators at risk. The European guidance issued to supporting the implementation of the 2013 directive assumed that concentrations critical to predators (or consumers) are most likely to occur in fish, with a theoretical trophic level (TL) of 4 in freshwaters and 5 in coastal waters (E.C. 2014). The guidance also recommended standardizing concentrations to a lipid content of 5% for hydrophobic substances, which bioaccumulate by lipid partitioning. For mercury and perfluorooctane sulfonate (PFOS), which do not accumulate by partitioning to lipids, it was recommended to standardize concentrations to a default dry weight fraction of 26%. As biota EQS values are based on either food consumption benchmarks or the secondary poisoning of piscivorous species, PS analysis should target either fillets or whole fish. Conversion factors based on regressions between fillet and whole-body concentrations can be used to translate measurements from fillet to whole-body matrix and vice versa (E.C. 2014 and citations therein).

Chemical monitoring on biota in water bodies can target species at a TL lower than 4. This situation is likely to occur in many cases, for practical (species abundance) and ethical (vertebrate sacrifice) considerations, in which case it is necessary to adjust the monitoring results to the required TL and lipid contents at which EQSs are set.

The 2013 directive authorized alternative monitoring options to fish, as long as such options provided an equivalent level of protection. To frame this possibility, the guidance document suggested implementing a tiered approach using alternative biota, such as wild crustaceans or bivalve molluscs, or other matrices, including caged

organisms (E.C. 2014). All these approaches are aimed at targeting sites where fish sampling would be fully justified by a risk of exceeding the EQSs, while also sparing time and resources and avoiding unnecessary fish sampling elsewhere. They all require predicting PS concentrations from field measurements in a theoretical standard fish having a TL of 4 and a lipid content of 5% (or a dry weight content of 26%). When alternative biota are used at this screening tier, the prediction of PS concentrations in the standard fish is based on the trophic magnification factor (TMF) equation, which describes the diet-weighted average biomagnification factor of chemical residues across a food web (Borgå et al. 2012).


Using caged invertebrates carries numerous advantages over sampling wild invertebrates. Besides making fieldwork more efficient, caged invertebrates allow controlling biotic confounding factors while measuring directly the bioavailable fraction of the contamination and provide robust, spatially-comparable results (Alric et al. 2019; Besse et al. 2013; Ciliberti et al. 2017). Among the freshwater invertebrate species, *Gammarus* spp. has emerged as suitable species for this kind of monitoring. Wild gammarids have been shown to accumulate trace elements (Filipović Marijić et al. 2016; Lebrun et al. 2014) and various persistent organic chemicals (Babut et al. 2017; Tlili et al. 2012), while caged gammarids have been successfully used for monitoring trace elements and organic chemicals (Besse et al. 2012; Besse et al. 2013; Lebrun et al. 2015). Nevertheless, strictly speaking, caged organisms do not belong to local food webs and so they cannot be assigned an actual TL.

Our purpose here is to describe a tiered approach for assessing compliance with EQSs for biota as per the European guidance requirements, using caged gammarids and selected TMFs, and to test this approach with one PS, the perfluorooctane sulfonate (PFOS) on riverine sites in France. To our knowledge, the tiered approach suggested in the European guidance (E.C. 2014) has never been tested in the real world at large spatial scale. Such a purpose entails addressing specific issues raised by the use of caged gammarids, such as the appropriate TL for these organisms partially disconnected from local food webs. As the ability to suitably predict either PS concentrations in fish or EQS exceedance depends also of the choices made for parameter values involved in these predictions, various options are explored and discussed.

## 2 Methods

### 2.1 Tiered approach concept

Figure 1 below summarizes the proposed monitoring strategy for PSs in biota. It involves caged organisms at the first tier (screening) with PS concentrations measured in fish at tier #2 if tier #1 results predict EQS exceedance. Conservative options should be preferred at tier #1, in order to minimize type II errors (or false negatives).



**Figure 1 – Schematic flow-chart of the biota monitoring approach using caged organisms**

#### 2.1.1 Tier #1 – screening monitoring sites for probable compliance with biota EQSs

Caging is carried out according to the French standard method “Water quality – In situ *Gammarus* caging for measuring bioaccumulation of chemical substances” (AFNOR 2019). Briefly, pools of male gammarids of a specific size class ( $\approx 2\text{mm}$ ) and originating from a unique wild population from a (pristine) reference site are introduced at monitored sites in polypropylene containers pierced with 1mm holes, thus allowing water, suspended particles and small organisms to flow through. The devices are left on site for three weeks, and gammarids are not fed during this period. At the end of exposure, the cages are retrieved, packed in cool boxes maintained at 4°C and sent to the laboratory within 24 hours. Gammarids are then pooled and processed for analysis. One measured concentration ( $C_{i\text{gam}}$ ) is thus available for each PS at each site.

Predicted concentrations in fish ( $C_{i\ fish}$ ) are estimated according to Equation 1, adapted from the European guidance for biota monitoring (E.C. 2014):

$$C_{i\ fish} = C_{i\ gam} \times TMF_i^{(TL_{fish}-TL_{gam})}$$

**Equation 1**

where  $TMF_i$  is the TMF selected for priority substance  $i$ , and  $TL_{fish}$  and  $TL_{gam}$  point to trophic levels (TLs) assigned to fish and caged gammarids, respectively.  $C_i$  designates the predicted whole-body concentration in fish ( $C_{i\ fish}$ ) or the measured concentration in gammarids ( $C_{i\ gam}$ ). Note that this equation is consistent with the generic trophic magnification model, which for hydrophobic substances is based on concentrations in lipids (Burkhard et al. 2013; Kidd et al. 2019).

Note too that estimating TLs is not strictly required, provided that  $\delta^{15}N$  measurements are available (Kidd et al. 2019), in which case TLs in Equation 1 are replaced by the respective  $\delta^{15}N$  values divided by 3.4‰, i.e. the mean trophic fractionation of  $\delta^{15}N$  (Post 2002).

A mean TL based on homologous local species is assigned to caged gammarids. As mentioned above, a value of 4 should be used for  $TL_{fish}$  in freshwaters so as to predict EQS exceedance. Compliance to EQS will be checked by comparing the geometric mean of measured concentrations (and the geometric standard deviation) or the concentration in a pooled sample against the EQS (E.C. 2014).

As few of the fish species selected for monitoring PSs in biota are expected to present the theoretical TL of 4, measured concentrations need to be adjusted at this TL using Equation 2 (E.C. 2014), which is derived from the above-mentioned trophic magnification model:

$$C_{TL-adjusted} = C_{measured} * TMF^{(4-TL_x)}$$

**Equation 2**

where  $x$  is the species monitored and  $C$  are the concentrations.  $TL$  and  $TMF$  were defined previously.

As stated above,  $EQS_{biota}$  generally refer to organisms that have a mean lipid content of 5%. Measured concentrations should therefore be adjusted to this lipid content prior to being compared to  $EQS_{biota}$ . For PSs such as mercury or PFOS that do not accumulate via partitioning to lipids, the guidance recommends by default an adjustment to a standard dry weight of 26% (Equation 3; E.C. 2014):

$$conc_{adjusted} = conc_{measured} \times \frac{0.26}{measured\ dry\ weight}$$

### Equation 3

#### 2.1.2 Tier #2 – Confirming exceedance of biota EQSs by sampling and analysing target fish species

Fish would be collected and analyzed at sites where predicted  $C_i$  fish would exceed one or several biota EQSs. The guidance document (E.C. 2014) states that two species, representing different habitats and feeding conditions and different bioaccumulation capacities, should be selected. In France, this has led to selecting barbel (*Barbus barbus*) and chub (*Squalius cephalus*). Roach (*Rutilus rutilus*) could be used instead of barbel at sites (catchments) where barbel is rare or absent.

The abovementioned guidance leaves the options open for analyzing individuals or pools, while outlining advantages and disadvantages of both. The option currently selected in France is to analyze one pool of about ten individuals of the same species and same size class per sampling event. Compliance with EQSs would follow the same rules as explained above.

## 2.2 Study design

Fifteen sites were selected from the current French river monitoring network. Selection criteria included biota availability and anthropogenic pressures in order to get a gradient of contamination by PSs.

Caged gammarids were exposed at these 15 sites throughout France for three weeks as per the existing guideline (AFNOR 2019), at three successive periods in 2018 (June, September and November). Site characteristics are presented in Table S1 of the Supplementary material (SM).

Fish were caught by electrofishing between the end of June and the end of November 2018, targeting either chub (*Squalius cephalus*), barbel (*Barbus barbus*) or roach (*Rutilus rutilus*). They were euthanatized on site with an excess dose of eugenol, in adherence with current ethical rules. Fifteen pools of 8 to 11 individuals of the same species and belonging to the same size class (Table S2 in SM) were put together and taken to the laboratory in cool boxes for dissection and analysis.

## 2.3 Analysis

Details on analytical protocols are provided in SM, section II.

### 2.3.1 Isotopic ratios

Finely ground gammarid samples of about 0.5 mg dry weight (dw) each were placed and weighed in tin capsules, then injected into a helium flow, with a limited quantity of oxygen (O<sub>2</sub>) into a combustion unit (950 °C). Gases passed first through a copper oven (650°C) so as to reduce nitrogen oxides into nitrogen (N<sub>2</sub>) and capture the excess O<sub>2</sub>, and then into a cold tube containing magnesium perchlorate to trap the remaining water. N<sub>2</sub> and CO<sub>2</sub> gases were finally separated on a chromatographic column and injected into an isotopic mass spectrometer. Results are expressed as δ<sup>13</sup>C or δ<sup>15</sup>N, i.e. the ratio of [<sup>13</sup>C]/[<sup>12</sup>C] or [<sup>15</sup>N]/[<sup>14</sup>N] in the sample to that of a reference sample (SM, section II.1). Isotopic ratio measurement in fish samples (i.e. fillets) followed the same procedure as that applied to caged gammarids. For both δ<sup>13</sup>C and δ<sup>15</sup>N, standard deviations of the value for gammarids and fish were calculated on two measurements, when the gap between these two values was <0.60 δ. Above this gap, a third measurement was done and accounted for in the standard deviation (SD) calculation. SD values ranged between 0.01 ‰ and 0.43 ‰, and between 0.01 ‰ and 1.16 ‰ for δ<sup>13</sup>C and δ<sup>15</sup>N respectively.

### 2.3.2 PFOS

PFOS was analyzed in gammarids and fish tissues, i.e. fillets and “rest of body” (ROB), following a modified Aquaref MA46 method (Aquaref 2013). Samples were first freeze-dried and finely ground. Internal standards and methanol (10 mL) were then added to an aliquot (1 g) of the resulting powder. Next followed a purification process involving activated carbon, evaporation and re-dissolution in acetonitrile. PFOS was analyzed by high-pressure liquid chromatography on a C18 column followed by tandem mass spectrometry (electro-spray ionization, negative mode). Injection blanks, calibration, method controls and methods blanks were used throughout the analytical process so as to check for absence of system contamination, trueness and signal drift. The limit of quantification (LQ) was determined according to French Standard NF T90-210 (details in SM, section II.2). Briefly, first, two candidate LQs were estimated based on the signal-to-noise ratio, then 20 blank tissue samples were spiked at the concentration level of the two candidate LQs (10 samples for each candidate LQ), and then analyzed. The candidate LQ was validated if the average determined concentration, including its uncertainty, was between 40% and 160% of the spiked target concentration.

## 2.4 TMF selection

TMFs are obtained by regression between the chemical concentrations in organisms of a food-web and their respective TLs (Borgå et al. 2012); in principle, all concentrations refer to whole-body. The TMFs used for the

tests were taken from previous studies on French riverine sites (Simonnet-Laprade et al. 2019a; Simonnet-Laprade et al. 2019b); while they came from field studies in a context relevant to the purpose of the present study, they were comparable to the TMFs retrieved from a literature review and screened according to Kidd et al. (2019). The TMFs determined in French rivers are compiled in Table S4, and TMFs from the literature survey are compiled in Figure S1 and Table S5 in SM. Two TMF values were selected for the predictive tests: the median (TMF<sub>50</sub>) and third quartile (TMF<sub>75</sub>) derived from the six TMFs for French riverine sites (Table S4 in SM). While TMF<sub>50</sub> is *a priori* neutral in terms of risk, TMF<sub>75</sub> represents a more conservative option, both for the adjustment of concentrations measured in fish and for the prediction based on concentrations in caged gammarids. TMF<sub>75</sub> was preferred to an even more conservative option such as the 90<sup>th</sup> percentile due to the limited number of TMFs available.

## 2.5 Prediction of fish contamination on the basis of caged organisms and TMFs – test design

The PFOS concentrations predicted in fish muscle (fillet) according to Equation 1 were compared against measured concentrations in this tissue (scenario #1). The predictions were based on PFOS concentrations in caged gammarids (September campaign), TMF values, and  $\delta^{15}\text{N}$  measured in caged gammarids ( $\delta^{15}\text{N}_{\text{gam}}$ ) and fish ( $\delta^{15}\text{N}_{\text{fish}}$ ). The slope of the regression between fillet and whole-body concentrations was used to convert the predicted whole-body concentrations to predicted fillet concentrations (Equation 4, derived from Equation 1; details in SM, section III); this conversion stems from the fact that the EQS for PFOS was derived from a benchmark targeting human health and fish consumption. The predictions were deemed correct if the concentration ranges were similar.

$$C_{\text{fish-fillet}} = \frac{C_{\text{gam}} \times \text{TMF}^{((\delta^{15}\text{N}_{\text{fish}} - \delta^{15}\text{N}_{\text{gam}})/3.4)}}{\text{slope}}$$

### Equation 4

where  $C_{\text{fish-fillet}}$  is predicted PFOS concentration in fish fillet,  $C_{\text{gam}}$  is measured PFOS concentration in gammarids, and *slope* is slope of the regression between PFOS concentrations measured in fillets and in whole fish.


Scenario #2 aimed to check whether the predicted concentrations were correct in terms of EQS exceedance. Prediction of EQS exceedance entailed adjusting the predicted whole-body concentrations in fish to TL=4 before estimating the corresponding concentrations in fillets (Equation 5, derived from Equation 2).

$$C_{ff-TL-adj} = \frac{C_{gam} \times TMF^{(4-TL_{gam})}}{slope}$$

**Equation 5**

where  $C_{ff-TL-adj}$  is predicted PFOS concentration in fillet of a fish at TL=4 according to Equation 2, using TLs collected from FishBase (Froese and Pauly 2019). This predicted PFOS concentration is compared against the measured PFOS concentrations adjusted to TL=4 according to Equation 2.  $TL_{gam}$  is the TL assigned to caged gammarids, assuming they hold a similar trophic position to that of wild populations.

Both scenarios are summarized in Figure 2.



**Figure 2 – Testing process for checking the reliability of Tier#1 (grey boxes: measurements; white boxes: modelled predictions, or data from external sources).**

Several options were used in each scenario for TMF values (TMF<sub>50</sub>, TMF<sub>75</sub>), for slope of the regression between fillet and whole-body concentrations in fish, and for TLs (fish and gammarids). For PFOS concentrations in gammarids, we used either the results from the September campaign (close to fish capture) or the arithmetic mean of two or three campaigns.

**2.6 Data processing and statistics**

Relationships between fillet and whole-body concentrations were assessed by Theil-Sen regression with using Pro-UCL 5.1 software (USEPA 2016), to include censored data in the datasets with Mann-Kendall tests applied prior to these regressions in order to check for the trend.


Multiple sets were compared using non-parametric Kruskal-Wallis tests followed by post-hoc Dunn tests (Pro-UCL 5.1 software or XLStat v19.4). Significance was set at  $\alpha=0.05$  in all tests.

### 3 Results

Gammarids were successfully exposed in the field and retrieved at three periods of the year; 43 samples were collected as two samples were lost at the same site (September and November campaigns). Survival rates ranged between 16% and 72% in June, 25% and 56% in September, and 42% and 66% in November, but enough quantities were collected sufficient for analysing PFOS and isotopic ratios in all samples. Fifteen fish samples were collected (Table S2 in SM): 13 chub, one barbel and one roach. All the data presented in the next section is available online (Babut et al. 2020; Miede et al. 2020).

#### 3.1 Isotopic ratios

Ranges  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values for caged gammarids from June (campaign #1) to November (campaign #3) 2018 are reported in Figure 3. Both isotopic ratios were significantly higher (for all sites pooled and almost systematically at each site) in November than in the two other campaigns. These variations added to the seasonal variations of isotopic ratios in gammarids at the site providing organisms for caging, i.e. a shift towards lower  $\delta^{13}\text{C}$  values in spring and towards slightly higher  $\delta^{15}\text{N}$  values in autumn/winter (SM, section V – Figure S2). Variations during caging was calculated on initial (T0) values in June, September and November (SM, section V).



**Figure 3 – Distributions of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in caged organisms at 15 sites in June (1), September (2) and November (3) 2018**

Both isotopic ratios changed in caged gammarids throughout the field exposure period.  $\delta^{13}\text{C}$  changes were variable among sites in June and September, with no discernible consistent pattern (Figure S2 in SM). In November,  $\delta^{13}\text{C}$  values became less negative during field exposure at almost all sites.  $\delta^{15}\text{N}$  values during field exposure increased at all sites but two in all three campaigns (Figure S3). These changes in isotopic ratios suggest that the caged gammarids exploited local food sources during field exposure, leading to a slightly higher  $\delta^{15}\text{N}$  value (mean gain  $\approx 1\text{‰}$ ) at most sites. The very nature of the food items consumed by caged gammarids also appears quite different in November, as shown by the positive variations of  $\delta^{13}\text{C}$  values contrasting with the more diverse variations in the June and September campaigns.

In fish,  $\delta^{13}\text{C}$  ranged between  $-28.46\text{‰}$  and  $-22.21\text{‰}$ , whereas  $\delta^{15}\text{N}$  varied between  $9.10\text{‰}$  and  $16.77\text{‰}$ . The lowest  $\delta^{15}\text{N}$  values were observed at sites 09 and 10, which are presumably less subject to anthropogenic pressures. The highest  $\delta^{15}\text{N}$  value was at site 05, which is located downstream from a densely populated watershed hosting numerous industries, as well as intensive agriculture.

Trophic levels in wild *Gammarus* spp. ranged from 1.26 to 2.91 (Figure S4 and Table S6 in SM, data from Hette-Tronquart and Belliard (2018)). In scenario #2, we used both the 75<sup>th</sup> percentile (2.28) and the median (2.00).

## 3.2 Contamination ranges

### 3.2.1 Caged organisms

PFOS was quantified in 80 % to 87% of caged organism samples from June and September campaigns respectively, but only 47% of samples from the November campaign. Concentrations ranges were nevertheless similar among the three campaigns, reaching  $22.7 \text{ ng g}^{-1} \text{ ww}$ ,  $22.5 \text{ ng g}^{-1} \text{ ww}$  and  $17.3 \text{ ng g}^{-1} \text{ ww}$  in June, September and November respectively.

PFOS levels in gammarids were fairly variable through time at each site, and no specific pattern could be identified (Figure S5 part A in SM).

### 3.2.2 Fish

PFOS was quantified in 77% of fish fillet or ROB samples. PFOS concentrations in fillet ranged from <LQ to  $250 \text{ ng g}^{-1} \text{ ww}$ . For 11 out of 15 pooled fish samples, PFOS was quantified in both fillets and ROB; for the four remaining

pools, PFOS concentrations were <LQ in fillets, or ROB, or both. For these four samples, it was therefore impossible to determine whole fish concentrations as detailed in SM, section III. Weighted mean concentrations in whole fish thus ranged from undetermined to 553 ng g<sup>-1</sup> ww. The current EQS (9.1 ng g<sup>-1</sup> ww), which was derived from a benchmark for human food consumption (EFSA 2008) and thus applies to fillet, was exceeded at two sites while a third site displayed a concentration at the EQS (SM, Figure S5 part B). Note that this classification is not a consistent compliance check, as no adjustment was made according to TL or dry weight content.

A Theil-Sen regression including all 15 sample pairs yielded a slope of 2.202, with a 95% confidence interval from 1.800 to 2.671 (intercept – 1.675;  $p \approx 0.00004$ ; SM, section III). This slope is very close to the value obtained with a linear regression accounting for uncensored values only (slope of  $2.20 \pm 0.01$ ; intercept –  $0.318 \pm 0.665$ ;  $R^2 = 1.000$ ,  $p < 0.0001$ ), although this regression was strongly influenced by the extremely high concentrations measured in fish from site 05. The Theil-Sen regression appears thus more robust. This result is also close to the slope found in a study on the same fish species (barbel and chub) in a few French rivers, but different from the slopes found in Lake Geneva (SM, Table S3).

### 3.3 Prediction of PFOS concentrations in fish

#### 3.3.1 Comparison of predicted and measured PFOS concentrations (scenario #1)

Predicted concentrations in fish fillets based on concentrations in caged gammarids from the September campaign are reported in Table 1. This prediction was impossible for sites 01 and 11, where PFOS was <LQ in caged gammarids. As expected, predicted concentrations were lower when based on TMF<sub>50</sub> than on TMF<sub>75</sub>, while using the lower bound of the confidence interval of the slope led to higher predicted concentrations, and vice versa.

Site	Predicted concentrations calculated with:						Measured concentrations (fish fillet)
	Median slope		Upper bound (CI 95%) of slope		Lower bound (CI 95%) of slope		
	TMF <sub>75</sub>	TMF <sub>50</sub>	TMF <sub>75</sub>	TMF <sub>50</sub>	TMF <sub>75</sub>	TMF <sub>50</sub>	
01	nd	nd	nd	nd	nd	nd	3.88
02	19.6	15.1	16.17	12.42	23.10	17.74	10.5
03	11.6	9.15	9.53	7.54	13.62	10.76	8.89
04	8.60	7.08	7.08	5.83	10.11	8.33	252
05	310	203	256	168	365	239	7.86
06	16.1	12.7	13.2	10.5	18.9	14.9	3.45
07	5.43	4.34	4.47	3.57	6.39	5.10	3.48
08	6.59	5.30	5.43	4.36	7.75	6.23	<LQ
09	4.37	3.55	3.60	2.93	5.14	4.18	6.03
10	3.71	3.33	3.06	2.74	4.37	3.92	<LQ
11	nd	nd	nd	nd	nd	nd	<LQ
12	4.11	3.16	3.38	2.60	4.83	3.72	4.64
13	24.6	18.9	20.3	15.5	28.9	22.2	3.43
14	41.8	32.4	34.4	26.7	49.2	38.1	7.97
15	22.8	17.6	18.8	14.5	26.8	20.6	3.81

**Table 1 – Predicted and measured PFOS concentrations (ng g<sup>-1</sup> ww) in fish fillets based on the September campaign (nd: not determined).**

Sites 04 and 05 are clearly outliers, with either predicted concentrations well below the measured ones, or vice versa. In the case of site 04, this discrepancy is due to the very high PFOS concentration in fish, which contrasts with a concentration in the lower range of PFOS concentrations in gammarids in the September campaign. In the case of site 05, the very high predicted PFOS concentration results from the combination of a high caged gammarid PFOS concentration and a high  $\delta^{15}\text{N}$  in fish.

Six other sites (02, 06, 08, 13, 14, 15) gave predicted concentrations well above the measured ones, whatever the TMF used, except for the variant combining TMF<sub>50</sub> and the upper bound of the slope confidence interval at site 02. Four sites (03, 07, 09, 12) showed near overlap between measured and predicted levels. For the three remaining sites (01, 10, 11), PFOS was <LQ in gammarids, making it impossible to predict concentrations in fish, but the corresponding measured PFOS concentrations in fish were either low or <LQ. In summary, predicted concentrations were almost correct at seven sites (01, 03, 07, 09, 10, 11, and 12) out of 15. Most of the incorrect predictions were overestimations that were slightly improved by applying TMF<sub>50</sub> instead of TMF<sub>75</sub>.

When the 95 % upper confidence limit of the regression slope was applied instead of the median slope to convert whole-fish to fillet concentrations, predicted concentrations were lower. Nevertheless, this did not substantially improve overall prediction accuracy. Predicted concentrations at sites 02, 06, 08, 13, 14 and 15 remained well

above the measured concentrations. Predicted concentrations at sites 03 and 07 remained close to measured concentrations, while at sites 09 and 12 they were slightly more underestimated than previously. Since the intercept of the regression between fillet and whole-fish concentrations was negative, introducing it in the prediction would increase the predicted concentrations, thus increasing the overestimation for sites 02, 06, 08, 13, 14 and 15. We therefore did not explore further this option in any scenarios. Similar conclusions can be drawn when applying the lower bound of the slope confidence interval.

### 3.3.2 Predicted PFOS concentrations vs EQS exceedance (scenario #2)

Four sets of concentrations adjusted to the TL at which the EQS is set were calculated based on Equation 2 and Equation 3, using either TMF<sub>75</sub> or TMF<sub>50</sub> (Table 2), and TLs obtained from FishBase (Froese and Pauly 2019), and with or without adjustment to mean dry weight contents (Table 2). The TL values varied from 2.7 (*S. cephalus*) at sites 01, 03, 04, 05, 06, 07, 08, 09, 10, 11, 13, 14 and 15, to 3.0 (*R. rutilus*) at site 12 and 3.1 (*B. barbuis*) at site 02. Weighted mean dry weight contents for fish are reported in SM, Table S2. These calculations led to higher predicted PFOS concentrations in fish fillets at all sites. Consequently, the number of samples exceeding the EQS of 9.1 ng g<sup>-1</sup> ww rose to 12, whatever the TMF applied (Table 2). We kept only the concentrations unadjusted to dry weight contents for the sake of comparison with predicted concentrations.

Site	Measured PFOS conc. (fillet)	Adjusted PFOS concentrations (fillet)			
		No adjustment to dry weight content		Adjusted to dry weight content	
		TMF <sub>75</sub>	TMF <sub>50</sub>	TMF <sub>75</sub>	TMF <sub>50</sub>
01	3.88	16.0	13.4	19.6	16.4
02	10.5	28.1	24.9	27.9	24.7
03	8.89	36.7	30.8	44.2	37.1
04	252	1039	872	1323	1110
05	7.86	32.4	27.2	41.9	35.2
06	3.45	14.3	12.0	17.6	14.8
07	3.48	14.4	12.0	17.3	14.5
08	<LQ	nd	nd	nd	nd
09	6.03	24.9	20.9	29.4	24.7
10	<LQ	nd	nd	nd	nd
11	<LQ	nd	nd	nd	nd
12	4.64	13.8	12.1	15.0	13.1
13	3.43	14.1	11.9	16.0	13.4
14	7.97	32.9	27.6	34.0	28.5
15	3.81	15.7	13.2	18.3	15.4

**Table 2 – Measured/adjusted PFOS concentrations in fish fillet (nd: not determined)**

The selected adjusted-measured concentrations were compared against predicted concentrations (example in Table 3, based on concentrations in gammarids and the median slope of the regression between fillet and whole-

body concentrations). Type II errors (or false negatives) occurred when the predicted concentration was below the EQS, while the adjusted concentration was above; the opposite situation yielded type I errors (false positives).

In this example, seven to nine fish samples were predicted to exceed the EQS for PFOS depending on the TMF applied, resulting in zero to one type I and four to five type II errors. The type II errors occurred mainly at sites where measured concentrations in gammarids were low or <LQ.

Site	TMF <sub>75</sub>			TMF <sub>50</sub>		
	Err. I	Err. II	Overall	Err. I	Err. II	Overall
01	0	1	1	0	1	1
02	0	0	0	0	0	0
03	0	0	0	0	1	1
04	0	0	0	0	0	0
05	0	0	0	0	0	0
06	0	0	0	0	0	0
07	0	1	1	0	1	1
08	0	0	0	0	0	0
09	0	1	1	0	1	1
10	1	0	1	0	0	0
11	0	0	0	0	0	0
12	0	1	1	0	1	1
13	0	0	0	0	0	0
14	0	0	0	0	0	0
15	0	0	0	0	0	0
<b>Total</b>	<b>1</b>	<b>4</b>	<b>5</b>	<b>0</b>	<b>5</b>	<b>5</b>

**Table 3 – Type I and type II errors (0= correct; 1= error; the third and sixth columns report whether the prediction of EQS exceedance was correct, i.e. 0, or erroneous, i.e. 1) in scenario #2: variant using a single field campaign with caged gammarids, and the median slope of the regression between fillet and whole body concentrations.**

Using the 95% upper confidence limit of the slope instead of the central value yielded the same number of type II errors when using TMF<sub>75</sub>, and no type I errors. In the TMF<sub>50</sub>-based variant, overall number of errors rose to 6, due to type II errors only. Conversely, using the 95% lower confidence limit of the slope yielded similar results to using the median slope, with four type II errors and one type I error for TMF<sub>75</sub> and TMF<sub>50</sub>.

Table 4 reports the results for scenario #2 where several variants of each parameter of Equation 5 were combined. For  $C_{gam}$ , we used either the concentrations from one campaign, or the means of two or three campaigns. For the slope, we tested the central value, and the 95% bounds of its confidence interval. For  $TL_{gam}$ , we took either the 75<sup>th</sup> percentile or the median of the TL distribution among 93 sites in French rivers (SM, Table

S6 and Figure S4, derived from Hette-Tronquart and Belliard (2018). Median  $TL_{gam}$  was introduced only twice, after having identified the most efficient combination of the other parameters (TMF,  $C_{gam}$  and slope).

TMF	$C_{gam}$	slope	$TL_{gam}$	N Type I	N Type II	Correct
TMF <sub>50</sub>	1 campaign (Sep)	lower bound	75 <sup>th</sup> percentile	1	4	10
TMF <sub>50</sub>	1 campaign (Sep)	median	75 <sup>th</sup> percentile	0	5	10
TMF <sub>50</sub>	1 campaign (Sep)	upper bound	75 <sup>th</sup> percentile	0	6	9
TMF <sub>50</sub>	2 campaigns (June-Sep)	lower bound	75 <sup>th</sup> percentile	1	4	10
TMF <sub>50</sub>	2 campaigns (June-Sep)	median	75 <sup>th</sup> percentile	0	5	10
TMF <sub>50</sub>	2 campaigns (June-Sep)	upper bound	75 <sup>th</sup> percentile	0	5	10
TMF <sub>50</sub>	3 campaigns	lower bound	75 <sup>th</sup> percentile	0	4	11
TMF <sub>50</sub>	3 campaigns	median	75 <sup>th</sup> percentile	0	5	10
TMF <sub>50</sub>	3 campaigns	upper bound	75 <sup>th</sup> percentile	0	5	10
TMF <sub>75</sub>	1 campaign (Sep)	lower bound	75 <sup>th</sup> percentile	1	4	10
TMF <sub>75</sub>	1 campaign (Sep)	median	75 <sup>th</sup> percentile	1	4	10
TMF <sub>75</sub>	1 campaign (Sep)	upper bound	75 <sup>th</sup> percentile	0	4	11
TMF <sub>75</sub>	2 campaigns (June-Sep)	lower bound	75 <sup>th</sup> percentile	1	1	13
TMF <sub>75</sub>	2 campaigns (June-Sep)	lower bound	median	2	0	13
TMF <sub>75</sub>	2 campaigns (June-Sep)	median	75 <sup>th</sup> percentile	1	3	11
TMF <sub>75</sub>	2 campaigns (June-Sep)	upper bound	75 <sup>th</sup> percentile	0	5	10
TMF <sub>75</sub>	3 campaigns	lower bound	75 <sup>th</sup> percentile	1	1	13
TMF <sub>75</sub>	3 campaigns	lower bound	median	2	0	13
TMF <sub>75</sub>	3 campaigns	median	75 <sup>th</sup> percentile	0	3	12
TMF <sub>75</sub>	3 campaigns	upper bound	75 <sup>th</sup> percentile	0	5	10

**Table 4 – Results obtained in scenario #2 according to Equation 5 parameter variants.  $C_{gam}$  is the concentration measured in caged gammarids, slope is the slope of the regression between fish fillet and whole-body concentrations, and  $TL_{gam}$  is the trophic level assigned to caged gammarids.**

The TMF<sub>50</sub>-based variants yielded four to six type II errors and no more than 11 correct predictions out of 15. Results were more contrasted with TMF<sub>75</sub>, with between zero and five type II errors and few type I errors, as with TMF<sub>50</sub>. The best predictions were obtained with the association of TMF<sub>75</sub>, mean  $C_{gam}$  from two or three campaigns, the lower bound of the slope confidence interval, and the median value of  $TL_{gam}$  distribution. An equal number of correct predictions was obtained with the 75<sup>th</sup> percentile of  $TL_{gam}$  distribution, two to three campaigns and the lower bound of the slope confidence interval, but with one type II error instead of zero. Environmental protection imperatives makes it is less worrying to overpredict an EQS exceedance (i.e. to commit a type I error) than to underpredict, so the variant with no type II error was slightly better.

## 4 Discussion

Exploring the effective validity of the proposed tiered approach concept entails two things: (i) addressing whether using caged organisms, especially gammarids, is a relevant solution, and (ii) examining the terms of Equation 5 (or Equation 4 if the purpose is to predict concentrations in fish without considering EQS exceedance). The deployment of caged gammarids has already been shown to be a robust biomonitoring tool (Besse et al. 2012; Besse et al. 2013), so the question here is more about finding a sensible proxy for the TL assigned to them and justifying it (which is why this point is included in the global discussion on the terms of Equation 5). We also discuss the first term of Equation 5, i.e. the measured  $C_{fish-tillet}$  adjusted to a standard fish, as this adjustment, and accordingly the gap to predicted concentrations, is sensitive to the values of TMF,  $TL_{fish}$  and dry weight content.

### 4.1 Adjustment to a standard fish

The adjustment recommended in the EU guidance relies on three parameters, i.e. TMF, TL for fish, and dry weight content which is presented as a proxy for total protein content (E.C. 2014).

Dry weight contents of fish pools in this study ranged from 20.1% to 26.2 % (Table S2 in SM), with a median value of 21.6%. Consequently, adjusting to the default dry weight content of 26% (E.C. 2014) increased the adjusted concentrations by 3.5% to 29%, except at the site where the actual dry weight content of the fish sample reached 26.2%. Therefore, the default dry weight content seems excessively high in our case, in contrast with a previous German study that found a relatively limited effect of standard dry weight content on adjusted PFOS concentrations in bream pools (Flidner et al. 2016). Based on our data, a standard (mean) dry weight content of 22% appears more appropriate.

Furthermore, PFOS accumulation is driven not by total protein content but by specific proteins (Jones et al. 2003; Luebker et al. 2002; Ng and Hungerbühler 2013). These specific proteins (chiefly serum albumin and liver fatty acid-binding proteins) are not expressed to the same extent across fish species (Ng and Hungerbühler 2013). Therefore, (i) the adjustment based on a proxy for total protein content does not seem justified, and (ii) the principle of this adjustment itself appears questionable in the case of PFOS. For these reasons, the measured/adjusted concentrations in scenario #2 were used without adjustment to dry weight contents.

TLs from FishBase (Froese and Pauly 2019) may also lack specificity compared to more geographically-focused values. Based on a script developed for calculating the TL distribution of freshwater fish species in France (Hette-

Tronquart and Belliard 2018) and the related database, we observed that median TL values for the three species studied here are higher than the respective mean TLs in FishBase (SM, Table S7). This would in turn lead to lower measured/adjusted concentrations, and possibly, to a different diagnosis of compliance with the EQS. Furthermore, using for instance the 25<sup>th</sup> percentile of the monitored species TL distributions would have the opposite effect, which could be deemed safer, as it would lower the probability of misclassifying the sites. It would therefore be advisable to complete the European guidance on these aspects.

## 4.2 Concentrations in gammarids

Gammarids were successfully exposed in the field and retrieved at three periods of the year, so the current standard protocol (AFNOR 2019) safely yields adequate materials for monitoring bioaccumulative PSs. Nevertheless, the rate of quantification was lower in the November campaign, in which the isotopic signatures of caged gammarids pointed to important changes in their food sources. We tested three options for the term  $C_{gam}$ , i.e. data from (a) one single campaign, or mean concentrations from (b) two or (c) three campaigns. The selection of the September campaign for option (a) was justified by its temporal proximity to the fish sampling campaign and a better PFOS quantification rate. Nevertheless, options (b) and (c) yielded better results for both scenarios #1 and #2. Due to the limited quantification rate in the November campaign,  $C_{gam}$  were not significantly different between options (b) and (c). The key conclusion here is that multiple campaigns would be more representative of real fish exposure.

## 4.3 Trophic magnification factors

TMFs are ecosystem-specific metrics, meaning they can vary across ecosystems (Borgå et al. 2012). Conversely, it seems almost impossible to precisely characterize the ecosystems at all sites of a nationwide monitoring network. From a nationwide assessment perspective, a unique TMF value applicable at all sites seems preferable in practice, but challenging to confidently select (Kidd et al. 2019). We carefully reviewed existing TMFs for PFOS in aquatic ecosystems (Table S4 in SM), and selected a set of TMFs derived from French rivers. All these TMFs were obtained in shallow freshwater ecosystems, and involved similar fish species. From this set, we derived two TMF values, i.e. the median (TMF<sub>50</sub>) and the third quartile (TMF<sub>75</sub>). According to Equation 5, TMF<sub>75</sub> is more representative of a worst-case scenario, leading to higher predicted concentrations. Using the 90<sup>th</sup> percentile of TMFs could be considered even more relevant, i.e. closer to the worst possible case, but as there were few TMF values available (Table S5 in SM), TMF<sub>75</sub> appears more robust. Nonetheless, the influence of TMF value on the fit

between predicted and measured concentrations (scenario #1) was moderate, leading to between 10% and 34% variations in predicted concentrations (all variants taken together). In scenario #2, using TMF<sub>50</sub> instead of TMF<sub>75</sub> did not substantially change the distribution between wrong and correct predictions, which seems logical: as mentioned before the TMF has also an influence on the measured/adjusted concentration, so we compared predicted against measured/adjusted concentrations based on identical TMFs on both sides of the equation. However, in this scenario #2, no variant with TMF<sub>50</sub> reached less than four type II errors (out of 15 sites).

#### 4.4 Trophic level of caged *Gammarus* spp. and fish

In the tiered approach framework, caged organisms substitute for their local/wild homologues, in order to lower the influence of confounding factors (e.g. nutrients) on PS bioaccumulation while ensuring inter-site comparability. Caged organisms are thus implicitly assumed to occupy a similar trophic position to their wild homologues.

According to the variations of both  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  during their exposure in the field, caged gammarids consumed local food items, which supports their use for monitoring bioaccumulative PSs, despite the fact that caged organisms are less connected to local food webs than their wild homologues. Moreover, caged gammarids showed very little influence of local (anthropic) nitrogen sources, unlike fish, which showed large variations in  $\delta^{15}\text{N}$ . This observation also argues for applying a TL derived from the TL distribution in wild gammarids. Indeed, here caged organisms were supplied by a wild population living in a pristine site, and their  $\delta^{15}\text{N}$  value varied little during exposure in the field (SM, Figure S3). They were thus still close to their initial trophic position even though they could not represent the actual TL of wild gammarids at each site. Instead, they could represent a generic population.

A robust determination of TLs in fish would not be applicable in a nationwide monitoring network at tier #1, and may even remain impracticable at tier #2. Thus, the application of fixed TL values appears relevant in practice. Again, generic TL values derived from the TL distribution in each fish species could be sufficient for checking EQS compliance, allowing to compare the severity of contamination at a large spatial scale, while limiting the effect of confounding factors. Nevertheless, the use of TLs derived from the actual TL distributions in e.g. large watersheds or ecoregions should be preferred to TLs from FishBase.

#### 4.5 Relationships between fish fillet and whole-body concentrations (conversion factor)

In the case of PFOS, the EQS refers to a human-health safety limit, which makes it important to consider the relationship between the concentrations in fillet and in whole fish. Three options were examined for the slope of the regression, namely the central value (median) and the lower and upper bounds of the slope CI. Neither the median nor the upper bound provided a good predictive accuracy in scenario #1, and the numbers of type I and type II errors were almost identical in scenario #2. The best result was obtained with the lower bound of the slope confidence interval, which tends to increase the predicted concentrations. Combined with  $TMF_{75}$  and mean  $C_{gam}$  (from two or three campaigns), it resulted in only two errors, either two type I and no type II, or one of each type. Because these parameter value choices both concur to increase predicted concentrations, it seems likely that caged gammarids tended to underestimate PFOS exposure for fish in the case of scenario #2.

One important limitation of this study was that the proportion of sites presumably above the EQS was rather high, meaning that we could not check whether the approach is as equally valid for predicting compliant situations.

## 5 Conclusion

For PFOS at least, the proposed tiered approach could work properly. Predictions at tier #1 were quite satisfactory when applying a combination of high TMF (i.e.  $TMF_{75}$ ), generic TLs, and the lower bound of the slope confidence interval. However, the number of sites was relatively limited, and the dataset was biased towards EQS exceedance. Further validation of the tiered approach for checking the compliance with EQSs for biota are still needed and should involve more sites covering a larger PFOS contamination gradient, as well as other PSs that behave differently to PFOS. Moreover, improvements are also needed for assigning appropriate TLs for the targeted fish species in the different catchments of concern.

### List of abbreviations and acronyms

EQS	Environmental Quality Standard, as defined in the Water Framework Directive
LQ	Limit of quantification
PFOS	Perfluorooctane sulfonate (CAS #1763-23-1), a mixture of linear and branched isomers
PS	Priority substance, as defined in the Water Framework Directive
TL	Trophic level

TMF	Trophic Magnification Factor
WFD	Water Framework Directive, i.e. Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy

### Declarations

- Ethics approval and consent to participate
  - Fishing (capture, transport, and disposition) authorization for scientific or monitoring purposes: granted to “Office Français de la Biodiversité” (OFB) regional offices by prefectural decrees from the local authorities. Fish were euthanized with an appropriate dose of eugenol.
  - Consent to participate: not applicable.
- Consent for publication: all authors approved the final manuscript and agreed to its submission to *Environmental Sciences Europe*.
- Availability of data and material
- Competing interests: the authors declare no competing interest
- Funding: fieldwork and analyses were given grant support by the French Agency for Biodiversity (AFB; Office Français de la Biodiversité, OFB since January 2020) as part of the prospective surveillance network programme for 2018-2019, part B.
- Authors' contributions
  - Marc BABUT: study design, data processing, writing
  - Benoit J.D. FERRARI: method design, writing, reviewing.
  - Patrick JAME: isotopic ratio analysis and interpretation
  - Azziz ASSOUMANI and François LESTREMAU: PFOS analysis, data processing
  - Nicolas HETTE-TRONQUART: interpretation, writing
  - Cecile MIEGE: study design and coordination, data management, reviewing.
  - Olivier PERCEVAL: study design, writing, reviewing.
- Acknowledgements: we thank the staff of BIOMAE (*Biosurveillance de la qualité de l'eau et des milieux aquatiques*, Chateau-Gaillard, France), and Nicolas DELORME and Laura GARNERO (INRAE, RIVERLY, Ecotoxicology laboratory) for their proactive assistance for fieldwork and handling of samples. Erik BONJOUR and Anthony ANCHISI managed isotopic ratio analysis. We also thank Glen McCULLEY (Metaform Langues, Clermont-Ferrand, France) for copy-editing the manuscript.

### References

- AFNOR (2019) Qualité de l'eau — Engagement in situ de gammarees pour la mesure de la bioaccumulation de substances chimiques vol XP T 90-721. <https://www.afnor.org/en/purchase-a-standard/>
- Alric B et al. (2019) Multisubstance Indicators Based on Caged Gammarus Bioaccumulation Reveal the Influence of Chemical Contamination on Stream Macroinvertebrate Abundances across France *Environ Sci Technol* 53:5906-5915 doi:10.1021/acs.est.9b01271
- Aquaref (2013) Composés perfluorés (chaîne linéaire en C8) - Méthode d'analyse dans les biotes. [https://www.aquaref.fr/system/files/Aquaref\\_2013\\_D1b\\_INERIS\\_MA46\\_PFCs\\_biotes\\_VF.pdf](https://www.aquaref.fr/system/files/Aquaref_2013_D1b_INERIS_MA46_PFCs_biotes_VF.pdf)
- Babut M, Jame P, Anchisi A, Bonjour E, Delorme N, Miege C (2020) Isotopic ratios in freshwater gammarids (caged) and wild fish from 15 continental sites (2018) and one reference site (2019), VERSION PROVISOIRE edn. Portail Data INRAE. <https://doi.org/10.15454/7GBGUQ>
- Babut M et al. (2017) Per- and poly-fluoroalkyl compounds in freshwater fish from the Rhône River: influence of fish size, diet, prey contamination and biotransformation *Sci Tot Environ* 605-606:38-47 doi:10.1016/j.scitotenv.2017.06.111
- Besse J-P, Geffard O, Coquery M (2012) Relevance and applicability of active biomonitoring in continental waters under the Water Framework Directive *TrAC-Trend Anal Chem* 36:113-127 doi:10.1016/j.trac.2012.04.004
- Besse JP, Coquery M, Lopes C, Chaumot A, Budzinski H, Labadie P, Geffard O (2013) Caged *Gammarus fossarum* (Crustacea) as a robust tool for the characterization of bioavailable contamination levels in continental waters: Towards the determination of threshold values *Wat Res* 47:650-660 doi:10.1016/j.watres.2012.10.024

- Borgå K et al. (2012) Trophic magnification factors: Considerations of ecology, ecosystems and study design *Integr Environ Assess Manag* 8:64-84 doi:10.1002/ieam.244
- Burkhard LP, Borgå K, Powell DE, Leonards P, Muir DCG, Parkerton TF, Woodburn KB (2013) Improving the Quality and Scientific Understanding of Trophic Magnification Factors (TMFs) *Environ Sci Technol* 47:1186–1187 doi:10.1021/es305253r
- Ciliberti A et al. (2017) Caged Gammarus as biomonitors identifying thresholds of toxic metal bioavailability that affect gammarid densities at the French national scale *Wat Res* 118:131-140 doi:10.1016/j.watres.2017.04.031
- E.C. (2008) Directive 2008/105 of The European Parliament and of the Council on environmental quality standards in the field of water policy and amending Directive 2000/60/EC.
- E.C. (2014) Common Implementation Strategy for the Water Framework Directive (2000/60/EC) - Guidance Document No. 32 on Biota monitoring (The implementation of EQS<sub>BIOTA</sub>) under the Water Framework Directive. European Union. doi:10.2779/833200
- E.P., E.C. (2000) Directive 2000/60/EC of the European Parliament and of the Council of 23 October 2000 establishing a framework for Community action in the field of water policy vol 2000/60, 22 Dec. 2000 edn. Official Journal of the European Communities,
- E.P., E.C. (2013) Directive 2013/39/EU of the European Parliament and of the Council of 12 August 2013 amending Directives 2000/60/EC and 2008/105/EC as regards priority substances in the field of water policy Text with EEA relevance vol 2013/39, 24 Aug. 2013 edn. Official Journal of the European Communities,
- EFSA (2008) Perfluorooctane sulfonate (PFOS), perfluorooctanoic acid (PFOA) and their salts Scientific Opinion of the Panel on Contaminants in the Food chain *EFSA Journal* 6:653 doi:10.2903/j.efsa.2008.653
- Filipović Marijić V et al. (2016) Investigation of the soluble metals in tissue as biological response pattern to environmental pollutants (Gammarus fossarum example) *Chemosphere* 154:300-309 doi:10.1016/j.chemosphere.2016.03.058
- Fliedner A et al. (2016) Biota monitoring and the Water Framework Directive—can normalization overcome shortcomings in sampling strategies? *Environ Sci Pollut Res* 23:21927-21939 doi:10.1007/s11356-016-7442-2
- Froese R, Pauly D (2019) FishBase. World Wide Web electronic publication. [www.fishbase.org](http://www.fishbase.org), Accessed 21.06.2019
- Hette-Tronquart N, Belliard J (2018) Indicateurs du fonctionnement trophique des cours d'eau - approche par l'analyse des isotopes stables vol action 20.
- Jones PD, Hu W, De Coen W, Newsted JL, Giesy JP (2003) Binding of perfluorinated fatty acids to serum proteins *Environ Toxicol Chem* 22:2639-2649 doi:10.1897/02-553
- Kidd KA et al. (2019) Practical Advice for Selecting or Determining Trophic Magnification Factors for Application Under the European Union Water Framework Directive *Integr Environ Assess Manage* 15:266-277 doi:10.1002/ieam.4102
- Lebrun JD, Geffard O, Urien N, François A, Uher E, Fechner LC (2015) Seasonal variability and inter-species comparison of metal bioaccumulation in caged gammarids under urban diffuse contamination gradient: Implications for biomonitoring investigations *Sci Tot Environ* 511:501-508 doi:10.1016/j.scitotenv.2014.12.078
- Lebrun JD, Uher E, Tusseau-Vuillemin MH, Gourlay-Francé C (2014) Essential metal contents in indigenous gammarids related to exposure levels at the river basin scale: Metal-dependent models of bioaccumulation and geochemical correlations *Sci Tot Environ* 466-467:100-108 doi:10.1016/j.scitotenv.2013.07.003
- Luebker DJ, Hansen KJ, Bass NM, Butenhoff JL, Seacat AM (2002) Interactions of fluorochemicals with rat liver fatty acid-binding protein *Toxicology* 176:175-185 doi:10.1016/S0300-483X(02)00081-1
- Miege C, Assoumani A, Lestremay F, Boutet P, Recoura-Massaquant R, Jubeaux G, Babut M (2020) PFOS and PFOA concentrations in freshwater gammarids (caged) and wild fish from 15 continental sites (2018), VERSION PROVISOIRE edn. Portail Data INRAE. <https://doi.org/10.15454/LKZJR4>
- Ng CA, Hungerbühler K (2013) Bioconcentration of perfluorinated alkyl acids: How important is specific binding? *Environ Sci Technol* 47:7214-7223 doi:10.1021/es400981a
- Post DM (2002) Using stable isotopes to estimate trophic position: Models, methods, and assumptions *Ecology* 83:703-718 doi:10.2307/3071875
- Simonnet-Laprade C et al. (2019a) Evidence for the spatial variability of poly- and perfluoroalkyl substance (PFAS) trophic magnification in French rivers *Sci Tot Environ* 686:393-401 doi:10.1016/j.scitotenv.2019.05.461
- Simonnet-Laprade C et al. (2019b) Biomagnification of perfluoroalkyl acids (PFAAs) in the food web of an urban river: assessment of the trophic transfer of targeted and unknown precursors and implications *Environ Sci-Proc Imp* 21:1864-1874 doi:10.1039/c9em00322c
- Tilili K, Labadie P, Bourges C, Desportes A, Chevreuil M (2012) Bioaccumulation of polybrominated diphenyl ethers by the freshwater benthic amphipod *Gammarus pulex* *Arch Environ Contam Toxicol* 63:69-76 doi:10.1007/s00244-012-9751-x
- USEPA (2016) Statistical Software ProUCL 5.1.00 for Environmental Applications for Data Sets with and without Nondetect Observations. <https://www.epa.gov/land-research/proucl-software> Accessed 2017.01.15 2017

# Figures

Tier #1 (screening)




Figure 1

Schematic flow-chart of the biota monitoring approach using caged organisms





Figure 2

Testing process for checking the reliability of Tier#1 (grey boxes: measurements; white boxes: modelled predictions, or data from external sources).



**Figure 3**

Distributions of  $\delta^{13}\text{C}$  and  $\delta^{15}\text{N}$  values in caged organisms at 15 sites in June (1), September (2) and November (3) 2018

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TieredapproachFSIformatted.docx](#)