

Description and Genome Analysis of *Microvirga Antarctica* Sp. Nov., A Novel Pink-Pigmented Psychrotolerant Bacterium Isolated From Antarctic Soil

Lin Zhu

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology

Weiwei Ping

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology

Siyue Zhang

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology

Ya Chen

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology

Ying Zhang

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology

Jianli Zhang (✉ zhangjianli@bit.edu.cn)

Key Laboratory of Molecular Medicine and Biotherapy, School of Life Science, Beijing Institute of Technology <https://orcid.org/0000-0001-7340-6392>

Research Article

Keywords: Microvirga, Polyphasic taxonomy, psychrotolerant, ANI, genome analysis, functional enzymes

Posted Date: May 19th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-443907/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Antonie van Leeuwenhoek on October 25th, 2021. See the published version at <https://doi.org/10.1007/s10482-021-01674-9>.

Abstract

During the investigation of exploring potential psychrotolerant species from Antarctica soil, a novel pink-pigmented bacterium designated strain 3D7^T was isolated. Cells of the isolate were observed to be rod-shaped (0.7–0.9×1.0–2.2 μm), Gram-stain negative and non-motile. It was able to grow at 4–32 °C, pH 7.0–10.0 and in the presence of 0–3 % (w/v) NaCl. Phylogenetic analysis based on 16S rRNA gene sequences showed that strain 3D7^T belongs to the genus *Microvirga* and was most closely related to '*Microvirga brassicacearum*' CDVBN77^T (98.3 %), *Microvirga subterranea* DSM 14364^T (96.8 %), *Microvirga guangxiensis* 25B^T (96.5 %) and *Microvirga aerophila* DSM 21344^T (96.5 %). The predominant quinone was ubiquinone 10 (Q-10), and the major fatty acids were summed feature 8 (C_{18:1} ω7c and/or C_{18:1} ω6c) and C_{19:0} cyclo ω8c. The predominant polar lipids were phosphatidylcholine and phosphatidylethanolamine. The genomic DNA G + C content of strain 3D7^T was 63.5 mol%. Its genome sequence showed genes encoding phosphatases and lipases. Genetic machinery related to carbohydrate-active enzymes and secondary metabolites were also observed. The average nucleotide identity and digital DNA–DNA hybridization values based on whole genome sequences of strain 3D7^T and its closely related species were below the threshold range for species determination. Phenotypic, chemotaxonomic, phylogenetic and genomic analyses suggested that strain 3D7^T represents a novel species of the genus *Microvirga*, for which the name *Microvirga antarctica* sp. nov. is proposed. The type strain is 3D7^T (= CGMCC 1.13821^T = KCTC 72465^T).

Introduction

To survive in an extremely cold and oligotrophic habitat, Antarctic microorganisms have formed unique physiological and biochemical properties in the long-term natural selection evolution (Niederberger et al. 2008). Many strains have genetic machinery to degrade multiple compounds as a source of nutrients and can produce low-temperature (cold-active) enzymes, antibacterial and anti-cancer active substances (Zhang et al. 2004), which are valuable in many fields including environmental engineering, agriculture biotechnology, pharmaceutical industry and enzyme industry. Accordingly, we carried out a programme to explore potential sources of psychrotolerant species from Antarctic soil, a putatively novel strain 3D7^T of the genus *Microvirga* was isolated. The genus *Microvirga* was proposed by Kanso and Patel (2003), with *Microvirga subterranea* as the type species. It belongs to the family *Methylobacteriaceae* of the order *Rhizobiales*. At the time of writing, the genus *Microvirga* contains 18 species listed on LPSN (List of Prokaryotic Names with Standing in Nomenclature: www.bacterio.net) with validly published names. They are distributed widely in various ecological habitats, such as air (Weon et al. 2010), natural, domestic, and contaminated soils (Dahal et al. 2017; Tapase et al. 2017; Zhang et al. 2009; Zhang et al. 2019), geothermal water (Kanso et al. 2003), Tibet hot spring sediments (Liu et al. 2020), human stool (Caputo et al. 2016), nodules of native legumes and cowpea (Ardley et al. 2012; Radl et al. 2017; Safronova et al. 2017) and roots of rapeseed plants (Jiménez-Gómez et al. 2019). Most members of the genus *Microvirga* showed (middle) thermophilic. Some studies report the genetic potential of strains

classified within the *Microvirga* genus for arsenic oxidation (Tapase et al. 2017) and the production of pigments, amylolytic enzymes (Radl et al. 2017), phosphatases and exopolysaccharides (Jiménez-Gómez et al. 2019). Based on polyphasic taxonomic characterisation, we propose the description of *Microvirga antarctica* sp. nov., isolated as a novel psychrotolerant member of the genus *Microvirga* with phosphatase and lipase activities. Moreover, we obtained and analyzed the genome sequence of strain *M. antarctica* 3D7^T, showing its genetic potential for biotechnological industrial application.

Materials And Methods

Isolation and culture conditions

Strain 3D7^T was isolated from a soil sample collected from the surface of Deception Island (62° 55' 09" S, 60° 34' 46" W), Antarctica. The collected soil sample (0.9 g) was suspended in 8.1 mL sterile water and stirred for 30 min as a 10⁻¹ dilution solution, then diluted it to 10⁻² by gradient. 150 µL of the 10⁻² sample dilution was spread on Reasoner's 2A agar (R2A; AOBOX) medium (pH 7.5). After 7 days of incubation at 15 °C, representative colonies were picked and purified by streaking repeatedly. A pink-coloured isolate, designated strain 3D7^T, was picked up and subsequently purified by plate streaking. The purified strain was stored in 20 % (v/v) glycerol suspensions at -20°C.

DNA amplification and determination of 16S rRNA gene sequence

The total DNA of strain 3D7^T was isolated from cell suspension culture. The 16S rRNA gene was amplified by PCR with the bacterial universal forward primer 27F and reverse primer 1525R (Li et al. 2006). The products above were purified and sequenced by BGI (The Beijing Genomics Institute). After sequencing, the 16S rRNA gene sequence of strain 3D7^T was obtained and similarity searches were performed by using the EzBioCloud server (www.ezbiocloud.net/identify) (Yoon et al. 2017). The phylogenetic tree was constructed according to the neighbour-joining (NJ) algorithm (Saitou et al. 1987) and supported by the minimum-evolution (ME) (Rzhetsky et al. 1992) and maximum-likelihood (ML) algorithms (Felsenstein 1981) in the MEGA X program (Kumar et al. 2018). Kimura's two-parameter model was used to calculate the evolutionary distances (Kimura 1980). Bootstrap values were determined based on 1000 replications (Felsenstein 1985).

Genome sequencing, assembly and function analysis

The genome of strain 3D7^T was sequenced using the Illumina HiSeq systems with paired-end sequencing technology. The sequencing data was filtered to remove the sequences containing the adaptor and the low quality data, and the obtained clean data was used for subsequent analysis. The genome assembly was performed by SOAPdenovo (version 2.04) (Li et al. 2010; Li et al, 2008). The assembly results were submitted to the NCBI (www.ncbi.nlm.nih.gov). The function of coding genes in the assembled genome were annotated by Gene Ontology (GO) (Ashburner et al. 2000), Clusters of Orthologous Groups (COG) (Galperin et al. 2015) and Kyoto Encyclopedia of Genes and Genomes (KEGG) (Minoru et al. 2016). The

carbohydrate-active enzymes (CAZymes) were analyzed using HMMER annotation (Zhang et al. 2018), and the analysis of gene clusters related to secondary metabolites production was performed using antiSMASH 5.0 webserver (Blin et al. 2019).

DNA-DNA hybridization and genome-based phylogenetic analysis

The genomic information of related strains of the same genus or neighbouring genera was obtained from the EzTaxon and NCBI databases. Phylogenomic tree based on the whole genome sequences of strain 3D7^T and related species was constructed using the Composition Vector (CV) approach. Average nucleotide identity (ANI) based on the BLAST algorithm (ANIb) and the MUMmer ultra-rapid aligning tool (ANIm), as well as the correlation indexes of tetranucleotide signatures (Tetra) were calculated through the website of JSpeciesWS (<http://jspecies.ribohost.com/jspeciesws/>) (Richter et al. 2016). The orthoANLu values were estimated using the EzBioCloud web service (www.ezbiocloud.net/tools/ani) as described by Yoon et al. (2017). Digital DNA–DNA hybridization (dDDH) was conducted using the Genome-to-Genome Distance Calculator (GGDC; version 2.1) under the recommended Formula 2 (<http://ggdc.dsmz.de/distcalc2.php>) provided by the DSMZ website (Meier-Kolthoff et al. 2013).

Morphology, physiological and biochemical analysis

Growth tests were performed on Reasoner's 2A, tryptic soy agar (TSA; AOBX), nutrient agar (NA; AOBX) and Ancylobacter–Spirosoma Medium (ASM; glucose 1 g, peptone 1 g, yeast extract 1 g, agar 15 g). Growth at different temperatures (0, 4, 10, 15, 20, 25, 28, 30, 32, 35 and 40 °C) was observed on R2A for 7 days. Tolerance to different NaCl concentrations (0–6 %, at intervals of 0.5 %, w/v) and pH (4.0–12.0, at intervals of 0.5 unit) were tested in R2A broth. The pH of the basal medium was adjusted using the buffer system: pH 4.0–5.0: 0.1 M citric acid/0.1 M sodium citrate; pH 6.0–8.0: 0.1 M KH₂PO₄/0.1 M NaOH; pH 9.0–10.0: 0.1 M NaHCO₃/0.1 M Na₂CO₃; pH 11.0: 0.05 M Na₂HPO₄/0.1 M NaOH; pH 12.0: 0.2 M KCl/0.2 M NaOH (Xu et al. 2005). Cell morphology was observed by light microscope (BH-2; Olympus) and transmission electron microscope (JEM-1400, JEDL) after 3 days growth on R2A medium at 28°C. Gram stain reaction was performed according to the method described by Dong and Cai (2001). Anaerobic growth test (Zhang et al. 2019) was performed on R2A medium with 1 g pyrogalllic acid and 2 ml 10 % (w/v) NaOH in the plate, which was then sealed with Vaseline and growth detected for up to 7 days. Motility was tested in R2A medium containing 0.4 % (w/v) agar and using the hanging-drop technique as described by Bernardet et al. (2002). Catalase activity was determined by assessing bubble production in 3 % (v/v) H₂O₂, and oxidase activity was determined using 1 % (w/v) tetramethyl-p-phenylenediamine (Ohta et al. 1983). Sensitivity and resistance to antibiotics were performed on R2A plates using filter-paper discs containing different antibiotics (Hangzhou Microbial Reagent). Hydrolysis of starch, gelatin and Tween 20, 40, 60 and 80 was determined on R2A for incubation at 28°C for up to 7 days as described by Tindall et al. (2007). Other physiological properties and enzyme activities were determined using the API 50CH, API 20NE and API ZYM systems (bioMérieux) according to the manufacturers' instructions.

Chemotaxonomic analyses

Cellular fatty acids of strain 3D7^T and its reference strains were analysed by using colonies grown on R2A medium at 28 °C for 3 days. The fatty acid methyl ester mixtures were separated and analysed using the standard protocol of the Sherlock Microbial Identification System (MIDI Sherlock software package, version 6.0) (Kämpfer et al. 1996; Sasser 1990). For analyses of quinones and polar lipids, cells were collected at the exponential phase by centrifugation, then washed three times with sterilized water and freeze-dried. Isoprenoid quinones were extracted and purified by the methods of Collins (1985) and then analysed by reversed-phase HPLC. Polar lipids were extracted from freeze-dried cells and loaded onto thin-layer silica gel 60 plates (Merck). Two-dimensional migration was performed on each plate using chloroform–methanol–water (65:25:4, by vol.) as the first solvent and chloroform–acetic acid–methanol–water (80:15:12:4, by vol.) as the second one (Minni et al. 1979; Collins et al. 1980). Total polar lipids were detected by spraying with phosphomolybdic acid solution followed by heating at 110 °C for 10 min. Aminolipids were detected by spraying the plate with a 0.4 % (w/v) solution of ninhydrin in butanol saturated with water followed by heating at 105 °C for 10 min. Phospholipids were detected by spraying with the reagent of Dittmer and Lester.

Results And Discussion

Phylogenetic characteristics

The nearly complete 16S rRNA gene sequence of strain 3D7^T (1396 bp) was determined and compared with the corresponding sequences. It shared the highest 16S rRNA gene similarity to '*Microvirga brassicacearum*' CDVBN77^T (98.3 %), followed by *Microvirga subterranea* DSM 14364^T (96.8 %), *Microvirga guangxiensis* 25B^T (96.5 %), and *Microvirga aerophila* DSM 21344^T (96.5 %). The NJ analyses (Fig. 1) showed that strain 3D7^T shared a branching node with '*M. brassicacearum*' CDVBN77^T, which was highly consistent with ME tree (Fig. S1) and ML tree (Fig. S2). It was clear that strain 3D7^T was a member of the genus *Microvirga*.

Genome composition and DNA-DNA hybridisation

The draft genome sequence of strain 3D7^T was 4,518,469 bp in length with 221 contigs. The coverage, N50 and DNA G + C content were 410×, 431,466 bp and 63.5 mol%. The genome had 4321 protein-coding genes and 49 RNAs (Table 1). Genomic analyses showed that strain 3D7^T and '*M. brassicacearum*' CDVBN77^T yielded ANIb and dDDH values of 77.5 and 22.2 %, respectively. The ANI values between strain 3D7^T and other species of the genus *Microvirga* are detailed in Table 1, which are all below standard criteria for classifying strains as different species (95–96 %) (Kim et al. 2014). The dDDH values between strain 3D7^T and other species of the genus *Microvirga* are also detailed in Table 1, and they are far below the 70 % cut-off value generally recommended for species differentiation (Wayne et al. 1987). A genome-based phylogenetic tree is included in Fig. S3, which shows that strain 3D7^T is affiliated to the genus *Microvirga*. These data confirm that strain 3D7^T represents a novel species of the genus *Microvirga*.

Genome features and function prediction

Gene Ontology database analysis results showed that strain 3D7^T has 42 different types functions. Among them, there were 14 features related to cell component, 14 features related to molecular function, 14 features related to biological process (Fig. S4). Among the 20 general COG functional categories, the detailed distribution of genes was as follows: Amino acid transport and metabolism, 497 genes; Inorganic ion transport and metabolism, 293 genes; Energy production and conversion, 204 genes; Transcription, 202 genes; Carbohydrate transport and metabolism, 200 genes. Detailed information of the COG functional categories was presented in Fig. S5. KEGG metabolic pathways were classified according to the relationship between KO (KEGG ORTHOLOGY) and Pathway. Functional annotation of genes by comparisons against the manually curated KEGG GENES database revealed that there were 62 genes related to the biosynthesis of other secondary metabolites, 122 genes related to the biodegradation and metabolism of xenobiotics, 374 genes related to the metabolism of carbohydrates and 45 genes related to the metabolism of terpenoids and polyketides (Fig. S6). The genome annotations showed genes encoding for proteins with phosphatase activity, such as the enzymes alkaline phosphatase (EC 3.1.3.1), acid phosphatase (EC 3.1.3.2), inorganic triphosphatase (EC 3.6.1.25) and pyrophosphatase (EC 3.6.1.1). Some enzymes involved in the production of triglyceride lipases, including lysophospholipase (EC 3.1.1.5) and unidentified phospholipase were also observed. These capabilities were tentatively proven in physiological tests, with potential applications in the agriculture biotechnology, washing industry and low-temperature environment remediation.

Analysis of the genome sequence of strain 3D7^T showed 132 genes encoding different CAZymes in five different classes: glycoside hydrolases (GHs), enzymes that catalyze the hydrolysis of glycosidic linkage of glucoside—27 gene counts; glycosyltransferases (GTs), involved in the formation of glycosidic bonds—47 gene counts; carbohydrate esterases (CEs), which hydrolyze carbohydrate esters—35 gene counts; auxiliary activities (AAs), redox enzymes that act in conjunction with CAZymes—20 gene counts; polysaccharide lyases (PLs), which perform non-hydrolytic cleavage of glycosidic bonds—2 gene counts and carbohydrate-binding modules (CBMs)—1 gene count (Table S1). AntiSMASH output revealed four biosynthetic gene clusters (BGCs) involved in the secondary metabolism of the bacterium. One of those clusters encodes terpene BGC, which is related to the synthesis of isoindolinomycin. Other clusters encode an arylpolyene, a hserlactone and a terpene BGCs that are not described for the production of an already known molecule. These genetic characteristics indicated that strain 3D7^T may have biotechnological potential for the degradation of biomass and the pharmaceutical industry.

Phenotypic characteristics

Strain 3D7^T grew well on R2A agar and ASM agar, but grew weakly on TSA and NA. Colonies on R2A agar plate were light-pink, semi-transparent, smooth and round. Cells were Gram-stain-negative, aerobic, non-motile, rod-shaped, 1.0–2.2 µm long and 0.7–0.9 µm wide (Fig. S7). It was able to grow at 4–32 °C (optimum, 25–28 °C), pH 7.0–10.0 (optimum, 7.0–7.5) and in the presence of 0–3 % (w/v) NaCl (optimum without NaCl). These characteristics markedly differentiated strain 3D7^T from the first related

strain '*M. brassicacearum*' CDVBN77^T. Sensitive to penicillin (10 U), ampicillin (10 µg), chloramphenicol (30 µg), tetracycline (30 µg), streptomycin (10 µg) and neomycin (30 µg), but resistant to polymyxin B (300 IU), vancomycin (30 µg) and bacitracin (0.04 U). Strain 3D7^T hydrolysed Tween 20, 40, 60, and weakly hydrolysed Tween 80. It can hydrolyse aesculin, but not gelatin and tyrosine. Positive reaction for alkaline phosphatase, valine arylamidase and naphthol-AS-BI-phosphohydrolase, weakly positive for lipase (C14) and trypsin. These characteristics differentiated strain 3D7^T from '*M. brassicacearum*' CDVBN77^T and other closely related reference strains. More differential characteristics between strain 3D7^T and its closely related species in the genus *Microvirga* were given in Table 2, and the other detailed physiological and biochemical characteristics are present in the species description.

Chemotaxonomic characteristics

The major cellular fatty acids of strain 3D7^T (> 10 %) were summed feature 8 (C_{18:1} ω7c and/or C_{18:1} ω6c) (36.2 %) and C_{19:0} cyclo ω8c (21.7 %), which was similar to that of closely related species of the genus *Microvirga*. Minor qualitative and quantitative differences could be used to distinguish strain 3D7^T from the closest relatives of the genus *Microvirga*. Compared with '*M. brassicacearum*' CDVBN77^T, strain 3D7^T possessed higher amounts of C_{16:0}, summed feature 2 (C_{14:0} 3-OH and/or iso-C_{16:1} I) and summed feature 3 (C_{16:1} ω6c and/or C_{16:1} ω7c), and lower amounts of C_{18:0}, C_{18:0} 3-OH, C_{18:1} ω7c 11-methyl and feature 8 (C_{18:1} ω7c and/or C_{18:1} ω6c). C_{14:0} and C_{17:0} cyclo were detected in strain 3D7^T, but not detected in '*M. brassicacearum*' CDVBN77^T (Table 3). The predominant respiratory quinone of strain 3D7^T was Q-10, which was in good agreement with other species of the genus *Microvirga*. The polar lipids of strain 3D7^T consisted of phosphatidylcholine and phosphatidylethanolamine as the major component, plus one unidentified aminophospholipid, two unidentified amino lipids and three unidentified lipids (Fig. S8). Strain 3D7^T shared the same major polar lipids with most of the described species of the genus *Microvirga*.

In conclusion, all phenotypic, chemotaxonomic, phylogenetic and genomic analyses suggested that strain 3D7^T should be considered to represent a novel species of the genus *Microvirga*, for which the name *Microvirga antarctica* sp. nov. is proposed.

Description of *Microvirga antarctica* sp. nov.

Microvirga antarctica (ant.arc'ti.ca. L. fem. adj. *antarctica* southern, pertaining to the Antarctica, where the type strain was isolated)

Cells are Gram-stain-negative, aerobic, non-motile and rod-shaped (0.7–0.9×1.0–2.2 µm). Growth occurs on R2A agar and ASM agar, weakly on NA and TSA. Colonies are light-pink, semi-transparent, smooth, round and smaller than 1.0 mm in diameter after 3 days at 28 °C. Growth occurs at a range of 4–32 °C (optimum, 25–28 °C) and pH 7.0–10.0 (optimum, 7.0–7.5) and in the presence of 0–3 % (w/v) NaCl (optimum without NaCl). Oxidase, catalase and nitrate reduction are positive, but glucose fermentation,

arginine dihydrolase, indole production and urease are negative. Hydrolyses aesculin, Tween 20, 40 and 60, weakly hydrolyses Tween 80, but not starch. Assimilates L-arabinose, D-xylose, D-ribose and D-cellobiose, weakly assimilates D-fucose, D-glucose and D-melosanose. Positive for alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, acid phosphatase and naphthol-AS-BI-phosphohydrolase, weakly positive for lipase (C14), cystine arylamidase and trypsin. The major polar lipids are phosphatidylcholine and phosphatidylethanolamine. The predominant quinone is Q-10 and the major fatty acids are summed feature 8 (C_{18:1} ω7c and/or C_{18:1} ω6c) and C_{19:0} cyclo ω8c. The genomic DNA G + C content of the type strain is 63.5 mol%.

The type strain, 3D7^T (= CGMCC 1.13821^T = KCTC 72465^T), was isolated from a soil sample collected from Deception Island, Antarctica. The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain 3D7^T is MH561859. This Whole Genome Shotgun project of strain 3D7^T has been deposited at DDBJ/ENA/GenBank under the accession number JAGEMM000000000.

Declarations

Funding information

This research was supported by the National Key Research and Development Program of China (2016YFC0501302) and by the National Natural Science Foundation of China (NSFC, grant number 31070002).

Author's contributions ZJL and ZL designed research and project outline. ZL and ZY performed isolation, deposition and polyphasic taxonomy. ZL, CY and PWW performed genome analysis. ZL, PWW and ZSY drafted the manuscript. ZJL revised the manuscript. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest The authors declare that they have no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

1. Ardley JK, Parker MA, De Meyer SE, Trengove RD, O'Hara GW, Reeve WG, Yates RJ, Dilworth, MJ, Willems A, Howieson JG (2012) *Microvirga lupini* sp. nov., *Microvirga lotononidis* sp. nov. and *Microvirga zambiensis* sp. nov. are alphaproteobacterial root-nodule bacteria that specifically nodulate and fix nitrogen with geographically and taxonomically separate legume hosts. *Int J Syst Evol Microbiol* 62:2579–2588
2. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K, Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese JC, Richardson JE, Ringwald

- M, Rubin GM, Sherlock G (2000)** Gene Ontology: tool for the unification of biology. *Nature Genetics* 25:25–29
3. **Bernardet JF, Nakagawa Y, Holmes B, Flavobacteri ST (2002)** Proposed minimal standards for describing new taxa of the family *Flavobacteriaceae* and emended description of the family. *Int J Syst Evol Microbiol* 52:1049–1070
 4. **Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Weber T (2019)** antiSMASH 5.0: Updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res* 47: 81–87
 5. **Caputo A, Lagier JC, Azza S, Robert C, Mouelhi D, Fournier PE, Raoult D (2016)** *Microvirga massiliensis* sp. nov., the human commensal with the largest genome. *Microbiologyopen* 5:307–322
 6. **Collins MD (1985)** Isoprenoid quinone analysis in classification and identification. In: Goodfellow M and Minnikin DE (editors). *Chemical Methods in Bacterial Systematics*. London: Academic Press. 267–287
 7. **Collins MD, Jones D (1980)** Lipids in the classification and identification of coryneform bacteria containing peptidoglycan based on 2, 4-diaminobutyric acid. *J Appl Bacteriol* 48:459–470
 8. **Dahal RH, Kim J (2017)** *Microvirga soli* sp. nov., an alphaproteobacterium isolated from soil. *Int J Syst Evol Microbiol* 67:127–132
 9. **Desper R, Gascuel O (2004)** Theoretical foundation of the balanced minimum evolution method of phylogenetic inference and its relationship to weighted least-squares tree fitting. *Mol Biol Evol* 21:587–598
 10. **Dong XZ, Cai MY (2001)** *Determinative Manual for Routine Bacteriology*. Beijing: Scientific Press
 11. **Felsenstein J (1981)** Evolutionary trees from DNA sequences: A maximum likelihood approach. *J Mol Evol* 17:368–376
 12. **Felsenstein J (1985)** Confidence limits on phylogenies: An approach using the bootstrap. *Evolution* 39:783–791
 13. **Galperin MY, Makarova KS, Wolf YI, Koonin EV (2015)** Expanded microbial genome coverage and improved protein family annotation in the COG database *Nucleic Acids Res* 43:261–269
 14. **Jiménez-Gómez A, Saati-Santamaría Z, Igual JM, Rivas R, Mateos PF, García-Fraile P (2019)** Genome insights into the novel species *Microvirga brassicacearum*, a rapeseed endophyte with biotechnological potential. *Microorganisms*, 7(9).
 15. **Kämpfer P, Kroppenstedt RM (1996)** Numerical analysis of fatty acid patterns of coryneform bacteria and related taxa. *Can J Microbiol* 42:989–1005
 16. **Kanso S, Patel BK(2003)** *Microvirga subterranea* gen. nov., sp. nov., a moderate thermophile from a deep subsurface Australian thermal aquifer. *Int J Syst Evol Microbiol* 53:401–406
 17. **Kim M, Oh HS, Park SC, Chun J (2014)** Towards a taxonomic coherence between average nucleotide identity and 16S rRNA gene sequence similarity for species demarcation of prokaryotes. *Int J Syst Evol Microbiol* 64:346–351.

18. **Kimura M (1980)** A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16:111–120
19. **Kumar S, Stecher G, Li M, Knyaz C, Tamura K (2018)** MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 35:1547–1549
20. **Lefort V, Desper R, Gascuel O (2015)** FastME 2.0: a comprehensive, accurate, and fast distance-based phylogeny inference program: table 1. *Mol Biol Evol* 32:2798–2800
21. **Li R, Li Y, Kristiansen K, Wang J (2008)** Soap: short oligonucleotide alignment program. *Bioinformatics* 24:713–714
22. **Li R, Zhu HM, Ruan J, Qian WB, Fang XD, Shi ZB, Li YR, Li ST, Shan G, Kristiansen K, Li SG, Yang HM, Wang J, Wang J (2010)** De novo assembly of human genomes with massively parallel short read sequencing. *Genome Res* 20:265–272
23. **Liu ZT, Xian WD, Li MM, Liu L, Ming YZ, Jiao JY, Fang BZ, Xiao M, Li WJ (2020)** *Microvirga arsenatis* sp. nov., an arsenate reduction bacterium isolated from Tibet hot spring sediments. *Antonie Van Leeuwenhoek* 113:1147–1153
24. **Li WJ, Zhang YQ, Schumann P, Chen HH, Hozzein WN, Tian XP, Xu LH, Jiang CL (2006)** *Kocuria aegyptia* sp. nov., a novel actinobacterium isolated from a saline, alkaline desert soil in Egypt. *Int J Syst Evol Microbiol* 56:733–737
25. **Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M (2013)** Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60
26. **Minnikin DE, Collins MD, Goodfellow M (1979)** Fatty acid and polar lipid composition in the classification of *Cellulomonas*, *Oerskovia* and related taxa. *J Appl Bacteriol* 47:87–95
27. **Minoru K, Yoko S, Masayuki K, Miho F, Mao T (2016)** KEGG as a reference resource for gene and protein annotation. *Nucleic Acids Res* 1:457–462
28. **Niederberger TD, McDonald IR, Hacker AL, Soo RM, Barrett JE, Wall DH, Cary, SC (2008)** Microbial community composition in soils of Northern Victoria Land, Antarctica. *Enviro Microbiol* 10:1713–1724
29. **Ohta H, Hattori T (1983)** *Agromonas oligotrophica* gen. nov., sp. nov., a nitrogen-fixing oligotrophic bacterium. *Antonie van Leeuwenhoek* 49:429–446
30. **Radl V, Simões-Araújo JL, Leite J, Passos SR, Martins LM, Xavier GR, Rumjanek NG, Baldani JI, Zilli JE (2017)** *Microvirga vignae* sp. nov., a root nodule symbiotic bacterium isolated from cowpea grown in semi-arid Brazil. *Int J Syst Evol Microbiol* 64:725–730
31. **Richter M, Rosselló-Móra R, Oliver Glöckner F, Peplies J (2016)** JSpeciesWS: a web server for prokaryotic species circumscription based on pairwise genome comparison. *Bioinformatics* 32:929–931
32. **Rzhetsky A, Nei M (1992)** A simple method for estimating and testing minimum evolution trees. *Mol Biol Evol* 9:945–967

33. **Safronova VI, Kuznetsova IG, Sazanova AL, Belimov AA, Andronov EE, Chirak ER, Osledkin YS, Onishchuk OP, Kurchak ON, Shaposhnikov AI, Willems A, Tikhonovich IA (2017)** *Microvirga ossetica* sp. nov., a species of rhizobia isolated from root nodules of the legume species *Vicia alpestris* Steven. *Int J Syst Evol Microbiol* 67:94–100
34. **Saitou N, Nei M (1987)** The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
35. **Sasser M (1990)** *Identification of Bacteria by Gas Chromatography of Cellular Fatty Acids*, MIDI Technical Note 101. Newark, DE: MIDI Inc
36. **Tapase SR, Mawlankar RB, Sundharam SS, Krishnamurthi S, Dastager SG, Kodam KM (2017)** *Microvirga indica* sp. nov., an arsenite-oxidizing Alphaproteobacterium, isolated from metal industry waste soil. *Int J Syst Evol Microbiol* 67:3525–3531
37. **Tindall BJ, Sikorski J, Smibert RA, Krieg NR (2007)** Phenotypic Characterization and the Principles of Comparative Systematics. American Society of Microbiology
38. **Wayne LG, Brenner DJ, Colwell RR, Grimont PAD, Kandler O, Krichevsky MI, Moore LH, Moore WEC, RGE Murray, Stackebrandt E (1987)** Report of the Ad Hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Evol Microbiol* 37:463–464
39. **Weon HY, Kwon SW, Son JA, Jo EH, Kim SJ, Kim YS, Kim BY, Ka JO (2010)** Description of *Microvirga aerophila* sp. nov. and *Microvirga aerilata* sp. nov. isolated from air, reclassification of *Balneimonas flocculans* Takeda et al. 2004 as *Microvirga flocculans* comb. nov. and emended description of the genus *Microvirga*. *Int J Syst Evol Microbiol* 60:2596–2600
40. **Xu P, Li WJ, Tang SK, Zhang YQ, Chen GZ, Chen HH, Xu LH, Jiang CL (2005)** *Naxibacter alkalitolerans* gen. nov., sp. nov., a novel member of the family ‘*Oxalobacteraceae*’ isolated from China. *Int J Syst Evol Microbiol* 55:1149–1153
41. **Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017)** Introducing EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and whole-genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617
42. **Yoon SH, Ha SM, Lim J, Kwon S, Chun J (2017)** A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie van Leeuwenhoek* 110:1281–1286
43. **Zhang BT, Miao JL, Li GY, Cui KY (2004)** Research progress of polar microbial active substances. *Mar Sci* 28:58–63
44. **Zhang H, Yohe T, Huang L, Entwistle S, Wu P, Yang Z, Yin Y (2018)** dbCAN2: a meta server for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 46: 95–101
45. **Zhang J, Song F, Xin YH, Zhang J, Fang C (2009)** *Microvirga guangxiensis* sp. nov., a novel alphaproteobacterium from soil, and emended description of the genus *Microvirga*. *Int J Syst Evol Microbiol* 59:1997–2001
46. **Zhang XJ, Zhang J, Yao Q, Feng GD, Zhu HH (2019)** *Microvirga flavescens* sp. nov., a novel bacterium isolated from forest soil and emended description of the genus *Microvirga*. *Int J Syst Evol Microbiol* 69:667–671

47. Zhang Y, Zhuang JL, Pang HC, Wang YN, Li YY, Zhang JL (2019) *Paenibacillus lutes* sp. nov., isolated from soil. Int J Syst Evol Microbiol 69:2354-2359

Tables

Table 1. Genome-based comparisons of strain 3D7^T and the closely related type strains of the genus *Microvirga*

Strains: 1, 3D7^T (JAGEMM000000000); 2, '*M. brassicacearum*' CDVBN77^T (VCMV000000000); 3, *M. subterranea* DSM 14364^T (QQBB000000000); 4, *M. guangxiensis* 25B^T (jgi.1041447.1.00000); 5, *M. aerophila* DSM 21344^T (QOIO000000000)

Calculation	1	2	3	4	5
Genome size (bp)	4,457,992	5221427	5,147,802	4,721,732	6,003,377
G+C content (%)	63.5	62.3	65.1	61.4	62.1
Contigs	221	88	36	34	184
N50 length (bp) (scaffolds)	431,466	130,073	519,753	407,384	101,852
Number of RNAs	49	51	67	61	61
Protein-coding genes	4321	5244	4885	4520	6022
ANIb (%)	-	77.47	75.27	74.34	74.75
ANIm (%)	-	84.47	83.96	83.89	83.96
Tetra	-	0.978	0.896	0.944	0.936
OrthoANLu (%)	-	79.27	76.94	75.84	76.46
GGDC distance (%)*	-	22.2	20.7	19.9	20.2

*DDH estimate (identities/HSP length formula)

Table 2. Differential characteristics between strain 3D7^T and other closely related species of the genus *Microvirga*

Strains: 1, 3D7^T; 2, '*M. brassicacearum*' CDVBN77^T; 3, *M. subterranea* DSM 14364^T; 4, *M. guangxiensis* 25B^T; 5, *M. aerophila* DSM 21344^T. All data listed were from this study unless indicated. +, positive; -, negative; w, weakly positive

Characteristic	1	2	3	4	5
Isolation source	Soil	Plant	Thermal aquifer	Soil	Air
Growth media	R2A, ASM	R2A, YMA	R2A, Rouf's agar	LB, Rouf's agar, GYM agar	R2A, NA
Colony colour	Light pink	White	Light pink	Light pink	Light pink
Motility	-	+	+	-	-
Growth at:					
Temperature range (°C)	4–32	12-37	25–45	16–42	10–35
Optimum temperature	25–28	28	41	37	30
pH range	7.0–10.0	6-10	6.0–9.0	5.0–9.5	7.0–10.0
Optimum pH	7.0-7.5	7.0	7.0	7.0	7.0
NaCl concentration (%)	0–3	0-1.5	0–1	0–2	0–2
Hydrolysis of:					
Aesculin	+	+	-	-	-
Gelatin	-	-	+	-	-
Starch	-	-	-	+	+
Tyrosine	-	w	w	-	-
Urea	-	-	-	+	-
Oxidase	+	+	-	+	+
Nitrate reduction	+	+	+	+	-
Enzyme activity:					
Alkaline phosphatase	+	+	-	-	-
Valine arylamidase	+	-	-	w	-
Esterase lipase (C8)	+	+	+	+	-
Lipase (C14)	w	-	-	-	-
Trypsin	w	+	w	-	-
Naphthol-AS-BI-phosphohydrolase	+	+	-	-	+

DNA G+C content (mol%)	63.5	62.3 [§]	65.1 [*]	64.3 [#]	62.2 ^{&}
------------------------	------	-------------------	-------------------	-------------------	-----------------------

[§] Data from Jiménez-Gómez et al. (2019)

^{*} Data from Kanso and Patel (2003)

[#] Data from Zhang et al. (2009)

[&] Data from Weon et al. (2010)

Table 3. Fatty acid profiles of strain 3D7^T and the closely related species of the genus *Microvirga*

Strains: 1, 3D7^T; 2, '*M. brassicacearum*' CDVBN77^T; 3, *M. subterranea* DSM 14364^T; 4, *M. guangxiensis* 25B^T; 5, *M. aerophila* DSM 21344^T. Values are percentages of total fatty acids. All tests were carried out using similar methods. -, not detected

Fatty acids (%)	1	2	3	4	5
C _{14:0}	1.4	-	-	0.8	1.1
C _{16:0}	12.2	9.2	6.5	7.9	7.6
C _{17:0}	-	3.8	7.7	0.8	-
C _{17:0} cyclo	7.3	-	-	0.6	2.1
C _{17:1} ω6c	-	-	1.1	-	-
C _{17:1} ω8c	-	-	2.0	-	-
C _{18:0}	1.3	6.1	3.6	6.0	1.0
C _{18:0} 3-OH	1.2	1.7	-	1.9	0.8
C _{18:1} ω7c 11-methyl	3.6	4.2	-	-	-
C _{19:0} cyclo ω8c	21.7	24.3	17.4	11.1	13.8
C _{19:0} 10-methyl	-	-	-	0.6	-
C _{20:2} ω6,9c	-	-	-	-	-
Summed Features*					
Summed feature 2	4.3	2.2	3.5	1.9	4.3
Summed feature 3	6.3	4.5	1.5	2.2	6.9
Summed feature 8	36.2	39.3	53.9	63.6	59.2

*Summed Features are fatty acids that cannot be resolved reliably from another fatty acid using the chromatographic conditions chosen. The MIDI system groups these fatty acids together as one feature with a single percentage of the total. Summed feature 2 comprised C_{14:0} 3-OH and/or iso-C_{16:1} I; summed feature 3 contained C_{16:1} ω7c and/or C_{16:1} ω6c; summed feature 8 contained C_{18:1} ω7c and/or C_{18:1} ω6c

Figures




Figure 1

Neighbor-joining phylogenetic tree based on 16S rRNA gene sequences showing the relationships between strain 3D7T and the closely related species of the genus *Microvirga*. Numbers at nodes represent bootstrap percentages (>50 %) based on 1000 replicates. *Phyllobacterium salinisoli* LLAN61T was chosen as an outgroup. Bar represents 0.01 substitutions per nucleotide position

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMaterials.pdf](#)