

# Soil Biotic and Abiotic Traits as Driven Factors for Site Quality of Araucaria Angustifolia Plantations

Tancredo Souza (✉ [tancredo\\_agro@hotmail.com](mailto:tancredo_agro@hotmail.com))

Universidade de Coimbra <https://orcid.org/0000-0001-8729-5478>

Mário Dobner Jr.

Federal University of Santa Catarina: Universidade Federal de Santa Catarina

Djalma Eugênio Schmitt

Federal University of Santa Catarina: Universidade Federal de Santa Catarina

Lucas Jónatan Rodrigues da Silva

Federal University of Santa Catarina: Universidade Federal de Santa Catarina

Júlia Carina Niemeyer

Federal University of Santa Catarina: Universidade Federal de Santa Catarina

Ketlin Schneider

Federal University of Santa Catarina: Universidade Federal de Santa Catarina

---

## Research

**Keywords:** Cambisols, Endangered tree species, Subtropical ecosystems, Soil organic horizons, Soil food web, Soil trophic structure

**Posted Date:** May 4th, 2021

**DOI:** <https://doi.org/10.21203/rs.3.rs-445199/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Biologia on January 29th, 2022. See the published version at <https://doi.org/10.1007/s11756-021-00988-3>.

# Abstract

**Purpose:** The role of soil biotic and abiotic factors in crucial ecosystem services such as primary production, organic matter dynamics, nutrient cycling, and soil biota community structure in the *Araucaria* ecosystem remains poorly quantified. We aimed to understand how the site quality affects the development of organic horizons, root growth, soil chemical properties, and the entire soil biota community in even aged and monospecific *Araucaria angustifolia* plantations.

**Methods:** We collected soil monoliths to describe layers of organic matter and the complex soil food web into these layers. We determined soil pH, soil moisture, total nitrogen, available P, and total organic carbon into each layer (litter, F-layer, H-layer, and A horizon), the biomass of fine roots, the community structure of soil biota, arbuscular mycorrhizal fungi, and nematodes, as well as the microbial biomass carbon.

**Results:** In the high-quality site, there was significantly higher organic matter formation, nutrient cycling (N and P), root growth, soil moisture, soil biota diversity, arbuscular mycorrhizal fungi, nematodes, and microbial activity evaluated by the microbial biomass carbon compared to the low-quality site.

**Conclusions:** High-quality sites promote the development of organic horizons, root growth on superficial layers that provide plant nutrient release, the A horizon nutrient contents, and the entire soil biota community in monospecific *Araucaria angustifolia* plantations located on humid subtropical Cambisols. This creates a positive plant-soil feedback that maintains soil quality and increases primary production, nutrient cycling, and habitat and food for the soil food web.

## 1. Introduction

It is well documented that biotic and abiotic factors significantly influence above- and belowground ecosystem primary production through positive plant-soil feedback (Bennett and Klironomos 2018). Even after decades of intensive research, it remains unclear and difficult to quantify such crucial ecosystem services in monodominance ecosystems such as *Araucaria angustifolia* (Bert.) O. Kuntze plantations (Marchioro et al. 2020). The role of soil biotic and abiotic factors remain far from being understood, primarily because of a wide range of complex compartments into soil ecosystems and a lack of scientific studies gathered from long-term field experiments (Bowsher et al. 2018; Santos et al. 2018; Souza et al. 2019). Ecosystem services, such as primary production, organic matter dynamics, nutrient cycling, and soil biota activity are among the most important services that promote soil quality and generate positive soil-plant feedback (Tateno et al. 2017). The ecological significance of these services is attributed to their characteristics to promote habitat and energy that enable soil biota and microbiota to create positive soil-plant feedback into the soil ecosystem (Souza and Freitas 2018). For instance, primary production is known to increase litter deposition, thus stimulating organic horizon development past over the years (Santos et al. 2018; Santos-Heredia et al. 2018). This first pathway creates a habitat for soil biota and microbiota; thus, they act by transforming and decomposing the organic matter through mechanisms such as a “priming effect”. Finally, a solid soil food web is created promoting soil biota community

structure, microbial growth, and plant uptake in a way similar as to the soil quality and nutrient content hypothesis described by Souza and Freitas (2018) and Melo et al. (2019). Therefore, the abundance and diversity of soil organisms may, in turn, be regulated by primary production, further modulating plant-soil feedback (Tateno et al. 2017; Bennett and Klironomos 2018).

Despite such positive plant-soil feedback between the soil organic horizon, plants and functionally diverse groups of soil biota, arbuscular mycorrhizal fungi, and soil nematodes, most of the experimental findings on the role of abiotic and biotic factors are often based on only one of these properties, gathered from pot experiments, or often in the absence of adult tree plant species (Bennet and Klironomos 2018; Gebremikael et al. 2016; Santos et al. 2018; Souza et al. 2019; Zhang et al. 2018). Such studies have been used by soil ecologists in soil biology and have increased our understanding of the role of soil biotic and abiotic factors in the processes of organic horizon formation, nutrient cycling, primary production, and soil food web (Ojeda et al. 2018; Yang et al. 2018). However, these findings do not consider the whole potential of the soil biotic and abiotic characteristics as driven factors for site quality and are far from representing the reality in the field. Yet, long-term studies in the presence of adult tree plant species considering these aspects are rare (Rasmussen et al. 2019).

After roughly 100 years of timber exploitation during the 20th century, *A. angustifolia* is currently defined as a critically endangered species (Thomas 2013; Marchioro et al. 2020) and only in-line plantations can be economically used after governmental permission. Besides this historic interest in its timber, and the recognized productive potential (Nutto et al. 2005; Dobner Jr. et al. 2019), the soil bio-chemical factors that influence the growth of *A. angustifolia* are still little known. Moreover, due to the lack of knowledge about the ecology of the *A. angustifolia*, mistakes were made firstly in selecting sites for establishing monospecific stands of this plant species (Breuninger et al. 2008) and secondly in managing them (Seitz 1986; Nutto et al. 2005).

This study aimed to understand the role of *Araucaria* site quality on the development of organic horizons, root growth, soil chemical properties, and the entire soil biota community. To this end, we analyzed soil monoliths, the soil's main chemical properties, soil biota and microbiota community structure, and microbial biomass carbon for two types of *A. angustifolia* plantations (e.g., low- vs. high-quality sites). With this approach, we sought to shed light on how site quality influences the formation of organic matter in humid-subtropical highland soil and the macro- and microbiota that provide ecosystem services into this condition. We hypothesized that sites with high quality are the ones characterized as follows: high biological activity in soil and with high cation exchange capacity resulting in an increased release of available nutrients as well as a concomitant increase in fine roots production and plant growth, thus creating a positive plant-soil feedback.

## 2. Material And Methods

### Species description

*Araucaria angustifolia* belongs to the family Araucariaceae. *A. angustifolia* is named as Araucaria, Paraná pine, Brazilian pine, and candelabra tree (Breuninger et al. 2008). An evergreen endangered tree to 40 m with thick, tough, and triangular-like leaves and a long straight trunk. These leaves are broad at the base. However, they are razor-sharp at the edges and tip (Hoogh and Dietrich 1979). *A. angustifolia* used to cover an area of 233,000 km<sup>2</sup> from Brazilian territory. Nowadays, it has lost 97% of its original area to logging, agriculture and Pinus plantations (Dobner Junior et al. 2019). It is tolerant of most soil types. It grows best in well drained, slightly acidic soils inside subtropical climate (e.g., with abundant rainfall more than 1,200 mm). In Brazil, its seeds (called pinhão) are used as a food resource and medicine for the regional farmers, and it played an important role for the small population of natives in the past.

## Study area location

The study sites are geographically located at the Florestal Gateados Enterprise in Campo Belo do Sul, Santa Catarina, highlands of Southern Brazil (Table 1). The mean annual temperature of the area is + 15°C and receives a mean annual precipitation of 1,750 mm. The experimental area was dominated by soybean in the 80's following a conventional farming system. The soil type in the experimental area was classified as Cambisols (WRB 2006). Köppen's classification defines the climate of the experimental area as humid-subtropical (Cfb) type (Alvares et al. 2013).

Table 1  
General description of the studied sites. with and altitude of 1,135 m above sea level

Site quality	Latitude	Longitude	Altitude (m)	Slope (%)	No. plots	No. soil samples	No. monoliths
Low	S27°51'14"	O50°49'21"	1,200	12	25	100	100
High	S28°01'50"	O50°51'18"	1,135	6	25	100	100

## Reconnaissance survey

A preliminary discussion was held with the managers of the Florestal Gateados Enterprise in order to get general information about the monospecific *A. angustifolia* plantation that covers a total area of 530 ha. Subsequently, a reconnaissance survey was conducted across the monospecific stand to have an overall impression about the study area. At this point, site quality was selected only in terms of dominant height, i.e., according to the accumulated productivity. The relationship between dominant height ( $h_{100}$ ) and age as a measure of site quality (Skovsgaard and Vanclay 2008) was obtained from 460 sample plots (500 m<sup>2</sup>). Stands were selected as closest as possible to 30 years of age, thus representing a well-established stand. Then, two sites adjacent to each other were selected for further study (Table 2). In terms of dominant height, at the age of 30 years, an 8-m difference was verified, which, when translated to mean annual increment (MAI) in volume, represents 14 to 31 m<sup>3</sup> ha<sup>-1</sup> yr<sup>-1</sup>. Both stands were not thinned and thus allowed robust comparisons in terms of growth and yield.

Table 2

Site characteristics including age, site index (SI) stocking (N), quadratic mean diameter at breast height ( $d_g$ ) dominant diameter at breast height ( $d_{100}$ ), dominant height ( $h_{100}$ ), basal area (G), standing volume (V), and mean annual increment (MAI) in volume.

Site quality	Age Years	SI <sup>a</sup>	N trees ha <sup>-1</sup>	$d_g$ cm	$d_{100}$ cm	$h_{100}$ m	G m <sup>2</sup> ha <sup>-1</sup>	V m <sup>3</sup> ha <sup>-1</sup>	MAI m <sup>3</sup> ha <sup>-1</sup> yr <sup>-1</sup>
Low	31	16	2,281	17.7	28.0	14.2	56.6	420.2	13.6
High	30	26	1,488	26.0	40.2	22.0	79.0	922.7	30.9
Amplitude	11– 51		60 – 2,560	11– 50	17– 78	7– 23	4–53	19– 556	-

<sup>a</sup> Site index, the dominant height at an index age of 40 years, according to the classification proposed by Schneider et al. (1992).

## Scheme of sample plot

A systematic sampling approach was implemented to conduct the field study. A total of twenty-five plots, 20 × 25 m were established in each site. The first plot was laid out systematically using a compass, 250 m away from the edge to avoid an edging effect. The transect lines (five transects per site) were made along the centre of each studied site and 50 m away from each other. Soil samples were taken in the centre of the plots at the 0–20 and 20–40 cm soil depths. The samples were homogenized, air-dried, and organic residues were removed manually. Then, soil samples were dried in an oven at 60°C, sifted in a 2-mm mesh sieve, and subjected to analyses. Clay content was 540 and 405 g kg<sup>-1</sup> at the 0–20 and 20–40 cm of soil depths, respectively. Soil pH (H<sub>2</sub>O) was 5.6 to 6.3 at 0–20 cm, and 5.2 to 5.7 at the 20–40 cm soil depth; CEC values at soil pH were 16.4 and 19.3 cmolc kg<sup>-1</sup> at the 0–20 and 20–40 cm soil depths, respectively. Total organic carbon and available P (Mehlich 1) ranged from 20.6 to 14.5 g kg<sup>-1</sup> and from 3.8 to 1.4 mg kg<sup>-1</sup>, at the 0–20 and 20–40 cm soil depths, respectively (EMBRAPA 1997; Tedesco et al. 1995). The inventories were conducted in July and December 2019.

## Data analysis

### Soil organic matter formation

To characterize the litter compartment and layers of organic matter at intermediate stages of decomposition (F-layer and H-layer), four soil monoliths by each studied plot were collected accordingly to Fassbender (1993). Before collecting the soil monoliths, an area of 20 × 20 cm on the soil surface was delimited for separately sampling the litter layer. After that, we extract soil monoliths with the following dimensions 20 × 20 × 20 cm. Next, we wrapped them with plastic film and transported all the monoliths

with minimal disturbance until analysis. During our analysis, the monoliths were dissected into the litter, F-layer, H-layer, and A horizon. We considered the F-layer to be the material composed of partly decomposed litter, the H-layer the material with well-decomposed litter, and the A horizon composed exclusively of mineral material (Toutain 1987). The ratio of organic matter layers was calculated using the following equation:  $ROML_i = dm_i/T_m$ , where  $dm_i$  is the dry mass of each layer (e.g., litter, F-layer, H-layer, and A horizon), and  $T_m$  are both the total dry mass of the soil monolith (20 × 20 × 20 cm) and dry biomass of litter.

## Fine roots

To estimate fine root (diameter: <2 mm) dry biomass, we collected roots from the soil samples of each layer (e.g., litter, F-layer, H-layer, and A horizon) during the monoliths processing described above. Fine roots in these layers were washed using a 0.5-mm nylon mesh bag. We sorted fine roots into living and dead roots based on morphology and condition. Only living roots were considered to estimate dry biomass. Fine roots included both tree and herbaceous species because it was difficult to distinguish between these precisely. Fine root dry biomass (g) was determined after drying the samples for 48 h at 70°C. Samples of each layer (litter, F-layer, H-layer, and A horizon) from monoliths were air-dried and passed through a 2-mm sieve.

## Soil chemical properties

We determined soil pH in a suspension of soil and distilled water (1:2.5 ratio) (Black 1965). Total soil nitrogen and soil organic carbon were estimated according to the methodology described by Okalebo et al. (1993). Phosphorus ( $P_{sbe}$ ) was determined using the Olsen's P protocol (Olsen et al. 1954). The soil moisture was measured by the gravimetric method, where a fresh soil sample was weighed, oven-dried until no further mass loss, and then reweighed (Black 1965).

## Soil fauna

At the end of the winter and summer, we sampled four soil monoliths (20 × 20 × 20 cm) to extract and characterize the soil fauna community per studied plot (e.g., low- and high-productivity sites), and collected the organisms manually using metal clips. They were stored in containers with 70% alcohol until identification as recommended in Tropical Soil Biology and Fertility (Anderson and Ingram 1989). These were later counted and identified under a stereoscopic microscope, at the level of a major taxonomic group. The term "taxonomic group" was used in the soil macroarthropod study, meaning either a Class, as Order or even Family, to comprise a set of individuals with a similar life form. The communities were characterized based on the following parameters: (a) richness and (b) Shannon Diversity Index (H) (Shannon and Weaver 1949). We assessed the frequency of occurrence of each taxonomic group by both studied sites. In addition, we classified the taxonomical groups according to

their functional groups as described by Souza and Freitas (2018). The frequency of occurrence was calculated using the following equation:  $FO_i = n_i/N$ , where  $n_i$  is the number of times an organism was observed, and  $N$  is the total number of organisms observed from each studied ecosystem.

## Arbuscular mycorrhizal fungi and soil nematodes

To sample the spores of arbuscular mycorrhizal fungi and soil nematodes, we sampled undisturbed soil cores ( $n = 4$  per studied plot and 300 g of soil each core), wrapped them, and stored them with minimal disturbance until specimen's extraction as recommended by Souza and Freitas (2018). For AMF extraction, spores and sporocarps from the field were extracted by the wet sieving technique (Gerdemann and Nicolson 1963) followed by sucrose centrifugation (Jenkins 1964). Initially, the extracted spores were examined in water under a dissecting microscope and they were separated based on morphological characteristics. Subsequently, they were mounted in polyvinyl alcohol in lacto-glycerol (PVLG) with and without the addition of Melzer's reagent (Walker et al. 2007). Species identification was based on the descriptions provided by Schenck and Perez (1987), and by consulting the online AMF collection of the Department of Plant Pathology, the University of Agriculture in Szczecin, Poland (<http://www.agro.ar.szczecin.pl/~jblaszkowski/>) and the International Culture Collection of Arbuscular Mycorrhizal Fungi Database—INVAM (<http://invam.caf.wvu.edu>). The AMF communities were characterized based on the following parameters: (a) richness and (b) Shannon Diversity Index (H) (Shannon and Weaver 1949). We assessed the frequency of occurrence of each taxonomic group at both studied sites. For soil nematodes, we used the method described by Buchan et al. (2013), to separate free-living nematodes from soil components (e.g., organic matter and clay). We counted the soil nematodes under a binocular microscope. Next, the soil nematodes were fixed with a 4% hot (70°C) formaldehyde solution. Finally, nematode identification using trophic groups was carried out according to Yeates et al. (1993).

## Microbial biomass carbon

Soil samples were put into pots. They were brought to and maintained at ca. 50% water-filled porosity and incubated at 18°C for 45 days. The amount of distilled water was based on the bulk density and initial moisture content of the soil. Water reposition was calculated weekly using a mass balance of each pot. Four replications from each studied site were sampled after 5, 15, 30, 45, and 60 days of incubation. Microbial biomass carbon ( $C_{mic}$ ) was determined using the fumigation-extraction protocol described by Vance et al. (1987). We divided the soil (20 g of fresh soil per pot) into fumigated and non-fumigated controls. The  $C_{mic}$  was extracted with 40 mL of 0.5 M  $K_2SO_4$  and stored at -18°C until analysis. Organic carbon contents of the extracts were determined by the rapid dichromate oxidation method described by Okalebo et al. (1993).

# Statistical analysis

Before analysis, all the variables were tested for normality (e.g., by Shapiro-Wilk test) and homoscedasticity (e.g., by the Bartlett test), and log transformations were applied to meet both required criteria. To find possible spatial autocorrelation, we used the Moran.I function (Gittleman and Kot 1990). We did not detect any relationship between the variables and the sampling points, indicating spatial independent samples. Soil properties, soil biota, and microbiota community composition, and microbial biomass carbon were analyzed with a non-parametric paired t-test followed by the Monte Carlo test (1000 permutations). The dissimilarities between the site quality (e.g., by Bray-Curtis distance measure) were analyzed using non-metric multidimensional scaling (NMDS), which provided a graphical ordination of the variables that when presenting a measure of stress less than 0.01, indicate an excellent fit of the model (Zuur et al. 2007). It also enables us to reduce the number of the variables used to determine which abiotic or biotic variable explained most of the variation in the productivity sites (Oksanen et al. 2013). All analyses were run using R 3.4.0 statistical software (R Core team 2018).

## 3. Results

The ratio of litter and organic matter layers (e.g., F-layer and H-layer) on soil monoliths was significantly higher in the high-quality site ( $t = 11.54, p < 0.01$ ;  $t = 17.33, p < 0.001$ ;  $t = 21.45, p < 0.001$ , respectively), while the A horizon was significantly higher in the low-quality site ( $t = 20.56, p < 0.001$ ). There were significant differences on total nitrogen ( $t = 21.28, p < 0.001$ ), and P by sodium bicarbonate extraction ( $t = 20.19, p < 0.001$ ) between low- and high- quality sites, reaching the lowest average of total nitrogen and P by sodium bicarbonate extraction ( $P_{sbe}$ ) in the A horizon. For all studied layers (litter, F-layer, H-layer, and A horizon), the total N and  $P_{sbe}$  were higher at high-quality site than at the low-quality site. Differently than observed for the soil layers, at least numerically, we did not find any significant differences between the studied sites on total organic carbon ( $t = 2.07, p = 80.798$ ) by analyzing soil monoliths. Overall, litter presented 206% and 6% more total nitrogen and  $P_{sbe}$  at the high-quality site than at the low-productivity one. For the A horizon, total nitrogen and available  $P_{sbe}$  tended to be higher (135% and 166%) at the high-quality site than at the low- quality one. Fine root dry biomass was significantly higher in litter ( $t = 11.34, p < 0.01$ ) and the A horizon ( $t = 14.82, p < 0.01$ ) at the high-quality site than at the low-productivity one. Whereas, for the H-layer, we found the significant highest fine root dry biomass at low-quality site ( $t = 14.27, p < 0.01$ ). The soil pH was not observably different for the litter at both low- and high-quality sites. There were significant differences between sites for the soil pH of the F-layer ( $t = 8.13, p < 0.05$ ), H-layer ( $t = 8.02, p < 0.05$ ) and A horizon ( $t = 7.45, p < 0.05$ ). For moisture, we found significant differences between low- and high-quality sites in the litter ( $t = 17.38, p < 0.01$ ), H-layer ( $t = 15.37, p < 0.01$ ), and A horizon ( $t = 19.81, p < 0.01$ ) (Table 3).

Table 3

Variation in the soil productivity site on the ratio of organic matter layers (%), total nitrogen ( $\text{g kg}^{-1}$ , b), available P ( $\text{mg kg}^{-1}$ ), total organic carbon ( $\text{g kg}^{-1}$ , d), fine roots dry biomass (g), soil pH (1:2.5 v:v), and soil moisture (%)

Sites <sup>a</sup>	Ratio of organic matter layers (%)	Total nitrogen ( $\text{g kg}^{-1}$ )	Available P ( $\text{mg kg}^{-1}$ )	Total organic carbon ( $\text{g kg}^{-1}$ )	Fine root dry biomass (g)	Soil pH (1:2.5 v:v)	Moisture (%)
Litter							
Low	1.4 ± 0.3	3.01 ± 0.02	10.2 ± 0.5	89.3 ± 0.2	1.9 ± 0.4	5.0 ± 0.5	37.9 ± 5.4
High	3.5 ± 0.5**	9.57 ± 0.09**	15.3 ± 0.4**	90.5 ± 0.1	47.4 ± 2.2**	5.0 ± 0.5	70.2 ± 11.2**
F-layer							
Low	5.0 ± 0.5	2.56 ± 0.22	5.5 ± 0.1	41.4 ± 0.2	11.8 ± 0.9	4.7 ± 0.1	11.0 ± 1.2
High	7.4 ± 0.7**	6.77 ± 0.17**	13.5 ± 0.3**	45.5 ± 0.3*	11.0 ± 0.8	6.6 ± 0.2*	11.0 ± 1.0
H-layer							
Low	19.1 ± 1.2	1.99 ± 0.43	4.3 ± 0.3	39.7 ± 0.4*	44.9 ± 5.4**	4.7 ± 0.3	14.6 ± 1.3
High	36.5 ± 2.3**	3.97 ± 0.19*	10.5 ± 0.2**	35.6 ± 0.2	21.0 ± 4.3	6.3 ± 0.2*	22.9 ± 3.6**
A horizon							
Low	74.5 ± 5.1**	0.99 ± 0.17	3.7 ± 0.2	25.7 ± 0.5	16.5 ± 3.9	4.1 ± 0.1	3.6 ± 0.6
High	52.6 ± 3.7	2.01 ± 0.05*	6.9 ± 0.5*	30.5 ± 0.4*	34.6 ± 0.9**	5.9 ± 0.3*	15.9 ± 0.2**
<sup>a</sup> Litter height (Low: +3.5–0.0 cm; High: +4.2–0.0 cm); F-layer depth (Low: 0.0–2.7 cm; High: 0.0–3.2 cm); H-layer depth (Low: 2.7–16.5 cm; High: 3.2–17.1 cm); A horizon depth (Low: 16.5–20.0 cm; High: 17.1–20.0 cm); *, ** significant at 5% and 1% by paired t-test, respectively.							

Our results about soil fauna revealed the general effects of soil quality on the average fauna abundance, occurrence of functional groups, richness, and diversity. The high-quality site had the highest richness and diversity. In this site, we identified exclusively soil organisms from Blattidae, Scarabaeidae, Spirobolidae, Enchytraeidae, Acaridae, Acerentomidae, Filistatidae, Scutigerae, Forficulidae, and Procampodeidae. The results from functional groups showed a similar pattern for ecosystem engineers, litter transformers, microregulators, and predators, with the high-productivity site having the highest values of soil fauna abundance and the exclusive presence of some families of these groups (Table 4).

Table 4

Soil fauna abundance (of individual monoliths), richness, diversity (Shannon's index), and functional groups at low- and high-productivity sites.

Taxonomic group / Family	Site Quality		Functional groups
	Low	High	
Blattodea / Blattidae	-	2	Ecosystem engineer
Blattodea / Termitidae	1	8**	Ecosystem engineer
Hymenoptera / Formicidae	17**	9	Ecosystem engineer
Coleoptera / Scarabaeidae	-	3	Litter transformers
Diplopoda / Spirobolidae	-	2	Litter transformers
Enchytraeidae	-	17	Litter transformers
Acari / Acaridae	-	1	Microregulator
Collembola / Isotomidae	2	27**	Microregulator
Protura / Acerentomidae	-	3	Microregulator
Araneae / Filistatidae	-	8	Predator
Chilopoda / Scutigerae	-	3	Predator
Dermoptera / Forficulidae	-	5	Predator
Strepsiptera / Myrmecolacidae	2	-	Predator
Diplura / Procamptodeidae	-	3	Herbivore
Larvae of Coleoptera	6	-	Herbivore
Richness	5	13**	-
Diversity (Shannon index)	1.12	2.15**	-
Soil monolith dimensions: 20 × 20 × 20 cm; Functional groups classification was done according to Souza and Freitas (2018); ** Significant at 1% by paired t-test.			

Collectively, we identified eight arbuscular mycorrhizal fungi (AMF) species in all samples. Eight AMF species were found at the high-quality site, and four at the low-quality one. There were differences between the AMF frequencies of occurrence between low- and high-quality sites. Across all the samples, *Funneliformis mosseae* was the most abundant identified AMF species. *Acaulospora morrowiae*, *C. pellucida*, *G. gigantea*, and *R. coralloidea* were AMF species found exclusively at the high-productivity site (Table 5).

Table 5  
Frequency of occurrence (%) of arbuscular mycorrhizal fungi at low- and high-productivity sites.

Arbuscular mycorrhizal fungi species	Site Quality	
	Low	High
<i>Acaulospora morrowiae</i> Spain & N.C. Schenck	-	12.33
<i>Cetraspora pellucida</i> (T.H. Nicolson & N.C. Schenck) Oehl, F.A. Souza & Sieverd.	-	8.03
<i>Claroideoglossum claroideum</i> (N.C. Schenck & G.S. Sm.) C. Walker & A. Schüßler	16.30**	9.63
<i>Funneliformis mosseae</i> (T.H. Nicolson & Gerd.) C. Walker & A. Schüßler	53.60**	25.39
<i>Glomus sp.</i>	14.90*	10.90
<i>Gigaspora gigantea</i> (T.H. Nicolson & Gerd.) Gerd. & Trappe	-	5.34
<i>Racocetra coralloidea</i> (Trappe, Gerd. & I. Ho) Oehl, F.A. Souza & Sieverd.	-	21.33
<i>Rhizophagus intraradices</i> (N.C. Schenck & G.S. Sm.) C. Walker & A. Schüßler	15.20**	7.05
Richness	4	8**
Diversity (Shannon index)	1.19	1.94*
*, ** Significant at 5% and 1% by paired t-test, respectively.		

The abundance and functional composition of nematodes in the low-quality site were dissimilar (36 vs. 76 ind. g<sup>-1</sup> soil) to the high-quality site. In the site with low-quality, we did not find Carnivores and Omnivores. The total abundance and abundances of bacterivores, carnivores, omnivores, and fungivores were significantly higher ( $p < 0.01$ ) at the high-quality site than at the low one. Only, the abundance of herbivores was significantly higher ( $p < 0.01$ ) at the low-quality site than at the high one (Fig. 1A). Microbial biomass carbon showed similar dynamics between low- and high-quality sites. The highest values were at 5 days of incubation, and then they started to decline until reaching a stable line after 45 and 30 days for high- and low-quality sites, respectively. In our study,  $C_{mic}$  was lower at the low-quality site than at the high- one and was significantly different ( $p < 0.001$ ) (Fig. 1B).

According to the NMDS analysis, the soil productivity sites were significantly dissimilar (Fig. 4). The ordination of the soil chemical characteristics (e.g., ratio of A horizon, ratio of H-layer, P contents in A horizon, F-layer, and H-layer, and N content in litter), soil biota abundance (e.g., Formicidae, and Isotomidae), soil microbiota abundance (e.g., *F. mosseae*, and Bacterivores) and microbial biomass carbon in each productivity site had a good fit (stress value = 0.18). Formicidae, the ratio of the A horizon, and *F. mosseae* were highly correlated with the low-productivity site, whereas Isotomidae, Bacterivores, microbial biomass carbon, litter N content, P content in the A horizon, F-layer, and H-layer, and the ratio of H-layer were highly correlated with the high-productivity site. Formicidae, the A horizon, *F. mosseae*,  $P_{A-H}$ ,  $C_{mic}$ , Bacterivores, Isotomidae,  $N_L$ , H-layer,  $P_{F-L}$ , and  $P_{H-L}$  explained 87.2, 79.8, 0.94, 0.90, 0.91, 0.94, 0.94, 0.95, 0.86, 0.82, and 0.76 of the variation in the productivity sites (Fig. 2).

## 4. Discussion

The soil quality of even-aged *A. angustifolia* plantations on the highlands of Southern Brazil impacts the soil biotic and abiotic properties. Our results indicate a promotion in soil organic matter formation (e.g., litter, F- and H-layers), plant nutrient release (e.g., total N and  $P_{sbe}$ ), and fine root production at the litter and A horizon of high-quality sites. This can be explained by the high productivity amplitude observed in *A. angustifolia* plantations in southern Brazil. While high productive sites and stands are available, with mean annual increments  $\sim 25 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$  (Nutto et al. 2005), or even  $> 30 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$  as was the case for the evaluated one, there are also sites and stands where productivity is negligible by Brazilian standards ( $< 10 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$ ). Unfortunately stands with low productivity are way more common than highly productive ones. In low-quality sites, the low cover rate leads to decrease both the soil organic matter, and nutrient cycling by the exposure of soil to wind and rainfall. This is the reason for the low commercial interest in *A. angustifolia* and why many *A. angustifolia* plantations were converted into other land uses. It is important to note that the site quality classification proposed by Schneider et al. (1992) and employed for the classification of the 530 ha of plantations regarded for stand selection, shows that the productivity level of *A. angustifolia* can be even lower than the low-quality stand evaluated in the present study ( $14 \text{ m}^3 \text{ ha}^{-1} \text{ yr}^{-1}$ ). On the other hand, the studied high-quality site had a dominant height at age of 30 years beyond the values given by Schneider et al. (1992), indicating that this site delivered an impressive production even beyond those authors' best stands. Recent results of *A. angustifolia* genetic breeding started in the 1970s by EMBRAPA are now delivering promising perspectives (Silva et al. 2018; Sousa and Aguiar 2012), which will probably be the genetic base of a new plantation wave. Nevertheless, this new enhanced genetic material will only deliver its maximum potential if accompanied by a deeper understanding of site quality. Thus, the abiotic and biotic characterization of the studied sites delivers a wider understanding of which factors are the drivers for the productivity of *A. angustifolia* plantations over a long period (30 years), as discussed in detail as follows. The decrease in soil organic matter formation and nutrient cycling is consistent with the findings of previous studies for cultivated tree species in the highland of southern Brazil (Hoogh and Dietrich 1979; EMBRAPA 1988; Horst et al. 2018; Koehler et al. 2010).

As we originally hypothesized, the quality level of the soil was explained by the biotic and abiotic properties, which were likely caused by an increase in the soil organic matter formation (e.g., by providing habitat), plant nutrient release (e.g., by improving N and P contents on litter, F-layer, and H-layer) as described by Eslamdoust and Sohrabi (2018), which promoted soil biota, arbuscular mycorrhizal fungi, and soil nematode abundance (Gebremikael et al. 2016; Moreira et al. 2007). However, differently than hypothesized, a soil depth of  $\geq 1$  m is not a crucial factor for *A. angustifolia* or, at least, it could be compensated for by other soil properties, such as N and P content and soil biota abundance and activity. Here we presented important evidence on the effects of soil history and quality sites on a variety of compartments into the soil profile in a long-term subtropical *Araucaria* ecosystem. Our results support the evidence that soil quality promotes the formation of layers of organic matter at the intermediate stage of decomposition (F- and H-layer), and nutrient cycling, which may influence the soil food web. Most of the variables analyzed in this study (e.g., layers of organic matter, nutrient cycling, and fine roots production) responded positively to the high-productivity site (Mishra et al. 2019). Productivity level (e.g., high- and low-quality sites) was an important variable to understand all studied variables, indicating that site quality influences the nutrient cycling, fine root production, and soil biota diversity and activity.

Site differences considering soil biodiversity were discussed in previous studies performed around the world (Bennett and Klironomos 2018; Beretta-Blanco et al. 2019). Our study provided evidence that high-productivity sites, when compared to low-productivity sites, had the highest values of organic matter layers, total N,  $P_{sbe}$ , fine root production, and soil moisture. Considering all these variables, we may consider that ecological processes, such as litter deposition, soil organic matter formation, primary production, and nutrient cycling, were dissimilar between the studied sites, indicating robust associations regardless of the soil quality on high-productivity sites and land degradation on low-productivity sites (Bini et al. 2013; Huangfu et al. 2019; Mauda et al. 2018; Roy et al. 2018; Mishra et al. 2019). This indicates that the abundance of soil biota, arbuscular mycorrhizal fungi, and functional-groups of soil nematodes might be playing an important role in high-productivity site sustainability (Gebremikael et al. 2016; Souza et al. 2019). The high-productivity site showed a better-quality residue and, consequently, the cycling of nutrients is faster in this environment, which can influence the productivity of *A. angustifolia*. In addition, the higher N and P contents in the monoliths may be due to the greater accumulation of residues in the high-quality site, since the litter in reforestation of *A. angustifolia* may be higher than the litter in native forests of southern Brazil (Bini et al. 2013). The N content in the araucaria residue is  $13 \text{ g kg}^{-1}$  (Pereira et al. 2013), which can bring about 200 kg of N to the soil per hectare.

Results on the high-productivity site were significant for soil biota abundance (e.g., Formicidae, Isotomidae, and Termitidae), arbuscular mycorrhizal fungi (e.g., *C. claroideum*, *Funneliformis mosseae*, *Glomus* sp., and *Rhizoglosum intraradices*), functional-groups of soil nematodes (e.g., bacterivores, carnivores, fungivores, herbivores, and omnivores), microbial biomass carbon, and ecological indexes (e.g., richness and diversity by soil fauna and AMF). These results emphasize the influence of site quality on soil biodiversity, which in turn affected productivity levels in a subtropical *Araucaria* ecosystem (Gebremikael et al. 2016; Forstall-Sosa et al. 2020; Souza et al. 2019). Essentially, we wanted to

understand how dissimilar sites (e.g., chosen by their productivity levels) promotes the soil food web in a long-term field experiment, considering a monodominance of *Araucaria angustifolia*. A pioneer study considering site quality for *A. angustifolia* was conducted by Hoogh and Dietrich (1979). These authors reported that the best growth performance was obtained in soils where *A. angustifolia* was planted immediately after the clear-cutting of the native forest. To our knowledge, our study is the first report in a subtropical *Araucaria* ecosystem showing the role of nutrient cycling, soil fauna diversity, AMF diversity, an abundance of soil nematodes, and microbial activity in sustain a positive plant-soil feedback as we found in the high-productivity site. The significant differences between the studied sites for soil properties may influence the soil biota community structure and function, consequently altering nutrient cycling and plant productivity (Jones et al. 2019; Romanowicz et al. 2016). Allied to this, it is also important to maintain soil moisture, adequate contents of plant-nutrients, and a constant supply of residues promoted by litter deposition, which is common in the *A. angustifolia* ecosystem.

The results observed in this study revealed that there were significant differences between the studied sites on soil biodiversity. Therefore, we may consider that the soil food web of the study sites is completely different as proposed by Anyango et al. (2020). According to Roy et al. (2018) and Parsons et al. (2020), soil organisms (e.g., here considering macro- and microbiota) may influence soil health, plant growth, and litter palatability. These authors also report that litter palatability may attract a high diversity of soil fauna functional groups (e.g., ecosystem engineers, and decomposers), which in turn may fuel higher trophic levels (e.g., bacterivores, fungivores, omnivores, and predators). These results agree with previous works (Beretta-Blanco et al. 2019; Gebremikael et al. 2016; Mauda et al. 2018; Mishra et al. 2019) that reported high litter deposition and rhizodeposition as the main factors increasing nutrient availability, microbial activity, and abundance of higher trophic levels as observed in the high-productivity site. By changing these two processes in the rhizosphere of *A. angustifolia*, a quality site may be improved over the years, generating high productivity levels because of increased release of plant-available nutrients (total N, and  $P_{sbe}$ ). These conditions would be responsible to promote soil nematode and microbial activity, especially by microbial biomass carbon, as well as microbivorous nematodes (e.g., bacterivores and fungivores (Gebremikael et al. 2016; Impastato and Carrington et al. 2020; Jo et al. 2020). Our hypothesis that a high-productivity site presents high biological activity resulting in an increased release of available nutrients and a concomitant increase in fine roots production in a specific layer, and plant growth, and thus, creating a positive plant-soil feedback was supported. Overall, soil fauna community composition in the high-productivity site was characterized by i) a high abundance of ecosystem engineers (e.g., Termitidae) and predators (e.g., Isotomidae), which give us evidence of sites with a high degree of bioturbation, organic matter decomposition, and biological control (Forstall-Sosa et al. 2020; Melo et al. 2019); ii) a low abundance of symbionts (e.g., e.g., *Claroideiglomus claroideum*, *Funneliformis mosseae*, *Glomus* sp., and *Rhizoglomus intraradices*) which reflect a high content of plant-available nutrients, creating an independence of *A. angustifolia* plants in this site (Souza et al. 2019); and iii) a high abundance of higher trophic levels of soil nematodes, which indicate that soil nematode community are promoting nutrient cycling, plant biomass, net N, and net P (Rosenfield and Müller 2020). These three characteristics created a positive effect in the trophic structure by promoting some important

ecological processes such as nutrient cycling, biological control, mutualism, parasitism, and soil organic matter formation (Barel et al. 2019; Cuassolo et al. 2020; Ge et al. 2019; Jo et al. 2020; Zhang et al. 2018).

## 5. Conclusions

High-quality sites promote the development of organic horizons (e.g., F- and H-layers), root growth on superficial layers which provide plant nutrient (e.g., P and N) release, soil chemical properties of the A horizon (e.g., contents of P and total N) and the entire soil biota community (e.g., abundance, richness, and diversity) in monospecific *Araucaria angustifolia* plantations located at humid subtropical Cambisols. High-quality sites are accompanied by high activity of fauna and microorganisms in the soil that can promote the nutrient cycling process, thus creating positive plant-soil feedback. Future studies could be carried out to assess which abiotic factors are more important in *A. angustifolia* plantations at different crop ages and if microorganisms could be inoculated to seedlings or sites to improve their productivity level. The results of this study contribute to a deeper view of plant-soil feedback influencing site quality that, in turn, may improve the interest in establishing new plantations of this endangered tree species. Besides the commercial and economic motivations, this would be also an important in-situ conservation strategy.

## Declarations

## Acknowledgments

We thank Florestal Gateados for field support and to support data about the history and soil management of the experimental area.

## Author's contributions

We declare that all the authors made substantial contributions to the conception, design, acquisition, analysis, and interpretation of the data. All the authors participate in drafting the article, revising it critically for important intellectual content, and finally, the authors gave final approval of the version to be submitted to the *Forest Ecosystems*.

## Compliance with Ethical Standards

Conflict of Interest: The authors declare that they have no conflict of interest.

## References

1. Alvares CA, Stape JL, Sentelhas PC, Moraes G, Leonardo J, Sparovek G (2013) Köppen's climate classification map for Brazil. *Meteorol Zeitschrift* 22:711-728. <https://doi.org/10.1127/0941->

2. Anderson JN, Ingram JSI (1993) Tropical soil biology and fertility: A handbook of methods. CAB International, Wallingford
3. Anyango JJ, Bautze D, Fiaboe KKM, Lagat ZO, Muriuki AW, Stöckli S, Riedel J, Onyambu GK, Musyoka MW, Karanja EN, Adamtey N (2020) The impact of conventional and organic farming on soil biodiversity conservation: a case study on termites in the long-term farming systems comparison trials in Kenya. BMC Ecology. <https://doi.org/10.1186/s12898-020-00282-x>
4. Barel JM, Kuyper TW, Boer W, Deyn GB (2019) Plant presence reduces root and shoot litter decomposition rates of crops and wild relatives. Plant Soil 438:313- 327. <https://doi.org/10.1007/s11104-019-0398107>
5. Bennett JA, Klironomos J (2018) Mechanisms of plant-soil feedback: interactions among biotic and abiotic drivers. New Phytologist. <https://doi.org/10.1111/nph.15603>
6. Beretta-Blanco A, Pérez O, Carrasco-Letelier L (2019) Soil quality decrease over 13 years of agricultural production. Nutr Cycl Agroecosyst 114:45-55. <https://doi.org/10.1007/s10705-019-09990-3>
7. Bini D, Santos CA, Carmo KB, Kishino N, Andrade G, Zangaro W, Nogueira MA (2013) Effects of land use on soil organic carbon and microbial process associated with soil health in Southern Brazil. European Journal of Soil Biology 55:117-123. <https://doi.org/10.1016/j.ejsobi.2012.12.010>
8. Black CA (1965) Methods of soil analysis, Part 2. In: Black CA (Ed.) Agronomy Monograph No. 9. American Society of Agronomy, Madison, pp 771-1572
9. Bowsher AW, Evans S, Tiemann LK, Friesen ML (2018) Effects of soil nitrogen availability on rhizodeposition in plants: a review. Plant Soil, 423:59-85. <https://doi.org/10.1007/s11104-017-3491-1>
10. Breuninger M, Einig W, Magel E, Cardoso E, Hampp R (2008) Mycorrhiza of Brazil pine (*Araucaria angustifolia* [Bert. O. Ktze.]). Plant Biol 2:4-10. <https://doi.org/10.1055/s-2000-9177>
11. Buchan D, Gebremikael MT, Ameloot N, Sleutel S, De Neve S (2013) The effect of free-living nematodes on nitrogen mineralisation in undisturbed and disturbed soil cores. Soil Biology & Biochemistry 60:142-155. <https://doi.org/10.1016/j.soilbio.2013.01.022>
12. Cuassolo F, Villanueva VD, Modenutti B (2020) Litter decomposition of the invasive *Potentilla anserina* in an invaded and non-invaded freshwater environment of North Patagonia. Biol Invasions, 22:1055-1065. <https://doi.org/10.1007/s10530-019-02155-x>
13. Dobner Jr M, Paixão CA, Costa EA, Finger CAG (2019) Effect of site and competition on diameter growth of *Araucaria angustifolia*. Floresta 49(4):717-724. <https://doi.org/10.5380/rev.v49i4.58161>
14. EMBRAPA (1988): Empresa Brasileira de Pesquisa Agropecuária. Centro Nacional de Pesquisas Florestais, CNPF: Zoneamento ecológico para plantios florestais no estado de Santa Catarina. Curitiba, 113 p.
15. EMBRAPA (1997) Centro Nacional de Pesquisa em solos. Manual de métodos de análise de solo. Rio de Janeiro, 212 p.

16. Eslamdoust J, Sohrabi H (2018) Carbon storage in biomass, litter, and soil of different native and introduced fast-growing tree plantations in the South Caspian Sea. *J For Res* 29(2):449-457. <https://doi.org/10.1007/s11676-017-0468-5>
17. Fassbender HW (1993) Modelos edafologicos de sistemas agroforestales. 2 ed. Centro Agronómico Tropical de Investigación y Enseñanza, Turrialba.
18. Forstall-Sosa KS, Souza TAF, Lucena EO, Silva SAI, Ferreira JTA, Silva TN, Santos D, Niemeyer JC (2020) Soil macroarthropod community and soil biological quality index in a green manure farming system of the Brazilian semi-arid. *Biologia* In press. <https://doi.org/10.2478/s11756-020-00602-y>
19. Ge T, Luo Y, He X (2019) Quantitative and mechanistic insights into the key process in the rhizodeposited carbon stabilization, transformation and utilization of carbon, nitrogen, and phosphorus in paddy soils. *Plant Soil* 445:1-5. <https://doi.org/10.1007/s11104-019-04347-9>
20. Gebremikael MT, Steel H, Buchan D, Bert W, Neve S (2016) Nematodes enhance plant growth and nutrient uptake under C and N-rich conditions. *Scientific Reports* <https://doi.org/10.1038/srep32862>
21. Gerdemann JW, Nicolson TH (1963) Spores of mycorrhizal *Endogone* species extracted from soil by wet sieving and decanting. *Transactions of the British Mycological Society* [https://doi.org/10.1016/S0007-1536\(63\)80079-0](https://doi.org/10.1016/S0007-1536(63)80079-0)
22. Gittleman JL, Kot M (1990) Adaptation: statistics and a null model for estimating phylogenetic effects. *Syst Zool* 39:227-241. <https://doi.org/10.2307/2992183>
23. Hoogh RJ, Dietrich AB (1979) Relação crescimento-sítio de *Araucaria angustifolia* (Bert) O. Ktze. em povoamentos plantados. *Silvicultura. Anais do 3º Congresso Florestal Brasileiro* 2(14):34-40.
24. Horst TZ, Dalmolin RSD, Ten Caten A, Moura-Bueno JM, Cancian LC, Pedron FA, Schenato RB (2018) Edaphic and Topographic factors and their relationship with dendrometric variation of *Pinus taeda* L. in a high-altitude subtropical climate. *Rev Bras Ciênc Solo* 42:1-16. <https://doi.org/10.1590/18069657rbcs20180023>
25. Huangfu C, Hui D, Qi X, Li K (2019) Plant interaction modulate root litter decomposition and negative plant-soil feedback with an invasive plant. *Plant Soil*, 437:179-194. <https://doi.org/10.1007/s11104-019-03973-7>
26. Impastato CJ, Carrington ME (2020) Effects of plant species and soil history on root morphology, arbuscular mycorrhizal colonization of roots, and biomass in four tallgrass prairie species. *Plant Ecol* <https://doi.org/10.1007/s11258-019-00997-y>
27. Jenkins W (1964) A rapid centrifugal-flotation technique for separating nematodes from soil. *Plant disease reporter* 48:692.
28. Jo I, Friedley JD, Frank DA (2020) Rapid leaf litter decomposition of deciduous understory shrubs and lianas mediated by mesofauna. *Plant Ecol*. <https://doi.org/10.1007/s11258-019-00992-3>
29. Jones DL, Cooledge EC, Hoyle FC, Griffiths RI, Murphy DV (2019) pH and exchangeable aluminum are major regulators of microbial energy flow and carbon use efficiency in soil microbial communities. *Soil Biology and Biochemistry* 138. <https://doi.org/10.1016/j.soilbio.2019.107584>

30. Marchioro CA, Santos KL, Siminski A (2020) Present and future of the critically endangered *Araucaria angustifolia* due to climate change and habitat loss. *Forestry*  
<https://doi.org/10.1093/forestry/cpz066>
31. Mauda EV, Joseph GS, Seymour CL, Munyai TC, Foord SH (2018) Changes in landuse alter ant diversity, assemblage composition and dominant functional groups in African savannas. *Biodivers Conserv* 27:947-965. <https://doi.org/10.1007/s10531-017-1474-x>
32. Melo LN, Souza TAF, Santos D (2019) Cover crop farming system affect macroarthropods community diversity of Caatinga, Brazil. *Biologia* <https://doi.org/10.2478/s11756-019-00272-5>
33. Mishra G, Marzaioli R, Giri K, Pandey S (2019) Soil quality assessment across different stands in tropical moist deciduous forests of Nagaland, India. *J For Res* 30(4):1479-1485.  
<https://doi.org/10.1007/s11676-018-0720-8>
34. Moreira M, Baretta D, Tsai SM, Gomes-da-Costa SM, Cardoso EJBN (2007) Biodiversity and distribution of arbuscular mycorrhizal fungi in *Araucaria angustifolia* forest. *Sci Agri* 64:396-399.  
<https://doi.org/10.1590/S0103-90162007000400010>
35. Nutto L, Spathelf P, Rogers R (2005) Managing diameter growth and natural pruning of Parana pine, *Araucaria angustifolia* (Bert.) O Ktze., to produce high value timber. *Annals of Forest Science* 62:163-173. <https://doi.org/10.1051/forest:2005008>
36. Ojeda JJ, Caviglia OP, Agnusdei MG (2018) Vertical distribution of root biomass and soil carbon stocks in forage cropping systems. *Plant Soil* 423:175-191. <https://doi.org/10.1007/s11104-017-3502-8>
37. Okalebo JR, Gathia KW, Woomer PL (1993) Laboratory methods of plant and soil analysis: A working manual. Technical bulletin No 1, Soil Science Society East Africa, Nairobi
38. Oksanen J et al (2013) Vegan: community ecology package. <http://CRAN.R-project.org/package=vegan>.
39. Olsen S, Cole C, Watanabe F, Dean L (1954) Estimation of available phosphorus in soils by extraction with sodium bicarbonate (USDA, circ. 939).
40. Parsons SE, Kermer LM, Frank SD (2020) Effects of native and exotic congeners on diversity of invertebrate natural enemies, available spider biomass, and pest control services in residential landscapes. *Biodiversity and Conservation* 29:1241-1262. <https://doi.org/10.1007/s10531-020-01932-8>
41. Pereira JM, Baretta D, Bini D, Vasconcellos RLF, Cardoso EJBN (2013) Relationships between microbial activity and soil physical and chemical properties in native and reforested *Araucaria angustifolia* forests in the state of São Paulo, Brazil. *Rev Bras Ciênc Solo* 37:572-586.  
<https://doi.org/10.1590/S0100-06832013000300003>
42. R Core Team (2018) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria <https://www.R-project.org/>
43. Rasmussen PU, Bennet AE, Tack AJM (2019). The impact of elevated temperature and drought on the ecology and evolution of plant-soil microbe interactions. *Journal of Ecology*.

<https://doi.org/10.1111/1365-2745.13292>

44. Romanowicz KJ, Freedman ZB, Upchurch RA, Argiroff WA, Zak DR (2016) Active microorganisms in forest soil differ from the total community yet are shaped by the same environmental factors: the influence of pH and soil moisture. *FEMS Microbiology Ecology* 92.  
<https://doi.org/10.1093/femsec/fiw149>
45. Rosenfield MF, Müller SC (2020) Plant traits rather than species richness explain ecological processes in subtropical forests. *Ecosystems* 23:52-66. <https://doi.org/10.1007/s10021-019-00386-6>
46. Roy S, Roy MM, Jaiswal AK, Baitha A (2018) Soil arthropods in maintaining soil health: Thrust areas for sugarcane production systems. *Sugar Tech* <https://doi.org/10.1007/s12355-018-0591-5>
47. Santos FM, Balieiro FC, Fontes MA, Chaer GM (2018) Understanding the enhanced litter decomposition of mixed-species plantations of *Eucalyptus* and *Acacia mangium*. *Plant Soil* 423:141-155. <https://doi.org/10.1007/s11104-017-3491-7>
48. Santos-Heredia C, Andresen E, Zárate DA, Escobar F (2018) Dung beetles and their ecological functions in three agroforestry systems in the Lacandona rainforest of Mexico. *Biodivers Conserv* 27: 2379-2394. <https://doi.org/10.1007/s10531-018-1542-x>
49. Schenck NC, Perez Y (1987) Manual for the identification of VA mycorrhizal fungi, Second edn. International Culture Collection of VA Mycorrhizal Fungi (INVAM), University of Florida, Gainesville
50. Schneider PR, Finger CAG, Hoppe JM (1992) Produção da *Araucaria angustifolia* (Bert.) O. Ktze. na região do Planalto Médio do Estado do Rio Grande do Sul. *Ciência Florestal* 2(1):99-118.  
<https://doi.org/10.5902/19805098278>
51. Seitz RA (1986) Erste Hinweise für die waldbauliche Behandlung von Araukarienwäldern. *Annales des Sciences Forestières* 43: 327-338.
52. Shannon CE, Weaver W (1949) The mathematical theory of communication. University of Illinois Press, Urbana
53. Silva JR, Santos W, Moraes MLT, Shimizu JY, Sousa VA, Aguiar AV (2018) Seleção de procedências de *Araucaria angustifolia* (Bert.) O. Kuntze para produção de madeira e pinhão. *Scientia Forestalis* 46(120):519-531. <https://doi.org/10.18671/scifor.v46n120.01>
54. Skovsgaard JP, Vanclay JK (2008) Forest site productivity: A review of the evolution of dendrometric concepts for even-aged stands. *Forestry* 81(1):13-30. <https://doi.org/10.1093/forestry/cpm041>
55. Sousa VA, Aguiar AV (2012) Programa de melhoramento genético de araucária da Embrapa Florestas: situação atual e perspectivas. *EMBRAPA Florestas, Documentos* 237, 40 p.
56. Souza TAF, Freitas H (2018) Long-Term Effects of Fertilization on Soil Organism Diversity. In: Gaba S, Smith B, Lichtfouse E (eds) *Sustainable Agriculture Reviews* Springer, Cham, 28, pp 211-247.  
[https://doi.org/10.1007/978-3-319-90309-5\\_7](https://doi.org/10.1007/978-3-319-90309-5_7)
57. Souza TAF, Santos D, Andrade LA, Freitas H (2019) Plant-soil feedback of two legume species in semi-arid Brazil. *Brazilian Journal of Microbiology*. <https://doi.org/10.1007/s42770-019-00125-y>

58. Tateno R, Taniguchi T, Zhang J, Shi W-Y, Zhang J-G, Du S, Yamanaka N (2017) Net primary production, nitrogen cycling, biomass allocation, and resource use efficiency along a topographical soil water and nitrogen gradient in a semi-arid forest near an arid boundary. *Plant Soil* 420:209-222. <https://doi.org/10.1007/s11104-017-3390-7>
59. Tedesco MJ, Gianello C, Bissani C, Bohnen H, Volkweiss SJ (1995) *Análise de solo, plantas e outros materiais*. Porto Alegre: UFRGS/FA/DS.
60. Thomas P (2013) *Araucaria angustifolia*. The IUCN Red List of Threatened Species 2013: e.T32975A2829141. <https://dx.doi.org/10.2305/IUCN.UK.2013-1.RLTS.T32975A2829141.en>
61. Toutain F (1987) Les litières: siège de systems interactifs et moteur de ce interactions. *Revue du Écologie et Biologie du Sol* 24:231-242
62. Vance FD, Brookes PC, Jenkinson DS (1987) An extraction method for measuring soil microbial biomass C. *Soil Biol Biochem* 19(6):703-707.
63. Walker C, Vestberg M, Demircik F, Stockinger H, Saito M, Sawari H, Nishmura I, Schüßler A (2007) Molecular phylogeny and new taxa in the Archaeosporales (Glomeromycota): *Ambispora fennica* gen. sp. nov., Ambisporaceae fam. nov., and emendation of *Archaeospora* and Archaeosporaceae. *Mycol Res* 111:137-153. <https://doi.org/10.1016/j.mycres.2006.11.008>
64. WRB - IUSS Working Group (2006) World Reference Base for Soil. World Soil Resources Reports. Rome, FAO
65. Yang B, Zhang W, Xu H, Wang S, Xu X, Fan H, Chen HYH, Ruan H (2018) Effects of soil fauna on leaf litter decomposition under different land uses in eastern coast of China. *J For Res* 29(4):973-982. <https://10.1007/s11676-017-0521-5>
66. Yeates GW, Bongers T, Goede RGM, Freckman DW, Georgieva SS (1993) Feeding habits in soil nematode families and genera – an outline for soil ecologists. *Journal of Nematology* 25: 315-331
67. Zhang W, Wang C, Lu T, Zheng Y (2018) Cooperation between arbuscular mycorrhizal fungi and earthworms promotes the physiological adaptation of maize under a high salt stress. *Plant Soil* 423:125-140. <https://doi.org/10.1007/s11104-017-3481-9>
68. Zuur AF, Ieno EN, Smith GM (2007) Principal coordinate analysis and non-metric multidimensional scaling. In: *Analysing ecological data. Statistics for Biology and Health*. Springer Science, NY, USA, pp 259–264. [https://doi.org/10.1007/978-0-387-45972-1\\_15](https://doi.org/10.1007/978-0-387-45972-1_15)

## Figures





Figure 1

Mean nematode abundance of each trophic group (A) and microbial biomass carbon (B) at low- and high-quality sites



**Figure 2**

Site quality dissimilarities base on soil chemical characteristics (soil organic matter layers, and nutrient cycling), soil biota and microbiota abundance, microbial C biomass (Cmic) plotted as non-metric multidimensional scaling (NMDS) of the dataset from low- and high-quality sites. The quality of sites is represented by the following symbols: low = gray triangles; and high = dark triangles. Vector length and direction represent only probability values less than 0.01 ( $p < 0.01$ ). A horizon (A-H) = Ratio of A horizon (%); L = Litter; F-L and H-L = F- and H-layer