

**Distinct role of 5'UTR sequences in dendritic trafficking of BDNF mRNA:
additional mechanisms for the BDNF splice variants spatial code.**

Andrea COLLIVA and Enrico TONGIORGI *

Department of Life Sciences, University of Trieste, 34127 Trieste, Italy.

*Corresponding author: Enrico Tongiorgi

Department of Life Sciences (Q building), University of Trieste

Via Licio Giorgieri, 5 - 34127 Trieste, Italy

Tel: +39 040 558 8724 - E-mail: tongi@units.it

Fax: +39 040 558 2443

Abstract

The neurotrophin Brain-derived neurotrophic factor (BDNF) is encoded by multiple bipartite transcripts. Each BDNF transcript is composed by one out of eleven alternatively spliced exons containing the 5'untranslated region (UTR), and one common exon encompassing the coding sequence (CDS) and the 3'UTR with two variants (short and long). In neurons, BDNF mRNA variants have a distinct subcellular distribution, constituting a "spatial code", with exon 1, 3, 5, 7 and 8 located in neuronal somata, exon 4 extending into proximal dendrites, and exon 2 and 6 reaching distal dendrites. We previously showed that the CDS encodes constitutive dendritic targeting signals (DTS) and that both the 3'UTR-short and the 3'UTR-long contain activity-dependent DTS. However, the role of individual 5'UTR exons in mRNA sorting remains unclear. Here, we tested the ability of each different BDNF 5'UTRs to affect the subcellular localization of the green fluorescent protein (GFP) reporter mRNA. We found that exon 2 splicing isoforms (2a, 2b, and 2c) induced a constitutive dendritic targeting of the GFP reporter mRNA towards distal dendritic segments. The other isoforms did not affect GFP-mRNA dendritic trafficking. Through a bioinformatic analysis, we identified five unique cis-elements in exon 2a, 2b, and 2c which might contribute to building a DTS. This study provides additional information on the mechanism regulating the cellular sorting of BDNF mRNA variants.

Keywords: Neurotrophic Factors; BDNF mRNA; RNA Localization; Neuronal Dendrites; Transport Mechanism, Dendritic Targeting Elements, Fragile-X protein family, hnRNPs, stress granules, P-bodies.

Introduction

The neurotrophin Brain-derived neurotrophic factor (BDNF) is a key plasticity factor and its local translation at synapses is required for long term potentiation (LTP) (1). BDNF is a highly conserved gene with a complex structure. In rodents, a common coding sequence (CDS) is spliced to one out of eleven upstream different 5'UTR alternative exons. Moreover, the 3'UTR contains two different polyadenylation sites which produce either a long or a short version of the 3'UTR, raising the resultant transcripts to a total of 22 variants (2). Following a series of *in vivo* (3–5) and *in vitro* studies (3,6–8) we and other groups have shown that in hippocampal pyramidal neurons, BDNF transcripts containing either exon 1, 3, 5, 7, or 8 are preferentially located in the soma, while transcripts with exon 4 extend into the proximal dendrites, and transcripts with exon 2 or 6 reach the distal dendritic compartment in an activity-dependent manner. Interestingly, in dentate gyrus (DG) hippocampal neurons *in vivo*, also the exon 3 transcript is targeted to the distal dendrites following *status epilepticus* (5). Based on these studies, we proposed that the differential subcellular distribution of BDNF transcripts endows neurons with a “spatial code” for a segregated expression of BDNF governed by the transcripts localized in the different subcellular compartments (3,9). Notably, variations in the local availability of the different BDNF transcripts in different cellular districts induce spatially restricted structural changes in the dendritic arbor (6,10,11).

The quest for the mechanisms regulating BDNF mRNA transport in dendrites has led to the discovery that the BDNF coding sequence (CDS) encodes a signal for constitutive dendritic targeting that requires the RNA-binding protein Translin (3). However, the localization of BDNF mRNA in dendrites or its restriction to the soma depends on the combination of the CDS with the different 5'UTR sequences. The alternatively spliced 5'UTR exons act as selectivity signals that modify the constitutive dendritic targeting mediated by the coding sequence (CDS) of BDNF. Indeed, exon 1 and exon 4 cause the retaining of BDNF mRNA to the first 30% of the dendrite, while exons 2a, 2b, 2c and 6 allow its transport into the distal dendritic compartment (3,6). In addition, both short and long 3' UTRs, display dendritic targeting in response to different stimuli: BDNF mRNAs with short 3' UTR respond *in vitro* to depolarization and NT3, or to seizures *in vivo*, and require CPEB-1, -2 and ELAV-2, -4 (8,12). Instead, dendritic targeting of the long 3'UTR is triggered by neuronal activity or BDNF, and requires CPEB-1, ELAV-1, -3, -4, FXR2, and FMR1 (8,12).

Despite these findings, one fundamental question remained unanswered: do the different 5'UTR splice variants contain targeting signals able to induce the sorting of BDNF mRNA to dendrites? The present study builds upon the above-cited works to evaluate how the different BDNF 5'UTR affect the subcellular localization of the green fluorescent protein (GFP) reporter mRNA, and identifies through a bioinformatic analysis the cis-acting elements present in the different BDNF 5'UTR variants.

Results and Discussion

To investigate whether the eleven 5'UTR splicing isoforms of rat BDNF encode any constitutive or activity-dependent targeting signals (DTS), we cloned each 5'UTR region upstream the GFP gene, used as reporter transcript (for details see: Materials and methods). Each chimaeric construct was transfected separately in cultured pyramidal hippocampal neurons at 6 days in vitro (DIV6) and then processed for *in situ* hybridization and densitometric analysis at DIV7. Figure 1A shows a representative *in situ* hybridization obtained using an antisense probe against GFP. The central panel of figure 1A shows the result of the application of the “Straighten” plug-in of ImageJ on the neuron shown at the top, leading to a linearized dendrite which was used for the densitometric analysis. The red line represents the region of interest in which the mean gray value was measured to evaluate the mRNA levels in the different compartments. The specificity of the *in situ* hybridization labeling was determined by using a sense probe for the GFP coding sequence on transfected neurons or a GFP antisense probe in non-transfected neurons. In both cases, the *in situ* signal obtained is indistinguishable from the background (Fig. 1A, bottom panel).

In agreement with our previous studies (3,6), *in situ* labeling for the GFP mRNA was found only in the soma and proximal dendrites of transfected neurons, as confirmed by densitometric analysis (Fig. 1B, bottom panel; GFP ctrl). Gray bars represent the plot of the mean gray values of control (=untreated) neurons, binned in 10 μm compartments starting from soma up to the first 100 μm of the apical dendrite. In parallel cultures, GFP-transfected neurons were treated for 3 hours with a 10 mM KCl solution to assess if GFP mRNA is targeted to dendrites in response to enhanced neuronal activity. Densitometric analysis on

KCl-treated transfected neurons confirmed that GFP RNA distribution was unaffected by KCl-induced depolarization (Fig. 1B, white bars; GFP KCl).

Using the same approach, we evaluated the distribution of chimaeric BDNF 5'exons-GFP transcripts in untreated or depolarized neurons (row data are provided in *Supplementary File#1*). Exon 1 and exon 4, which were previously shown to inhibit the constitutive dendritic targeting of the BDNF CDS (3), did not alter the distribution of the GFP reporter mRNA, displaying a distribution similar to GFP mRNA alone (Fig. 1C and Fig. 1D). Exon 2 and 6 were previously demonstrated to be permissive for BDNF CDS-mediated dendritic targeting (3,6). Here, we found that only exon 2 variants induced a significant dendritic targeting of the GFP reporter transcript (Fig. 2A, B and C; * = $p < 0.05$). These three exon 2 variants resulted to encode for a constitutively active DTS, because their dendritic localization was not affected by KCl-induced depolarization (Fig. 2A, B and C, white bars). Instead, exon 6 did not affect the GFP chimaeric transcript localization with respect to GFP alone, neither in unstimulated nor in depolarized conditions (Fig. 2D). We then evaluated the localization of the other BDNF 5'UTR splicing variants that were previously less characterized regarding their contribution to BDNF mRNA sorting. Exon 3-GFP chimaeric mRNA distribution in unstimulated condition was not significantly different with respect to GFP alone (Fig. 3A, gray bars) but the depolarization induced a weak, not statistically significant, transport into dendrites (Fig. 3A, white bars). Exon 5 and exon 7-GFP displayed a distribution similar to the GFP reporter transcript, in both conditions analyzed (Fig. 3B and C respectively). Exon 8-GFP resulted to be less abundant in dendrites of untreated neurons compared to the GFP transcript, especially in the first 40 μm (Fig. 3D, gray bars, exon 8-GFP CTRL; * $p < 0.05$). KCl depolarization significantly increased this transcript with respect to the untreated conditions in the soma, but not in dendrites (Fig. 3D, white bars, exon 8-GFP KCl; #, $p < 0.05$).

To identify the RBP binding sites present in each BDNF 5'splicing isoform sequences, we performed a bioinformatic analysis on each BDNF 5'UTR variant (FASTA sequences in *Supplementary File#2*) using the RBPDB database (<http://rbpdb.cabr.utoronto.ca>) containing 757 different RNA recognition motifs, and the RBPmap database (<http://rbpmap.technion.ac.il>) containing 114 human/mouse and 51 *Drosophila melanogaster* experimentally defined motifs. Each BDNF variant showed numerous putative RBP recognition sites (Table 2; Full details of recognition sequences and p values

are reported in *Supplementary File#3*). Searching for motifs that could account for the constitutive DTS of exons 2a, 2b, and 2c, we focused on cis-elements common to all three variants but absent in the other BDNF 5'exons. We identified five sites unique to exon 2a, 2b and 2c, which are recognized by the RBPs: Heterogeneous Nuclear Ribonucleoprotein A1 (HNRNPA1), Poly(A) Binding Protein Cytoplasmic 3 (PABPC3), Small Nuclear Ribonucleoprotein U1 Subunit 70 (SNRNP70), Zinc Finger CCCH-Type Containing 10 (ZC3H10), Zinc Finger CCHC-Type And RNA Binding Motif Containing 1 (ZCRB1) (highlighted in green in Table 2). All five recognition elements occur only once in each exon 2 sequence. Even if the algorithm indicated that the recognition motif for HNRNPA1 was present twice, given the large overlap of the two sequences, we considered them as a single motif. HNRNPA1 was previously identified as a component of neuronal mRNA-transporting granules by pull-down experiments with anti-Kinesin5 (KIF5) immunoprecipitating antibodies, together with several other RBPs involved in mRNA trafficking and translational regulation (13). Therefore, it represents a strong candidate for being involved in the constitutive DTS for exons 2a, 2b, and 2c. Interestingly, in all exon 2 variants the recognition sequence for HNRNPA1 largely overlaps with the binding sequence for HNRNPA2/B1, another key factor in mRNA trafficking and translational regulation (14). This recognition sequence is located in the central position of the RNA molecule and is quite distant from other regions that are characterized by long stem-loop structures containing recognition sites for other regulatory proteins (Fig. 4). HNRNPA2/B1 can form multimers when bound to the scaffolding protein tumor-overexpressed gene (TOG), which in turn, binds to multiple HNRNPA2 molecules (14). Interestingly, we found one recognition site for HNRNPA2/B1 not only in all three exon 2 variants but also in exon 6, the other prominent BDNF mRNA with dendritic localization (3,15) and exon 3, which we found to be enriched in the dendrites of hippocampal DG granule cells following *status epilepticus* (5) (Table 2, highlighted in orange). Since, another recognition site for HNRNPA2/B1 is also present in the CDS of BDNF (*data not shown*), a possible cooperative effect between multiple HNRNPA2/B1 molecules and HNRNPA1 is conceivable.

The other four unique RBP recognition sites present in exon 2 variants are likely involved in mRNA dendritic targeting and/or translational repression. PABP3 is a less studied member of the poly(A)-binding protein family with a slightly lower affinity to poly(A) as compared to the best-characterized isoform PABP1 and belongs to the mRNA 3'End Processing Factors required for the process of polyadenylation involved in mRNA translation (16). As PABP1

was shown to regulate the nuclear export of mRNAs towards the cytoplasm (17), and formation of P-bodies (18) it is plausible that also PABP3 may play a role either in cytoplasmic mRNA trafficking or translational regulation, or both. SNRNP70 belongs to the family of small nuclear nucleoproteins that bind to the spliceosomal U1 RNA, forming the core of the spliceosome. SNRNP70 is essential for recognition of the pre-mRNA 5' splice-site and the subsequent spliceosome assembly. It has been shown that this protein is able to inhibit polyadenylation via direct interaction with the poly(A)-polymerase (PAP) (19). Interestingly, recent studies have shown that SNRNP70 participates in the formation of detergent-insoluble aggregates in the brain of patients affected by Alzheimer's disease together with other well-characterized RNA-binding proteins such as TAR DNA-binding protein 43 (TDP-43) and fused in sarcoma (FUS), for which we found binding sites in exon 3 and exon 6, but not in exon 2 variants (20). ZC3H10 is a CCCH zinc finger protein and belongs to a family of RNA-binding proteins that function as RNA metabolism regulators (21). Its role in mRNA trafficking is unclear.

Another interesting information that we obtained from the bioinformatic analysis is related to the finding of recognition sequences for FXR1 in exon 1, and FXR2 and FMRP in exon 4 and 6. We have previously shown that these proteins provide signals for somatic retention of BDNF mRNA both *in vitro* and *in vivo*, and this retention signal is released by electrical activity, or BDNF application (12,22). Another observation pertains to the presence in exons 1, 2a,b,c, 3, and 5 of recognition sites for CPEB2 and CPEB4 which, in this context, do not appear generating any DTS. We previously showed that these recognition sites are also present in both short and long 3' UTRs where they participate in activity-dependent dendritic targeting of BDNF mRNA (8,12). The CPEB proteins are involved in multiple functions and their role is depending upon the reciprocal distance of their recognition sites and upon the distance of each site from the 3' end (8). Regarding the BDNF 5'UTRs, we favour the hypothesis that CPEB2-CPEB4 might be involved in the formation of P-bodies and stress granules together with another prominent stress granule protein, namely TIA1 (23), whose recognition site is also present in exons 1, 2a,b,c, 3, and 5. Of note, the recognition sites for these RBPs are clustered within one distinctive stem-loop structure that is present in all exon 2 variants (Fig. 4). This stem-loop structure additionally harbours the recognition sites for HNRNPC/CL1 and HUR (also known as ELAV-like protein 1; see Fig.4) that are also involved in the formation of P-bodies and stress granules (24) and are part of the 3'UTR signals involved in the dendritic trafficking of BDNF mRNA (12).

Our bioinformatic analysis of the BDNF 5' sequences identified only a few putative binding sites for RNA binding proteins (RBPs) involved in mRNA trafficking. This is in agreement with the current literature, reporting that most of the cis-elements found in the 5'UTRs are involved in mRNA alternative splicing, stability, and translation, while the regulatory sequences for mRNA trafficking are preferentially located in the CDS or in the 3'UTR (25,26). However, some important exceptions to this rule were reported in the literature. For instance, the 5'UTR of the BC1 untranslated RNA encodes a DTS organized in a stem-loop structure bound by the HNRNPA2 protein (27). In conclusion, following a systematic analysis of the different BDNF 5'UTR variants, we can exclude an active role for exon 1, 3, 4, 5, 6, 7, and 8 in dendritic targeting of BDNF mRNA. In addition, we provide evidence that exon 2 variants encode a constitutively active dendritic targeting, possibly involving an HNRNPA1 and HNRNPA2/B1-mediated interaction between 5'UTRs and CDS of BDNF to regulate BDNF mRNA dendritic trafficking.

Materials and Methods

Animal treatment. Animals were treated according to the institutional guidelines in compliance with national laws (D.lgs. 26/2014, art.1/04), the European Council Directive 86/609, and NIH Guide for the care and use of laboratory animals. The present study has been carried out under project authorization N. NO88TON18 from the Italian Ministry of Health. Animals were housed in groups under standard conditions (12:12 h light/dark cycle, ambient temperature 23°C, access to food and water *ad libitum*).

Primary hippocampal neuronal cultures. Primary rat hippocampal neuronal cultures were prepared from P0-P1 Wistar rat pups as previously described (28). In brief, hippocampi were dissected and collected into tubes with ice-cold Hanks balanced salt solution (HBSS; 4.2 mM NaHCO₃, 12 mM Hepes, 33 mM D-glucose, 2 mM Kinurenic acid and 0.95% Hank's salt powder, all purchased from Sigma). Tissue was digested for 8 minutes with trypsin 0.25% (Euroclone) in HBSS solution, then an equal volume of Dulbecco's Modified Eagle Medium (DMEM; Euroclone) supplemented with penicillin/streptomycin (Euroclone) and 10% fetal bovine serum (FBS, Euroclone) was added to stop enzymatic digestion. Digested tissue was centrifuged at room temperature for 5 minutes at 800 rpm, the supernatant was removed and replaced with pre-warmed DMEM

+ 10% FBS. Tissue was homogenized by pipetting, the homogenate was filtered through a 0.40 µm cell strainer (Falcon DB). Cells were plated at a density of 1×10^5 cells in a 24 multi-well plate (Sarstedt) on 12 mm cover glasses (Sacco) pre-treated with 0.1% poly-ornithine (Sigma) and coated with 2% Matrigel (BD science). After one hour, the culture medium was replaced with fresh pre-warmed Neurobasal (Invitrogen) supplemented with B-27 (Invitrogen) and penicillin-streptomycin. Medium supplemented with 0.5 µM Cytosine β-D-arabinofuranoside (Sigma) was replaced at days in vitro (DIV) 2 to stop non-neuronal cell proliferation. Half of the medium was changed twice a week.

Cloning and generation of BDNF 5'UTRs exons-pEGFP-N1 constructs. The different 5'UTR isoforms (exon 1, 2a, 2b, 2c, 3, 4, 5, 6, 7, 8 – See *Supplementary File #1*) of rat BDNF gene were cloned into a pEGFP-N1 vector (Clontech) to obtain the expression of the chimaeric transcript 5'UTR BDNF exon-GFP coding sequence. Total RNA was extracted from adult rat brain using TRIzol (Thermofisher) and retro-transcribed to cDNA using Superscript-II transcriptase (Promega). Forward primers for 5'UTR exons cloning introduced a XhoI restriction site into the amplicon, while reverse primers introduced an AgeI restriction site. Amplification of the specific exon sequences were obtained using Phusion high-fidelity DNA polymerase (Finnzymes). Primers and PCR conditions are reported in Table 1. Amplified sequences were isolated on agarose gel using GenElute Gel extraction kit (Sigma) and cloned into pEGFP-N1 vector digested with XhoI and AgeI restriction enzymes (New England Biolabs) and ligase reaction was performed using T4-DNA ligase (New England Biolabs).

Constructs transfection. Rat hippocampal cultured neurons were transfected at DIV6 with 2 µl of Lipofectamine 2000 (Invitrogen) and 1 µg of chimaeric plasmid DNA (exon 1, 2a, 2b, 2c, 3, 4, 5, 6, 7, or 8-GFP constructs). After one hour, medium with transfectant reagent was removed and replaced with fresh medium. After 24 hours from transfection, cells were depolarized for 3 hours with a isosmotic solution containing 10 mM KCl, 1.8 mM CaCl₂, 1.6 mM MgSO₄, 100 mM NaCl, 26 mM NaHCO₃, 1mM NaH₂PO₄, 30 mM HEPES and 0.7% D-Glucose (all reagents were from Sigma). Treated and untreated cells were fixed in a 4% paraformaldehyde (PFA) in PBS at room temperature for 15 minutes.

In situ hybridization. *In situ* hybridization was performed as previously described (29). Cells were fixed for 15 min at RT in 4% PFA in PBS, washed in PBS 0.1% Tween20

(Sigma) (PBST), and permeabilized in absolute ethanol for 15 min at -20°C. After rehydration with increasing concentration of PBST (50% ethanol/50% PBST, 30% ethanol/70% PBST, and finally PBST), cells were hybridized with approximately 50-100 ng/cover slip of antisense or sense probes for GFP or BDNF coding sequence. Before probe hybridization, coverslips were equilibrated at 55°C in hybridization buffer containing 20 mM Tris-HCl pH 7.5 (Sigma), 300 mM NaCl (Sigma), 1mM EDTA (Sigma), 0.5 mg/ml polyadenylic acid (Sigma), 0.5 mg/ml salmon sperm (Labtek Eurobio), 1X Denhardt's solution, 100 mM Dithiothreitol (DTT, Sigma) and 50% deionised formamide (Sigma). After 1 hour, the hybridization buffer was removed and replaced with the hybridization mix (hybridization buffer + 10% dextran sulfate, Sigma) containing probes. Coverslips were incubated overnight at 55°C. Then, hybridization mix was removed, coverslips were washed with 2X Sodium Saline Citrate (Sigma) buffer (150 mM NaCl, 15mM C₆H₅Na₃O₇) containing 0.1% Tween20 (SSCT 2X) and 50% formamide at 55°C twice, then with SSCT 2X at 55°C once and finally in SSCT 0.1X at 60°C twice. Coverslips were incubated with PBST 5% fetal bovine serum (FBS; Euroclone) for 1 hour at room temperature to block unspecific binding of primary antibodies, then incubated with anti-digoxigenin alkaline phosphatase-conjugated antibody (Roche) in PBST 5% FBS for 2 hours at room temperature. After washing with PBST to remove the excess of unbound antibody, hybridized probes were detected by developing *in situ* signal with 70 mg/ml 4-nitroblue tetrazolium (NBT, Labtek Eurobio) and 50 mg/ml 5-bromo-4-chloro-3-indolyl-phosphate (BCIP, Sigma) in a buffer solution containing 100 mM Tris-HCL pH 9.5, 50 mM MgCl₂, 100 mM NaCl and 1 mM Levamisole (Sigma). Developing was carried out in the dark and time of incubation was determined empirically. The reaction was stopped by removing developing buffer and replacing it with a stop solution containing 10 mM Tris-HCl pH 8.0 and 1 mM EDTA pH 8.0. Coverslips were washed with PBS and deionized water and then mounted with Mowiol (Sigma).

Digoxigen-labeled probes synthesis. Probe synthesis was performed as previously described (3). The open reading frame of GFP was subcloned into pBluescript and digoxigenin (DIG)-labelled probes were synthesized with DIG-RNA labeling kit (Roche Diagnostics) using linearized plasmids as templates, according to the manufacturer's instructions. GFP sense and antisense probe were fragmented through alkaline hydrolysis in a carbonate buffer (40 mM NaHCO₃, 60 mM Na₂CO₃, pH 10.2) to increase probe penetration into cells.

Densitometric analysis on non-radioactive *in situ* hybridization Non-radioactive *in situ* hybridization for GFP coding sequence was analyzed by viewing stained cultures under bright-field illumination as previously described (3) using a Nikon AXM1200 digital camera on a Nikon E800 Microscope with interference contrast-equipped lens (60x magnification) and then analyzed with the image analysis program ImageJ (NIH), using the “Straighten” plugin, on apical dendrites of *bonafide* pyramidal neurons identified by morphological criteria. Dendrites were traced conservatively, starting from the base of dendrite after soma and up to the point in which *in situ* labeling was distinguishable. The background level for each image was determined by evaluation of the mean gray value in the distal portion of apical dendrites of transfected neurons in which the *in situ* labeling was absent. At least 50 neurons from 3 or 4 independent cultures were analyzed for GFP, exon 1, 2a, 2b, 2c, 4, and 6 densitometric analysis, while at least 30 neurons from 2 - 4 independent cultures were analyzed for exon 3, 5, 7 and 8.

Bioinformatic Analysis. The research of putative cis-elements for RNA Binding Proteins on rat BDNF 5'UTR sequence was performed using RBPDB database (<http://rbpdb.ccb.utoronto.ca>). A threshold of 0.8 was set, then FASTA sequences were scanned by the algorithm. The 5'UTR two-dimensional conformation was investigated with the platform RNAfold from the University of Vienna (<http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi>) (30,31).

Statistical Analysis. Statistical data analysis and graph generation were carried out using Sigma Plot 11 software (Systat Software Inc.). The 5'UTR-GFP densitometries were analyzed with Krustal-Wallis ANOVA on ranks for exon-GFP untreated or depolarized versus GFP control or depolarized condition in all exon-GFP experiments.

List of abbreviations

BDNF: Brain-derived neurotrophic factor;

CDS: coding sequence;

CTRL: control;

GFP: green fluorescent protein;

LTP: long term potentiation;

mRNA: messenger ribonucleic acid;
RBP: RNA-Binding Protein;
HNRNPA1: Heterogeneous Nuclear Ribonucleoprotein A1;
PABPC3: Poly(A) Binding Protein Cytoplasmic 3;
SNRNP70: Small Nuclear Ribonucleoprotein U1 Subunit 70;
ZC3H10: Zinc Finger CCCH-Type Containing 10;
ZCRB1: Zinc Finger CCHC-Type And RNA Binding Motif Containing 1;
UTR: untranslated region;
TOG: tumor overexpressed gene;
HNRNPA2/B1: Heterogeneous Nuclear Ribonucleoprotein A2/B1;
HNRNPA2: Heterogeneous Nuclear Ribonucleoprotein A2;
PAP: poly(A)-polymerase;
P-bodies: processing bodies;
FMR1: fragile X mental retardation 1;
FXR1: FMR1 Autosomal Homolog 1;
HBSS: Hanks balanced salt solution;
DMEM: Dulbecco's Modified Eagle Medium;
FBS: fetal bovine serum;
PCR: polymerase chain reaction;
RT: room temperature;
PBS: phosphate-buffer saline;
PBST: phosphate-buffer saline + Tween20;
SSC: Sodium Saline Citrate;
SSCT: Sodium Saline Citrate + Tween20;
BCIP: 5-bromo-4-chloro-3-indolyl-phosphate;
NBT: 4-nitroblue tetrazolium;
EDTA: ethylene-diamino tetra-acetate;
DIG: digoxigenin;
A1CF: APOBEC1 complementation factor;
ANKHD1: Ankyrin repeat and KH domain-containing protein 1
BRUNOL4: Bruno-like 4
BRUNOL5: Bruno-like 5
BRUNOL6: Bruno-like 6
CNOT4: CCR4-NOT transcription complex, subunit 4

CPEB2: Cytoplasmic polyadenylation element-binding protein 2
CPEB4: Cytoplasmic polyadenylation element-binding protein 4
CUG-BP: CUG triplet repeat, RNA binding protein
ENOX1: Ecto-NOX Disulfide-Thiol Exchanger 1
ESRP2: Epithelial Splicing Regulatory Protein 2
HNRNPA1: Heterogeneous Nuclear Ribonucleoprotein A1
HNRNPA2B1: Heterogeneous Nuclear Ribonucleoprotein A2/B1
HNRNPC/CL1: Heterogeneous nuclear ribonucleoprotein C-like 1
HNRNPF: Heterogeneous Nuclear Ribonucleoprotein F
HNRNPH1: Heterogeneous Nuclear Ribonucleoprotein H1
HNRNPH2: Heterogeneous Nuclear Ribonucleoprotein H2
HNRNPK: Heterogeneous Nuclear Ribonucleoprotein K
HNRPL: Heterogeneous Nuclear Ribonucleoprotein L
HNRPLL: Heterogeneous Nuclear Ribonucleoprotein L-Like
HNRNPM: Heterogeneous Nuclear Ribonucleoprotein M
HNRNPU: Heterogeneous Nuclear Ribonucleoprotein U
HuR: human antigen R
IGF2BP2: Insulin-Like Growth Factor 2 mRNA Binding Protein 2
KHDRBS1: KH RNA Binding Domain Containing, Signal Transduction Associated 1
KHDRBS3: KH RNA Binding Domain Containing, Signal Transduction Associated 3
LIN28A: Lin-28 Homolog A
MATR3: Matrin 3
MBNL1: Muscleblind-Like Splicing Regulator 1
MSI1: Musashi RNA Binding Protein 1
NOVA1: RNA-binding protein Nova-1
PABPC3: Poly(A) Binding Protein Cytoplasmic 3
PCBP1: Poly(RC) Binding Protein 1
PCBP2: Poly(RC) Binding Protein 2
PCBP3: Poly(RC) Binding Protein 3
PTBP1: Polypyrimidine tract-binding protein 1
RALY: RNA-binding protein Raly
RBFox1: RNA Binding Fox-1 Homolog 1
RBM24: RNA Binding Motif Protein 24
RBM28: RNA Binding Motif Protein 28

RBM38: RNA Binding Motif Protein 38
RBM4: RNA Binding Motif Protein 4
RBM6: RNA Binding Motif Protein 6
RBM41: RNA Binding Motif Protein 41
RBM42: RNA Binding Motif Protein 42
RBM5: RNA Binding Motif Protein 5
RBM6: RNA Binding Motif Protein 6
RBM8A: RNA Binding Motif Protein 8A
RBMS1/RBMS3: RNA Binding Motif Single Stranded Interacting Protein 1/ RNA Binding Motif Single Stranded Interacting Protein 3
SAMD4A: Sterile Alpha Motif Domain Containing 4A
SFPQ: Splicing Factor Proline And Glutamine Rich
SNRNP70: Small Nuclear Ribonucleoprotein U1 Subunit 70
SRSF1: Serine And Arginine Rich Splicing Factor 1
SRSF10: Serine And Arginine Rich Splicing Factor 10
SRSF2: Serine And Arginine Rich Splicing Factor 2
SRSF3: Serine And Arginine Rich Splicing Factor 3
SRSF5: Serine And Arginine Rich Splicing Factor 5
SRSF7: Serine And Arginine Rich Splicing Factor 7
SRSF9: Serine And Arginine Rich Splicing Factor 9
TARDBP: TAR DNA-binding protein
TIA1: Tia1 cytotoxic granule-associated RNA binding protein
TRA2A: Transformer 2 Alpha Homolog
TRA2B: Transformer 2 Beta Homolog
TUT1: Terminal Uridylyl Transferase 1, U6 SnRNA-Specific
U2AF2: U2 Small Nuclear RNA Auxiliary Factor 2
YBX1: Y-Box Binding Protein 1
ZC3H10: Zinc Finger CCCH-Type Containing 10
ZC3H14: Zinc Finger CCCH-Type Containing 14
ZCRB1: Zinc Finger CCHC-Type And RNA Binding Motif Containing 1
ZNF638: Zinc Finger Protein 638

Declarations

Ethics approval and consent to participate: The present study has been carried out under project authorization N. NO88TON18 from the Italian Ministry of Health.

Consent for publication: Not applicable

Availability of data and materials: All data generated or analyzed during this study are included in this published article and its supplementary information files.

Competing interests: The authors declare that they have no competing interests

Funding: This study has been supported by the Italian Ministry of the University (MUR) grant SYNACTIVE PRIN_2017HPTFFC.

Authors' contributions: AC carried out the experiments, analyzed the data and revised the manuscript. ET conceived the project, analyzed and interpreted the data, wrote and revised the manuscript. Both authors read and approved the final manuscript.

Acknowledgments: Not applicable.

Supplementary Files

Supplementary File#1 – Row data of the densitometric analysis of the in situ hybridization for the different 5'UTR BDNF exons

Supplementary File#2 - FASTA sequences of the 5'UTR BDNF exons used in the bioinformatic analysis

Supplementary File#3 – Recognition sequences and p values of the bioinformatic analysis of each 5'UTR BDNF exon

Literature citations

1. Pang PT, Lu B. Regulation of late-phase LTP and long-term memory in normal and aging hippocampus: role of secreted proteins tPA and BDNF. *Ageing Res Rev.* 2004 Nov;3(4):407–30.
2. Aid T, Kazantseva A, Piirsoo M, Palm K, Timmusk T. Mouse and rat BDNF gene structure and expression revisited. *J Neurosci Res.* 2007 Feb 15;85(3):525–35.
3. Chiaruttini C, Vicario A, Li Z, Baj G, Braiuca P, Wu Y, et al. Dendritic trafficking of BDNF mRNA is mediated by translin and blocked by the G196A (Val66Met) mutation. *Proc Natl Acad Sci.* 2009 Sep 22;106(38):16481–6.
4. Baj G, D'Alessandro V, Musazzi L, Mallei A, Sartori CR, Sciancalepore M, et al. Physical Exercise and Antidepressants Enhance BDNF Targeting in Hippocampal CA3 Dendrites: Further Evidence of a Spatial Code for BDNF Splice Variants. *Neuropsychopharmacology.* 2012 Jun;37(7):1600–11.
5. Baj G, Del Turco D, Schlaudraff J, Torelli L, Deller T, Tongiorgi E. Regulation of the spatial code for BDNF mRNA isoforms in the rat hippocampus following pilocarpine-treatment: A systematic analysis using laser microdissection and quantitative real-time PCR. *Hippocampus.* 2013 May;23(5):413–23.
6. Baj G, Leone E, Chao MV, Tongiorgi E. Spatial segregation of BDNF transcripts enables BDNF to differentially shape distinct dendritic compartments. *Proc Natl Acad Sci.* 2011 Oct 4;108(40):16813–8.
7. Aliaga EE, Mendoza I, Tapia-Arancibia L. Distinct subcellular localization of BDNF transcripts in cultured hypothalamic neurons and modification by neuronal activation. *J Neural Transm.* 2009 Jan;116(1):23–32.
8. Oe S, Yoneda Y. Cytoplasmic polyadenylation element-like sequences are involved in dendritic targeting of BDNF mRNA in hippocampal neurons. *FEBS Lett.* 2010 Aug 4;584(15):3424–30.
9. Tongiorgi E. Activity-dependent expression of brain-derived neurotrophic factor in dendrites: Facts and open questions. *Neurosci Res.* 2008 Aug;61(4):335–46.
10. Maynard KR, Hobbs JW, Sukumar M, Kardian AS, Jimenez DV, Schloesser RJ, et al. Bdnf mRNA splice variants differentially impact CA1 and CA3 dendrite complexity and spine morphology in the hippocampus. *Brain Struct Funct.* 2017 Sep;222(7):3295–307.
11. O'Neill KM, Donohue KE, Omelchenko A, Firestein BL. The 3' UTRs of Brain-Derived Neurotrophic Factor Transcripts Differentially Regulate the Dendritic Arbor. *Front Cell Neurosci.* 2018 Mar 7;12:60.
12. Vicario A, Colliva A, Ratti A, Davidovic L, Baj G, Gricman Ł, et al. Dendritic targeting of short and long 3' UTR BDNF mRNA is regulated by BDNF or NT-3 and distinct sets of RNA-binding proteins. *Front Mol Neurosci* [Internet]. 2015 Oct 29 [cited 2020 Jul 9];8. Available from: <http://journal.frontiersin.org/Article/10.3389/fnmol.2015.00062/abstract>

13. Kanai Y, Dohmae N, Hirokawa N. Kinesin Transports RNA. *Neuron*. 2004 Aug;43(4):513–25.
14. Carson JH, Gao Y, Tatavarty V, Levin MK, Korza G, Francone VP, et al. Multiplexed RNA trafficking in oligodendrocytes and neurons. *Biochim Biophys Acta BBA - Gene Regul Mech*. 2008 Aug;1779(8):453–8.
15. Chiaruttini C, Sonogo M, Baj G, Simonato M, Tongiorgi E. BDNF mRNA splice variants display activity-dependent targeting to distinct hippocampal laminae. *Mol Cell Neurosci*. 2008 Jan;37(1):11–9.
16. Feral C. Human testis expresses a specific poly(A)-binding protein. *Nucleic Acids Res*. 2001 May 1;29(9):1872–83.
17. Afonina E, Stauber R, Pavlakis GN. The Human Poly(A)-binding Protein 1 Shuttles between the Nucleus and the Cytoplasm. *J Biol Chem*. 1998 May 22;273(21):13015–21.
18. Frydryskova K, Masek T, Borcin K, Mrvova S, Venturi V, Pospisek M. Distinct recruitment of human eIF4E isoforms to processing bodies and stress granules. *BMC Mol Biol*. 2016 Dec;17(1):21.
19. Gunderson SI, Polycarpou-Schwarz M, Mattaj JW. U1 snRNP Inhibits Pre-mRNA Polyadenylation through a Direct Interaction between U1 70K and Poly(A) Polymerase. *Mol Cell*. 1998 Jan;1(2):255–64.
20. Diner I, Hales CM, Bishof I, Rabenold L, Duong DM, Yi H, et al. Aggregation Properties of the Small Nuclear Ribonucleoprotein U1-70K in Alzheimer Disease. *J Biol Chem*. 2014 Dec 19;289(51):35296–313.
21. Fu M, Blackshear PJ. RNA-binding proteins in immune regulation: a focus on CCCH zinc finger proteins. *Nat Rev Immunol*. 2017 Feb;17(2):130–43.
22. Louhivuori V, Vicario A, Uutela M, Rantamäki T, Louhivuori LM, Castrén E, et al. BDNF and TrkB in neuronal differentiation of Fmr1-knockout mouse. *Neurobiol Dis*. 2011 Feb;41(2):469–80.
23. Luo Y, Na Z, Slavoff SA. P-Bodies: Composition, Properties, and Functions. *Biochemistry*. 2018 May;57(17):2424–31.
24. Hubstenberger A, Courel M, Bénard M, Souquere S, Ernoult-Lange M, Chouaib R, et al. P-Body Purification Reveals the Condensation of Repressed mRNA Regulons. *Mol Cell*. 2017 Oct;68(1):144-157.e5.
25. Doyle M, Kiebler MA. Mechanisms of dendritic mRNA transport and its role in synaptic tagging: Mechanisms of dendritic mRNA transport. *EMBO J*. 2011 Aug 31;30(17):3540–52.
26. Glock C, Heumüller M, Schuman EM. mRNA transport & local translation in neurons. *Curr Opin Neurobiol*. 2017 Aug;45:169–77.

27. Muslimov IA, Iacoangeli A, Brosius J, Tiedge H. Spatial codes in dendritic BC1 RNA. *J Cell Biol.* 2006 Nov 6;175(3):427–39.
28. Baj G, Patrizio A, Montalbano A, Sciancalepore M, Tongiorgi E. Developmental and maintenance defects in Rett syndrome neurons identified by a new mouse staging system in vitro. *Front Cell Neurosci* [Internet]. 2014 [cited 2020 Jul 10];8. Available from: <http://journal.frontiersin.org/article/10.3389/fncel.2014.00018/abstract>
29. Colliva A, Maynard KR, Martinowich K, Tongiorgi E. Detecting Single and Multiple BDNF Transcripts by In Situ Hybridization in Neuronal Cultures and Brain Sections. In: Duarte CB, Tongiorgi E, editors. *Brain-Derived Neurotrophic Factor (BDNF)* [Internet]. New York, NY: Springer New York; 2018 [cited 2020 Jul 9]. p. 27–53. (*Neuromethods*; vol. 143). Available from: http://link.springer.com/10.1007/7657_2018_14
30. Gruber AR, Lorenz R, Bernhart SH, Neubock R, Hofacker IL. The Vienna RNA Websuite. *Nucleic Acids Res.* 2008 May 19;36(Web Server):W70–4.
31. Lorenz R, Bernhart SH, Höner zu Siederdisen C, Tafer H, Flamm C, Stadler PF, et al. ViennaRNA Package 2.0. *Algorithms Mol Biol.* 2011 Dec;6(1):26.

FIGURE LEGENDS

Figure 1. Trafficking of GFP, exon 1-GFP, and exon 4-GFP transcripts in hippocampal neurons. A) Representative image of an *in situ* hybridization against the GFP coding sequence on one hippocampal pyramidal neuron transfected with pEGFP-N1 plasmid before (upper panel) and after straightening (central panel). Bottom panels represent two control *in situ* hybridization on pEGFP-N1 transfected neurons using GFP coding sequence sense probe (left) and on untransfected neurons using antisense probe. Scale bar: 10 μm . B-D) Densitometric analysis on *in situ* hybridization of apical dendrites of neurons transfected with GFP (panel B), exon 1- GFP (panel C), and exon 4-GFP (panel D) untreated (gray boxes) or treated with 10 mM KCl for 3 hours (white boxes). Below each graph, representative straightened neurons labeled by *in situ* hybridization are reported. Lines in graphs C and D represent the plot of the median gray value of GFP densitometric analysis untreated (solid line) and after KCl stimulation (dashed line). Scale bar: 10 μm .

Figure 2. Trafficking of exon 2a-GFP, exon 2b-GFP, exon 2c-GFP, and exon 6-GFP transcripts in hippocampal pyramidal neurons. A-D) Densitometric analysis on *in situ* hybridization of apical dendrites of pyramidal neurons transfected with exon 2a-GFP (panel A), exon 2b-GFP (panel B), exon 2c-GFP (panel C) and exon 6-GFP untreated (gray boxes) or treated with 10 mM KCl for 3 hours (white boxes). Below each graph, representative straightened neurons labeled by *in situ* hybridization are reported. Lines in graphs represent the plot of the median gray value of GFP densitometric analysis untreated (solid line) and after KCl stimulation (dashed line). Scale bars: 10 μm . *, $p \leq 0.05$ exon-GFP ctrl respect to GFP control and exon-GFP KCl vs GFP KCl

Figure 3. Trafficking of exon 3-GFP, exon 5-GFP, exon 7-GFP, and exon 8-GFP transcripts in hippocampal neurons. A-D) Densitometric analysis of *in situ* hybridization on apical dendrites of neurons transfected with exon 3-GFP (panel A), exon 5-GFP (panel B), exon 7-GFP (panel C) and exon 8-GFP untreated (gray boxes) or treated with 10 mM KCl for 3 hours (white boxes). Below each graph, representative straightened neurons labeled by *in situ* hybridization are reported. Lines in graphs represent the plot of the median gray value of GFP densitometric analysis untreated (solid line) and after KCl stimulation (dashed line). Scale bars: 10 μm . *, $p \leq 0.05$ exon-GFP ctrl respect to GFP control and exon-GFP KCl vs GFP KCl. #, $p \leq 0.05$ exon-GFP ctrl vs exon-GFP KCl.

Figure 4: Predicted folding of exon 2 mRNA variants A-C) Predicted secondary structures using RNAfold from the University of Vienna (<http://rna.tbi.univie.ac.at/cgi-bin/RNAWebSuite/RNAfold.cgi>) of *BDNF* mRNA 5'UTR splicing variants exon 2a (A), 2b (B) and 2c (C). The overall secondary structures of each exon are reported on the right. The red insets on the left highlight a common region displaying a putative binding site for two clusters of RNA binding proteins: CPEB2/CPEB4, HNRNPC/CL1, HuR, TIA1 cluster (green curved line) and HNRP2AB1 cluster (orange straight line).

Table 1: Primers and PCR conditions for BDNF 5'UTR cloning

Exon	Primer Forward	Primer Reverse	PCR conditions			
			Step	Temperature	Time	
Exon I	AATTCTCGAGTAA AGCGGTAGCCGG CTGGTGCAGG	GTCTACCGGITT TGCTGTCCTGGA GACTCAGTGTC	Denaturation	98° C	10 sec	31 cycles
			Annealing	57° C	20 sec	
			Extension	72° C	40 sec	
Exon IIa	GATCCTCGAGGC TTTGGGAAAGCC ATCCGCACGTGAC	GCCACCGGTCT GGATGAAGTACT ACCACCTCGGAC	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	35 sec	
Exon IIb	GATCCTCGAGGC TTTGGCAAAGCC ATCCGCACGTGAC	GTTACCGOTOC AGCTTGCCAAGA GTCTATTCCAG	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	35 sec	
Exon IIc	GATCCTCGAGGC TTTGGCAAAGCC ATCCGCACGTGAC	GAAACCGGTCTTCTTT GCGGCTTACACCACC CCGGTGGCTAG	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	35 sec	
Exon III	AGGCTCGAGGCC CTCACGATTCTCG CTGGATAG	GCTAACCGGTCT GGGCTCAATGA AGCATCCAGCCC	Denaturation	98° C	15 sec	31 cycles
			Annealing	55° C	15 sec	
			Extension	68° C	20 sec	
Exon IV	AAITCTCGAGACC CACTTTCCATTACCGG AGG	GACTACCGGTCTCAGTC ACTACTTGCAAAGTA AACATCAAGGC	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	20 sec	
Exon V	GACTCTCGAGAAACCAT AACCCCGCACACTCTGT GTAGTTTCATTGTGTGT TCG	GACTACCGGTCTTCC CGCACCTTCCCGCAC CACAGAGCTAGAAAA AGCGAACACAC	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	15 sec	
Exon VI	GACTCTCGAGCCAATC GAAGCTCAACCGAAGA GC	GATCACCGGTCTCAG GGTCCACACAAAGCT CTCGG	Denaturation	98° C	10 sec	31 cycles
			Annealing	57.5° C	20 sec	
			Extension	72° C	15 sec	
Exon VII	GACTCTCGAGCACTGTC ACCTGCTTTCTAGGGAG TATTACC	GTACACCGGTCTCCC GGATGAAAGTCAAAA CTTTCACCTTCTCTGG AGG	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	20 sec	
Exon VIII	GTCACCTCGAGGTTATAG AGTTGGATGCAAGCGTA ACCCG	GCATACCGGTGACAC CATTTTCAGCAATCGTT TGTTTCAGCTCC	Denaturation	98° C	10 sec	31 cycles
			Annealing	58° C	20 sec	
			Extension	72° C	35 sec	

Table 2. Identified RNA-binding protein cis-recognition motifs in rodents BDNF 5'UTRs. In green are labeled the RBPs unique to Exons 2a,2b,2c. In orange, the HNRNPA2B1 and in blue, the RBPs of the Fragile-X family.

Exon1	Exon2a	Exon2b	Exon2c	Exon3	Exon4	Exon5	Exon6	Exon7	Exon8
A1CF(1)	A1CF(1)	A1CF(1)	A1CF(1)						
ANKHD1(1)			ANKHD1(1)						
	BRUNOL4/L5(2)	BRUNOL4/L5(2)	BRUNOL4/L5(2)	BRUNOL4/L5(2)		BRUNOL4/L5(1)	BRUNOL4/L5(2)		BRUNOL4/L5(4)
						BRUNOL6(2)	BRUNOL6(1)		BRUNOL6(3)
		CNOT4(2)	CNOT4(2)						
CPEB2(3)	CPEB2(3)	CPEB2(3)	CPEB2(3)	CPEB2(11)		CPEB2(1)			
CPEB4(3)	CPEB4(3)	CPEB4(3)	CPEB4(3)	CPEB4(14)		CPEB4(1)			
CUG-BP(6)	CUG-BP(2)	CUG-BP(2)	CUG-BP(2)	CUG-BP(6)	CUG-BP(9)	CUG-BP(3)	CUG-BP(9)		CUG-BP(5)
ENOX1(1)									
			ESRP2(1)						
					FMR1(1)		FMR1(1)		
FXR1(1)									
					FXR2(2)		FXR2(1)		
				FUS(1)	FUS(6)		FUS(8)		
	HNRNPA1(2)	HNRNPA1(2)	HNRNPA1(2)						
	HNRNPA2B1(3)	HNRNPA2B1(3)	HNRNPA2B1(3)	HNRNPA2B1(1)	HNRNPA2B1(2)		HNRNPA2B1(1)		
HNRNPC/CL1(6)	HNRNPC/CL1(4)	HNRNPC/CL1(5)	HNRNPC/CL1(5)	HNRNPC/CL1(16)		HNRNPC/CL1(1)			HNRNPC/CL1(1)
HNRNPF(1)	HNRNPF(4)	HNRNPF(4)	HNRNPF(4)						HNRNPF(5)
						HNRNPH1(1)			
				HNRNPH2(3)	HNRNPH2(2)	HNRNPH2(1)	HNRNPH2(1)		
		HNRNPK(1)							
									HNRPL(2)
HNRPLL(2)									HNRPLL(4)
HNRNPM(1)		HNRNPM(1)	HNRNPM(2)	HNRNPM(3)					
				HNRNPU(1)					
HUR(3)	HUR(1)	HUR(1)	HUR(2)	HUR(16)					HUR(2)
IGF2BP2(1)									
		KHDRBS1(1)	KHDRBS1(1)						
									KHDRBS3(1)
		LIN28A(2)	LIN28A(2)						
									MATR3(1)
MBNL1(8)	MBNL1(2)	MBNL1(9)	MBNL1(9)	MBNL1(14)	MBNL1(11)	MBNL1(3)	MBNL1(13)	MBNL1(3)	MBNL1(9)
MS1(1)									
NOVA1(6)				NOVA1(1)					NOVA1(1)
	PABPC3(1)	PABPC3(1)	PABPC3(1)						
PCBP1(1)									
				PCBP2(2)					
	PCBP3(1)	PCBP3(3)	PCBP3(3)	PCBP3(1)					
					PPRC1(1)				
PTBP1(24)		PTBP1(18)	PTBP1(18)	PTBP1(25)		PTBP1(7)			PTBP1(16)
RALY(5)	RALY(3)	RALY(3)	RALY(3)	RALY(16)		RALY(1)			PTBP1(4)
									RBFOX1(4)
RBM24(1)	RBM24(2)	RBM24(2)	RBM24(2)	RBM24(1)		RBM24(1)	RBM24(1)		RBM24(2)
		RBM28(1)	RBM28(1)						
	RBM38(2)	RBM38(2)	RBM38(2)			RBM38(2)	RBM38(1)		RBM38(4)
					RBM4(2)	RBM4(2)	RBM4(5)		
	RBM6(1)								
RBM41(1)									
									RBM42(1)
									RBM6(1)
					RBM8(1)				
					RBM8A(1)		RBM8A(2)		
RBMS1/RBMS3(1)									
			SFPQ(1)	SAMD4A(1)	SAMD4A(2)		SAMD4A(2)		
									SFPQ(1)
	SNRNP70(1)	SNRNP70(1)	SNRNP70(1)						
SRSF1(2)				SRSF1(1)	SRSF1(1)		SRSF1(1)	SRSF1(1)	SRSF1(1)
SRSF10(1)	SRSF10(2)		SRSF10(2)						
SRSF2(2)	SRSF2(2)	SRSF2(6)	SRSF2(20)	SRSF2(3)	SRSF2(6)		SRSF2(6)		
SRSF3(20)	SRSF3(6)	SRSF3(15)	SRSF3(10)	SRSF3(10)	SRSF3(4)	SRSF3(3)	SRSF3(6)	SRSF3(10)	SRSF3(7)
SRSF5(19)		SRSF5(2)	SRSF5(5)	SRSF5(4)	SRSF5(5)	SRSF5(2)		SRSF5(8)	SRSF5(9)
SRSF7(6)	SRSF7(1)	SRSF7(1)	SRSF7(1)		SRSF7(1)		SRSF7(1)		SRSF7(1)
									SRSF9(1)
TARDBP(15)	TARDBP(4)	TARDBP(4)	TARDBP(4)	TARDBP(13)	TARDBP(5)	TARDBP(10)	TARDBP(8)		TARDBP(16)
TIA1(4)	TIA1(3)	TIA1(3)	TIA1(3)	TIA1(13)		TIA1(1)			
									TRA2A(2)
TRA2B(14)	TRA2B(3)	TRA2B(3)	TRA2B(7)	TRA2B(2)		TRA2B(1)		TRA2B(5)	TRA2B(10)
									TUT1(1)
U2AF2(4)	U2AF2(3)	U2AF2(7)	U2AF2(7)	U2AF2(13)		U2AF2(2)			U2AF2(1)
YBX1(1)									
	ZC3H10(1)	ZC3H10(1)	ZC3H10(1)						
ZC3H14(2)	ZC3H14(1)	ZC3H14(1)	ZC3H14(1)	ZC3H14(14)					
	ZCRB1(1)	ZCRB1(1)	ZCRB1(1)						
				ZNF638(2)		ZNF638(1)			