

Three-hour post-ERCP amylase level: a useful indicator for early prediction of post-ERCP pancreatitis

Ze-Hui Lv

First Affiliated Hospital of Dalian Medical University

Da-Qing Kou

First Affiliated Hospital of Dalian Medical University

Shi-Bin Guo (✉ gsb@dl.cn)

First Affiliated Hospital of Dalian Medical University

Research article

Keywords: Endoscopic Retrograde Cholangiopancreatography, Serum amylase, Pancreatitis

Posted Date: February 10th, 2020

DOI: <https://doi.org/10.21203/rs.2.13822/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Gastroenterology on April 20th, 2020. See the published version at <https://doi.org/10.1186/s12876-020-01254-7>.

Abstract

Background: To evaluate the value of the 3-hour post-ERCP serum amylase level for early prediction of post-ERCP pancreatitis (PEP). **Method:** A study of 206 patients performed ERCP was analysed. The patients with PEP were recorded. ROC curves were used to statistically analyze the data. **Results:** PEP occurred in 21 patients (10.19%). The 3-hour post-ERCP pancreatic amylase level was used as the test variable, and the PEP occurrence as the state variable to plot ROC curve. The area under the curve (AUC) was 0.816, and was statistically significant ($P < 0.001$). The standard error (SE) was 0.0507, the 95% confidence interval (CI) was 0.756-0.866, and the optimal cut-off value was 351U/L (sensitivity 76.19%, specificity 83.24%, positive likelihood ratio 4.55, negative likelihood ratio 0.29, Youden index 59.43%). The ROC curves were plotted for both serum amylase and lipase respectively. The areas under the ROC curves were statistically significant ($P < 0.001$). The area under the ROC curve for the 3-hour post-ERCP lipase was 0.778, the 95% confidence interval was 0.673-0.862, and optimal cut-off value was 1834 U/L. The area under the ROC curve for the 3-hour post-ERCP serum amylase was 0.780, and the 95% confidence interval was 0.676-0.864. The optimal cut-off is 380U/L, and there was no statistically significant difference between the two for diagnostic accuracy. According to gender, there was no statistically significant difference in the diagnostic accuracy. In the male group, 436 U/L serum amylase provided the greatest diagnostic accuracy with sensitivity(SE) of 70.5%, specificity(SP) of 89.2%, positive predictive value (PPV) 87.5%, and negative predictive value (NPV) 78.1%. Whereas, in the female group, 357U/L serum amylase provided the greatest diagnostic accuracy with sensitivity of 76.9%, specificity of 81.2%, positive predictive value of 80.4%, negative predictive value of 77.9%. **Conclusions:** 1. The 3-hour post-ERCP serum amylase level is a useful measurement for predicting post-ERCP pancreatitis. 2. There was no significant difference between serum amylase and lipase 3-hour post-ERCP for predicting PEP. 3. There was no statistically significant difference between male and female using the 3-hour post-ERCP serum amylase level to predict PEP. For female, the optimal cut-off value was 357 U/L, whereas male 436U/L.

Background

Since first reported by McCunne in 1968[1], ERCP has been widely used in diagnosis and treatment of pancreatic and biliary diseases, especially in the treatment of common bile duct stones. Compared to traditional surgery, it is safer, more effective, with less damage and shorter hospitalization, and has benefits when repeated performance for stone extraction is required[2]. However, its postoperative complications are sometimes difficult to avoid[3]. Post-ERCP pancreatitis (PEP) is one of the most common complications after ERCP procedure. It is reported that the incidence of PEP is 1.6-15.7%[4]. Although 90% are mild or moderate[5], 1% may develop acute necrotizing pancreatitis[6]. Since discharge and readmission after pancreatitis results in worse outcome[7, 8], early diagnosis and timely treatment is very important[9, 10]. So far, the determination of serum amylase is still the most commonly used indicator for the diagnosis of PEP, but there is still no clear standard for the early prediction of PEP at different time points and different levels of amylase[11, 12]. In this study, the ROC curve of the diagnostic test was used to evaluate the value of early prediction of PEP by 3-hour post-ERCP serum amylase level.

Methods

Patients

From January 2011 to November 2016, a total of 546 patients were diagnosed or treated with ERCP at the First Affiliated Hospital of Dalian Medical University. The study was conducted in compliance with the Helsinki Declaration and in accordance with local legislation, was approved by the Ethics Committee of First Affiliated Hospital, Dalian Medical University. (Ethics References No: YJ-KY-FB-2019-01). Written informed consent was obtained from all of the patients or their relatives before the procedure.

The inclusion and exclusion criteria were as follows:

Inclusion criteria: (1) preoperative serum amylase and lipase levels were normal; (2) age ≥ 18 years; (3) serum amylase level was measured 3 hours after ERCP; (4) patients have not used trypsin inhibitor before the diagnosis of PEP.

Exclusion criteria: (1) preoperative diagnosis of acute and chronic pancreatitis (2) abnormal renal function (serum creatinine $> 92\mu\text{mol} / \text{L}$) (3) pregnant women

According to the inclusion and exclusion criteria above, a total of 206 patients were enrolled, including 92 males and 114 females, among them, 84 cases were simultaneously detected for serum lipase.

The diagnosis of pancreatitis after ERCP is based on the consensus reached by Cotton et al[4, 13], and the international consensus on the classification of acute pancreatitis in Atlanta in 2012[14].

(1) Acute pancreatic abdominal pain within 24 hours after ERCP;

(2) Serum amylase more than 3 times the upper limit of normal within 72 hours after ERCP(normal value is 30-110U/L) or lipase greater than 3 times the upper limit of normal within 96h after ERCP(normal reference range 23-300U/L) ;

(3) Contrast-enhanced CT, MRI, abdominal ultrasound showing acute pancreatitis changes (pancreatic enlargement, exudation, necrosis and other AP signs); having two of the three criteria will lead to a diagnosis of PEP.

(4) The necessity for new or continued hospitalization for at least 2 nights.

1.2 Equipment Description

Duodenoscope JF-260, Olympus Optical Corporation, Tokyo, Japan), guide wire (Hydra Jagwire 0.035 inch, Boston Scientific Microvasive. Cork, Ireland), triple lumen sphincterotome (Papillotome, ENDO-FLEX GmbH, Germany), CRE balloon catheter (Boston Scientific Microvasive, Cork, Ireland), retrieval balloon catheter (Extractor Three Lumen Retrieval Balloon, Boston Scientific Microvasive. Cork, Ireland), Dormia

basket (Web™ extraction basket, Wilson-Cook Medical Inc. Winston-Salem, North Carolina, United States), Percuflex Biliary Stent (Boston Scientific Corporation, One Boston Scientific Place, Natick, MA 01760-1537, USA), nasal biliary drainage tube (nasobil.Sonde, ENDO-FLEX GmbH, Germany), WallFlex Biliary RX Fully Covered Stent System (Boston Scientific Corporation, One Boston Scientific Place, Natick, MA 01760-1537, USA), Mechanical lithotripsy (BML-4Q; Olympus Optical, Tokyo, Japan).

Some ERCP procedures were accomplished under ECG monitoring. **Tetracaine** was given for local anesthesia of the pharynx. The patients were sedated and relieved pain by intramuscular administration of diazepam and meperidine. 20 mg of butyl scopolamine bromide was injected intramuscularly prior to the procedure to inhibit duodenal peristalsis.

A sphincterotome with a guide wire was used for selective cannulation. Difficulties in selective cannulation including stenosis and sclerosis of the papillary, incarceration of common bile duct (CBD) stone at papilla, and periampullary diverticulum—especially the papilla is located at bottom of the diverticulum.

For some cases difficulty in bile cannulation, precut through pancreatic duct was applied. For cases with incarceration of CBD stone at papilla, needle knife was used for precut sphincterotomy. For papilla located at the bottom of diverticulum, it was exposed by eversion diverticulum through biopsy forceps, or fixation by metal clip, or submucosal injection of saline.

For patients with CBD stones, small endoscopic sphincterotomy (EST) alone or EST combined with endoscopic papillary balloon dilation (EPBD) was performed. For biliary tract benign stricture or biliary fistula, ERBD was given. For biliary or pancreatic malignant tumor, a self-expanding metal stent was placed.

Mechanical lithotripsy was used for large common bile duct stones. However, if patient was in poor condition and cannot tolerate long procedure time, just ERBD was placed for drainage and to relieve symptoms. And a next ERCP was performed for removal of cholelithiasis three months later.

Serum amylase, lipase, biochemical liver function and renal function were detected by Johnson & Johnson's VITROS FS5.1 automatic biochemical analyzer.

1.4 statistical methods

According to the inclusion and exclusion criteria above, a total of 206 patients were enrolled, and each patient's progress note was reviewed by the medical record system. Sudden pancreatitis-like mid-abdominal pain within 24 hours after ERCP was marked positive, otherwise negative; combined with CT, serum amylase, lipase levels, comprehensive estimation was made according to the PEP diagnostic criteria .

Statistical methods: 1. Descriptive analysis of the enrolled patients, a preliminary understanding of the basic situations of the study objects. 2. The ROC curve was plotted according to the 3-hour post-ERCP

serum amylase level of the enrolled patients and PEP diagnosis. According to the area under the ROC curve and the 95% confidence interval of the area, the accuracy of the tests was evaluate based on statistical tests. The Youden index was calculated according to the sensitivity and specificity of each level and the optimal cut-off value was obtained. The sensitivity, specificity, positive and negative predictive value, positive and negative likelihood ratio, Youden index of different cut-off sites were calculated, comprehensively evaluated the accuracy and predictive value of the predictive diagnostic method. 3. According to gender, they were divided into two groups, and the ROC curves were drawn and analyzed separately aimed to find best cut-off in different gender. As a same lab test whose diagnosis accuracy be showed as AUC in ROC should have no difference between genders or other classifications of subjects. So we use Z test to test the reliability of data then obtain the optimal diagnostic cut-off value according to these two curves. 4. Matched-pair analysis of ROC curves in patients who were simultaneously tested for both serum amylase levels and lipase levels 3 hours after ERCP. According to the two test Methods, the ROC curves were drawn and statistically tested, and the difference test (Z test) was performed to compare and analyze the diagnostic accuracy and diagnostic value in the two groups.

Statistical software: Medcalc was applied to perform matched-pair analysis. The ROC curve of serum amylase of different gender patients were plotted using SPSS software, and then the difference test was performed according to the standard error and the area under the curve. The Z value was manually calculated, and the P value was looked up in the table .

Related formula:

Sensitivity: Se; Specificity: Sp

$$LR(+) = \frac{Se}{1-Sp}$$

Positive likelihood ratio: (1)

$$LR(-) = \frac{1-Se}{Sp}$$

Negative likelihood ratio: (2)

$$PPV = \frac{p * Se}{p * Se + (1-p) * (1-Sp)} \times 100\%$$

Positive predictive value: (3)

$$NPV = \frac{(1-p) * Sp}{(1-p) * Sp + p * (1-Se)} \times 100\%$$

Negative predictive value: (4)

$$YI = Se + Sp - 1$$

Youden Index: (5)

Results

1. Basic situations:

A total of 562 patients who underwent diagnostic or therapeutic ERCP in our hospital between January 2011 to November 2016. The total number of patients who met the criteria for enrollment was 206, 92 males and 114 females. Among the 206 patients, endoscopic sphincterotomy (including EST alone and EST combined with EPBD) was performed in 136 patients; endoscopic nasobiliary drainage (ENBD) was performed in 140 patients; endoscopic retrograde biliary drainage (ERBD) was performed in 36 patients (Table 1). The median duration from the procedure to discharge was 7 days (min = 1d, max = 13d). A total of 21 patients (10.19%) were diagnosed with PEP, 2 patients (0.97%) had upper gastrointestinal bleeding, and 3 patients (1.45%) had post-ERCP infection. No case of perforation.

The average amylase level was 224.44 U/L in the non-PEP group, with a median of 115.00 U/L. The average amylase level in the PEP group was 688.90 U/L, with a median of 456 U/L. The serum amylase levels in the PEP group and the non-PEP group after ERCP were positively skew distribution, and the results showed that the serum amylase levels 3-hour after ERCP were significantly higher than those of the non-PEP group (non-PEP group stem width = 100; PEP group stem width = 1000).

2. Analyses of ROC curve of amylase level 3-hour after ERCP and PEP diagnosis in 206 patients.

The non-parametric method was used in this paper, the serum amylase level 3-hour post-ERCP as the test variable, the PEP occurrence as the state variable and the ROC curve was drawn. The result is shown in Fig. 1 (the solid line is the drawn ROC curve, and the dotted line is 95% confidence interval), In tables 2, the area under the ROC curve was 0.816, the diagnostic value is good, the standard error was 0.0507, the 95% confidence interval was 0.756-0.866, and the significance test Z value was 6.235, $P < 0.0001$. The area under the ROC curve was statistically significant, and the optimal cut-off value was 351 U/L. The sensitivity, specificity, positive and negative likelihood ratio, positive and negative predictive value, and Youden index of different cut-off values are shown in Table 3:

As seen from the Table 4, when 351U/L was taken as the optimal cut-off value, a better positive predictive value (81.90) could be obtained, the Youden index was the highest (59.43%), and the diagnosis was the most accurate at this time. Meanwhile, the positive and the negative likelihood ratio were both high (4.55, 0.29), indicating that the possibility of diagnosis or elimination of the disease was high. When taking 105U/L (close to the standard upper limit of normal), the negative likelihood ratio reached 0.21, which indicates that this is an ideal diagnostic criterion with a favorable negative predictive value and has certain clinical reference significance.

2.3 ROC curve matched-pair testing for the 3-hour post-ERCP amylase level between male and female, and further analysis according to gender

In order to investigate whether different genders groups need different diagnostic criteria, ROC curve was applied to evaluate this issue in our study. 206 patients were divided into two groups according to gender,

and the ROC curves were plotted respectively(Fig. 2, male; Fig. 3, female). The results are shown as follows:

The ROC curves drawn for the two groups both had statistical significance ($P=0.001$ for both groups), the area under the curve for the male group was 0.859, the standard error was 0.065, and the area under the curve for the female group was 0.794, and the standard error was 0.068 (Table 5,6). It indicated that the accuracies of the two tests were good; the two groups were independent samples, suitable for comparison of independent ROC curves to test the statistical significance. The Z value of the hypothesis test was 0.69098, less than 1.96, $P>0.05$, which indicated there was no statistical difference between the two groups. We know that the ROC curve is not affected by the prevalence rate. The results indicate that although the prevalence rates were different in the two groups, there was no statistical difference in the accuracy of predicting PEP by amylase level 3-hour post-ERCP between the two groups, which also verifies the validity of our study.

The corresponding diagnostic values according to the optimal cut-off values are shown in the following Table 7: In the male group, 436 U/L serum amylase provided the greatest diagnostic accuracy with sensitivity of 70.5%, specificity of 89.2%, positive predictive value of 87.5%, and negative predictive value of 78.1%. Whereas, in the female group, 357U/L serum amylase provided the greatest diagnostic accuracy with sensitivity of 76.9%, specificity of 81.2%, positive predictive value of 80.4%, negative predictive value of 77.9%. The NPV in the male group was higher than that of the female group, which indicated that the proportion of PEP in the male group was also higher than that of the female group. This was consistent with the objective situation that women have a higher incidence of PEP than men.

2.4. Matched-pair analysis of the 3-hour post-ERCP serum amylase and lipase levels

The ROC curve matched-pair testing was used to compare the serum amylase and serum lipase 3 hours after ERCP. A total of 83 patients with both the serum amylase and lipase levels 3-hour post-ERCP were selected. Among them, 15 patients were diagnosed as PEP. The ROC curves were drawn (Fig. 4) and the results were compared as follows: The ROC curves of the two groups were statistically significant ($P < 0.001$). The area under the ROC curve of lipase levels 3-hour post-ERCP was 0.778, with a 95% confidence interval of 0.673-0.862; The area under the ROC curve of amylase 3-hour post-ERCP was 0.780, and the 95% confidence interval was 0.676-0.864, indicating that the accuracy of predicting PEP was high (Table 8). Matched-pair testing was performed between the two groups, and the results showed that there was no statistically significant difference in the accuracy of predicting PEP (Z score 0.0598, $P=0.9523$, $P>0.05$) (Table 9), and the negative predictive value of lipase 3-hour post-ERCP was not high (0.688) (Table 10).

Discussion

ERCP has become an important method in clinical diagnosis and treatment for biliary and pancreatic diseases. However, its complications should not be ignored, especially post-ERCP pancreatitis (PEP). As the most common post-ERCP complication, PEP has restricted the development of ERCP. The average

incidence of PEP is reported to be 3.5% in the literatures, but in high-risk groups (such as previous PEP, difficulty in cannulation of the bile duct, etc.), the incidence can be as high as 30%[5]. Although most cases of PEP were mild[5], they recovered after conservative treatment in less than 72 h, in some severe cases, the length of hospital stay can be extended, the costs can be increased, multiple organ failure may occur, even death[6]. Since it is difficult to avoid PEP even if the procedure is performed by an endoscopic expert, early detection and timely treatment for the PEP are very important. Serum amylase measurement technology is simple, easy to obtain for low price with high sensitivity and early appearance. Although the level of amylase cannot estimate the severity of pancreatitis, it is still one of the most commonly used indicators for diagnosis of PEP currently. However, there is still no clear criteria for various cut-off serum amylase levels at different time points in prediction and diagnosis of PEP[9, 11, 15-18].

The role of serum amylase levels in predicting the occurrence of PEP after ERCP has been recognized[19]. Takayoshi Nishino[16] et al have demonstrated a good negative predictive value of the serum amylase levels 4-hour post-ERCP for PEP(AUC 0.91, $p=0.007$) through a retrospective study of 1631 patients. Kapetanios et al[20] also have demonstrated that the 2-hour post-ERCP amylase cut-off value 3 times the upper limit of normal exhibited an NPV and a PPV for PEP of 95% and 32% respectively, which indicates the 2-hour post-ERCP serum amylase is valuable for eliminating the diagnosis of PEP. However, it can not be used for early detection of the PEP. Studies have shown that serum amylase levels at 6, 8, and 12 h after ERCP have a good predictive effect on PEP, and the accuracy of prediction is higher over time, but the significance of early prediction has been lost[21]. The studies by Sutton et al[12] suggest that serum amylase value 4-hour after ERCP greater than 5 times the upper limit of normal (ULN), or serum amylase value greater than 2.5 times the ULN and pancreatic duct visualization are good predictors of PEP.

Our study showed that the serum amylase level 3-hour after ERCP has a good accuracy in the prediction of PEP. The cut-off at 351U/L provides the optimal diagnostic accuracy, when setting the cut-off at 105U / L (close to the standard upper limit of normal), the negative likelihood ratio can reach 0.21, indicating that this is a valuable predictive value for eliminating PEP.

Studies have confirmed that female gender is one of the important risk factors for pancreatitis after ERCP[22], so whether different genders need to use different predictive diagnostic criteria is one of the purposes of this study. By using the ROC curve matched-pair testing, the results showed that the two groups have the best diagnostic accuracy at different cut-off values. For female, the optimal cut-off value was 357U/L, whereas male 436U/L. This conclusion may provide help for clinicians. In addition, nearly 25% of patients with type 2 diabetes have elevated levels of lipase and/or amylase without the symptoms of acute pancreatitis[23]. Whether we need to establish different cut-off values for the diagnosis of PEP to obtain a better predictive effect for those patients, is still to be validated by more clinical studies.

Serum lipase is a specific diagnostic indicator of pancreatitis. Its activity is maintained longer than amylase, and there is no other source in the blood[24], so it is considered to be more sensitive than serum amylase (95%;79%). The British guidelines for the diagnosis and treatment of acute pancreatitis in 2005

indicated that the pancreas is the only source of lipase and is superior to amylase in terms of sensitivity, specificity and accuracy[25]. And the American guidelines for the diagnosis and treatment of acute pancreatitis in 2006 also had similar views[26]. The European Guidelines for Digestive Endoscopy in 2014 clearly indicated that serum amylase $<1.5 \times \text{ULN}$ or serum lipase $<4 \times \text{ULN}$ monitored 2-4 hours after ERCP provide a high NPV for PEP, and the patients can be discharged without considering the risk of PEP (recommended level B)[19], which suggests that serum lipase has a high negative predictive value for PEP. Gumaste et al[27] pointed out based on an analysis of 170 patients with abdominal pain, the negative predictive value for acute pancreatitis was 98%. However, we emphasized the early (3-hour) lipase and amylase levels after ERCP in this study. And the results showed that in terms of comparison the predictive value for PEP between serum lipase and amylase 3-hour post-ERCP, lipase was not better than amylase. It is well known that serum lipase starts to rise at 24-48h and reaches a peak at 72-96 hours and last for 7-10d. This seems to explain why the early lipase monitoring value is not significantly better than that of amylase. Based on the actual situations in China, the predictive value of the joint monitoring needs to be further verified, and the economic benefits of the two tests need further cost-benefit analysis.

However, this was a retrospective study in a single center, and small sample was another limitation for the study. More cases are required to confirm the value of the 3-hour post-ERCP serum amylase level for early prediction of post-ERCP pancreatitis in the future.

Conclusions

In summary, we have come to the following conclusions:

The 3-hour post-ERCP serum amylase level is a useful measurement for early predicting post-ERCP pancreatitis . With the 3-hour post-ERCP serum amylase level 1-1.5 times upper limit of normal, the negative predictive value is high.

There was no significant difference between serum amylase and lipase 3-hour post-ERCP for predicting PEP.

There was no statistical difference between male and female using the 3-hour post-ERCP serum amylase level for prediction PEP. For female, the optimal cut-off value was 357U/L, whereas male 436U/L.

List Of Abbreviations

post-ERCP pancreatitis PEP

area under the curve AUC

standard error SE

confidence interval CI

sensitivity SE
specificity SP
positive predictive value PPV
negative predictive value NPV
Endoscopic Retrograde Cholangiopancreatography ERCP
Endoscopic Sphincterotomy EST
Endoscopic Nasobiliary Drainage ENBD
Endoscopic Retrograde Biliary Drainage ERBD
Endoscopic Papillary Balloon Dilatation EPBD
Endoscopic Retrograde Pancreatic Drainage ERPD
Endoscopic Metal Biliary Endoprosthesis EMBE
upper limit of normal ULN

Declarations

Ethics and Consent to participate

The study was conducted in compliance with the Helsinki Declaration and in accordance with local legislation, was approved by the Ethics Committee of First Affiliated Hospital, Dalian Medical University. (Ethics References No: YJ-KY-FB-2019-01). Informed consent was not acquired as this was a retrospective study.

Consent for Publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

No funding was obtained for this study

Author information

Affiliations

1. Department of Gastroenterological Endoscopy, the First Affiliated Hospital of Dalian Medical University, Dalian 116011, Liaoning Province, China. Ze-Hui Lv & Shi-Bin Guo
2. Department of Clinical Laboratory, the First Affiliated Hospital of Dalian Medical University, Dalian 116011, Liaoning Province, China. Da-Qing Kou

Contributions

G SB designed the study; L ZH and K DQ collected and analyzed the data; All authors were involved in writing and editing the manuscript. L ZH and K DQ have equal contribution. All authors have read and approved the manuscript.

Corresponding author

Correspondence to Shi-Bin Guo

Acknowledgements

We would like to acknowledge the contribution of the colleagues of the department of Endoscopy, Surgery, First Affiliated Hospital of Dalian Medical University.

Availability of data and material

Data and material was sent in attachment. Data for the analyses are available from the corresponding author on request.

References

1. McCune WS, Shorb PE, Moscovitz H: **Endoscopic cannulation of the ampulla of vater: a preliminary report.** *Ann Surg* 1968, **167**(5):752-756.
2. Meng W, Leung JW, Zhang K, Zhou W, Wang Z, Zhang L, Sun H, Xue P, Liu W, Wang Q *et al*: **Optimal dilation time for combined small endoscopic sphincterotomy and balloon dilation for common bile duct stones: a multicentre, single-blinded, randomised controlled trial.** *Lancet Gastroenterol Hepatol* 2019.
3. Kim HG, Cheon YK, Cho YD, Moon JH, Park DH, Lee TH, Choi HJ, Park SH, Lee JS, Lee MS: **Small sphincterotomy combined with endoscopic papillary large balloon dilation versus sphincterotomy.** *World J Gastroenterol* 2009, **15**(34):4298-4304.
4. Cotton PB, Garrow DA, Gallagher J, Romagnuolo J: **Risk factors for complications after ERCP: a multivariate analysis of 11,497 procedures over 12 years.** *Gastrointest Endosc* 2009, **70**(1):80-88.
5. Dumonceau JM, Andriulli A, Deviere J, Mariani A, Rigaux J, Baron TH, Testoni PA, European Society of Gastrointestinal E: **European Society of Gastrointestinal Endoscopy (ESGE) Guideline: prophylaxis**

- of post-ERCP pancreatitis. *Endoscopy* 2010, **42**(6):503-515.
6. Elmunzer BJ: **Preventing Postendoscopic Retrograde Cholangiopancreatography Pancreatitis.** *Gastrointest Endosc Clin N Am* 2015, **25**(4):725-736.
 7. Cote GA, Lynch S, Easler JJ, Keen A, Vassell PA, Sherman S, Hui S, Xu H: **Development and Validation of a Prediction Model for Admission After Endoscopic Retrograde Cholangiopancreatography.** *Clin Gastroenterol Hepatol* 2015, **13**(13):2323-2332 e2321-2329.
 8. Freeman ML, DiSario JA, Nelson DB, Fennerty MB, Lee JG, Bjorkman DJ, Overby CS, Aas J, Ryan ME, Bochna GS *et al.*: **Risk factors for post-ERCP pancreatitis: a prospective, multicenter study.** *Gastrointest Endosc* 2001, **54**(4):425-434.
 9. Minakari M, Sebghatollahi V, Sattari M, Fahami E: **Serum amylase and lipase levels for prediction of postendoscopic retrograde cholangiopancreatography pancreatitis.** *J Res Med Sci* 2018, **23**:54.
 10. Gottlieb K, Sherman S, Pezzi J, Esber E, Lehman GA: **Early recognition of post-ERCP pancreatitis by clinical assessment and serum pancreatic enzymes.** *Am J Gastroenterol* 1996, **91**(8):1553-1557.
 11. Lee YK, Yang MJ, Kim SS, Noh CK, Cho HJ, Lim SG, Hwang JC, Yoo BM, Kim JH: **Prediction of Post-Endoscopic Retrograde Cholangiopancreatography Pancreatitis Using 4-Hour Post-Endoscopic Retrograde Cholangiopancreatography Serum Amylase and Lipase Levels.** *J Korean Med Sci* 2017, **32**(11):1814-1819.
 12. Sutton VR, Hong MK, Thomas PR: **Using the 4-hour Post-ERCP amylase level to predict post-ERCP pancreatitis.** *JOP* 2011, **12**(4):372-376.
 13. Cotton PB, Lehman G, Vennes J, Geenen JE, Russell RC, Meyers WC, Liguory C, Nickl N: **Endoscopic sphincterotomy complications and their management: an attempt at consensus.** *Gastrointest Endosc* 1991, **37**(3):383-393.
 14. Banks PA, Bollen TL, Dervenis C, Gooszen HG, Johnson CD, Sarr MG, Tsiotos GG, Vege SS, Acute Pancreatitis Classification Working G: **Classification of acute pancreatitis–2012: revision of the Atlanta classification and definitions by international consensus.** *Gut* 2013, **62**(1):102-111.
 15. Xu LH, Qian JB, Gu LG, Qiu JW, Ge ZM, Lu F, Wang YM, Li YM, Lu HS: **Prevention of post-endoscopic retrograde cholangiopancreatography pancreatitis by epinephrine sprayed on the papilla.** *J Gastroenterol Hepatol* 2011, **26**(7):1139-1144.
 16. Thomas PR, Sengupta S: **Prediction of pancreatitis following endoscopic retrograde cholangiopancreatography by the 4-h post procedure amylase level.** *J Gastroenterol Hepatol* 2001, **16**(8):923-926.
 17. Ito K, Fujita N, Noda Y, Kobayashi G, Horaguchi J, Takasawa O, Obana T: **Relationship between post-ERCP pancreatitis and the change of serum amylase level after the procedure.** *World J Gastroenterol* 2007, **13**(28):3855-3860.
 18. Sultan S, Baillie J: **What are the predictors of post-ERCP pancreatitis, and how useful are they?** *JOP* 2002, **3**(6):188-194.
 19. Dumonceau JM, Andriulli A, Elmunzer BJ, Mariani A, Meister T, Deviere J, Marek T, Baron TH, Hassan C, Testoni PA *et al.*: **Prophylaxis of post-ERCP pancreatitis: European Society of Gastrointestinal**

Endoscopy (ESGE) Guideline - updated June 2014. *Endoscopy* 2014, **46**(9):799-815.

20. Kapetanios D, Kokozidis G, Kinigopoulou P, Xiarchos P, Antonopoulos Z, Progia E, Kitis G: **The value of serum amylase and elastase measurements in the prediction of post-ERCP acute pancreatitis.** *Hepatogastroenterology* 2007, **54**(74):556-560.
21. Katsinelos P, Kountouras J, Paroutoglou G, Beltsis A, Mimidis K, Zavos C: **Intravenous N-acetylcysteine does not prevent post-ERCP pancreatitis.** *Gastrointest Endosc* 2005, **62**(1):105-111.
22. Zamri Z, Razman J: **Comparison of complication outcomes in acute pancreatitis following ERCP and conservative management at UKM medical centre: a six years retrospective study.** *Clin Ter* 2012, **163**(6):467-471.
23. Davison LJ: **Diabetes mellitus and pancreatitis—cause or effect?** *J Small Anim Pract* 2015, **56**(1):50-59.
24. Papachristos A, Howard T, Thomson BN, Thomas PR: **Predicting post-endoscopic retrograde cholangiopancreatography pancreatitis using the 4-h serum lipase level.** *ANZ J Surg* 2018, **88**(1-2):82-86.
25. Working Party of the British Society of G, Association of Surgeons of Great B, Ireland, Pancreatic Society of Great B, Ireland, Association of Upper GI SoGB, Ireland: **UK guidelines for the management of acute pancreatitis.** *Gut* 2005, **54 Suppl 3**:iii1-9.
26. Banks PA, Freeman ML, Practice Parameters Committee of the American College of G: **Practice guidelines in acute pancreatitis.** *Am J Gastroenterol* 2006, **101**(10):2379-2400.
27. Gumaste VV, Roditis N, Mehta D, Dave PB: **Serum lipase levels in nonpancreatic abdominal pain versus acute pancreatitis.** *Am J Gastroenterol* 1993, **88**(12):2051-2055.

Tables

Due to technical limitations the Tables are available as a download in the Supplementary Files.

Figures

Fig.1





Figure 1

ROC curve of amylase level 3-hour after ERCP and PEP diagnosis in 206 patients

Fig.2




Diagonal segments are produced by ties.

Figure 2

ROC curve of amylase level 3-hour after ERCP and PEP diagnosis in male group

Fig.3



Diagonal segments are produced by ties.

Figure 3

ROC curve of amylase level 3-hour after ERCP and PEP diagnosis in female group

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [data.xls](#)
- [Table26.pdf](#)