Physical Fitness and Cardiometabolic Risk in Seven to Nine Year Old Children

Gerardo Weisstaub (✉ gweiss@inta.uchile.cl)

Universidad de Chile Instituto de Nutricion y Tecnologia de los Alimentos <https://orcid.org/0000-0003-3938-4972>

Maria Angelica Gonzalez

University of Talca

Antonio Garcia Hermoso

USACH

Gabriela Salazar

INTA

Research article

Keywords: Physical fitness, Cardio-metabolic risk, Muscular strength, Obesity

Posted Date: July 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-45298/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Obesity and lower physical fitness levels in children are related to the development of cardiometabolic risk. The objective of the study was to determine the relationship of cardio-metabolic risk and physical fitness in school children from Santiago de Chile.

Methods

Physical fitness was assessed as cardiorespiratory fitness and muscle strength. Fitness was measured using the six-minute-walk-test and muscle strength was assessed by hand grip and standing lung jump. Overweight (≥ 1 SD) and obesity (≥ 2 SD) were determined by body mass index. A cross sectional study was done in 452 children (185 boys and 267 girls), age range (7–9 years). Cardio-metabolic-risk (serum glucose, triglycerides, HDL, insulin) and waist for height ratio was expressed as z score. Data was analyzed using bivariate analysis and multiple and logistic regression.

Results

A 21% were obese and 27% overweight. Children with high body fat and low cardiorespiratory fitness and muscle strength had an OR of 6.8-fold (IC 95% 3.4 to 13.9) cardiometabolic risk z-score, compared to those most fit.

Conclusion

School children with higher body fat and lower physical fitness had increased cardio-metabolic-risk-score. No significant difference in cardio-metabolic risk in the non-obese children was found independent of their physical fitness.

Introduction

The prevention and treatment of cardiovascular disease and its risk factors are commonly focused on the adult population. However, in recent years, evidence has revealed that these risk factors can be observed in early life.^[1, 2] Childhood obesity in Chilean children has tripled in recent decades, reaching 23.9% in 2017 ^[3]. Obesity has been shown to be associated with a high prevalence of hypercholesterolemia (27%) and high insulin levels during fasting (42% of all obese children have abnormal values) ^[4]. Childhood obesity is also associated with an increased risk of cardiovascular disease in adulthood ^[5]. The sum of risk factors, assessed using a cardiometabolic risk (CMR) score, is considered a more appropriate tool for defining disease risk in children ^[6, 7]. Inappropriate eating patterns and a lack of regular moderate to

vigorous physical activity [8] are important contributors to the early onset of CMR. Sixty percent of the world's population does not have an adequate level of physical activity in terms of preventing noncommunicable disease [9,10]. Chilean children are predominantly sedentary, given that only 20.2% of them meet the physical activity recommended guidelines [11, 12, 13]. Similarly, in northeast England, sedentary behaviour was found in 78% to 82% of 7- to 9-year-old children [14]. In school children and adolescents, this problem has further consequences if we consider that only 9% of school-age Chilean children have acceptable physical fitness levels [15]

Cardiorespiratory and muscle fitness are inversely associated with the risk factors for cardiovascular diseases [16]. Physical fitness assessments are feasible measures that could help improve the monitoring of paediatric health status [17]. The independent biological effect of cardiorespiratory fitness and muscle strength on cardiometabolic risk has been demonstrated in adolescents [18,19].

Artero et al, evaluated the relationship between metabolic risk and physical fitness in 709 adolescents. They found that muscular fitness was negatively associated with clustered metabolic risk independent of cardiorespiratory fitness ($\beta = -0.249$, $p < 0.001$). Independent of muscular fitness, an inverse association was also found between cardiorespiratory fitness and clustered metabolic risk ($\beta = 0.264$, $p < 0.001$) [20]. Steene-Johannessen J, et al, measured in 2818 children 9- and 15-year-olds physical fitness and metabolic risk. They also encountered that muscle fitness was negatively associated with clustered metabolic risk, independent of cardiorespiratory fitness, and after adjustment for age, sex, and pubertal stage ($\beta = -0.112$, $P < 0.001$). [21] [22]. Additionally, there is scarce information in this group age. The objective of the study is to determine the relationship of cardiometabolic risk and physical fitness in school children from Santiago de Chile.

Methods

This study is cross-sectional; the sample was drawn from children enrolled in the "Growth and Obesity Chilean Cohort Study", which assesses the association of early growth and development with adiposity and metabolic risk.[23].

We randomized by using random numbers, included healthy 7- to 9-year-old schoolchildren from the original study, both girls and boys, who had provided a blood sample in 2009. According to the objective of this project, fitness tests were performed in 2010. Parents or guardians signed a previously approved informed consent form by the Institute of Nutrition and Food Technology Ethics Committee for research in human subjects. The final sample size was 452 children (267 girls and 185 boys).

Weight was measured with an electronic scale (Body Composition Analyzer TANITA BC-418, USA) with a 0.1 kg accuracy and 220 kg maximum measurement. Children were measured in underclothing, while placing their feet in the centre of the scale. Height measurements were performed with a portable stadiometer (SECA 222 ®) with a range between 0 and 200 cm and a 1 mm precision using the Frankfurt standard [24]. Nutritional status was determined using body mass index (BMI) and BMI-Z. Overweight and

obesity were defined as BMI-z ≥ 1 and $2 \geq$ SD respectively (WHO 2007) [25,26]. Waist circumference was measured with an inextensible tape (SECA ®) [27]. All variables were measured twice for each child. The measurements agreed within 0.5 cm for height and waist circumference and 20 g for weight. Finally, waist circumference and height were used to calculate the waist-to-height ratio (WHtR), and central obesity was defined as having a WHtR ≥ 0.5 [28].

The body composition of a child was classified in tertiles of BMI or WHtR. According to the results, body fat mass was classified as low, medium and high.

Identification of pubertal status (Tanner stages I–II) was assessed by study professionals using the method described by Tanner and Whitehouse. [29].

Blood samples were obtained between 08.30 and 10.30 after an 8-h overnight fast. Glucose, insulin and blood lipids were assessed after 8 h of fasting. Ten millilitres of venous blood were collected, and serum glucose was assessed using a commercial kit by the GOD-PAP (Clinical Chemistry Applied SA) enzymatic colorimetric method. Insulin was measured by RIA (RIA DCP Diagnostic Products Corporation, LA, USA). Glucose intolerance was defined by a fasting glucose ≥ 100 mg/dL [30]. Basal insulin sensitivity was estimated by homeostatic model assessment - HOMA (fasting insulin (mIU/dl) * fasting glucose (mmol/l)/22.5) [31]. Hyperinsulinemia was defined as baseline insulin ≥ 10 mIU / dl in children with Tanner I or II [32]. Insulin resistance was diagnosed if HOMA ≥ 2.1 , which corresponds to > 75th percentile of the observed quartile in Chilean children [33]. Cholesterol, HDL and triglycerides were determined by dry analytical methodology (Vitros, Johnson & Johnson, Clinical Diagnostics Inc.) using the USA population as a reference (Non-HDL cholesterol > 145 mg/dL; HDL < 40 mg/dL and triglycerides > 130 mg/dL) [34]. The remaining blood was eliminated.

The CMR score was defined by the sum of the Z score of the WHtR-z and blood lipid related variables: (Glycaemia-z, Insulin-z and Triglycerides-z – HDL-z)/5, as previously reported [35]. These variables were chosen because they are commonly used in adults as criteria to diagnose metabolic syndrome [36, 37].

Physical fitness was assessed as cardiorespiratory fitness and muscle strength. Cardiorespiratory fitness was measured using a sub-maximal test, the six-minute walk test (6MWT) [38, 39]. The 6MWT, hand grip strength and standing long jump without momentum have been shown to be valid, reliable, and feasible for health monitoring purposes at the population level [34, 42]. The 6MWT results were expressed as the maximal distance (metres) divided by height because leg length was not assessed [40]. The value was also standardized using the z score.

Muscle strength was assessed using hand grip and standing long jump. Handgrip was measured using an adjustable hand grip digital dynamometer (Baseline 12-0286 ®; 100 g accuracy), and the results were expressed in kg. The hand grip was performed twice by each child, by squeezing the dynamometer as hard as possible, for at least two seconds. One minute of recovery between squeezes was measured (alternating the right and left hands). Finally, hand grip strength was divided by body weight (kg) to take

into account body size differences. The best result of two attempts was registered [41]. The best value was also standardized using the z score.

The standing long jump was performed by starting in a semi-squatting position, then jumping forward. This test was performed on a 2-m long non-slippery surface, free of obstacles. A cross mark was made to identify the initial position from which the child jumped. A piece of chalk was used to mark the landing point for each of the two attempts; the distance in cm served to register the best jump [42]. The best value was also standardized using the z score¹⁶.

The results for muscle strength are expressed as a z score and were calculated based on standard values of hand grip and standing long jump from this study (hand grip/ weight z + standing long jump -z) [43].

The physical fitness z score was defined as 6MWT/height -z score + (hand grip/ weight z score + standing long jump -z score). The z score data were continuous.

Statistical analysis

The normality of the data for each variable was checked using the Shapiro-Wilk test and graphical methods. Statistical normality was tested using both statistical (Shapiro Wilk test) and graphical methods (normal probability plots). The majority of outcomes were normally distributed except hand grip, hand grip/weight, triglycerides, insulin, HOMA and body mass index.

Based on the distribution of the variables, the Mann-Whitney test or a t test was used to analyse independent samples. Preliminary analyses showed significant interactions between sex, physical fitness and cardiometabolic variables; therefore, combined association analyses were performed with boys and girls together to increase the statistical power of the results.

The effect of muscle strength and cardiorespiratory fitness on the CMR was also studied using two-way ANCOVA by dividing groups by muscle strength and cardiorespiratory fitness tertiles. The interaction between CMR-z score and fatness categories (low and high BMI or WHtR) was assessed. The post hoc analysis of differences between groups was performed using a Bonferroni adjustment to test for significance and was adjusted by age and maturation stage. The respective physical fitness values for age groups were segregated by tertiles since no reference values are available. Finally, the odds ratio (OR) and 95% CI of having a high CMR were examined by combined physical fitness variables (i.e., cardiorespiratory fitness and muscle strength). A multinomial logistic regression adjusted by age, sex and maturation stage was used. All analyses were performed using SPSS software, version 21.0. Statistical significance was set at $p < 0.05$.

Results

Descriptive statistics for the studied children are shown.

Table 1 shows that girls were significantly older than boys and that boys had slightly higher blood glucose than girls ($p < 0.05$).

Table 1
Description of anthropometric variables and cardiometabolic risk.

	Boys (n = 185)	Girl (n = 267)	Total (n = 452)	p^1
Age (years)	7.75 ± 0.5	7.88 ± 0.4	7.8 ± 0.5	0.002*
Weight (kg)	27.9 ± 7.6	28.2 ± 8.1	28 ± 8.1	0.2
Height (cm)	126.6 ± 5.6	127.1 ± 5.8	126.9 ± 5.7	0.7
BMI (kg/m ²)	17.1 (3.2)	17.4 (3.9)	17.3 (3.8)	0.2
Waist-to-height ratio	0.47 ± 0.05	0.48 ± 0.08	0.5 ± 0.07	0.1
Body fat (%)	27,6 ± 7,9	32,2 ± 6,7	29,8 ± 7,6	0.01
Glycaemia (mg/dl)	90.2 ± 6.2	88.7 ± 6.5	89.3 ± 6.4	0.002*
Triglyceride (mg/dl)	88.0 (51.5)	87.0 (49)	87 (49.8)	0.6
High-density lipoprotein (mg/dl)	49.6 ± 14.4	50.1 ± 12.3	49.9 ± 13.2	0.3
Insulin (uU/dl)	5.10 (0.6)	5.20 (0.7)	5.2 (0.7)	0.1
HOMA	1.14 (0.2)	1.13(0.2)	1.1 (0.2)	0.7
Cardiometabolic risk z-score	-0.04 ± 0.6	-0.02 ± 0.6	-0.02 ± 0.6	0.7
Mean ± SD or median and (interquartile range), ¹ Differences between boys and girls were examined with Student's t test for independent observations with equal variances (variables with normal distribution) or with the Mann-Whitney nonparametric test. BMI was calculated as weight (kg)/height (cm) ²				

Table 2 shows the CMR factors by sex and nutritional status. Significant differences in CMR score and WHtR were found in both sexes; higher values for both variables were observed in overweight children compared to normal weight children ($p < 0.001$). Overweight girls had significantly higher triglycerides, insulin and HOMA than normal weight girls ($p < 0.001$).

Table 2
Description of cardiometabolic risk factors by nutritional status.

	Boys (n = 185)			Girls (n = 267)		
	Normal (n = 93)	OvOb (n = 92)	p	Normal (n = 141)	OvOb (n = 126)	p
WC/Height	0.45 ± 0.03	0.50 ± 0.06	< 0.001	0.45 ± 0.04	0.53 ± 0.05	< 0.001
Glycaemia (mg/dl)	89.5 ± 5.9	90.9 ± 6.4	0.06	88 ± 6.7	89.4 ± 5.7	0.06
TG (mg/dl)	84 (55.5)	91.5(46.7)	0.16	79 (41)	101(47.2)	< 0.001
HDL (mg/dl)	49.8 ± 14.3	49.3 ± 14.6	0.92	51.2 ± 12.6	48.9 ± 11.9	0.12
Insulin (uU/dl)	5.10(0.75)	5.20(0.75)	0.29	5.10(0.5)	5.30(0.8)	0.004
HOMA	1.14 (0.20)	1.17(0.21)	0.06	1.12 (0.16)	1.15(0.22)	< 0.001
CMR-z	-0.19 ± 0.54	0.15 ± 0.63	< 0.001	-0.26 ± 0.54	0.20 ± 0.57	< 0.001
Mean ± SD or median and (interquartile range); BMI , body mass index; WC/height waist circumference / height; TG , triglycerides; HDL , high-density lipoprotein; HOMA , homeostasis model assessment; CMR z , cardiometabolic risk score, OvOb , overweight and obese						

Table 3 shows the physical fitness results by nutritional status and sex. Compared to obese children, the normal nutritional status group had better muscle strength and cardiorespiratory fitness in both sexes in most tests ($p < 0.05$). Notably, on the hand grip strength test, children with overweight and obesity had better records than normal weight children, but when adjusted for weight, the results were reversed. Boys showed higher values of muscle strength and cardiorespiratory fitness than girls ($p < 0.05$). The findings suggest that increasing at least moderate-intensity physical activity and decreasing sedentary time, decreases cardiometabolic risk in children.

Table 3
Description of physical fitness variables by nutritional status.

	Boys (n = 185)			Girls (n = 267)		
	Normal (n = 93)	OvOb (n = 92)	p ¹	Normal (n = 141)	OvOb (n = 126)	p
Standing long jump (cm)	116.9 ± 7.5	111.2 ± 4.9	0.02	106.2 ± 15.6	101.0 ± 3.5	< 0.005
Hand grip (kg)	11.1 (2.4)	12.5 (4.0)	< 0.001	10.6 (2.3)	11.4 (3.2)	< 0.001
Hand grip/weight	0.45 (0.1)	0.39 (0.1)	< 0.001	0.44 (0.1)	0.40 (0.1)	< 0.001
6MWT (m)	625.2 ± 53.7	615.9 ± 44.9	0.2	615.8 ± 52.8	609.2 ± 42.4	0.2
6MWT/height (m)	503.6 ± 49.8	477.6 ± 53.3	0.002	493.1 ± 48.7	474.9 ± 48.6	< 0.001
Muscle strength-z score	1.14 ± 2.1	-0.01 ± 1.7	< 0.001	0.34 ± 1.9	-1.10 ± 1.6	< 0.001
6MWT/height- z score	0.43 ± 1.2	-0.20 ± 1.3	0.002	0.17 ± 1.2	-0.30 ± 1.2	< 0.001
Mean ± SD or median and (interquartile range), OvOb , overweight and obese 6MWT Six-minute walk test; ¹ Differences in nutritional status were examined with Student's t test for normal distributions or with the Mann-Whitney non-parametric test for non-normal observations.						

Figure 1a shows the CMR score tertiles in relation to physical fitness and fat categories (BMI and WHtR). In subjects with a low fat mass tertile, there was no difference in the CMR risk by muscle strength and cardiorespiratory fitness tertiles. Whereas in children with a high fat mass tertile, higher muscle strength or cardiorespiratory fitness was associated with a lower CMR than that of other groups. The post hoc analysis found significant differences between the low and high tertiles ($p < 0.05$), and CMR was lower for the higher tertile. Muscular strength has a relevant role in attenuating the association between physical inactivity and metabolic risk in children, in a comparison of active and sedentary children.

The post hoc analysis found significant differences between the low and high tertiles ($p < 0.05$), and CMR was lower for the higher tertile.

Figure 1b shows the CMR score according to fat categories and muscle strength tertiles. In subjects with a low fat mass tertile, there were no differences in CMR according to cardiorespiratory fitness tertile. In high-fat tertile children, the group with higher cardiorespiratory fitness had lower CMR values than the low

tertile cardiorespiratory fitness group. The results of Fig. 1a and 1b were adjusted by age, sex and maturation status.

Table 4 shows the odds ratio (OR) of the relationship between high CMR and the combined association of physical fitness (cardiorespiratory fitness and muscle strength). The analysis was performed with the high tertile of cardiorespiratory fitness and the high tertile of muscle strength as references. The results show that a low tertile of cardiorespiratory fitness and a low tertile of muscle strength are associated with a 6.8 times higher CMR (95% CI, 3.4 to 13.9) compared to the high tertiles for both of these factors. In addition, a low tertile of muscle strength and a medium tertile of cardiorespiratory fitness had a high CMR (OR = 3.5 Finally, a medium tertile of cardiorespiratory fitness and a medium tertile of muscle strength resulted in a high CMR (OR = 2.6 95% CI, 1.2 to 5.5; p = 0.003).

Table 4
Combined effects of physical fitness parameters on high cardiometabolic risk

Cardiorespiratory fitness level	Muscle strength	Cardiometabolic risk (OR and 95% CI)
High	High	reference
	Medium	1.6 (0.8 to 3.3)
	Low	2.4 (0.9 to 4.9)
Medium	High	1.5 (0.7 to 3.1)
	Medium	2.6 (1.2 to 5.5)*
	Low	3.5 (1.4 to 8.8)*
Low	High	0.6 (0.2 to 2.0)
	Medium	4.1 (2.8 to 8.4)*
	Low	6.8 (3.4 to 13.9)*
OR significant * All data were compared to higher muscle strength and higher cardiorespiratory fitness.		

Discussion

Our results show that both muscle strength and cardiorespiratory fitness are inversely associated with the CMR-z score. In another study in adolescents, it was stated that continuous CMR-z scores are practical tools and are accurate and easy to apply [44]. Although fatness (BMI tertile) was the factor that explained CMR, overweight children with better physical fitness had a significantly lower CMR. This suggests that body composition is a critical determinant of metabolic health, as other studies have shown in adolescents [45,46]. On the other hand, children, who increase their adiposity levels as they enter puberty, despite remaining normal weight are at risk of developing cardiometabolic risk factors. [47]

Our results show that overweight children had greater hand grip strength than eutrophic children. Ervin et al, found a similar result in a sample of 1224 youth aged 6 to 15 years, wherein grip strength was significantly higher in obese children and adolescents than in eutrophic children. However, when hand grip strength was adjusted by weight, eutrophic children achieved better results, a fact that suggests that static hand strength correlates more with lean mass than with total weight or BMI [48]. Children with low grip strength had higher systolic blood pressure, low-density lipoprotein cholesterol, higher metabolic syndrome score (0.36), compared with those with moderate/high grip strength [49]. Cohen et al, has shown that when analyses were stratified by sex, handgrip strength was negatively associated with cardiometabolic z-score in girls. Furthermore, young people in the lowest strength quartile were three times more likely to have elevated metabolic risk than those in the highest strength quartile. [50].

No difference was found by sex or nutritional status for the 6MWT, but when the 6MWT was adjusted by height, the maximal distance was significantly different ($p < 0.05$). This adjustment by height was done to minimize the bias that body size has on the walked distance (correlation $r = 0.17$) (data not shown). The test was validated by Li et al, where the correlation of the 6MWT with size had an $r = 0.45$ ($p < 0.05$). The same author has shown that this test is valid, easy to perform and economical as an indicator of sub-maximum capacity, compared to cardiorespiratory fitness or other stressful tests [51]. Moura BP et al, found that light sedentary physical activity can be considered an effective alternative to reduce sedentary behavior as improves the cardiometabolic health indicators in the children [52].

The first multinomial logistic regression analysis revealed that muscle strength and cardiorespiratory fitness explained CMR. However, in the overweight group of schoolchildren, it was observed that children with better muscle strength and cardiorespiratory fitness had a lower CMR. These results coincide with other works in adolescents and adults, which were attributed to changes in muscle metabolism, as decreased visceral fat mass provides protection [16, 53]. In similar studies, muscular strength has shown a stronger association with CMR score than cardiorespiratory fitness has [20,21]. Similar findings in previous work in children and adolescents have shown an independent association of muscle strength with insulin sensitivity [15] and metabolic disease risk [54]. Agostinis-Sobrinho et al [55], in another study in adolescents, indicated that persistent greater and increasing muscle strength is associated with lower levels of cardiovascular risk. The results of Schmidt et al [56] strengthen the findings that cardiorespiratory fitness has a strong association with CMR.

Several studies have shown that CMR is higher in overweight children and adolescents with low muscle strength [7, 49]. The results of this study suggest that muscle strength in schoolchildren may confer additional benefits attributed to cardiorespiratory fitness. Most obese children are reluctant to participate in aerobic training; therefore, power exercise may be more attractive and better tolerated [57].

Hand grip strength and the standing long jump have proven to be a good validity criterion, compared to laboratory tests [34,58], suggesting that they are appropriate measures of muscle strength, especially in the first years of schooling.

Among the strengths of our study, we confirm that a risk score composed of metabolic risk factors can be used to investigate the cardiorespiratory fitness and muscle strength [16, 18, 30, 45]. Such variables indicators may better reflect the health status of participants than the assessment of individual risk factors [59].

Limitations

However, our results should be interpreted with care. The CMR score is specific to the sample under study and is based on the assumption that each component is equally weighted in the prediction of CMR. The 6MWT is a submaximal test to assess cardiorespiratory fitness, and it is not comparable with the standard aerobic test, i.e., the Course-Navette test. Finally, to evaluate muscle strength, only hand grip strength and standing long jump were measured, and other tests could have been chosen^{36, 46}. However, these tests are easy to apply and well tolerated in children of young ages, whose results are reported in this study.

Conclusion

Muscle strength and cardiorespiratory fitness are associated independently and inversely with CMR score. These results confirm the importance of stimulating physical activity to prevent metabolic risk at an early age, especially in sedentary children or those with high weights. In the future, longitudinal studies are needed to further find the independent and combined effects of cardiorespiratory fitness and muscle strength in the prevention of cardiovascular disease risk.

Abbreviations

BMI	Body Mass Index
CMR	Cardio-metabolic risk
6MWT	Six-minute walk test

Declarations

Conflict of interest The authors declare that they have no conflict of interest.

Declarations

Ethics approval and consent to participate: Ethical approval: Research involving human participants. All procedures performed were in accordance with the ethical standards as laid down in the 1964 Helsinki declaration. Informed consent: Parents or guardians previously signed the informed consent forms approved by INTA's Committee for research in human subjects.

Consent for publication

- Availability of data and materials: The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request
- Competing interests: The authors declare that they have no conflict of Interest
- Funding: There is no funding source
- Authors' contributions

GW designed the study, review of the literature, writing and revised the manuscript, approval of the final manuscript

MAG review of the literature and statistical analysis the data, approval of the final manuscript.

AGH statistical analysis the data and approval of the final manuscript

GS supervise the data collection, review of the literature, writing and revised the manuscript, approval of the final manuscript.

Acknowledgements: We thank to Algerina Anziani (IRMS laboratory technician)

References

- 1 Perez-Escamilla R, Lutter CK, Rabadan-Diehl C, Rubinstein A, Calvillo A, Corvalán C, et al: Prevention of childhood obesity and food policies in Latin America: from research to practice. *Obes Rev* 2017, 19 Suppl 2:28-38.
- 2 Pacheco LS, Blanco E, Burrows R, Reyes M, Lozoff B, Gahagan S: Early Onset Obesity and Risk of Metabolic Syndrome Among Chilean adolescents. *Prev Chronic Dis* 2017, 14:E93.
- 3 Junta de Auxilio Escolar y Becas de Chile (JUNAEB). Prevalencia de Obesidad en escolares de Primero Básico. Mapa Nutricional. 2017. [Available Dic 2019 in <https://www.junaeb.cl/wp-content/uploads/2013/03/Mapa-Nutricionalpresentacio%CC%81n2.pdf>
- 4 Vasquez FD, Corvalan CL, Uauy RE, Kain JA: Anthropometric indicators as predictors of total body fat and cardiometabolic risk factors in Chilean children at 4, 7 and 10 years of age. *Eur J Clin Nutr* 2017, 71:536-543.
- 5 Cote AT, Harris KC, Panagiotopoulos C, Sandor GG, Devlin AM: Childhood obesity and cardiovascular dysfunction. *J Am Coll Cardiol* 2013, 62:1309-19.
- 6 Cesa CC, Sbruzzi G, Ribeiro RA, Barbiero SM, de Oliveira Petkowicz R, Eibel B, Machado et al: Physical activity and cardiovascular risk factors in children: meta-analysis of randomized clinical trials. *Prev Med* 2014, 69:54-62.

- 7 García-Hermoso A, Ramírez-Campillo R, Izquierdo M: Is Muscular Fitness Associated with Future Health Benefits in Children and Adolescents? A Systematic Review and Meta-Analysis of Longitudinal Studies. *Sports Med* 2019, 49:1079–1094.
- 8 Schuler G, Adams V, Goto Y: Role of exercise in the prevention of cardiovascular disease: results, mechanisms, and new perspectives. *Eur Heart J* 2013, 34:1790-9.
- 9 Organización Mundial de la Salud. Estrategia Mundial sobre Régimen Alimentario, Actividad Física y Salud. La inactividad física. [available Dic 2019
http://www.who.int/dietphysicalactivity/factsheet_inactivity/es/index.html]
- 10 Sirico F, Bianco A, D'Alicandro G, Castaldo C, Montagnani S, Spera R, Di Meglio F, Nurzynska D (2018) Effects of Physical Exercise on Adiponectin, Leptin, and Inflammatory Markers in Childhood Obesity: Systematic Review and Meta-Analysis. *Child Obes* 14:207-217.
- 11 Aguilar-Farias N, Miranda-Marquez S, Sadarangani KP, Martino-Fuentealba P, Cristi-Montero C, Carcamo et al. Results from Chile's 2018 Report Card on Physical Activity for Children and Youth. *J Phys Act Health* 2018, 15(S2):S331-S332.
- 12 Díaz-Martínez X, Petermann F, Leiva AM, Garrido-Méndez A, Salas-Bravo C, Martínez MA, et al: Association of physical inactivity with obesity, diabetes, hypertension and metabolic syndrome in the chilean population. *Rev Med Chil* 2018, 146:585-595.
- 13 Telama R, Yang X, Leskinen E, Kankaanpää A, Hirvensalo M, Tammelin T, et al. Tracking of physical activity from early childhood through youth into adulthood. *Med Sci Sports Exerc* 2014, 46:955-962.
- 14 Basterfield L, Pearce MS, Adamson AJ, Fray JK, Parkinson KN, Wright CM, Reilly JJ: Gateshead Millennium Study Core Team. Physical activity, sedentary behavior, and adiposity in English children. *Am J Prev Med* 2012, 42:445-51.
- 15 Report SIMCE Physical Education. Eight grade. Ministry of Education. Unity of Curriculum and Evaluation. Santiago, Chile. 2015. (Available Dic 2019 www.simce.cl)
- 16 Delgado Floody PA, Caamaño Navarrete F, Palomino-Devia C, Jerez-Mayorga D, et al: Relationship in obese Chilean schoolchildren between physical fitness, physical activity levels and cardiovascular risk factors. *Nutr Hosp* 2019, 36:13-19.
- 17 Lang JJ, Larouche R, Tremblay MS: The association between physical fitness and health in a nationally representative sample of Canadian children and youth aged 6 to 17 years. *Health Promot Chronic Dis Prev Can* 2019, 39:104-111.
- 18 Agostinis-Sobrinho C, García-Hermoso A, Ramírez-Vélez R, Moreira C, Lopes L, Oliveira-Santos J, et al: Longitudinal association between ideal cardiovascular health status and muscular fitness in adolescents: The LabMed Physical Activity Study. *Nutr Metab Cardiovasc Dis* 2018, 28:892-899.

- 19 Mintjens S, Menting MD, Daams JG, van Poppel MNM, Roseboom TJ, Gemke RJ:Cardiorespiratory Fitness in Childhood and Adolescence Affects Future Cardiovascular Risk Factors: A Systematic Review of Longitudinal Studies. *Sports Med* 2018, 48:2577-2605.
- 20 Artero EG, Ruiz JR, Ortega FB, et al. Muscular and cardiorespiratory fitness are independently associated with metabolic risk in adolescents: the HELENA study. *Pediatr Diabetes*. 2011;12(8):704–712.
- 21 Steene-Johannessen J, Anderssen SA, Kolle E, Andersen LB: Low muscle fitness is associated with metabolic risk in youth. *Med Sci Sports Exerc*. 2009;41(7):1361–1367.
- 22 Myers J, Kokkinos P, Nyelin E: Physical Activity, Cardiorespiratory Fitness, and the Metabolic Syndrome. *Nutrients*. 2019;11(7):1652.
- 23 González L, Corvalán C, Pereira A, Kain J, Garmendia ML, Uauy R: Early adiposity rebound is associated with metabolic risk in 7-year-old children. *Int J Obes (Lond)* 2014, 38:1299-304.
- 24 Capon T: Standardised anatomical alignment of the head in a clinical photography studio. A comparison between the Frankfort Horizontal and the natural head position. *J Vis Commun Med* 2016, 39:105-111.
- 25 de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J: Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ* 2017, 85:660-7.
- 26 National Center for Health Statistics. Growth curves. CDC growth Charts for United States: Methods and development. 2000. (Available Janury 2020 in www.cdc.gov/growthcharts).
- 27 Gomez-Campos R, Andruske CL, Hespanhol J, Torres JS, Arruda M, Duarte-Rocha C, Cossio-Bolaños MA: Waist Circumferences of Chilean Students: Comparison of the CDC-2012 Standard and Proposed Percentile Curves. *Int J Environ Res Public Health* 2015, 9:7712-7724.
- 28 Asif M, Aslam M, Altaf S: Evaluation of anthropometric parameters of central obesity in Pakistani children aged 5-12 years, using receiver operating characteristic (ROC) analysis. *J Pediatr Endocrinol Metab* 2018, 31:971-977.
- 29 Tanner JM, Whitehouse RH: Clinical longitudinal standards for height, weight, height velocity, weight velocity, and stages of puberty. *Arch Dis Child* 1976, 51: 170–179.
- 30 American Diabetes Association: Diagnosis and Classification of Diabetes Mellitus. *Diabetes Care* 2017, 40:s11-s24.
- 31 Radziuk J: Homeostatic Model Assessment and Insulin Sensitivity /Resistance. *Diabetes* 2014, 63:1850-1854.

- 32 Moschonis G, Karatzi K, Polychronopoulou MC, Manios Y: Waist circumference, trunk and visceral fat cutoff values for detecting hyperinsulinemia and insulin resistance in children: the Healthy Growth Study. *Eur J Nutr* 2016, 55:2331-4.
- 33 Burrows R, Correa-Burrows P, Reyes M, Blanco E, Albala C, Gahagan S: Healthy Chilean Adolescents with HOMA-IR ≥ 2.6 Have Increased Cardiometabolic Risk: Association with Genetic, Biological, and Environmental Factors. *J Diabetes Res* 2015, 783296.
- 34 Smith AJ, Turner EL, Kinra S: Universal Cholesterol Screening in Childhood: A Systematic Review. *Acad Pediatr* 2016, 16:716-725.
- 35 Corvalán C, Uauy R, Stein AD, Kain J, Martorell R: Effect of growth on cardiometabolic status at 4 y of age. *Am J Clin Nutr*. 2009; 90(3):547–555.
- 36 Mokha J, Srinivasan S, DasMahapatra P, Fernandez C, Chen W, Xu J, Berenson G: Utility of waist-to-height ratio in assessing the status of central obesity and related cardiometabolic risk profile among normal weight and overweight/obese children: The Bogalusa Heart Study. *BMC Pediatrics* 2010, 10:73.
- 37 Kamel M, Smith BT, Wahi G, Carsley S, Birken CS, Anderson LN: Continuous cardiometabolic risk score definitions in early childhood: a scoping review. *Obes Rev* 2018, 19:1688-1699.
- 38 ATS Statement Guidelines for the Six-Minute Walk Test. *Am J Respir Crit Care Med* 2002, 166:111-117.
- 39 Li A, Yin J, Hung S, Tsang T, Wong E, Fok T, Ng P: Standard Reference for the Six-Minute-Walk Test in Healthy Children Aged 7 to 16 years. *Am J Respir Crit Care Med* 2007, 176:174-80.
- 40 Oliveira AC, Rodrigues CC, Rolim DS, Souza AA, Nascimento OA, Jardim JR, Rozov T: Six-minute walk test in healthy children: is the leg length important? *Pediatr Pulmonol* 2013, 48:921-6.
- 41 Ervin RB, Fryar CD, Wang CY, Miller IM, Ogden CL: Strength and body weight in US children and adolescents. *Pediatrics* 2014, 134:e782-e789.
- 42 Castro-Piñero J, Ortega F, Artero E, Girela-Rejon M, Mora J, Sjostron M, Ruiz J: Assessing muscular strength in youth: usefulness of standing long jump as a general index of muscular fitness. *J Strength Cond Res* 2010, 24:1810–1817.
- 43 Andersen LB, Harro M, Sardinha LB, Froberg K, Ekelund U, Brage S, Anderssen SA: Physical activity and clustered cardiovascular risk in children: a cross-sectional study (The European Youth Heart Study). *Lancet* 2006, 22:299-304.
- 44 Šebeková K, Gurecká R, Csongová M, Koborová I, Šebek J: Estimation of the proportion of metabolic syndrome-free subjects on high cardiometabolic risk using two continuous cardiometabolic risk scores: a cross-sectional study in 16- to 20-year-old individuals. *Eur J Pediatr* 2019, 178:1243-1253.

- 45 Vásquez FD, Corvalán CL, Uauy RE, Kain JA: Anthropometric indicators as predictors of total body fat and cardiometabolic risk factors in Chilean children at 4, 7 and 10 years of age. *Eur J Clin Nutr* 2017, 71:536-543.
- 46 Ervin RB, Fryar CD, Wang CY, Miller IM, Ogden CL: Strength and body weight in US children and adolescents. *Pediatrics* 2014, 134: e782-e789.
- 47 Van Hulst A, Ybarra M2, Mathieu ME, Benedetti A, Paradis G, Henderson M: Determinants of new onset cardiometabolic risk among normal weight children. *Int J Obes (Lond)*. 2020; 44:781-789.
- 48 . Wong SL: Grip strength reference values for Canadians aged 6 to 79: Canadian Health Measures Survey, 2007 to 2013. *Health Rep* 2016, 27:3-10.
- 49 [1] Laitinen TT, Saner C, Nuotio J, et al: Lower grip strength in youth with obesity identifies those with increased cardiometabolic risk]. *Obes Res Clin Pract*. 2020;S1871-403X(19)30792-6.
- 50 Cohen DD, Gómez-Arbeláez D, Camacho PA, et al: Low muscle strength is associated with metabolic risk factors in Colombian children: the ACFIES study. *PLoS One*. 2014; 9(4): e93150.
- 51 [1]. Li A, Yin J, Yu C, Tsang T, So H, Wong E, et al: The six-minute walk test in healthy children: reliability and validity. *Eur Respir J* 2005, 25:1057-1060.
- 52 Moura BP, Rufino RL, Faria RC2: Amorim PRS. 2Effects of isothermal substitution of sedentary behavior with light-intensity or moderate-to-vigorous physical activity on cardiometabolic markers in male adolescents. *PLoS One*. 2019; 14 (11):e0225856.
- 53 Isasi CR, Strizich GM, Kaplan R, Daviglus ML, Sotres-Alvarez D, Vidot DC: The association of cardiorespiratory fitness with cardiometabolic factors, markers of inflammation, and endothelial dysfunction in Latino youth: findings from the Hispanic Community Children's Health Study/Study of Latino Youth. *Ann Epidemiol* 2018, 28:583-589.
- 54 Castro-Piñero J, Perez-Bey A, Cuenca-Garcia M, Cabanas-Sanchez V, Gómez-Martínez S, Veiga OL: Cut Points for Early Assessment of Cardiovascular Risk in Children and Adolescents. *J Pediatr* 2018, S0022-3476(18)31466-5.
- 55 Agostinis-Sobrinho C, Ruiz JR, Moreira C, Lopes L, Ramírez-Vélez R, García-Hermoso A, Mota J, Santos R: Changes in muscular fitness and its association with blood pressure in adolescents. *Eur J Pediatr* 2018, 177:1101-1109.
- 56 Schmidt MD, Magnussen CG, Rees E, Dwyer T, Venn AJ: Childhood fitness reduces the long-term cardiometabolic risks associated with childhood obesity. *Int J Obes (Lond)* 2016, 40:1134-1140.
- 57 Thivel D, Ring-Dimitriou S, Weghuber D, Frelut ML, O'Malley G: Muscle Strength and Fitness in Pediatric Obesity: a Systematic Review from the European Childhood Obesity Group. *Obes Facts* 2016, 9:52-63.

58 Laurson KR, Saint-Maurice PF, Welk GJ, Eisenmann JC: Reference Curves for Field Tests of Musculoskeletal Fitness in U.S. Children and Adolescents: The 2012 NHANES National Youth Fitness Survey. *J Strength Cond Res* 2017, 31:2075-2082.

59 Segura-Jiménez V, Parrilla-Moreno F, Fernández-Santos JR, Esteban-Cornejo I, Gómez-Martínez S, Martínez-Gomez D, et al: Physical fitness as a mediator between objectively measured physical activity and clustered metabolic syndrome in children and adolescents: The UP&DOWN study. *Nutr Metab Cardiovasc Dis* 2016, 26:1011-1019.

Figures

Figure 1

Cardiometabolic risk z-score by fat tertiles categories (a), cardiorespiratory fitness (b) and muscular strength (c). * p<0.05 between high and low tertiles. In subjects with low fat, there was no difference in the CMR risk as muscle strength tertiles; whereas in children with high fat, higher muscle strength showed lower CMR compared to other groups with lower muscle strength.