

Serum Albumin and Activities of Daily Living in Chinese Centenarians: A Cross-Sectional Study

Ke Han

Chinese PLA General Hospital

Shengshu Wang

Chinese PLA General Hospital

Wangping Jia

Chinese PLA General Hospital

Wenzhe Cao

Chinese PLA General Hospital

Miao Liu

Chinese PLA General Hospital

Shanshan Yang

Chinese PLA General Hospital

Jianhua Wang

Chinese PLA General Hospital

Yao He (✉ heyao301@163.com)

Research article

Keywords: activities of daily living, serum albumin, centenarians, prevalence, China

Posted Date: January 9th, 2020

DOI: <https://doi.org/10.21203/rs.2.13830/v2>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on June 30th, 2020. See the published version at <https://doi.org/10.1186/s12877-020-01631-7>.

Abstract

Background Objective deterioration in activities of daily living (ADL) exists in older people, particularly grows with age. While as the criterion standard of positive aging and longevity, little information that focuses on centenarians is available. The objective of this study is to examine the relationship between serum albumin and activities of daily living among centenarians in long-lived areas. Methods design—population-based cross-sectional study; setting—a complete sample study in Hainan, the longest-lived areas of China; subjects—Chinese older adults aged 100 and above (N=1002); measurements—serum albumin, basic and instrumental activities of daily living (BADL and IADL). Results Of 1002 participants included in the analysis 287(28.64%) were BADL disability, and 648(64.67%) were IADL disability. The median serum albumin level was 38.5 g/L (interquartile range, 36.2-41.3). In multivariable analyses controlling for socio-demographic characteristics, lifestyle characteristics and other influential factors, low albumin in the bottom quartile were associated with BADL disability (OR=5.994;95% CI: 3.542-10.143) and IADL disability (OR=4.398; 95% CI, 2.661-7.269). Serum albumin concentration showed a significant inverse linear relationship with ADL disability (P for trend <0.001). In sex-stratified analyses, the association was more pronounced in women. And we found a nonlinear relationship between serum albumin and IADL disability in female centenarians (P=0.025) from restricted cubic splines. Conclusions Low levels of albumin were associated with the decline in ADL function in centenarians, and this association was gender-specific and more prominent in the female.

Background

The global population is aging. Worldwide, the proportion of the population aged 60 years and older increased from 9.2% in 1990 to 11.7% in 2013 and will continue to increase, and account for 21.1% of the world's population by 2050 [1]. The population of very old adults aged 85 years and older is increasing at a faster rate than other groups; particularly, the number of centenarians has increased exponentially worldwide [2]. Centenarians are said to represent the criterion standard of positive aging and longevity [3]. Basic research regarding longevity in the population has arisen as an essential area in the field of gerontology and aims to solve the problems imposed by population aging on health and social security systems.

Activities of daily living (ADLs) are a core set of essential activities for individuals to live independently and is an important index that determines the quality of life and life expectancy [4]. An increase in the number of centenarians may indicate a higher proportion of disabilities and poor health in the older population, and eventually increase the burden of medical care and treatment [5]. Objective deterioration of functional status exists among centenarians, most of whom depend on help from family or professional aides in their ADLs [6]. Many factors influence the impairment of ADLs in older adults, with nutritional biomarkers being significant [7].

Serum albumin (Alb) is the most abundant plasma protein in the human body with critical biological functions. Among older people, hypoalbuminemia is a prognostic indicator of various complications,

disability, and mortality [8-10]. Age increment leads to lower Alb levels and a higher risk of hypoalbuminemia [11]. Given the potentially unique pattern of vulnerability, centenarians deserve more attention.

Previous studies have shown that a strong relationship exists between Alb level and ADL in the elderly population [8, 10-19]. A few of them have focused on the oldest old populations [15-17]. Only one study of 420 persons aged ≥ 95 years provided data regarding Chinese longevous persons [16]. Further studies with large samples comprising both community-dwelling and institutionalized centenarians are required to understand the association clearly.

Therefore, we assessed the distribution characteristics of Alb levels and ADL disability and investigated the association between Alb and ADL impairment among centenarians using cross-sectional data obtained from a complete sample of a centenarian cohort from Chinese longest-lived areas.

Methods

Study population

We assessed cross-sectional data collected from 2014–2016 in the China Hainan Centenarian Cohort Study (CHCCS) [20], a longitudinal observational study that targeted the population aged 100 years or older in Hainan, China. The CHCCS is based on a complete sample of the Hainan community-dwelling and institutionalized centenarian population conducted by an interdisciplinary research team. The study protocol was approved by the institutional review board of the Chinese People's Liberation Army General Hospital (Beijing, China). All study participants provided written informed consent prior to enrollment.

According to the household register provided by the Civil Affairs Bureau in 2014, there were 1811 living centenarians in total. After a rigorous age validation process, we established contact with 1473 eligible individuals. We also excluded individuals who were not conscious and could not perform ADLs or participate in the questionnaire interview, physical health examination, and blood tests. Finally, 1002 centenarians aged 100–116 years were enrolled during recruitment and baseline data were collected between July 2014 and December 2016.

Activities of daily living

We conducted face-to-face interviews to assess the ability to perform ADL and inspections through relatives or caregivers to ensure effective and reliable investigations. ADLs consist of two components: basic ADLs (BADLs) and instrumental ADLs (IADLs). The Barthel index (BI) and Lawton scale were used to evaluate BADLs and IADLs, respectively.

The BI was developed in 1965 and later modified as a scoring technique that measures the patient's performance in 10 activities of daily life (eating, grooming, bathing, dressing, bowel and bladder care, toilet use, ambulation, transfers, and stair climbing) [21, 22]. Each item is scored 0–5 points, 0–10 points, or 0–15 points according to different evaluation results and summed to a total of 0–100 points. The total

BI score is used to classify the individuals' level of dependence as total (0–20), severe (21–60), moderate (61–90), or slight (91–99). A total score of 100 indicates that the patient is independent of assistance from others [23]. In this study, we defined BADL disability as BI \leq 60 points [24].

The Lawton IADL scale is an appropriate instrument used to assess independent living skills that are considered more complex than the BADLs (using the phone, shopping, preparing food, housekeeping, doing laundry, using transportation, handling medications, and handling finances) [25, 26]. Each skill measured by the scale requires some degree of both cognitive and physical function [27]. It takes 10–15 minutes to complete with a summary score ranging from 0–8. The total score of the scale sorts IADL dependence into 4 levels: severe (\leq 2), moderate (3–5), slight (6–7), and none (8). In this study, we defined IADL disability as a Lawton IADL score \leq 2.

Serum albumin

Experienced nurses collected 8 ml of blood from each participant in a fasting state using four vacutainer tubes (2 ml each). All blood samples were transported to the clinical laboratory in a chilled transport container (4°C) and assayed within 4 hours. Comprehensive metabolic panel examinations were performed using a fully automatic biochemical autoanalyzer (Cobas 8000; Roche Products Ltd, Basel, Switzerland). Alb level was categorized into two groups according to the cutoff point of hypoalbuminemia (35 g/L) [28].

Covariates

For the interview and examination, participants were visited at their residences or nearby clinics by a well-trained interdisciplinary medical group. Using a face-to-face interview in the appropriate regional dialect, the interviewers recorded detailed information on a standardized structured questionnaire. Socio-demographic characteristics assessed included age, gender, education, ethnicity, marital status, educational level, and type of residence. We also considered lifestyle characteristics such as smoking status, alcohol drinking status, and weekly exercise. Standing height and weight were measured using a manufactured instrument, and the body mass index (BMI) was calculated by dividing the body weight by the height (in kg/m²). Participants with a BMI <18.5 kg/m² were categorized as underweight [29]. Respondents were asked to report whether they had been diagnosed and treated by a doctor for any specific medical conditions. The presence of heart disease or stroke was self-reported. The presence of hypertension was defined by a self-report of high blood pressure, and/or sitting systolic blood pressure >140 mmHg and/or diastolic blood pressure >90 mmHg [30]. Similarly, diabetes was defined by self-report and/or a fasting blood glucose concentration of \geq 7.0 mmol/L [31]. Impaired renal function was defined by self-report and/or a glomerular filtration rate <60 ml·min⁻¹/1.73 m² [32].

Statistical analysis

The results are presented as medians and interquartile ranges or means \pm standard deviations for continuous variables, and as n (%) for categorical variables. Differences among groups were evaluated

using the Mann–Whitney *U* test or the chi-squared test, as appropriate. Multivariable logistic regression models were constructed to evaluate the associations between Alb and BADL disability and IADL disability. ADL disability (BI \leq 60) and IADL disability (Lawton IADL score \leq 2) were used as outcomes. Alb level considered dichotomously (35 g/L as the cutoff point) was used as the independent variable. Also, stratified analyses were performed to assess the association between Alb and ADL disability among different sexes. In the multivariable analyses, we considered socio-demographic characteristics, lifestyles, morbidity, and other influential factors including BMI classification, hemoglobin (Hb), total cholesterol (TC) level, and serum 25-hydroxyvitamin D (25OHD) as covariates. All statistical analyses were performed using SPSS Statistics version 24.0 (IBM Corporation, Armonk, NY, USA). All tests were two-sided and performed at a 5% significance level.

Results

Baseline characteristics

A total of 1002 centenarians were included in this study, including 822 women, who accounted for 82.04% of the total population. The ages of the participants ranged from 100–116 years, with a median age of 102 years (interquartile range [IQR], 101–104). The participants were mainly of Han ethnicity, illiterate, widowed/divorced/unmarried, and living with their families. The proportions of participants who smoked, drank alcohol, and exercised weekly were all lower than 20%. The median Alb level was 38.5 g/L (IQR, 36.2–41.3) for the entire sample. Participants who had a lower level of Alb had lower levels of Hb, TC, and 25OHD, and a lower prevalence of hypertension (Table 1).

Prevalence of ADL disability

The higher Alb group had a significantly lower rate of BADL and IADL disability than did the lower Alb group (Table 1). The prevalence of BADL disability was 28.64%, and that of IADL disability was 64.67%. Only 19 individuals were completely independent regarding IADL, and accounted for 1.9% of the population.

Association of serum albumin level with BADL disability and IADL disability

Multivariable logistic regression analyses revealed a significant association between Alb level and ADL disability. In model 3 (which controlled for socio-demographic characteristics, lifestyle characteristics, morbidities, and other covariates), low Alb level (<35 g/L) was found to be significantly associated with ADL disability (BADL: odds ratio [OR]=3.119, 95% confidence interval [CI]: 2.089–4.656; IADL: OR=3.687, 95% CI: 2.187–6.213). The effect was weakened but in the same direction when the covariates were gradually incorporated (Table 2).

Stratified analyses by sex

Among male centenarians, all logistic regression models suggested that the association between Alb and BADL disability was no more significant in the comparison between the two Alb level groups ($P>0.05$).

Regarding IADL disability among male adults, we only found a significant relationship in the unadjusted model (OR=2.606, 95% CI: 1.164–5.834). Among female centenarians, the associations were significant in all models, and the ORs were higher than those calculated for the entire population (Table 3).

Discussion

Prior to our study, only one study had assessed the association between Alb level and ADL among Chinese centenarians [17]. Our data were derived from the 2014–2016 cross-sectional data of the CHCCS, which included a complete sample of a large population in Hainan province. Hainan province has one of the highest life expectancies (76.3 years in 2010) and the highest proportion of centenarians (18.75/100 000) in China [33]. Therefore, our study may have some implications for positive aging and longevity.

As a clinical routinely-tested indicator and modifiable risk-related factor, Alb is important and imperative for providing potential clues for public interventions and long-term clinical care. Among 1002 Hainan centenarians assessed in this study, the average Alb level was 38.5 g/L (IQR, 36.2–41.3). This is lower than that identified in another Chinese longevity cohort with an average age of 97 (IQR, 96–98) years, which showed a mean albumin level of 42.4 ± 4.6 g/L [9]. The difference may be related to the age distribution characteristics of the subjects and differences in laboratory testing. A previous study showed that albumin concentration is related to aging and gradually decreases by 0.08–0.17 g/L per year, with levels tending to decline faster among males than among females [10]. In a study that included 95 Japanese centenarians, the mean Alb concentration was 36.0 ± 4.0 g/L among women and 36.0 ± 5.0 g/L among men [17]. The Alb concentration among centenarians was close to the lower limit of the adult reference value (35.0–50.0 g/L). Even in the high Alb group (with levels >35.0 g/L), a significant proportion of the elderly was ADL disabled. Thus, we have reason to doubt the utility of using this traditional reference in the very old population, which is in line with the findings of Kuzuya et al [34]. There is a need to establish a reference standard for healthy elderly populations, not only of Alb but also other significant serological indicators.

In this study, 287 (28.64%) centenarians were identified as having BADL disability. The prevalence of BADL disability was lower than that reported among older people and centenarians in similar studies regarding the population aged ≥ 90 years in other countries [35, 36]. Approximately 50% of participants were classified as having BADL disability in the baseline data of a local longevity cohort study, which investigated 433 long-lived individuals aged ≥ 95 years [9]. In the Chinese Longitudinal Healthy Longevity Survey, a nationwide survey of a randomly-selected half of the counties and cities in 22 of the 31 provinces, covering about 85% of the total population of China, 37.1% of the sample had BADL disability at the baseline [37, 38]. One possible reason for the discrepancy is that diseases or factors of early death are delayed or absent among centenarians. Comparatively, long-lived individuals tend to live in good functional states. Besides, different versions of rating scales with different scoring criteria were used, and different definitions of disability were used among the studies.

BADL functions are essential for an individual's self-care, whereas IADL functions are more concerned with higher-level skills in specific situations of social living. In our study, only 19 (1.9%) individuals had complete IADL independence according to the total score of the Lawton IADL scale. This means that almost every centenarian had lost at least one aspect of advanced functional independence necessary for daily living. The combination of all IADL items explains a larger portion of the difference in physical and psychological dimensions of quality of life compared to IADL items [39]. The performance of IADL in this sample is worse than that of the younger older population reported in similar studies [39-41]. This may be due to the fact that the incidence rate of ADL disability increases with age [42]. Advanced functions in daily life are important aspects of mental health and quality of life among older people. Thus, there is a need to pay attention to the impact of such a poor IADL in the very old.

The main finding of our study was that low levels of Alb were associated with ADL disability. Similar results have been reported in previous cross-sectional studies [12, 13, 16] and cohort studies [14, 18]. However, a few studies have reported negative results [43]. Alb plays an essential physiological role in the maintenance of health, including maintaining plasma colloid osmotic pressure and transporting endogenous substances and exogenous drugs. Its correlation with ADL may be attributable to the following hypotheses. As an essential indicator of nutritional status, Alb has great prognostic significance in protein-energy malnutrition. Also, it is a significant marker of acute and chronic inflammatory responses. Both hypoalbuminemia [19], inflammation [44], and malnutrition [45] are associated with reduced muscle mass (sarcopenia) in older adults. The positive association of sarcopenia with falls and fractures in older adults [46] is exact, which would contribute to the decline of ADL. Moreover, Alb plays a vital role in the pathogenesis of organ or system dysfunction/failure [47]. Albumin-associated inflammatory state and the associated poor acute pathological recovery might explain the ADL disability. On the other hand, IADL disability is a multidimensional construct that includes physical and cognitive domains. Significant cognitive decline is typical among older people [48] and is strongly associated with difficulties in IADL [49]. Thus, the severe deterioration of IADL among centenarians is explainable. The mechanism and causal link between Alb and ADL are not precise and need to be assessed in further studies.

Sex differences existed in the association in our study. The negative results obtained among men are similar to those reported by Liu et al., who enrolled 97 extremely longevous individuals [16] and Okamura et al. who studied 797 younger old men [18]. The crude mortality rate of elderly males is higher than that of females [50]. This may lead to the earlier death of the older men with severe impairment of ADL status and mask the relationship between Alb and impaired ADL. Given the very advanced level of age, the sample size of our study is relatively sufficient. However, we still could not determine if the results obtained in the male group are due to this fact or the influence of the sample size. Therefore, more extensive research has to be conducted to understand the relationship entirely.

Several limitations in this study deserve mention. First, due to the cross-sectional design of the study, causal inference is limited. A longitudinal survey will further clarify the clinical significance and predictive effectiveness of Alb in the older population. Second, we excluded some individuals who were unable to

participate in interviews and physical examinations before the investigation. The excluded subjects may be more likely in a low level of physical and cognitive function, which may lead to a certain underestimation of the prevalence of ADL disability. Third, the self-reported results of the questionnaire may be biased toward older people with poor cognitive function. Fourth, due to the lack of systematic nutritional assessment, we were unable to verify the exact association between Alb and nutritional status among centenarians.

Conclusions

This study is the first to report on the levels of Alb and prevalence of ADL disability and to explore their association among Chinese centenarians. In this study, the concentration of Alb was low and the prevalences of BADL disability and IADL disability were 28.64% and 64.67%, respectively. Low levels of albumin were associated with a decline in ADL function among centenarians, and this association was gender-specific and more prominent among females. Paying attention to Alb levels among the elderly and properly caring and treating elderly persons with low Alb levels may significantly improve their physical function. However, there is a need to validate whether albumin-deficient interventions are ideal strategies for preventing disability development or improving the independence of older persons through appropriately designed randomized and controlled trials.

Abbreviations

ADL: Activities of daily living; BADL: basic activities of daily living; IADL: instrumental activities of daily living; Alb: Serum albumin; CHCCS: China Hainan Centenarian Cohort Study; BMI: body mass index; Hb: hemoglobin; TC: total cholesterol; 25HOD: serum 25-hydroxyvitamin D; IQR: interquartile range; 95% CI: 95% confidence intervals.

Declarations

Ethics approval and consent to participate

The study obtained ethics approval from the Ethics Committee of the Hainan branch of the Chinese People's Liberation Army General Hospital (No. 301hn11201601). All participants were informed about the study goals and signed informed consent.

Consent for publication

Not applicable.

Availability of data and material

Data can be obtained from the corresponding author upon reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

The study was supported by Beijing Nova Program (Z181100006218085), Opening Foundation of State Key Laboratory of Kidney Diseases (KF-01-115), Opening Foundation of National Clinical Research Center of Geriatrics (NCRCG-PLAGH-2017017), National Natural Science Foundation of China (81773502, 81703285, 81703308), Medical Big Data Fund of Chinese PLA General Hospital (2018MBD-029).

Authors' contributions

KH, SSW, and YH conceived and designed the project; KH, SSW, ML, SSY, JHW, WPJ, WZC, and YH collected the data; KH drafted and revised the paper; KH, SSW, WPJ, and WZC conducted the data analysis; ML, SSY, JHW, and YH supervised the analyses and suggested revisions of the paper. ML, SSY, JHW provided administrative, technical, or material support; All the authors have read and approved the final manuscript.

Acknowledgements

The authors wish to express gratitude to all the participants for their cooperation in this study.

References

1. United Nations, Department of Economic and Social Affairs, Population Division (2013). **World Population Ageing 2013**. ST/ESA/SER.A/348.
2. Teixeira L, Araújo L, Jopp D, Ribeiro O: **Centenarians in Europe**. *Maturitas* 2017, **104**:90-95.
3. Perls TT, Silver MH: **Living to 100 : lessons in living to your maximum potential at any age**. New York: Basic Books; 1999.
4. Covinsky K: **Aging, arthritis, and disability**. *Arthritis and rheumatism* 2006, **55**(2):175-176.
5. Andersen-Ranberg K, Schroll M, Jeune B: **Healthy centenarians do not exist, but autonomous centenarians do: a population-based study of morbidity among Danish centenarians**. *Journal of the American Geriatrics Society* 2001, **49**(7):900-908.
6. Richmond RL, Law J, Kay-Lambkin F: **Physical, mental, and cognitive function in a convenience sample of centenarians in Australia**. *Journal of the American Geriatrics Society* 2011, **59**(6):1080-1086.
7. Maeda K, Koga T, Akagi J: **Nutritional variables predict chances of returning home and activities of daily living in post-acute geriatric care**. *Clin Interv Aging* 2018, **13**:151-157.
8. Corti MC, Guralnik JM, Salive ME, Sorkin JD: **Serum albumin level and physical disability as predictors of mortality in older persons**. *Jama* 1994, **272**(13):1036-1042.
9. Liu Z, Zhong G, Li S, Deng W, Zhang Y, Qian D, Jin L, Wang X: **Use of serum albumin and activities of daily living to predict mortality in long-lived individuals over 95 years of age: a population-based**

- study.** *Age (Dordrecht, Netherlands)* 2015, **37**(4):9809.
10. Cabrerizo S, Cuadras D, Gomez-Busto F, Artaza-Artabe I, Marin-Ciancas F, Malafarina V: **Serum albumin and health in older people: Review and meta analysis.** *Maturitas* 2015, **81**(1):17-27.
 11. Salive ME, Corconi-Huntley J, Phillips CL, Guralnik JM, Cohen HJ, Ostfeld AM, Wallace RB: **Serum albumin in older persons: relationship with age and health status.** *Journal of clinical epidemiology* 1992, **45**(3):213-221.
 12. Aung KC, Feng L, Yap KB, Sitoh YY, Leong IY, Ng TP: **Serum albumin and hemoglobin are associated with physical function in community-living older persons in Singapore.** *J Nutr Health Aging* 2011, **15**(10):877-882.
 13. Onem Y, Terekeci H, Kucukardali Y, Sahan B, Solmazgul E, Senol MG, Nalbant S, Sayan O, Top C, Oktenli C: **Albumin, hemoglobin, body mass index, cognitive and functional performance in elderly persons living in nursing homes.** *Archives of gerontology and geriatrics* 2010, **50**(1):56-59.
 14. Kitamura K, Nakamura K, Nishiwaki T, Ueno K, Nakazawa A, Hasegawa M: **Determination of whether the association between serum albumin and activities of daily living in frail elderly people is causal.** *Environmental health and preventive medicine* 2012, **17**(2):164-168.
 15. Gondo Y, Hirose N, Arai Y, Inagaki H, Masui Y, Yamamura K, Shimizu K-i, Takayama M, Ebihara Y, Nakazawa S *et al.*: **Functional status of centenarians in Tokyo, Japan: developing better phenotypes of exceptional longevity.** *The journals of gerontology Series A, Biological sciences and medical sciences* 2006, **61**(3):305-310.
 16. Liu Z, Wang Y, Huang J, Chu X, Qian D, Wang Z, Sun X, Chen F, Xu J, Li S *et al.*: **Blood biomarkers and functional disability among extremely longevous individuals: a population-based study.** *The journals of gerontology Series A, Biological sciences and medical sciences* 2015, **70**(5):623-627.
 17. Nozaki H, Nohara Y, Ashitomi I, Zukeran R, Inafuku T, Akisaka M, Suzuki M: **Serum albumin level and activities of daily living in centenarians.** *Nihon Ronen Igakkai zasshi Japanese journal of geriatrics* 1998, **35**(10):741-747.
 18. Okamura T, Hayakawa T, Hozawa A, Kadowaki T, Murakami Y, Kita Y, Abbott RD, Okayama A, Ueshima H: **Lower levels of serum albumin and total cholesterol associated with decline in activities of daily living and excess mortality in a 12-year cohort study of elderly Japanese.** *Journal of the American Geriatrics Society* 2008, **56**(3):529-535.
 19. Baumgartner RN, Koehler KM, Romero L, Garry PJ: **Serum albumin is associated with skeletal muscle in elderly men and women.** *Am J Clin Nutr* 1996, **64**(4):552-558.
 20. He Y, Zhao Y, Yao Y, Yang S, Li J, Liu M, Chen X, Wang J, Zhu Q, Li X *et al.*: **Cohort Profile: The China Hainan Centenarian Cohort Study (CHCCS).** *Int J Epidemiol* 2018.
 21. Mahoney FI, Barthel DW: **FUNCTIONAL EVALUATION: THE BARTHEL INDEX.** *Maryland state medical journal* 1965, **14**:61-65.
 22. Wade DT, Collin C: **The Barthel ADL Index: A standard measure of physical disability?** *International disability studies* 1988, **10**(2):64-67.

23. Shah S, Vanclay F, Cooper B: **Improving the sensitivity of the Barthel Index for stroke rehabilitation.** *Journal of clinical epidemiology* 1989, **42**(8):703-709.
24. Chivite D, Formiga F, Corbella X, Conde-Martel A, Aramburu Ó, Carrera M, Dávila MF, Pérez-Silvestre J, Manzano L, Montero-Pérez-Barquero M: **Basal functional status predicts one-year mortality after a heart failure hospitalization in elderly patients - The RICA prospective study.** *Int J Cardiol* 2018, **254**:182-188.
25. Lawton MP, Brody EM: **Assessment of older people: self-maintaining and instrumental activities of daily living.** *The Gerontologist* 1969, **9**(3):179-186.
26. Graf C: **The Lawton instrumental activities of daily living scale.** *The American journal of nursing* 2008, **108**(4):52-62; quiz 62-53.
27. Ng TP, Niti M, Chiam PC, Kua EH: **Physical and cognitive domains of the Instrumental Activities of Daily Living: validation in a multiethnic population of Asian older adults.** *The journals of gerontology Series A, Biological sciences and medical sciences* 2006, **61**(7):726-735.
28. Gatta A, Verardo A, Bolognesi M: **Hypoalbuminemia.** *Internal and emergency medicine* 2012, **7 Suppl 3**:S193-199.
29. He Y, Lam TH, Jiang B, Li LS, Sun DL, Wu L, Liu M, Yang SS, Wang YY, Tobias DK *et al*: **Changes in BMI before and during economic development and subsequent risk of cardiovascular disease and total mortality: a 35-year follow-up study in China.** *Diabetes care* 2014, **37**(9):2540-2547.
30. Williams B, Mancia G, Spiering W, Agabiti Rosei E, Azizi M, Burnier M, Clement DL, Coca A, de Simone G, Dominiczak A *et al*: **2018 ESC/ESH Guidelines for the management of arterial hypertension.** *European Heart Journal* 2018, **39**(33):3021-3104.
31. **Diagnosis and Classification of Diabetes Mellitus.** 2010, **33**(Supplement 1):S62-S69.
32. Coresh J, Astor BC, Greene T, Eknoyan G, Levey AS: **Prevalence of chronic kidney disease and decreased kidney function in the adult US population: Third National Health and Nutrition Examination Survey.** *American journal of kidney diseases : the official journal of the National Kidney Foundation* 2003, **41**(1):1-12.
33. Wang L, Li Y, Li H, Holdaway J, Hao Z, Wang W, Krafft T: **Regional aging and longevity characteristics in China.** *Archives of gerontology and geriatrics* 2016, **67**.
34. Kuzuya M, Izawa S, Enoki H, Okada K, Iguchi A: **Is serum albumin a good marker for malnutrition in the physically impaired elderly?** *Clinical nutrition (Edinburgh, Scotland)* 2007, **26**(1):84-90.
35. Berlau DJ, Corrada MM, Peltz CB, Kawas CH: **Disability in the oldest-old: incidence and risk factors in the 90+ study.** *The American journal of geriatric psychiatry : official journal of the American Association for Geriatric Psychiatry* 2012, **20**(2):159-168.
36. Herr M, Arvieu JJ, Robine JM, Ankri J: **Health, frailty and disability after ninety: Results of an observational study in France.** *Archives of Gerontology and Geriatrics Archives of Gerontology and Geriatrics* 2016, **66**:166-175.
37. Zeng Y, Feng Q, Hesketh T, Christensen K, Vaupel JW: **Survival, disabilities in activities of daily living, and physical and cognitive functioning among the oldest-old in China: a cohort study.** *Lancet* 2017,

- 389(10079):1619-1629.
38. Li LW, Zhang J, Liang J: **Health among the oldest-old in China: which living arrangements make a difference?** *Social science & medicine (1982)* 2009, **68**(2):220-227.
 39. Gobbens RJ: **Associations of ADL and IADL disability with physical and mental dimensions of quality of life in people aged 75 years and older.** *PeerJ* 2018, **6**:e5425.
 40. Park J, Lee YJ: **Patterns of instrumental activities of daily living and association with predictors among community-dwelling older women: A latent class analysis.** *BMC Geriatr* 2017, **17**(1):158.
 41. Henning-Smith C, Shippee T, Capistrant B: **Later-Life Disability in Environmental Context: Why Living Arrangements Matter.** *The Gerontologist* 2018, **58**(5):853-862.
 42. Sjölund BM, Wimo A, Engström M, von Strauss E: **Incidence of ADL Disability in Older Persons, Physical Activities as a Protective Factor and the Need for Informal and Formal Care—Results from the SNAC-N Project.** *PLoS ONE* 2015, **10**(9):e0138901.
 43. Schalk BWM, Visser M, Deeg DJH, Bouter LM: **Lower levels of serum albumin and total cholesterol and future decline in functional performance in older persons: the Longitudinal Aging Study Amsterdam.** *Age and ageing* 2004, **33**(3):266-272.
 44. van Atteveld VA, Van Ancum JM, Reijnierse EM, Trappenburg MC, Meskers CGM, Maier AB: **Erythrocyte sedimentation rate and albumin as markers of inflammation are associated with measures of sarcopenia: a cross-sectional study.** *BMC geriatrics* 2019, **19**(1):233.
 45. Pierik VD, Meskers CGM, Van Ancum JM, Numans ST, Verlaan S, Scheerman K, Kruizinga RC, Maier AB: **High risk of malnutrition is associated with low muscle mass in older hospitalized patients - a prospective cohort study.** *BMC geriatrics* 2017, **17**(1):118.
 46. Yeung SSY, Reijnierse EM, Pham VK, Trappenburg MC, Lim WK, Meskers CGM, Maier AB: **Sarcopenia and its association with falls and fractures in older adults: A systematic review and meta-analysis.** *J Cachexia Sarcopenia Muscle* 2019, **10**(3):485-500.
 47. Arroyo V, García-Martínez R, Salvatella X: **Human serum albumin, systemic inflammation, and cirrhosis.** *Journal of hepatology* 2014, **61**(2):396-407.
 48. Hendrie HC, Albert MS, Butters MA, Gao S, Knopman DS, Launer LJ, Yaffe K, Cuthbert BN, Edwards E, Wagster MV: **The NIH Cognitive and Emotional Health Project. Report of the Critical Evaluation Study Committee.** *Alzheimer's & dementia : the journal of the Alzheimer's Association* 2006, **2**(1):12-32.
 49. Connolly D, Garvey J, McKee G: **Factors associated with ADL/IADL disability in community dwelling older adults in the Irish longitudinal study on ageing (TILDA).** *Disability and rehabilitation* 2017, **39**(8):809-816.
 50. Jacobs JM, Cohen A, Ein-Mor E, Stessman J: **Gender differences in survival in old age.** *Rejuvenation Res* 2014, **17**(6):499-506.

Tables

Table 1. Prevalence of risk factors for ADL disability at two levels of serum albumin among 1002 centenarians

Characteristics	Total(n=1002)	Serum albumin		P value
		<35g/L(n=174)	≥35g/L(n=828)	
Age, median(IQR),y	102.0(101.0-104.0)	103.0(101.0-105.0)	102.0(101.0-104.0)	0.003
Female,%	822(82.0)	140(80.5)	682(82.4)	0.551
Han ethnicity,%	883(88.1)	156(89.7)	727(87.8)	0.492
Illiterate,%	915(91.3)	162(93.1)	753(91.1)	0.325
Married,%	100(10.0)	16(9.2)	84(10.1)	0.704
Living with families,%	863(86.1)	159(91.4)	704(85.0)	0.027
Current smoker,%	35(3.5)	6(3.5)	29(3.5)	0.963
Current alcohol drinker,%	99(9.9)	14(8.1)	85(10.3)	0.002
Weekly exerciser,%	129(12.9)	11(6.3)	118(14.3)	0.005
BMI<18.5kg/m ² ,%	575(57.4)	118(67.8)	457(55.2)	0.005
Alb, median(IQR),g/L	38.5(36.2-41.3)	32.8(30.6-34.2)	39.4(37.7-41.7)	□0.001
Hb, median(IQR), g/L	103.0(114.0-123.0)	102.0(94.0-113.0)	116.9(107.0-125.0)	□0.001
TC, median(IQR), mmol/L	4.60(4.1-5.3)	4.0(3.4-4.6)	4.66(4.2-5.3)	□0.001
25OHD, median(IQR), ng/mL	21.6(16.9-27.8)	20.9(14.0-26.5)	21.6(17.4-28.1)	0.001
Diabetes,%	96(9.6)	18(10.3)	78(9.4)	0.706
Hypertension,%	757(75.6)	121(69.5)	636(76.8)	0.042
Heart disease,%	41(4.1)	5(2.9)	36(4.3)	0.255
Stroke,%	22(2.2)	1(0.6)	21(2.5)	0.153
Renal function impaired,%	299(29.8)	56(32.2)	243(29.4)	0.457
BADL disability,%	287(28.6)	94(54.0)	193(23.3)	□0.001

IADL disability,%	648(64.7)	151(86.8)	497(60.0)	0.001
-------------------	-----------	-----------	-----------	-------

Abbreviations: ADL, activities of daily living; IQR, interquartile range; BMI, body mass index; Alb, serum albumin; Hb, hemoglobin; TC, total cholesterol; 25OHD, serum 25-hydroxyvitamin D.

Table 2. Association between serum albumin level and BADL/IADL disability ^{a,b}

	Categorized variables of Alb	
	≥35g/L(n=828)	<35g/L(n=174)
BADL disability		
Crude model	Ref.	3.866(2.755,5.425)
Model 1	Ref.	3.868(2.739,5.462)
Model 2	Ref.	3.180(2.140,4.725)
Model 3	Ref.	3.119(2.089,4.656)
IADL disability		
Crude model	Ref.	4.372(2.760,6.928)
Model 1	Ref.	4.259(2.658,6.825)
Model 2	Ref.	3.712(2.208,6.242)
Model 3	Ref.	3.687(2.187,6.213)

^a Crude model: None adjusted;

Model 1: Adjusted for age, gender;

Model 2: Adjusted for age, gender, ethnicity, marriage, educational levels, residence type, smoking status, alcohol drinking status, weekly exercise;

Model 3: Adjusted for variables in Model 2 plus BMI classification, hemoglobin, total cholesterol, serum 25-hydroxyvitamin D and hypertension.

^b Described as OR (95%CI).

Table 3. Sex-stratified analyses of the association between serum albumin level and BADL/IADL disability ^{a,b}


	Male		Female	
	Categorized variables of Alb ≥35g/L	<35g/L	Categorized variables of Alb ≥35g/L	<35g/L
BADL disability				
Crude model	Ref.	2.023(0.902,4.537) *	Ref.	4.538(3.104,6.634)
Model 1	Ref.	2.084(0.894,4.856) *	Ref.	4.472(3.048,6.563)
Model 2	Ref.	2.398(0.859,6.692) *	Ref.	3.364(2.168,5.219)
Model 3	Ref.	2.366(0.832,6.726) *	Ref.	3.291(2.104,5.149)
IADL disability				
Crude model	Ref.	2.606(1.164,5.834)	Ref.	5.834(3.229,10.540)
Model 1	Ref.	2.117(0.911,4.922) *	Ref.	5.544(3.051,10.076)
Model 2	Ref.	1.669(0.620,4.497) *	Ref.	4.836(2.542,9.198)
Model 3	Ref.	1.557(0.568,4.267) *	Ref.	4.695(2.467,8.936)

^a The adjusted covariates in models are the same as in Table 2 except for gender.

^b Described as OR (95%CI).

*P > 0.05

Figures



The distribution of BADL is different from that of IADL by chi-square test. (P<0.001)

Figure 1

ADL distribution according to the score of the Barthel index and Lawton Scale. The distribution of BADL is different from that of IADL by chi-square test. ($P < 0.001$)




Figure 2

Restricted cubic splines of the relationship between serum albumin and the risk of ADL disability. The y-axis represents the odds ratio, and the x-axis is serum albumin concentration. The solid line shows the odds ratio estimate and the shaded part the 95% confidence intervals. All adjusted for age, ethnicity, marriage, educational levels, residence type, smoking status, alcohol drinking status, weekly exercise, BMI classification, hemoglobin, total cholesterol, serum 25-hydroxyvitamin D and hypertension. A: total BADL disability; B: male BADL disability; C: female BADL disability; D: total IADL disability; E: male IADL disability; F: female IADL disability (P for nonlinear relationship = 0.025).