

The predictive value of D-dimer test for venous thromboembolism during puerperium in women age 35 or older: A prospective cohort study

Wen Hu

Zhejiang University School of Medicine

Dong Xu

Zhejiang University School of Medicine

Juan Li

Zhejiang University School of Medicine

Cheng Chen

Zhejiang University School of Medicine

Yuan Chen

Zhejiang University School of Medicine

Fangfang Xi

Zhejiang University School of Medicine

Feifei Zhou

Zhejiang University School of Medicine

Xiaohan Guo

Zhejiang University School of Medicine

Baihui Zhao (✉ zhaobh@zju.edu.cn)

Zhejiang University School of Medicine

Qiong Luo (✉ luoq@zju.edu.cn)

Zhejiang University School of Medicine <https://orcid.org/0000-0002-0782-4462>

Research

Keywords: D-dimer, Venous thromboembolism, women age at 35 years or older, Puerperium

Posted Date: July 24th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-45646/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 16th, 2020. See the published version at <https://doi.org/10.1186/s12959-020-00241-y>.

Abstract

Background

This study aimed to investigate the predictive value of the D-dimer for venous thromboembolism (VTE) events during puerperium of women age at 35 years or older, as well as to identify other risk factors associated with the occurrence of VTE.

Methods

It was a prospective observational cohort study, from January 2014 to December 2018, which involved 12451 women age 35 or older who delivered after 28 weeks of gestation at Women's Hospital of Zhejiang University, School of Medicine. The maternal and fetal demographic characteristics, pregnancy complications, imaging finding and results of laboratory test within postpartum 24 hours including D-dimer, platelet and fibrinogen were collected for analyses.

Results

30(2.4‰) women were identified as VTE, including 1 PE event and 29 DVT events. The receiver operating characteristic (ROC) curve analysis suggested the best cutoff point for D-dimer level within postpartum 24 hours of women age 35 or older was 5.545 mg/L, with a specificity of 70.0% and a sensitivity of 75.4%. Besides, there was no statistical correlation between platelets and VTE, as well as between fibrinogen and VTE. On multivariate analysis, D-dimer \geq 5.50 mg/L (OR = 5.874, 95%CI: 2.678–12.886) and emergency cesarean section (OR = 11.965, 95%CI: 2.732–52.401) were independently associated with VTE in puerperium of women age 35 or older.

Conclusions

We concluded that D-dimer \geq 5.50 mg/L was an independent predictor of VTE in puerperium with maternal age 35 or older and D-dimer testing was a necessary examination for perinatal women.

Background

Venous thromboembolism (VTE) remains one of the leading causes of maternal mortality [1], taking the place of postpartum hemorrhage, which has been highly prevented and treated. The incidence is 4 to 5 times higher among pregnant and postpartum women than that of non-pregnant women [2]. Pregnancy is an acquired and independent risk factor for the development of VTE. Many other risk factors have been linked to VTE, such as advanced maternal age, thrombophilia, cesarean section, obesity, and a personal or family history of VTE [3, 4]. In recent years, maternal age at childbirth continues to increase worldwide, particularly in China, as a consequence of the changes in fertility concept and the adjustment of the birth

policy. Increasing maternal age is associated with the increasing incidence of pregnancy complications, which together lead to the increasing incidence of VTE.

Among the screening tools for VTE, D-dimer testing has been proved its reliability in non-pregnant individuals by several studies, with high sensitivity, high negative predictive value and moderate specificity [5]. For pregnant women, D-dimer concentration increased progressively during the pregnancy and peaked at the first postpartum day [6]. Some studies proved the predictive value of D-dimer test for pregnant related VTE by raising the cutoff value or finding a higher D-dimer reference range. D-dimer also has been shown to increase by patient age [7]. However, there is a lack of research on the predictive value of D-dimer for VTE in the women age 35 or older.

Therefore, we designed a prospective observational study to identify the incidence and risk factors of VTE during the postpartum period in women age 35 or older. Furthermore, we investigated the predictive value of coagulation markers including D-dimer, platelet and fibrinogen, and attempted to determine a suitable threshold for the assessment in postpartum period of older mothers.

Methods

Patients

We initiated the study in January 1st 2014 and continued until December 31st 2018, and we prospectively collected the data of women age 35 or older who gave birth after 28 weeks of gestation at Women's Hospital of Zhejiang University, School of Medicine. Women used anticoagulant or anti-platelet drugs before delivery or with incomplete clinical data were excluded from this study (Figure 1). We set $\alpha=0.05$ and $\beta=0.10$ (power=0.90), the sample size of this study was larger than the estimated sample size according to the reported incidence of pregnancy-related VTE [8]. All clinical variables were recorded, including age, body mass index (BMI), pregnancy times, parity, number of births, gestational weeks of delivery, fetal position, mode of delivery, fetal birth weight, pregnancy complications, postpartum hemorrhage and predictive biomarkers within postpartum 24 hours including D-dimer, platelet and fibrinogen. All biomarker values were obtained from the same laboratory affiliated to the hospital.

Clinical diagnosis of VTE

Imaging evidence was confirmed as the diagnostic criteria for VTE. DVT was diagnosed by upper and lower extremity venous color Doppler ultrasound and/or computed tomographic (CT) venography, and PE was diagnosed by CT pulmonary angiography.

Imaging examinations were required if the following conditions were present: (1) with suspicious symptoms of VTE, including pain or tenderness when move limbs, swelling of the limbs, measurement of inconsistencies in the circumference of the bilateral limbs, or unexplained dyspnea, chest pain or cough; or (2) with multiple high risk factors, and the clinician considered that the probability for VTE was great. Anticoagulation and antithrombotic therapy would be applied immediately when imaging examination

indicated the diagnosis of VTE. All the VTE patients were told to follow up in the vascular department after discharge. Other women were followed up in the communities and would be accessed by vascular specialist in case of suspicious VTE symptoms (Figure 1).

Laboratory assays

We performed laboratory tests including platelet count, fibrinogen and D-dimer. The detection of platelet count was measured by impedance (XN9000; Sysmex, Kobe, Japan). The detection of fibrinogen was measured by the solidification (Stago-R, Paris, France). The detection of D-dimer was measured by the latex-enhanced immunoturbidimetry (Stago-R, Paris, France) (normal reference range for non-pregnant adults is less than 0.5 mg/L).

Based on receiver operating characteristic (ROC) curve analysis, the best cutoff point for D-dimer level within postpartum 24 hours was 5.545 mg/L (Figure 2). For convenience in clinical practice, the predefined cutoff value for dichotomized variables of D-dimer was set at 5.50 mg/L.

Statistical analyses

Data inputting and statistical analysis were performed in SPSS 22.0 (IBM Corporation, New York, USA). Continuous variables were described as means \pm standard deviation. The continuous variables were compared by Student's T test. The difference in the categorical variables was compared through Chi square test, Yate's correction of continuity or Fisher's exact test. Furthermore, to estimate the risk factors of VTE, the forward stepwise multiple logistic regression was performed. The associations between biomarkers and VTE were expressed as ROC curve analysis. Statistical significance was set at $p < 0.05$.

Results

12451 women were enrolled in this study after screening (Fig. 1). In our cohort, 30 (2.4‰) women were identified as VTE, including 1 PE event and 29 DVT events. The DVT events included: 5 women with bilateral DVT of lower extremity, 11 women with DVT of right lower extremity and 13 with DVT of left lower extremity. VTE occurred at median of 3.5 days postpartum (range: 2–15 days).

All the D-dimer test results of VTE patients in this study exceeded the upper limit of the reference value (0.5 mg/L). The ROC curve analysis showed the best cutoff point for D-dimer level within postpartum 24 hours of women age 35 or older was 5.545 mg/L, with a specificity of 70.0% and a sensitivity of 75.4%. When the cutoff value was set at 6.475 mg/L, the specificity could increase to 80.0%, but the sensitivity would decrease to 53.3%; when the cutoff value was set at 9.875 mg/L, the specificity could increase to 90.0%, but the sensitivity would decrease to 36.7%. The AUCs of fibrinogen and platelets were close to 0.5, indicating that there was no statistical correlation between them and VTE (Fig. 2).

Table 1 shows a comparison of maternal and fetal characteristics between VTE and non-VTE groups. Age, heights, gestational weeks of delivery, neonatal weight, fibrinogen and platelet count were not

significantly associated with VTE. The average D-dimer within postpartum 24 hours with maternal age ≥ 35 in VTE group was significantly higher than that of non-VTE group (8.91 vs. 4.55 mg/L, $P < 0.001$).

Table 1

Comparison of general characteristics between VTE and non-VTE delivery women age ≥ 35 ($x \pm SD$)

	VTE (n = 30)	Non-VTE (n = 12421)	t value	P value
Age (years)	38.10 \pm 2.23	37.31 \pm 2.29	1.893	.058
Height (cm)	161.13 \pm 5.76	160.57 \pm 4.71	0.650	.516
BMI before pregnancy	21.65 \pm 2.96	21.05 \pm 2.91	1.139	.255
BMI before delivery	26.70 \pm 2.80	26.50 \pm 2.99	0.363	.717
Gain weight during pregnancy (kg)	13.07 \pm 3.19	14.04 \pm 4.94	-1.067	.286
Gestational age at delivery (weeks)	37.83 \pm 2.26	37.99 \pm 1.95	-0.447	.655
Neonatal birth weight(g)	3114.67 \pm 558.19	3203.16 \pm 604.89	-0.800	.423
Laboratory test results within postpartum 24 hours				
D-dimer (mg/L)	8.91 \pm 6.16	4.55 \pm 4.31	3.870	.001
Fibrinogen (g/L)	4.37 \pm 0.85	4.63 \pm 0.81	-1.706	.088
Platelet count ($10^9/L$)	172.00 \pm 40.50	187.82 \pm 53.53	-1.590	.112

The risk factors predisposing to VTE in puerperium of women age ≥ 35 were analyzed in Table 2. Mode of delivery, scared uterus, and D-dimer ≥ 5.50 mg /L were significantly associated with VTE in puerperium of older mothers (Table 2).

Table 2
Risk factors predisposing to VTE in puerperium of women age ≥ 35

Risk factors	VTE(%) (n = 30)	Non-VTE(%) (n = 12421)	χ^2 value	P value
Previous obstetric history				
0	11(36.67)	2938(23.65)	2.804	.094
≥ 1	19(63.33)	9483 (76.35)		
Parity				
1	30(100.00)	12048(97.00)		1.000 ^a
≥ 2	0(0.00)	373(3.00)		
Fetal position of singleton cases				
Head	28(93.33)	11406(94.67)		.385 ^a
Breech	1(3.33)	506(4.20)		
Transverse	1(3.33)	136(1.13)		
Mode of delivery				
Vaginal delivery	2(6.67)	4786(38.53)	25.792	< .001
Emergency cesarean section	16(53.33)	2399(19.31)		
Elective caesarean section	12(40.00)	5236(42.15)		
In vitro fertilization (IVF)	0(0.00)	678(5.46)	0.834	.361
Scared uterus	20(66.67)	5612(45.18)	5.58	.018
Relative cephalopelvic disproportion	0(0.00)	164(1.32)		1.000 ^a
Placenta previa	3(10.00)	612(4.93)	0.738	.390
Adherent placenta	4(13.33)	610(4.91)	2.910	.088
Fetal growth restriction	0(0.00)	182(1.47)		1.000 ^a
Premature birth	5(16.67)	1557(12.54)	0.165	.684
Macrosomia (birthweight ≥ 4000 g)	1(3.33)	789(6.35)	0.092	.762
Premature rupture of membranes	7(23.33)	2184 (17.58)	0.682	.409
Fetal distress	5(16.67)	1566(12.61)	0.155	.694

^a evaluated by Fisher's exact test.

Risk factors	VTE(%) (n = 30)	Non-VTE(%) (n = 12421)	χ^2 value	P value
Intrauterine infection	1(3.33)	139(1.12)		.288 ^a
Postpartum hemorrhage	3(10.00)	458(3.69)	1.809	.179
Anemia	9(30.00)	2654(21.37)	1.327	.249
Intrahepatic cholestasis of pregnancy	3(10.00)	398(3.20)		.071 ^a
Gestational diabetes mellitus	3(10.00)	2951(23.76)	3.130	.077
Hypertensive disorders of pregnancy	3(10.00)	924(7.44)	0.034	.853
Cardiac insufficiency	0(0.00)	20(0.16)		1.000 ^a
Uterine rupture	0(0.00)	84(0.68)		1.000 ^a
D-dimer \geq 5.545 mg /L	21(70.00)	3057(24.61)	33.130	< .001
D-dimer \geq 6.475 mg /L	16(53.33)	2480(19.97)	20.789	< .001
D-dimer \geq 9.88 mg /L	11(36.67)	1242(10.00)	20.661	< .001
^a evaluated by Fisher's exact test.				

A multivariate model using forward stepwise regression was constructed to identify the risk-factors associated with VTE in puerperium. D-dimer \geq 5.50 mg /L and emergency cesarean section were independently associated with VTE in puerperium (Table 3).

Table 3
Multivariate logistic regression of VTE risk factors during puerperium of women age \geq 35

Risk factors	P value	OR	95% CI
Emergency cesarean section	.001	11.965	2.732–52.401
D-dimer \geq 5.545 mg /L	< .001	5.874	2.678–12.886

Discussion

VTE was reported 1.0-1.8/1000 in women during pregnancy and puerperium[8]. Our study showed that the rate of VTE during puerperium of women age \geq 35 (2.4‰) was higher than that of younger mothers, which supported age was a risk factor for VTE, which had been proved by many other studies[3, 9–11]. It was considered that more cases of DVT events occurred in the left lower extremity, which was related to the more serious venous stasis of the left lower extremity caused by the compression of the pregnant

uterus[12]. But in our study, the proportion of DVT in the left lower extremity was only a little higher than that in the right. This may be due to the limitation of the small sample size.

The diagnostic value of D-dimer for pregnant related VTE is not clear up to now. For pregnant women, D-dimer concentration increased progressively during the pregnancy and peaked at the first postpartum day[6]. Most healthy pregnant women have higher D-dimer values during pregnancy and puerperium than the normal reference range[13]. A prospective study showed that in the first trimester, 84% women had normal D-dimer, in the second 33%, and by the third trimester only 1%, which suggests that D-dimer has no practical diagnostic use of VTE if the threshold for abnormal is used[14]. Guidelines from Royal College of Obstetricians and Gynaecologists recommended that D-dimer testing should not be performed in the investigation of acute VTE in pregnancy[15]. Guidelines from American College of Obstetricians and Gynecologists also recommended that the rise of D-dimer cannot predict VTE reliably[14, 16]. However, there still some studies supporting to perform D-dimer test and provide a higher threshold to increase the specificity of D-dimer without reducing the sensitivity [14, 17–20]. Actually, D-dimer test is still being used by obstetricians. If the D-dimer level was abnormally high, the need for prophylactic use of low molecular weight heparin (LMWH) was decided by the doctors according to their own experiences.

However, up to now, there are few studies on the correlation between D-dimer and VTE in delivery women age ≥ 35 . The prevention of VTE in delivery women age ≥ 35 is particularly important because the risk of VTE increases with age. A variety of acquired prothrombotic risk factors (e.g., autoimmune disorders, diabetes and infection) also gradually develop with aging. Concurrently, aging is associated with a variety of coagulation and hemostasis changes in general population [21]. In our study, a large sample of older mothers was observed and we initially found the predictive value of D-dimer test for VTE during puerperium in women age ≥ 35 . Although the specificity and sensitivity are not particularly high, the ROC curve analysis offered an even higher threshold of D-dimer in delivery women age ≥ 35 . Multivariate analysis also indicated that D-dimer ≥ 5.50 mg/L was independently associated with VTE in puerperium of older mothers. Therefore, we think it is necessary to perform the D-dimer test within postpartum 24 hours of women age ≥ 35 .

Elevated D-dimer level was not the only criterion for high risk of VTE. Our study revealed emergency cesarean section was another important independent risk factors of VTE in puerperium of women age ≥ 35 , which was consistent with previous studies. A meta-analysis found that the risk of VTE was four fold greater following cesarean section than following vaginal delivery, and was greater following emergency cesarean section than following elective cesarean section [4]. Other studies also revealed several independent risk factors of VTE in puerperium such as higher BMI, thrombophilia, multiple pregnancy, gestational diabetes, premature birth, anemia, chorioamnionitis, in vitro fertilization with ovarian hyperstimulation, cardiac diseases and postpartum hemorrhage [9]. But we didn't find these risk factors because these factors may be age related and the objects of our study were older mothers while theirs were delivery women of all ages. Based on the results of this study, we recommend the use of LMWH to prevent VTE when the level of D-dimer was higher than 5.50 mg/L of older mother, or the delivery mode of the older mother was emergency cesarean section.

The limitation of this study is that the number of VTE cases is insufficient, but this is consistent with the incidence of pregnancy-related VTE. Other limitations of this study include the effects of choice bias, for all the women in this study were at our hospital, and loss to follow-up bias, for we could hardly get the data of follow-up in communities and vascular department. Furthermore, this study did not discuss the predictive effect of D-dimer test for VTE during pregnancy and 24 hours after postpartum, which need further study.

Conclusion

In summary, this study calculated that D-dimer ≥ 5.50 mg /L was an independent factor associated with VTE in puerperium of women age ≥ 35 , which confirmed the predictive value of D-dimer test for older delivery women. Another independent risk factor of VTE in puerperium of women age ≥ 35 was emergency cesarean section. We believe our study provides a new reliable evidence for clinicians to focus on the emphasis risk factors for VTE of older delivery women, which was expected to reduce the incidence of VTE.

List Of Abbreviation

VTE venous thromboembolism

ROC receiver operating characteristic

BMI body mass index

CT computed tomographic

PE pulmonary embolism

DVT deep venous thrombosis

AUC area under the curve

LMWH low molecular weight heparin

Declarations

Ethics approval and consent to participate

This paper is approved by Medical Ethics Committee of Women's Hospital, Zhejiang University School of Medicine. Informed consent was not obtained because no patient information was published herein.

Consent for publication

Not applicable

Availability of data and materials

Not applicable

Competing interests

The authors declare that they have no competing interests.

Funding

The paper is supported by Natural Science Foundation of Zhejiang province, China (No. LQ20H040008 and LY20H040009), Medical and Health Research Projects of Zhejiang Province (No.201244300) and Key R & D plan of Zhejiang Province (No. 2018CD3010).

Authors' contributions

WH and DX wrote the manuscript. BZ and QL made critical revisions of the manuscript for important intellectual content and contributed to the study concept, design and implementation. JL, CC, YC and FX were responsible for data collection, input and correction. FZ and XG contributed to data statistics and analysis. All the authors discussed the first draft of the paper and put forward suggestions for revision. All authors read and approved the final manuscript.

Acknowledgements

Not applicable

References

1. Abe K, Kuklina EV, Hooper WC, Callaghan WM. Venous thromboembolism as a cause of severe maternal morbidity and mortality in the United States. *SEMIN PERINATOL*. 2019;43:200–4.
2. Heit JA, Kobbervig CE, James AH, et al. Trends in the incidence of venous thromboembolism during pregnancy or postpartum: a 30-year population-based study. *ANN INTERN MED*. 2005;143:697–706.
3. James AH, Jamison MG, Brancazio LR, Myers ER. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *AM J OBSTET GYNECOL*. 2006;194:1311–5.
4. Blondon M, Casini A, Hoppe KK, et al. Risks of Venous Thromboembolism After Cesarean Sections: A Meta-Analysis. *CHEST*. 2016;150:572–96.
5. Johnson ED, Schell JC, Rodgers GM. The D-dimer assay. *AM J HEMATOL*. 2019;94:833–9.
6. Wang M, Lu S, Li S, Shen F. Reference intervals of D-dimer during the pregnancy and puerperium period on the STA-R evolution coagulation analyzer. *CLIN CHIM ACTA*. 2013;425:176–80.
7. Haase C, Joergensen M, Ellervik C, Joergensen MK, Bathum L. Age- and sex-dependent reference intervals for D-dimer: evidence for a marked increase by age. *THROMB RES*. 2013;132:676–80.

8. Meng K, Hu X, Peng X, Zhang Z. Incidence of venous thromboembolism during pregnancy and the puerperium: a systematic review and meta-analysis. *J Matern Fetal Neonatal Med.* 2015;28:245–53.
9. Galambosi PJ, Gissler M, Kaaja RJ, Ulander VM. Incidence and risk factors of venous thromboembolism during postpartum period: a population-based cohort-study. *Acta Obstet Gynecol Scand.* 2017;96:852–61.
10. Sharma S, Monga D. Venous thromboembolism during pregnancy and the post-partum period: incidence and risk factors in a large Victorian health service. *Aust N Z J Obstet Gynaecol.* 2008;48:44–9.
11. Jensen TB, Gerds TA, Gron R, et al. Risk factors for venous thromboembolism during pregnancy. *Pharmacoepidemiol Drug Saf.* 2013;22:1283–91.
12. James AH, Tapson VF, Goldhaber SZ. Thrombosis during pregnancy and the postpartum period. *AM J OBSTET GYNECOL.* 2005;193:216–9.
13. Reger B, Peterfalvi A, Litter I, et al. Challenges in the evaluation of D-dimer and fibrinogen levels in pregnant women. *THROMB RES.* 2013;131:e183–7.
14. Kovac M, Mikovic Z, Rakicevic L, et al. The use of D-dimer with new cutoff can be useful in diagnosis of venous thromboembolism in pregnancy. *Eur J Obstet Gynecol Reprod Biol.* 2010;148:27–30.
15. Gynaecologists RCOO, editor. *Thromboembolic Disease in Pregnancy and the Puerperium: Acute Management. Green Top Guideline No.37b.* London: Royal College of Obstetricians and Gynaecologists; 2015.
16. ACOG Practice Bulletin No. Thromboembolism in Pregnancy. *OBSTET GYNECOL.* 2018;196:132:e1–17.
17. Parilla BV, Fournogerakis R, Archer A, et al. Diagnosing Pulmonary Embolism in Pregnancy: Are Biomarkers and Clinical Predictive Models Useful? *AJP Rep.* 2016;6:e160–4.
18. Murphy N, Broadhurst DI, Khashan AS, et al. Gestation-specific D-dimer reference ranges: a cross-sectional study. *BJOG.* 2015;122:395–400.
19. Tang J, Lin Y, Mai H, et al. Meta-analysis of reference values of haemostatic markers during pregnancy and childbirth. *Taiwan J Obstet Gynecol.* 2019;58:29–35.
20. Xu D, Cai SP, Xu JW, Liang C, He J. [Study on the dynamic changes of D-dimer during pregnancy and early puerperium]. *Zhonghua Fu Chan Ke Za Zhi.* 2016;51:666–71.
21. Favaloro EJ, Franchini M, Lippi G. Aging hemostasis: changes to laboratory markers of hemostasis as we age - a narrative review. *SEMIN THROMB HEMOST.* 2014;40:621–33.

Figures




Figure 1

Flow chart of indicating the patients included in and excluded from the study.




Figure 2

Receiver operating characteristic (ROC) curve for D-dimer, fibrinogen, and platelet count within postpartum 24 hours of women age ≥ 35 . AUC (ROC of D-dimer): 0.732 (P < .001); AUC (ROC of fibrinogen): 0.592 (P = .086); AUC (ROC of platelet count): 0.594 (P = .081).