

Global Disease Burden of Uncorrected Refractive Error Among Adolescents From 1990 to 2019

Zhenlan Yang (✉ zhenlan14@163.com)

Sun Yat-sen University

Guangming Jin

Sun Yat-sen University

Zijing Li

Sun Yat-sen University

Yunru Liao

Sun Yat-sen University

Xiang Gao

Sun Yat-sen University

Yichi Zhang

Sun Yat-sen University

Yuqing Lan

Sun Yat-sen University

Research Article

Keywords: uncorrected refractive error, adolescents, disability-adjusted life years, socioeconomic, urbanization, education

Posted Date: April 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-459379/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background To estimate the global disease burden of uncorrected refractive error (URE) among adolescents and assess the contributions of various risk factors to disability-adjusted life-years (DALYs) due to URE.

Methods Global, regional and country-level DALY numbers and rates due to URE among adolescents were acquired from the Global Burden of Disease Study 2019 database. Human Development Index (HDI), Socio-Demographic Index (SDI) and other country-level data were obtained from other open databases as potential indicators. Regression analysis was used to evaluate associations between DALY rates among adolescents and potential predictors.

Results Global DALYs due to URE among adolescents rose by 8% between 1990 and 2019 but moderately decreased by 4.8% during this period after adjusting for population size. Female adolescents showed higher DALY rates. DALY rates sharply increased from 5 to 9 years of age, then rose more slowly, reaching a plateau before 20 years of age. Country-level DALY rates in 2019 were positively associated with HDI, SDI, and urbanization rates but negatively correlated with primary school dropout rates. Higher disease burden of adolescents visually impaired from URE was associated with lower primary school dropout rates ($\beta=-0.257$, 95% CI -0.376 to -0.138 , $P<0.001$) and higher urbanization rates ($\beta=0.257$, 95% CI 0.067 to 0.256 , $P=0.001$).

Conclusions Higher socioeconomic status, urbanization rates and education levels are associated with a heavier disease burden of URE among adolescents. The findings of this study can provide a reference for policy making on resource allocation for URE prevention and control in teenagers.

Background

Uncorrected refractive error is an underappreciated but widespread public health problem that significantly diminishes both quality of life and productivity. Among the global population with vision impairment (VI) and blindness in 2015, URE resulted in moderate to severe VI in 116.3 million people and blindness in 7.4 million people. With an extremely high prevalence, URE ranks first among VI causes, and the affected population is expected to continually grow[1]. Being at the critical period of eye development, children and adolescents could be sensitive to environmental factors and at high risk of suffering from refractive disorders. Especially during the coronavirus disease 2019 (COVID-19) pandemic period, these school-aged children could be vulnerable to myopia owing to the home confinement policy[2].

Given the difficulties that people with URE face in activities of daily living and social functioning, it is not surprising that URE can be associated with lower quality of life[3] and productivity loss[4]. The adverse effects of URE can be more serious in children, whose poor-quality visual experience due to URE can lead to unreversible amblyopia[5] and inadequate knowledge acquisition from the outer world during development and maturation[6]. There is no doubt that good visual performance has a profound effect on children's cognitive and psychosocial development[7, 8].

Although there have been many studies on the epidemiology and etiology of refractive error, there are few evaluations quantifying its health burden. The Global Burden of Diseases Study (GBD study)[9] developed disability-adjusted life years (DALYs) to estimate the disease burden of many conditions, including VI due to URE. DALYs are determined by the number of disabled years weighted by the level of disability caused by a disability or disease. Previous research has examined the DALYs of URE among people of all ages[10]. However, an aged population was generally affected by presbyopia and the previous study failed to fully describe the URE burden among adolescents. As most URE in adolescents is preventable and controllable, in this study, we focused on the disease burden of URE in adolescents[11–13]. The aim of this study was to describe the temporal trends of the global disease burden of URE among adolescents and its distribution across different age brackets, sexes, and regions and to explore the risk factors for DALYs due to URE in adolescents, including national-level data on demographic, socioeconomic and educational factors.

Methods

Global disease burden of URE among adolescents

DALY data of URE were retrieved from the open database of the Global Burden of Disease 2019 Study in the Global Health Data Exchange (GHDx; <http://ghdx.healthdata.org/gbd-results-tool>), which aggregated DALYs due to 369 diseases and injuries for 204 countries and territories from 1990 to 2019. International Classification of Diseases (10th edition) codes H52–H52.7 were mapped to URE covering myopia, hypermetropia, astigmatism, etc. in the database. The methodology has been concretely illustrated in a previous study[9]. Overall, DALYs are calculated as the sum of years lived with disability (YLD) and years of life lost (YLL). As a nonfatal disease, URE's DALY number = YLD = the number of prevalent cases * the disability weight. The DALY rate means the number of cases per 100,000 population (controlling for population size).

The following data were extracted from the database for further analysis: (1) the global DALY numbers, rates and global prevalence numbers and rates among adolescents (0–20 years old) in 1990–2019, (2) global sex-specific DALY numbers and rates from 1990 to 2019 and their age distribution (four age groups: 1 to 4, 5 to 9, 10 to 14 and 15 to 19 years of age) in 2019; (3) global maps of DALY numbers and rates among adolescents in 2019, with the top ten countries ranked by DALY numbers and rates. Global maps were generated from a data visualization tool available from the GHDx, supported by the Institute for Health Metrics and Evaluation (IHME) (<https://vizhub.healthdata.org/gbd-compare/>); and (4) World Health Organization (WHO) income level, the World Bank income level and Socio-Demographic Index (SDI)-level regional DALY rates in 2019.

Country-level indicators

We assessed the relationship of country-specific DALY rates with several country-level demographic, socioeconomic and educational indicators derived from the following well-known open databases. 2019 SDI data were obtained from the GHDx open database (<http://ghdx.healthdata.org/>). The SDI is a

composite measure of a country's socio-demographic development that combines information on gross national income (GNI) per capita, life expectancy at birth, and average and expected years of schooling. The SDI ranges from 0 to 1, with a higher value indicating a higher degree of socioeconomic development. The Human Development Index (HDI), primary school dropout rates and urbanization rates were obtained from the open database of the United Nations Development Programme (<http://hdr.undp.org/en/data>). The HDI has four components: a life expectancy index, a mean years of schooling index, an expected years of schooling index, and an income index.

Statistical analysis

The outcomes included the time, age, sex, and geographic distribution of DALYs due to URE. The Wilcoxon signed rank test was used to compare sex differences in DALY numbers and rates. One-way analysis of variance (ANOVA) was used to compare DALY rates in different WHO regions. The Kruskal-Wallis H test was used to investigate differences in DALY rates across four income-based and five SDI-based country groups, followed by the Mann-Whitney U test for post hoc pairwise comparisons. Scatter plots, Pearson correlation and univariate linear regression analyses were performed to explore the relationships between diverse demographic, socioeconomic and educational variables and DALY rates of uncorrected refractive disorder at the country level. Indicators that were significant in univariate analyses were further applied to stepwise multiple linear regression analyses. Statistical analyses were performed using SPSS software (SPSS, Inc.; Chicago, IL, USA; version 25.0), with $P < 0.05$ considered statistically significant. Figures were drawn using GraphPad Prism software (version 7.0e, GraphPad Software; San Diego, CA, USA).

Results

Trends in the disease burden of URE from 1990 to 2019

According to the GBD 2019, the number of DALYs due to URE slightly increased by 8%, from 814,261.98 (95% confidence interval [CI] 502,307.00–1,239,889.76) in 1990 to 879,736.05 (95% CI 538,302.00–1,353,108.75) in 2019 (Fig. 1A). Conversely, the DALY rate exhibited a slight downward trend, decreasing by 4.8%, from 35.81 (95% CI 22.09–54.53) to 34.11 (95% CI 20.87–52.46) (Fig. 1B). In addition, we observed that the numbers and rates of URE prevalence have remained high and relatively stable over the past 30 years (Fig. 1C and D).

Global disease burden of URE by age and sex

Wilcoxon signed-rank tests showed that there were significant sex disparities, as total DALYs for women significantly exceeded those of men from 1990 to 2019 (for both DALY numbers and rates, $P < 0.001$) (Fig. 2A and B). Figure 2C and D showed the global DALY variation by sex in different age groups in 2019. The two sexes showed a similar trend in global DALY rates by age, sharply increasing from 5 to 9 years of age, then slowly rising, and reaching a plateau before 20 years of age. Wilcoxon signed rank tests

showed that there were no significant sex disparities in the global DALY rates for each age group ($P=0.068$).

Regional differences in global disease burden due to URE

The severity of disease burden due to URE varies worldwide (Fig. 3A and B). As expected, the countries with the largest populations had the highest DALY numbers (Fig. 3C). India had the greatest total number of DALY due to URE (186,533.31, 95% CI 114,179.97–287,138.60). After adjusting for population size, the DALY rate was highest in Oman (82.85, 95% CI 49.99–128.59) (Fig. 3D). One-way ANOVA and the Bonferroni correction for multiple comparisons indicated that the Eastern Mediterranean region had the heaviest disease burden compared to the remaining five WHO regions over the past 30 years ($P<0.001$), followed by the Region of the Americas. The African Region had the lightest burden. All time trends for rates in the six WHO regions remained steady from 1990 to 2019 (Fig. 4A). Each of the income-level and SDI-level regions were also analyzed. The DALY rates were highest in high-income regions and lowest in low-income regions (Fig. 4B). Middle SDI and low SDI regions ranked first and last, respectively, in terms of the DALY rates, while in recent years, high-middle SDI regions passed middle SDI regions and took first place (Fig. 4C).

Country-level disease burden due to URE and associations with national indicators

The scatter plots between the disease burden of URE and country-level indicators are presented in Fig. 5. Linear trends were observed between the disease burden of URE and SDI, HDI, primary school dropout rate and urbanization rate. Pearson correlation and linear regression analysis revealed that the country-level SDI, HDI and urbanization rate were significantly positively correlated with the DALY rates of URE, while primary school dropout rates were negatively correlated with DALY rates (Table 1 and Table 2). Upon further analysis, primary school dropout rates and HDI were the most influential indicators. In stepwise multiple linear regression analyses, higher urbanization rates ($\beta = 0.257$, 95% CI 0.067–0.256, $P = 0.001$) and lower primary school dropout rates ($\beta = -0.257$, 95% CI -0.376 to -0.138, $P < 0.001$) were associated with a higher disease burden resulting from URE (Table 2).

Table 1

Pearson correlation analysis of the relationship between country-level disease burden of URE and socioeconomic variables

Factors	Pearson correlation	
	r	P value
SDI	0.414	⊠0.001*
HDI	0.459	⊠0.001*
Primary school dropout rate (%)	-0.460	⊠0.001*
Urban (%)	0.400	⊠0.001*

P < 0.05 was considered statistically significant; SDI – Socio-Demographic Index, HDI – Human Development Index.

Table 2

Linear regression analysis of the relationship between country-level disease burden of URE and socioeconomic variables

Factors	Univariate linear regression			Multiple linear regression		
	Adjusted R square	β(95% CI)	P value	Adjusted R square	β(95% CI)	P value
SDI	0.167	30.548 (20.637 to 40.459)	⊠0.001*	-	-	-
HDI	0.206	41.17 (29.347 to 53.001)	⊠0.001*	-	-	-
Primary school dropout rate (%)	0.207	-0.355 (-0.463 to -0.247)	⊠0.001*	0.257	-0.257 (-0.376 to -0.138)	⊠0.001*
Urban (%)	0.155	0.231 (0.154 to 0.308)	⊠0.001*	0.257	0.162 (0.067 to 0.256)	0.001*

P < 0.05 was considered statistically significant; SDI – Socio-Demographic Index, HDI – Human Development Index.

Discussion

This study used DALYs for the first time to elucidate the disease burden associated with URE among adolescents, exploring temporal trends from 1990 to 2019 and the global distribution by age, sex, WHO regions, level of healthcare and socioeconomic factors. From 1990 to 2019, the world DALY numbers due to URE increased by 8%, while the world DALY rates slightly decreased over time after adjusting for increases in population. The global DALY rates of URE increased by age. Females had higher DALY rates than males of the same age. Regarding regional distribution, the disease burden in the Eastern

Mediterranean, high-income and middle SDI regions was highest. Further research revealed that the disease burden of URE was associated with HDI, SDI, primary school dropout rates and urbanization rate.

Refractive errors are the result of a mismatch between the axial length of the eye and its optical power, creating blurred vision. Eye growth, namely, emmetropization, is believed to be completed at the age of 13 years, while axial length growth in myopic eyes may continue[14]. Myopes typically exhibit early progression quickly, and the progression then slows down; in general, their refractive error eventually stabilizes during late adolescence, e.g., 15–21 years of age[15]. High incidence and high degrees of astigmatism were demonstrated to exist among children, especially newborns. As children grow older, the cornea flattens with significantly reduced astigmatism and stabilizes before adulthood[16]. Therefore, in this study, adolescence was defined as ages below 20 years.

The results showed fluctuation in the prevalence and disease burden of URE among adolescents from 1990 to 2019. The temporal variation in the disease burden of URE was largely in accordance with the variations observed in prevalence. However, compared to prevalence, DALYs are more suitable indicators to measure the influence of a disease on quality of life. The total DALY numbers due to URE increased by 8% from 1990 to 2019, while the DALY rates remained stable after adjusting for global population growth. The high disease burden of URE among adolescents persists, suggesting that current health policies to control URE have failed to alleviate the additional disease burden caused by increasing prevalence. According to Holden et al[17], the global myopic population will continue to expand and reach 4.758 billion (49.8% of the world population) by 2050, which means healthcare planners will face an even heavier burden of VI related to URE.

Sex inequality in the global burden of URE exists even among adolescents and has persisted since 1990, with females bearing a significantly higher burden of URE than males. Females had a higher burden of URE than males for all age groups in 2019, as has been reported previously[18]. Many studies have revealed a higher prevalence of refractive disorders among young females than in males. This may reflect different environmental risk factors, such as the tendency of girls to spend a greater number of hours engaged in near vision activities and significantly fewer hours outdoors than boys[19–22]. The longer life expectancy of females in most cultures could also contribute to a higher global burden of URE. Moreover, the inequality of social, cultural, and economic status between men and women is believed to reduce access to eye care services, including refractive correction for women[23]. However, the sex discrepancy of spectacle coverage has been reported as less extensive than expected[24, 25], which implies that the gender inequality of resource distribution may not be the primary cause.

The age-specific variation curve of DALY rates due to URE showed that the steepest increase in disease burden occurred in the 5–9 age group, followed by relatively smooth growth in the 10–14 age group; DALY rates then seemingly reached a plateau at 15–19 years of age. This trend was nicely consistent with the growth law of the ocular axis and myopia, which typically exhibits fast progression early at the age of 6–8 years, followed by slowing and eventual stabilization of refractive error in late

adolescence[14, 15, 26]. The increasing study burden among these school-age children may explain the growing prevalence rate of VI associated with URE[21, 27].

Regional inequality was also obvious, with a higher disease burden among adolescents in the WHO Eastern Mediterranean Region, Region of Americas and South-East Asia Region. Our results also revealed that regions with middle to high income and SDI have higher DALY rates. The HDI and SDI, two widely used indexes of socioeconomic development, were significantly positively correlated with VI due to URE in univariate linear regression. We compared the results of our disease burden analysis focused on adolescents to similar studies among individuals of all ages. Our results turned out to contrast with the findings of Lou et al., in which age-standardized DALY rates were inversely associated with HDI[10]. These differences could be attributed to the diverse spectrum of diseases at different ages. The aging process of ocular refractive structures, such as loss of lens power and lens opacity, can lead to presbyopia and crystalline source refractive error among older people, which constitutes a large part of VI due to URE. A total of 244 million uncorrected presbyopia cases among people aged < 50 years were associated with a potential productivity loss of US \$11.023 billion (0.016% of global GDP)[28]. The prevalence of presbyopia is estimated to be higher in regions with longer life expectancies, whereas a greater burden of VI resulting from uncorrected presbyopia occurs in less developed countries. The low amounts of available eye care resources and poor optical correction rates in these areas were proven to be the underlying causes[29]. Moreover, the quantity and quality of cataract surgery that could correct refractive errors to some extent were also positively associated with GDP per capita and HDI[30]. For children under 20 years of age, a recent meta-analysis by Hashemi et al. estimated global and regional prevalence figures of refractive errors: the estimated pooled prevalence of astigmatism (> 0.50 D, 14.9%) was higher than that of myopia (≤ -0.50 D, 11.7%) and hyperopia ($\geq +2.0$ D, 4.6%). The prevalence and severity of astigmatism and hyperopia often decrease during emmetropization, while myopia tends to exhibit fast progression in school-age children[31]. This process can be greatly influenced by environmental factors[13, 32]. Among these factors, education as an element of SDI and HDI plays an important role. The tendency for schooling to lead to an increased prevalence of myopia and VI has been documented in almost all major population groups[21, 27, 33]. This result agrees with our finding that primary school dropout rates were inversely correlated with the DALY rates due to URE in both Pearson correlation and univariate linear regression analyses.

Additionally, variations in other parameters, such as living environment, diet and lifestyle accompanying societal development may also contribute to the high prevalence of URE and VI among adolescents[13, 32]. Therefore, we explored the effects of primary school dropout rates and urbanization rates on URE burden with multiple linear regression analysis. A heavier disease burden due to URE in young people was suggested to occur among higher urbanized countries with lower primary school dropout rates. Potential mechanisms were enumerated as follows: first, primary and middle students could be more susceptible to myopia due to environmental factors such as heavy educational stress and fewer outdoor activities[21]. Second, findings from many studies indicated that the burden of myopia may be heavier among urban residents, which is consistent with our results[34–36]. Urban areas are usually characterized by more severe environmental pollution (less green space, ambient light exposure),

different lifestyles (lower levels of time outdoors and higher levels of indoor activities)[35] and heavy academic pressure, all of which may affect refractive errors. Recent interventional prospective studies have shown that encouraging outdoor activity and exposure to sunlight effectively prevents the development of myopia[11, 12]. Third, significant inequities in resources for refractive error correction still exist in developed countries with high urbanization rates[37, 38] and in urban areas of developing countries[39]. As a result, the need for refractive error correction in many people, especially migrants, remains unmet, although resources are adequate in these regions. This is partially attributable to socioeconomic barriers, such as low income, lower rates of health care coverage, fewer visits to health services, and language barriers[40, 41].

This study has several limitations. First, the accuracy of the disease burden information was affected the limitations of the data source; the data were not collected first-hand. Second, data from 204 countries and territories were obtained in the GBD database, and the completeness of the study was limited by the unavailability of data from the remaining countries and territories.

Despite the limitations mentioned above, the study findings can be useful for developing targeted strategies to address the severe VI resulting from URE. The estimated large disease burden reveals a challenging task for healthcare policymakers, and considerable efforts will be required to scale up refractive error services for adolescents.

Conclusions

In conclusion, the global VI burden of URE among adolescents remained stable without significant alleviation from 1990 to 2019. Older age and female sex were associated with a higher burden of URE. The steepest increase in DALY rates occurred in the 5–9 age group, namely, primary school students. High socioeconomic development status (indicated by HDI and SDI), low primary school dropout rates and high urbanization rates were associated with increasing DALY rates. This suggests that we should be aware of the growing disease burden of URE among adolescents along with the process of urbanization and universal education. These findings can provide information for policymakers to craft more targeted strategies to prevent and control child and adolescent VI due to URE.

Abbreviations

URE

uncorrected refractive error; DALYs:Disability-adjusted life years; YLDs:Years lived with disability; YLLs:Years of life lost; GBD:Global burden of disease; VI:visual impairment; IHME:Institute for Health Metrics and Evaluation; WHO:World health organization; HDI:Human development index; SDI:Socio-Demographic Index; GNI:gross national income; ANOVA:one-way analysis of variance.

Declarations

Ethics approval and consent to participate: Not applicable.

Consent for publication: Not applicable.

Availability of data and materials: The datasets generated during and/or analysed during the current study are available in the Global Health Data Exchange repository, <http://ghdx.healthdata.org/gbd-results-tool> and United Nations Development Programme repository, <http://hdr.undp.org/>.

Competing interests: The authors declare that they have no competing interests.

Funding: Not applicable.

Authors' contributions: YZ and YqL contributed conception and design of the work, ZY and GJ analyzed the data of the work and drafted the manuscript, ZL, YnL, XG took part in acquisition the data of the work and interpretation of data for the work. All authors had read the work and revised

it critically for important intellectual content. All authors had agreement to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Acknowledgements: This research was supported by the National Natural Science Foundation of China (Grant No:81700833).

References

1. **Trends in prevalence of blindness and distance and near vision impairment over 30 years: an analysis for the Global Burden of Disease Study.** *The Lancet Global health* 2021, **9**(2):e130-e143.
2. Wang J, Li Y, Musch DC, Wei N, Qi X, Ding G, Li X, Li J, Song L, Zhang Y *et al*: **Progression of Myopia in School-Aged Children After COVID-19 Home Confinement.** *JAMA ophthalmology* 2021.
3. Tahhan N, Papas E, Fricke TR, Frick KD, Holden BA: **Utility and uncorrected refractive error.** *Ophthalmology* 2013, **120**(9):1736-1744.
4. Smith TS, Frick KD, Holden BA, Fricke TR, Naidoo KS: **Potential lost productivity resulting from the global burden of uncorrected refractive error.** *Bulletin of the World Health Organization* 2009, **87**(6):431-437.
5. Shotton K, Powell C, Voros G, Hatt SR: **Interventions for unilateral refractive amblyopia.** *The Cochrane database of systematic reviews* 2008(4):Cd005137.
6. O'Neill DK, Astington JW, Flavell JH: **Young children's understanding of the role that sensory experiences play in knowledge acquisition.** *Child development* 1992, **63**(2):474-490.
7. Packwood EA, Cruz OA, Rychwalski PJ, Keech RV: **The psychosocial effects of amblyopia study.** *Journal of AAPOS: the official publication of the American Association for Pediatric Ophthalmology and Strabismus* 1999, **3**(1):15-17.

8. Birch EE, Castañeda YS, Cheng-Patel CS, Morale SE, Kelly KR, Beauchamp CL, Webber A: **Self-perception of School-aged Children With Amblyopia and Its Association With Reading Speed and Motor Skills.** *JAMA ophthalmology* 2019, **137**(2):167-174.
9. **Global burden of 369 diseases and injuries in 204 countries and territories, 1990-2019: a systematic analysis for the Global Burden of Disease Study 2019.** *Lancet (London, England)* 2020, **396**(10258):1204-1222.
10. Lou L, Yao C, Jin Y, Perez V, Ye J: **Global Patterns in Health Burden of Uncorrected Refractive Error.** *Investigative ophthalmology & visual science* 2016, **57**(14):6271-6277.
11. He M, Xiang F, Zeng Y, Mai J, Chen Q, Zhang J, Smith W, Rose K, Morgan IG: **Effect of Time Spent Outdoors at School on the Development of Myopia Among Children in China: A Randomized Clinical Trial.** *Jama* 2015, **314**(11):1142-1148.
12. Ngo CS, Pan CW, Finkelstein EA, Lee CF, Wong IB, Ong J, Ang M, Wong TY, Saw SM: **A cluster randomised controlled trial evaluating an incentive-based outdoor physical activity programme to increase outdoor time and prevent myopia in children.** *Ophthalmic & physiological optics : the journal of the British College of Ophthalmic Opticians (Optometrists)* 2014, **34**(3):362-368.
13. Harb EN, Wildsoet CF: **Origins of Refractive Errors: Environmental and Genetic Factors.** *Annual review of vision science* 2019, **5**:47-72.
14. Fledelius HC, Christensen AS, Fledelius C: **Juvenile eye growth, when completed? An evaluation based on IOL-Master axial length data, cross-sectional and longitudinal.** *Acta ophthalmologica* 2014, **92**(3):259-264.
15. **Myopia stabilization and associated factors among participants in the Correction of Myopia Evaluation Trial (COMET).** *Investigative ophthalmology & visual science* 2013, **54**(13):7871-7884.
16. Read SA, Collins MJ, Carney LG: **A review of astigmatism and its possible genesis.** *Clinical & experimental optometry* 2007, **90**(1):5-19.
17. Holden BA, Fricke TR, Wilson DA, Jong M, Naidoo KS, Sankaridurg P, Wong TY, Naduvilath TJ, Resnikoff S: **Global Prevalence of Myopia and High Myopia and Temporal Trends from 2000 through 2050.** *Ophthalmology* 2016, **123**(5):1036-1042.
18. Lou L, Liu X, Tang X, Wang L, Ye J: **Gender Inequality in Global Burden of Uncorrected Refractive Error.** *American journal of ophthalmology* 2019, **198**:1-7.
19. Saxena R, Vashist P, Tandon R, Pandey RM, Bhardawaj A, Menon V, Mani K: **Prevalence of myopia and its risk factors in urban school children in Delhi: the North India Myopia Study (NIM Study).** *PloS one* 2015, **10**(2):e0117349.
20. Ostadimoghaddam H, Fotouhi A, Hashemi H, Yekta A, Heravian J, Rezvan F, Ghadimi H, Rezvan B, Khabazkhoob M: **Prevalence of the refractive errors by age and gender: the Mashhad eye study of Iran.** *Clinical & experimental ophthalmology* 2011, **39**(8):743-751.
21. Wang J, Ying GS, Fu X, Zhang R, Meng J, Gu F, Li J: **Prevalence of myopia and vision impairment in school students in Eastern China.** *BMC ophthalmology* 2020, **20**(1):2.

22. Pärssinen O, Kauppinen M, Viljanen A: **The progression of myopia from its onset at age 8-12 to adulthood and the influence of heredity and external factors on myopic progression. A 23-year follow-up study.** *Acta ophthalmologica* 2014, **92**(8):730-739.
23. Courtright P, Lewallen S: **Why are we addressing gender issues in vision loss?** *Community eye health* 2009, **22**(70):17-19.
24. Fotouhi A, Hashemi H, Raissi B, Mohammad K: **Uncorrected refractive errors and spectacle utilisation rate in Tehran: the unmet need.** *The British journal of ophthalmology* 2006, **90**(5):534-537.
25. Marmamula S, Madala SR, Rao GN: **Prevalence of uncorrected refractive errors, presbyopia and spectacle coverage in marine fishing communities in South India: Rapid Assessment of Visual Impairment (RAVI) project.** *Ophthalmic & physiological optics : the journal of the British College of Ophthalmic Opticians (Optometrists)* 2012, **32**(2):149-155.
26. Hou W, Norton TT, Hyman L, Gwiazda J: **Axial Elongation in Myopic Children and its Association With Myopia Progression in the Correction of Myopia Evaluation Trial.** *Eye & contact lens* 2018, **44**(4):248-259.
27. Williams KM, Bertelsen G, Cumberland P, Wolfram C, Verhoeven VJ, Anastasopoulos E, Buitendijk GH, Cougnard-Grégoire A, Creuzot-Garcher C, Erke MG *et al.*: **Increasing Prevalence of Myopia in Europe and the Impact of Education.** *Ophthalmology* 2015, **122**(7):1489-1497.
28. Frick KD, Joy SM, Wilson DA, Naidoo KS, Holden BA: **The Global Burden of Potential Productivity Loss from Uncorrected Presbyopia.** *Ophthalmology* 2015, **122**(8):1706-1710.
29. Fricke TR, Tahhan N, Resnikoff S, Papas E, Burnett A, Ho SM, Naduvilath T, Naidoo KS: **Global Prevalence of Presbyopia and Vision Impairment from Uncorrected Presbyopia: Systematic Review, Meta-analysis, and Modelling.** *Ophthalmology* 2018, **125**(10):1492-1499.
30. Wang W, Yan W, Müller A, He M: **A Global View on Output and Outcomes of Cataract Surgery With National Indices of Socioeconomic Development.** *Investigative ophthalmology & visual science* 2017, **58**(9):3669-3676.
31. Hashemi H, Fotouhi A, Yekta A, Pakzad R, Ostadimoghaddam H, Khabazkhoob M: **Global and regional estimates of prevalence of refractive errors: Systematic review and meta-analysis.** *Journal of current ophthalmology* 2018, **30**(1):3-22.
32. Grzybowski A, Kanclerz P, Tsubota K, Lanca C, Saw SM: **A review on the epidemiology of myopia in school children worldwide.** *BMC ophthalmology* 2020, **20**(1):27.
33. Nickels S, Hopf S, Pfeiffer N, Schuster AK: **Myopia is associated with education: Results from NHANES 1999-2008.** *PloS one* 2019, **14**(1):e0211196.
34. Shapira Y, Mimouni M, Machluf Y, Chaiter Y, Saab H, Mezer E: **The Increasing Burden of Myopia in Israel among Young Adults over a Generation: Analysis of Predisposing Factors.** *Ophthalmology* 2019, **126**(12):1617-1626.
35. Guo Y, Liu LJ, Xu L, Lv YY, Tang P, Feng Y, Meng M, Jonas JB: **Outdoor activity and myopia among primary students in rural and urban regions of Beijing.** *Ophthalmology* 2013, **120**(2):277-283.

36. Lee YY, Lo CT, Sheu SJ, Lin JL: **What factors are associated with myopia in young adults? A survey study in Taiwan Military Conscripts.** *Investigative ophthalmology & visual science* 2013, **54**(2):1026-1033.
37. Ngo G, Trope G, Buys Y, Jin YP: **Significant disparities in eyeglass insurance coverage in Canada.** *Canadian journal of ophthalmology Journal canadien d'ophtalmologie* 2018, **53**(3):260-265.
38. Zhang X, Elliott MN, Saaddine JB, Berry JG, Cuccaro P, Tortolero S, Franklin F, Barker LE, Schuster MA: **Unmet eye care needs among U.S. 5th-grade students.** *American journal of preventive medicine* 2012, **43**(1):55-58.
39. Wang X, Yi H, Lu L, Zhang L, Ma X, Jin L, Zhang H, Naidoo KS, Minto H, Zou H *et al*: **Population Prevalence of Need for Spectacles and Spectacle Ownership Among Urban Migrant Children in Eastern China.** *JAMA ophthalmology* 2015, **133**(12):1399-1406.
40. Muñoz B, West SK, Rodriguez J, Sanchez R, Broman AT, Snyder R, Klein R: **Blindness, visual impairment and the problem of uncorrected refractive error in a Mexican-American population: Proyecto VER.** *Investigative ophthalmology & visual science* 2002, **43**(3):608-614.
41. Fiscella K, Franks P, Gold MR, Clancy CM: **Inequality in quality: addressing socioeconomic, racial, and ethnic disparities in health care.** *Jama* 2000, **283**(19):2579-2584.

Figures




Figure 1

Time trends of global health burden and prevalence of URE among adolescents in 1990–2019. Panel A. DALY numbers. Panel B. DALY rates. Panel C. Prevalence numbers. Panel D. Prevalence rates. Dashed lines represent 95% CIs; DALY, disability-adjusted life year; URE, uncorrected refractive error.




Figure 2

The distribution of global burden due to URE among adolescents by age and sex. Panel A. Sex-specific DALY numbers. Panel B. Sex-specific DALY rates. Panel C. Age-specific and sex-specific DALY numbers in 2019. Panel D. Age-specific and sex-specific DALY rates in 2019. DALY, disability-adjusted life year; URE, uncorrected refractive error.




Figure 3

Global map of the disease burden of adolescents visually impaired from URE. Panel A. DALY numbers. Panel B. DALY rates. Panel C. The 10 countries with the highest DALY numbers. Panel D. The 10 countries with the highest DALY rates. DALY, disability-adjusted life years; URE, uncorrected refractive error. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.




Figure 4

Global health burden distribution of URE among adolescents in different regions. Panel A. WHO regions. Panel B. Different SDI-level regions. Panel C. Different income-level regions. WHO, World Health

Organization; SDI, Socio-Demographic Index; URE, uncorrected refractive error.




Figure 5

Relationship between the disease burden of URE among adolescents and country-level indicators. Panel A. SDI. Panel B. HDI. Panel C. Primary school dropout rates. Panel D. Urbanization rates. HDI, Human Development Index; SDI, Socio-Demographic Index.