A Novel Association of Hbq India Trait with Sick Cell Anemia: A New Insight in Hemoglobinopathies

Rakesh Kumar Gupta (✉ rakesh.newjobi@gmail.com)

All India Institute of Medical Sciences - Raipur <https://orcid.org/0000-0001-9031-1109>

Kartavya Kumar Verma

All India Institute of Medical Sciences - Raipur

Gurmeet Singh

Jawahar Lal Nehru Hospital and Research Centre Bhilai

Research Article

Keywords: HbQ India, Hemoglobinopathy, sickle cell trait, HPLC.

Posted Date: July 1st, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-466525/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at SN Comprehensive Clinical Medicine on January 4th, 2022. See the published version at <https://doi.org/10.1007/s42399-021-01103-y>.

Abstract

HbQ India is a rare α -chain structural hemoglobinopathy usually asymptomatic and presents in the heterozygous form or co-inherited with β -thalassemia trait. Herein, we are reporting the third case of novel association of HbQ India with HbS trait hemoglobinopathy in a 30-years-old young male presented with chief complaints of yellowish discoloration of sclera since 5-years with raised serum bilirubin levels along with pedigree analysis of the family.

Introduction

HbQ India is a rare hemoglobinopathy [1]. Identification of these hemoglobinopathies is crucial for prognostication, treatment, genetic counselling and preventive strategies formulation. HbQ India (HbA1:c. 193 G > C), is an uncommon α -chain structural hemoglobin variant. The structural changes at codon 64 of the $\alpha 1$ -globin gene cause substitution of histidine for aspartic acid. HbQ India usually presents in the heterozygous form or co-inherited with β -thalassemia trait [2]. We are reporting the third case of HbQ India with HbS trait hemoglobinopathy in a 30-years-old young male presented with chief complaints of yellowish discoloration of sclera along with pedigree analysis of the family. To the best of our knowledge only two index cases of co-inherited, HbQ India/HbS, hemoglobinopathy has been reported previously by Parab et al [3] in the English literature.

Case Report

A 30-years-old young male presented with flu-like symptoms at COVID centre of our hospital. Patient's oro-pharyngeal and nasal swab samples were taken for the RTPCR which came positive for COVID-19 infection and he was admitted to the isolation ward. He also complained of persistent yellowish discoloration of the sclera since 5 years. In view of Chhattisgarh being endemic state for sickle cell anemia (SCA) and his belonging to 'Sahu community' which is a common ethnic group in the region for development of SCA his detailed past medical history was also inquired. It was found that patient had yellowish discoloration of sclera with raised bilirubin levels (total bilirubin: 6.56 mg/dl, direct bilirubin: 1.02 mg/dl and indirect bilirubin 5.54 mg/dl) since 5 years. In 2018, patient was diagnosed to have sickle cell trait along with mild splenomegaly. His sickling test was positive, while direct and indirect Coomb's tests were negative. Except mildly deranged bilirubin levels other liver function tests (LFT) were within normal limits. His hemogram showed slightly decreased RBCs count: $3.98 \times 10^6/\mu\text{l}$, however other parameters were normal. Peripheral blood smear showed predominantly normocytic normochromic RBCs. No schistocytes, pencil cells or nRBC were found. At present admission, a high performance liquid chromatography (HPLC) was done for characterization of the hemoglobinopathy which revealed two peaks of 41.8% and 9.1% at retention time 4.14 and 4.60 corresponding to HbS and HbQ India respectively suggestive of sickle cell - HbQ India hemoglobinopathy (Fig. 1). Again, LFT delineated high bilirubin levels (total bilirubin: 5.69 mg/dl, direct bilirubin: 0.85 mg/dl, indirect bilirubin: 4.84 mg/dl). We also did a pedigree analysis of the family (Fig. 2) by HPLC and found wife carrying normal Hb pattern, while elder daughter (3 years) was sickle cell trait and younger daughter (1 years) was also having sickle

cell - HbQ India hemoglobinopathy. His both daughters were asymptomatic with normal hemogram. Patient was symptomatically treated for the COVID-19 infection and discharged after 7 days, when his COVID-19 RTPCR report became negative. At 3 months of follow up, patient is otherwise healthy resumed his office, however still having persistent icterus.

Discussion

HbQ India is a very rare alpha-chain structural variant hemoglobinopathy with a prevalence of 0.4% in the Indian subcontinent [2]. Sachdev et al in their HPLC based study in northern India and Nepal, reported a prevalence of 0.2% for HbQ India hemoglobinopathy [4]. HbQ is prevalent in South East Asia and shows three structural variants which includes HbQ India (alpha 64 Asp to His), HbQ Thailand (alpha 74 Asp to His), and HbQ Iran (alpha 75 Asp to His). HbQ India is clustered in western and northern India and more common in the Sindhi and Punjabi communities [5]. The earlier reported indexed cases by Parab et al [3] were also belonging to Sindhi community but the present case is from Sahu community which is not identified as risk factor for the HbQ India hemoglobinopathy. Chhattisgarh is an endemic zone for sickle cell anemia, however only two cases of HbQ India with HbS trait hemoglobinopathy are reported from this part of country by Parab et al [3]. Phanasgaonkar et al [2] in their study reported 64 patients of HbQ India which included 36 HbQ India trait, 22 HbQ India- β thalassemia trait, 3 HbQ India- β thalassemia major and 3 cases of HbQ India homozygous. The literature about HbQ India is very limited and only one of the earlier published paper by Parab et al^[3] reported co-inheritance of HbQ India with HbS trait. So, this is a third case of HbQ India-HbS trait. HbQ India trait and HbQ India- β thalassemia trait patients are usually asymptomatic, sometimes show only mild hematological parameters derangement and diagnosed incidentally during screening. Parab et al in their index cases reported mild anaemia in both mother and daughter (Hb 10.1 g/dl and 10.7 g/dl respectively) with decreased mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC) [3]. However, authors have not discussed any details about clinical presentation and liver function test in their cases. In present cases, patient presented with a long term (about 5 years) persistent icterus with indirect hyperbilirubinemia, mild anemia and decreased total RBCs count. Daughter of the patient was asymptomatic and showed mild anemia with slightly decreased MCH. Panigrahi et al [6] in their series found jaundice in one patient, however which could be explained by concomitantly inherited G6PD deficiency. We assume that HbQ India in presence of co-inherited sickle cell trait is clinically symptomatic and more severe in comparison to other types of inheritance pattern and association. Since, this is a case report, the exact mechanism and pathophysiology of icterus further needs to be elucidated.

Mean corpuscular volume (MCV) and MCH levels reduces in patient showing co-inheritance of HbQ India with β thalassemia, however, in present case these parameters were within normal range.

Harrison et al [7] reported HbQ India levels varies from 13.6–24.4% and 7.4–9.0% in patients with HbQ India trait & HbQ India with β thalassemia trait respectively. In present case, HbQ level was 9.1% corresponding to an unknown post HbS peak at retention time 4.60 minute. Parab et al in their case reported HbQ India peak at 4.55 minute as well as a hybrid peak of HbS and HbQ India in HbC window at

retention time of 5.08 minute [3]. Harrison et al [7] also reported minor additional peaks (post HbQ, Split HbA2 and HbS). In present, case hybrid peak or minor peaks were not found.

To conclude, HbQ India is a very rare hemoglobinopathy usually inherited in heterozygous form alone or in common association with β thalassemia trait in Sindhi and Punjabi community and remain silent. This is a first case of HbQ India co-inherited with sickle cell trait in Sahu community with chronic persistent jaundice.

Declarations

Funding: None

Conflicts of interest/Competing interests: No conflicts of interest to be declared

Availability of data and material: Yes

Code availability: Not applicable

Authors' contributions:

1. **Dr Rakesh Kumar Gupta-** Concept, design, definition of intellectual content, Manuscript editing
2. Dr Kartavya Verma- Data acquisition, data analysis, literature search, manuscript preparation
3. Dr Gurmeet Singh- Manuscript editing, and manuscript review

Ethics approval: Not applicable

Consent to participate: Not applicable

Consent for publication: Taken

References

1. Rao S, Kar R, Gupta SK, Chopra A, Saxena R. Spectrum of haemoglobinopathies diagnosed by cation exchange-HPLC & modulating effects of nutritional deficiency anaemias from north India. Indian J Med Res. 2010;132:513–9.
2. Phanasgaonkar S, Colah R, Ghosh K, Mohanty D, Gupte S. Hb Q(India) and its interaction with beta-thalassaemia: a study of 64 cases from India. Br J Biomed Sci. 2007;64:160–3.
3. Parab S, Sakhare S, Sengupta C, Velumani A. Diagnosis of a novel hemoglobinopathy of compound heterozygosity of hemoglobin S/hemoglobin Q India. Clin Chim Acta. 2015;442:33–5.
4. Sachdev R, Dam AR, Tyagi G. Detection of Hb variants and hemoglobinopathies in Indian population using HPLC: report of 2600 cases. Indian J Pathol Microbiol. 2010;53:57–2.

5. Mutreja D, Tyagi S, Tejwani N, Dass J. Double heterozygous hemoglobin Q India/hemoglobin D Punjab hemoglobinopathy: Report of two rare cases. Indian J Hum Genet. 2013;19:479–2.
6. Panigrahi I, Bajaj J, Chatterjee T, Saxena R, Mahapatra M, Pati HP. Hb Q India: is it always benign? Am J Hematol. 2005;78:245–6.
7. Harrison A, Mashon RS, Kakkar N, Das S. Clinico-Hematological Profile of Hb Q India: An Uncommon Hemoglobin Variant. Indian J Hematol Blood Transfus. 2018;34:299–3.

Figures

Figure 1

High performance liquid chromatography analysis of patient and his daughter showing two peaks corresponding to HbS (black arrow) and HbQ India (red arrow) respectively.

Figure 2

Flowchart showing pedigree analysis of the patient's family.