

Effects of Elevated Atmospheric CO₂ and its Interaction with Temperature and Nitrogen on Yield of Barley (*Hordeum vulgare* L.): A Meta-analysis

Mekides Woldegiorgis Gardi (✉ mekides.gardi@uni-hohenheim.de)

University of Hohenheim: Universitat Hohenheim <https://orcid.org/0000-0002-3709-6718>

Bettina I.G Haussmann

University of Hohenheim

Waqas Ahmed Malik

University of Hohenheim

Petra Högy

University of Hohenheim

Research Article

Keywords: CO₂-enrichment, *Hordeum vulgare* L., Nitrogen, Temperature, Yield-variable

Posted Date: May 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-466735/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Plant and Soil on April 11th, 2022. See the published version at <https://doi.org/10.1007/s11104-022-05386-5>.

Abstract

Aims

The general aim of this meta-analysis is to synthesize and summarise the mean response of barley yield variables to elevated CO₂ (eCO₂) and its interaction with temperature and N fertilization.

Methods

The present study quantitatively synthesized the response of barley to eCO₂ and its interaction with temperature, and Nitrogen (N). A meta-analysis procedure was used to analyse five yield variables of barley extracted from 76 articles to determine the effect size and the magnitude in relation to eCO₂ and its interaction with temperature and N.

Results

CO₂ enrichment increased biomass (23.8%), grain number (24.8%), grain yield (27.4%), and thousand-grain weight (5.6%). However, responses to eCO₂ were affected by genotype, additional stress, and experimental conditions. In comparison, genotype "*Anakin*" shows the highest response of biomass (47.1%), while "*Genebank accessions*" had a higher grain number (46.1%) and grain yield (57.1%) under eCO₂. The maximal enhancement of barley yield was observed when plants grow under a combination of eCO₂ and higher nitrogen fertilizer (>100 kg ha⁻¹). Nevertheless, biomass (-12%), and grain yield (-17%) responses were lower when eCO₂ is combined with high temperature (>25 °C). It was further noted the response of barley yield to eCO₂ was higher in the growth chamber than in other CO₂ exposure methods. Moreover, comparing pot-rooted versus field-rooted barley plants, a higher response of biomass and grain yield was observed for pot-rooted plants.

Conclusions

Overall, results suggest that the maximal barley production under eCO₂ will be obtained in combination with high nitrogen fertilizer and optimal temperature (21-25 °C).

Introduction

Evidence indicates that atmospheric carbon dioxide (CO₂) concentration increased globally, from 280 ppm in the pre-industrial period to about 410 ppm in 2019 and it might increase to 550 ppm by 2050 (NOAA, 2020). CO₂ is the most important anthropogenic greenhouse gas (GHG) and it represented 72% of overall anthropogenic GHG emissions in 2018. The changes in CO₂ concentration and other GHG emissions are expected to increase air temperature by 2.5 to 4.5 °C at the end of the 21st century (IPCC, 2018). These environmental changes will have a substantial effect on crop growth and food supply in the future. At the same time, the total global food production has to increase by 25 to 70% within the next 40 years, to meet the food demand for the projected increase in the global population (Fróna, 2019; United

Nations, 2011). Alleviating future food security challenges will need precisely estimating crop production response to the ongoing increase of atmospheric CO₂, together with rising temperature, and soil fertility. The potential beneficial effect of elevated CO₂ (eCO₂) on plant growth and yield has been well documented (Ainsworth et al., 2002; Clausen et al., 2011; Curtis and Wang, 1998; Wang et al., 2013), however, the magnitude of this effect varies widely among species, genotypes, and growing conditions.

Barley (*Hordeum vulgare* L.) is one of the most important and extensively cultivated cereal crops worldwide for human nutrition and as animal feed. The production of barley in 2019 was about 150 × 10⁶ tonnes and barley has been cultivated in more than 100 countries worldwide (FAOSTAT, 2020). An increase in atmospheric CO₂ generally exerts beneficial effects on plant biomass by increasing net photosynthesis by 30 to 50% and reducing photorespiration (Drake et al., 1997; Poorter and Navas, 2003; Schapendonk et al., 2000). This has been studied for cereals including barley, wheat, rice, oat, and rye (Conroyac et al., 1994; Kimball et al., 2002; Long et al., 2006). For instance, in a meta-analysis comprising 79 crop and wild species, Jablonski et al. (2002) documented an average increase in yield of 28% due to eCO₂. A climate chamber experiment with 700 ppm CO₂ on barley reported an increment of grain yield by 54% compared to 400 ppm (Alemayehu et al., 2014). Moreover, Manderscheid and Weigel (2006) evaluated the effects of eCO₂ using free-air carbon dioxide enrichment (FACE) at 550 ppm, and obtained yield increases of 7–15%.

The projected increase in biomass and grain yield of C3 crops due to eCO₂ is affected by certain environmental factors such as rising air temperature (Jaggard et al., 2010; Weigel et al., 1994). In general, increased temperature is primarily linked with evapotranspiration, acceleration of plant development, and consequently shortening of developmental phases, leading to early maturation and decreased yields (Barnabás et al., 2008; Cox et al., 2000; Hansen et al., 2000; Högy et al., 2013; Mangelsen et al., 2011; Vara Prasad et al., 2002). Studies on six major crops including barley have indicated that increasing the seasonal average temperature by 1 °C results in a significant grain yield reduction by 4 to 10% (Barnabás et al., 2008; Hatfield et al., 2011). Similarly, Clausen et al. (2011) found grain yield reduction in well-watered barley of 14% compared with the same level of eCO₂ and ambient temperature, while a 53% reduction in grain yield was recorded by another study (Alemayehu et al., 2014).

Besides, the positive effect of eCO₂ on aboveground biomass and grain yield, which varied between 20 to 90% (Fangmeier et al., 1996; Fangmeier et al., 2000; Weigel et al., 1994) were modified by the influence of nitrogen fertilizer (Manderscheid et al., 2009; Thompson and Woodward, 2007). N influences plant growth mainly through seasonal variation in the total green leaf area index and the accumulated radiation absorbed (Amthor, 2001). At low N fertilizer, eCO₂ effects on the yield of barley (Fangmeier et al., 2000) and other cereals (Kim et al., 2003) are usually small. This is reinforced by findings of FACE studies with wheat and rice (Högy and Fangmeier, 2008; Kim et al., 2003). Even though the pattern of yield response to eCO₂ and its interactions with temperature and nitrogen are similar within C3 crops, distinctions are evident across species, genotypes, and growing conditions (Connor, 2002).

Previous meta-analytic studies on C3 crops, such as wheat, rice, and soybean, have provided insights into the extent of the effects of eCO₂ on yield variables (Ainsworth, 2008; Broberg et al., 2019; Feng et al., 2008). However, despite its importance, the effect of eCO₂ and its interaction with temperature and N fertilizer on barley production has not been quantitatively reviewed using meta-analysis. The objectives of the current meta-analysis are therefore two-fold (1) to synthesize and summarise the mean response of barley yield variables to eCO₂, temperature, and N fertilization and, (2) to determine whether different CO₂ exposure methods, rooting conditions, genotypes, or different stress treatments significantly alter the mean response of barley to eCO₂.

Materials And Methods

Database development

Peer-reviewed primary literature focusing on barley yield responses to eCO₂, temperature, and N were searched on Scopus, Science Direct, and Google Scholar. The following keywords were used to search the literatures in various combinations “barley”, “elevated CO₂”, “interaction”, “interactive effects”, “high temperature”, “warming” and “N fertilizer”. The search was intended to be comprehensive, including all relevant studies that were published between 1991 and 2020. The final database covered a total of 76 articles out of which 36 were on the response of barley to eCO₂, 22 on the interaction of eCO₂ with temperature, and 18 on the interaction of eCO₂ with N. To minimize biases the following four inclusion criteria were applied for including studies in the database: (1) the ambient CO₂ level has to be ≤ 450 ppm (intended to represent the past and the near future concentration) and the eCO₂ has to be ≥ 451 ppm (representing the CO₂ concentration after 2050); (2) at least one of the selected yield variables is evaluated; (3) the response means, and sample sizes (n) are reported directly in the text, table or can be indirectly derived from figures, and (4) the CO₂ exposure technique is specified.

To test the interaction effect of eCO₂ with temperature and nitrogen, we extracted data from studies that included CO₂, temperature, and nitrogen treatments over the entire experimental period. Observations were considered independent within individual studies, if measurements were taken on different CO₂ concentration levels, genotypes, or combinations with various environmental stress factors following previous meta-analysis studies (Ainsworth et al., 2002; Gurevitch and Hedges, 1999). For experiments involving additional stress factors, such as O₃ and drought, the mean values of the control group were used. The final database included studies on the responses of barley aboveground biomass, grain number, thousand-grain weight (TGW), grain yield, and harvest index (HI). For each variable, response means and sample size were recorded from the treatment and control groups for each observation.

The putative factors likely to affect the response to eCO₂ of barley yield variables were further partitioned into several categories, such as CO₂ exposure methods (growth chamber, greenhouse, open-top chamber, and Free-air CO₂ enrichment), CO₂ treatment levels (451-550, 551-650, 651-750 ppm), air temperature

level (<15, 15-20, 21-25, >25 °C), N fertilizer level (0-50, 51-100, 101-150, 151-200, >200 kg ha⁻¹), rooting conditions (pot and field), and genotypes (Table 1).

Meta-analysis

Meta-analysis commonly describes the extent of an experimental treatment mean (\bar{y}_T) relative to the control treatment mean (\bar{y}_C) (Ainsworth et al., 2002). To calculate the magnitude of CO₂, temperature and N treatment effect on the selected variables, we used, as the effect size, the log response ratio (expressed as mean yield ratio), the natural log of the ratio of the mean yield of the experimental to the control group. The log-transformation can make the data better approximate the normal distribution, reduce skewness and make non-linear relationships more linear. For each of the three variables, we calculated the natural logarithm of the ratio of the mean response of the experimental treatment (\bar{y}_T) to the mean response of the control treatment (\bar{y}_C) as, $r = \ln(\bar{y}_T/\bar{y}_C)$ and its a percentage change from the control ($[r-1] \times 100$). Thus, the expected mean percentage change is positive for $r > 1$ but negative for $r < 1$. Linear mixed models were fitted, assuming that differences among studies within a class are due to both sampling error and random variation. As variance or related parameters were not reported in several studies, unweighted analyses were performed for all the variables. However, resampling and bootstrapping techniques were used to obtain the confidence intervals of the mean effect size (Gurevitch and Hedges, 1999). For each categorical variable, between-group heterogeneity (QB) was examined. Means of different categorical variables were considered significantly different from each other if their 95% confidence intervals (CI) did not overlap (Ainsworth et al., 2002; Gurevitch and Hedges, 1999). All the analyses were performed in R statistical software (R Core Team 2019). The linear mixed model was fitted using the R library *lme4*.

Table 1	
Summary of barley genotypes included in the meta-analysis	
Genotypes	Genotype description
Genebank accession	Collection of genotypes
Alexis	Two-row German malting barley cultivar, heat tolerant
Arena	Two-row Australian modern spring barley cultivar, powdery mildew tolerant
Anakin	Modern spring barley cultivar
Atem	European modern spring cultivar, drought tolerant
Aura	Spring malting barley cultivar, disease-tolerant
Gairdner	Australian malting barley, moderately tolerant to powdery mildew
Gammer-Dansk	Old landrace
Scarlett	Two-row German spring malting barley
Spring_cultivars	A group of spring cultivars
Theresa	Winter barley cultivar

Results

CO₂ effect on barley yield variables

Across all the 76 studies, a significant enhancement of most of the measured yield variables was observed at elevated eCO₂ levels (i.e., 451-750 ppm) relative to aCO₂ (\leq 450 ppm). On average, 23.8% [CI: 18.0–27.8%] CO₂-induced enhancement of aboveground biomass of barley was observed compared to barley plants grown under aCO₂ (Figure 1). Besides, grain yield was 27.4% [CI: 18.5–36.2%] higher under eCO₂, mainly due to higher grain number (24.8% [CI: 17.7-31.9%]) and thousand-grain weight (5.6% [CI: 3.5-8.1%]), however, the response of harvest index was not affected by eCO₂ (Figure 1). Furthermore, the response of barley yield varied with eCO₂ concentration level such that the response of aboveground biomass (28.7%) was the highest under the eCO₂ concentration level of 551-650 ppm relative to 451-550 ppm. Also, the responses of grain yield, grain number, TGW, and HI were all significantly higher under the highest eCO₂ concentration level (651-750 ppm) relative to all the lower levels (Table 2 and Figure 2).

Table 2

Between-group heterogeneity (QB) for eCO₂ effect size across different categorical variables

Variables	No. of studies	Genotype	CO ₂ -treatment level	CO ₂ -exposure technique	Rooting condition	Stress	
						N	Temperature
AGB	58	23.4***	22.5***	14.3***	21.9***	21.5***	16.8***
GN	34	15.6***	22.4**	23.5*	20.4***	18.6**	14.7**
GY	54	29.2**	14.3 ^{ns}	22.8 ^{ns}	26.9***	24.5***	20.7*
TGW	44	4.7***	5.7 ^{ns}	1.0 ^{ns}	4.8 ^{ns}	15.6 ^{ns}	-6.2 ^{ns}
HI	44	1.9 ^{ns}	2.3*	4.6 ^{ns}	1.8 ^{ns}	5.4 ^{ns}	-

AGB: aboveground biomass, GN: grain number, GY: grain yield, TGW: thousand-grain weight. Significance level: P < 0.001 (***), P < 0.01 (**), P < 0.05 (*) and not-significant (ns).

Interactive effect of eCO₂ with N fertilization and temperature treatments

Although only a few studies reported the interactive effects of CO₂ with N or temperature, the response of barley yield variables to eCO₂ was significantly affected by both N fertilizer and temperature treatments (Table 2). The highest responses of aboveground biomass (31% [CI: 26-39%]), TGW (13% [CI: 8-16%]), and HI (20.4% [CI: 15.2-24.2%]) to eCO₂ were observed under the N level of 151-200 kg ha⁻¹, relative to lower N levels. Besides, an enhancement of grain yield by 24.5% [CI: 19.1-30%] and grain number by 31.7% [CI: 27.5-36.9%] were recorded for eCO₂ combined with an N level of 101-150 kg ha⁻¹. There was no significant difference in the response of barley yield variables to eCO₂ between plants grown under N levels of 0-50 kg ha⁻¹ and 51-100 kg ha⁻¹ (Figure 3). On the other hand, the highest enhancement of aboveground biomass by 28.4% [CI: 20.7-34.4%], grain yield by 28% [CI: 22-32%], grain number by 36.4% [CI: 29-41%], and TGW by 11% [CI: 5.6-16%] were observed for the combination of eCO₂ with a temperature level of 21-25 °C (Figure 4). However, the response of aboveground biomass and grain yield declined by -12% [CI: -18-(-5)%] and -17% [CI: -23-(-10)%], respectively, when eCO₂ was combined with a high-temperature level (>25 °C). The response of TGW was unaffected by variation in eCO₂ under the other temperature levels (Figure 4).

Genotypic variation

Barley genotypes had a statistically significant effect on the response of aboveground biomass, grain number, and grain yield across all eCO₂ treatment levels (Table 2). The genotypes showed an increase in aboveground biomass, averaged across eCO₂ treatments, of 14.7 to 47.1%. The highest increase in aboveground biomass of 47.1% [CI: 38.7-53.5%] was observed for genotype “*Anakin*” followed by “*Genebank accessions*” (46.5% [CI: 38.1-51.3%]) and “*Scarlett*” (33.5% [CI: 29.7-38%]). By contrast, the genotype “*Theresa*” showed the lowest increase in aboveground biomass of 14.7% [CI: 6.3-23.1%] under eCO₂. The highest increase in grain number of 46.1% [CI: 38.3-52.7%] and grain yield of 57.1% [CI: 43.6-64.5%] were observed for “*Genebank accessions*”. The lowest effect of eCO₂ on the response of grain number of 9% [CI: 3-14%] and grain yield of 10.8% [CI: 6.2-16%] was observed for “*Theresa*”. The responses of TGW and harvest index to eCO₂ did not differ among the nine genotypes (Figure 5)

CO₂ exposure methods and rooting conditions

The responses of barley yield (except for TGW) to eCO₂ were significantly affected by the four CO₂ exposure methods (Table 2). Aboveground biomass increased by 38% [CI: 26.9-47.7%] in the GC, 21.1% [CI: 16.7-26.3%] in OTC, and 12% [CI: 6-17%] in FACE under eCO₂ compared to aCO₂ (Figure 6). The highest increase in barley grain yield under eCO₂ (49.8% [CI: 40.1-55.1%]) was observed for the plants grown in the GC. In contrast, barley plants that were grown under OTC had only a 9% [CI: 3-14%] higher harvest index, while the response declined by -5.2% [CI: -10.3-0%] under OTC (Figure 6). Barley yield response was also significantly affected by rooting conditions. Higher responses of aboveground biomass (31% [CI: 26-36%]), grain number (30% [CI: 23-35%]), and grain yield (41% [CI: 34-47%]) were obtained for the plants grown in pots. However, the responses of TGW and harvest index of barley were not affected by rooting conditions (Table 2 and Figure 7).

Discussion

Responses of barley yield components to atmospheric CO₂ enrichment

This quantitative review synthesises the literature on barley yield as a function of eCO₂, temperature, and N fertilizer treatments and provides insights into contemporary paradigms. The rise in atmospheric CO₂ causes mostly positive effects on the total biomass of C₃ plants such as barley, by stimulating net photosynthesis and reducing photorespiration (Drake et al., 1997; Mitterbauer et al., 2017; Schapendonk et al., 2000). The average increase in aboveground biomass by 23.8% under eCO₂ is similar to other reported estimates (Manderscheid et al., 2014; Thompson and Woodward, 1994; Weigel and Manderscheid, 2012). The aboveground biomass and yield showed similar patterns of increase with increasing level of eCO₂. Plants grown under a CO₂ treatment level of 551-650 ppm showed the highest response in aboveground biomass (28.7%) compared to lower CO₂ concentrations (450-550 ppm). Other

studies have also reported a significant increase in aboveground biomass in barley of 38% and 24% at 550 and 650 ppm, respectively (Fangmeier et al., 2000; Weigel et al., 1994). On the other hand, a maximum increase of about 70 to 110% in aboveground biomass of two barley genotypes was reported under a eCO₂ concentration of 500 ppm (Weigel et al., 1994). FACE experiments with wheat and several crop types also reported about 12% enhancement in aboveground biomass under eCO₂ (Ainsworth and McGrath, 2010; Kimball et al., 2002; Weigel and Manderscheid, 2012). Similarly, a positive effect of eCO₂ on aboveground biomass of 37% in soybean (Ainsworth et al., 2002), 10% in ryegrass (Weigel and Manderscheid, 2012), and 21% in oilseed rape (Högy et al., 2010a) have been observed. A meta-analytic study of 79 crop and wild species also documented an average enhancement of biomass by 28% across all species due to eCO₂ (Jablonski et al., 2002).

Likewise, a positive effect of eCO₂ was recorded for grain yield (27.4%) averaged across all the studies, mainly due to the significant and positive effect of eCO₂ on grain number (27.4%). These results support findings from other studies on barley, that have reported an increase in grain yield induced by eCO₂ (Fangmeier et al., 2000; Högy et al., 2009; Weigel et al., 1994). Similarly, studies with other C3 crops, for instance, wheat (Högy et al., 2010b), soybean (Ainsworth et al., 2002), and oilseed rape (Högy et al., 2010a) have also noted significant increases in grain yield due to CO₂ enrichment of 10, 24, and 18%, respectively. In agreement with the findings from other studies, the response of TGW was not affected by eCO₂, which reflects its lower market value (Schmid et al., 2016; Weigel et al., 1994). Comparing the response of harvest index under different eCO₂ levels, the highest increase was observed for plants grown under the highest concentration level of 651-750 ppm. However, the response of harvest index of barley was not affected across all the CO₂ levels, similar to findings of other studies (Fangmeier et al., 1996; Palta and Ludwig, 2000).

The interactive effect of eCO₂, N fertilizer, and temperature

Aboveground biomass and grain yield responses were lower when eCO₂ was combined with low N level (<50 kg ha⁻¹) than with higher N levels. Nevertheless, the important interactive effects of N with eCO₂ have only rarely been studied. High positive effects of eCO₂ on the response of aboveground biomass (31%) and harvest index (20.4%) were observed under the higher N level (151-200 kg ha⁻¹). Accordingly, barley plants fertilized with 140 kg ha⁻¹ N had 13% more aboveground biomass than plants fertilized with 80 kg ha⁻¹ under eCO₂ (Fangmeier et al., 2000). In addition, a significant decrease in grain yield of cereal crops of between 10 and 22% was recorded under a combination of eCO₂ and 50 kg ha⁻¹ N compared to 100 kg ha⁻¹ N (Manderscheid et al., 2009). The response of harvest index to eCO₂ also slightly decreases under low N fertilizer (Fangmeier et al., 1996). The CO₂ enrichment effect on biomass and grain yield of barley and other cereals is negative when N accessibility is condensed compared with when it is not in accord with findings of other studies.

Even though eCO₂ had a positive effect on the yield production of barley, the effect was low under high temperatures (>25 °C). High temperature had negative impacts on aboveground biomass and grain yield compared with optimal temperatures (Högy et al., 2019; Usui et al., 2014; Wheeler et al., 1996). Consequently, aboveground biomass and grain yield decreased by -12% and -17%, respectively, when eCO₂ was combined with high temperatures (>25 °C). This might be due to the shorter duration of crop growth and development under high temperatures, and the perturbation of processes related to carbon assimilation (Stone, 2001). TGW and harvest index were not affected by the interaction of eCO₂ with temperature but TGW can decrease significantly with eCO₂ at high temperatures (Alemayehu et al., 2014; Högy et al., 2013).

Variation in the response of barley yield to eCO₂

Genotypic variation

The significant effect of genotype on barley yield response to eCO₂ was might be related to the varietal character of genotypes. The modern genotypes had higher aboveground biomass. Thus, the modern spring barley genotype "*Anakin*" had 47.1% more aboveground biomass under eCO₂ than the other genotypes. The genotype "*Genebank accessions*" had the highest grain number (46.1%) and its grain yield increased by 57.1% under eCO₂. Other studies have similarly found modern barley genotypes to higher responses than old ones (15-34%) under CO₂ enrichment (Alemayehu et al., 2014; Plessl et al., 2005; Schmid et al., 2016). Similarly, a stronger positive effect of eCO₂ on grain yield was observed for the modern genotype "*Mauri*" (released in 2004) than for old genotypes released between 1952 and 1996 (Franzaring et al., 2013). This suggests breeding for the exploitation of eCO₂ might enhance future crop production. The lowest response to eCO₂ was recorded for the genotype "*Gammel Dansk*", consistent with findings of other CO₂ enrichment studies (Clausen et al., 2011).

However, modern genotypes do not necessarily always perform better than new ones at higher CO₂ levels. For example, the aboveground biomass of an older wheat genotype (52%) can increase more than that of a new genotype (39%) in response to increasing CO₂ (Hay and Gilbert, 2001). Furthermore, older wheat genotypes can have higher grain yield than newer genotypes under certain experimental conditions (Franzaring et al., 2013). The responses of harvest index and TGW to eCO₂ did not differ among the nine study barley genotypes but TGW can vary widely among other barley genotypes (Weigel et al., 1994). For example, a climate chamber experiment with old and new malt barley genotypes recorded higher TGW for the modern genotype "*Bambina*" (Schmid et al., 2016). A review of the harvest index of cereals found that modern genotypes had a significantly higher harvest index than older ones (HAY, 1995). Other trials with barley, oat, and oilseed rape have also shown that plant genotypes react differentially to climate change (Clausen et al., 2011; Fangmeier et al., 2000; Johannessen et al., 2005).

Experimental conditions

The response of crop yield to CO₂ enrichment can be significantly altered by CO₂ exposure methods. But the effect of CO₂ exposure methods can vary with several factors, including yield variable and crop species. Thus, enhancement of grain yield at eCO₂ can be lower under FACE experiments than under enclosure methods (Long et al. 2006). Similarly, aboveground biomass yield can be greater under FACE experiments than under the OTC or growth chamber (Tubiello et al., 2007). But rice yield can be higher for plants grown in OTCs than with FACE (Broberg et al., 2019). Our meta-analysis shows that the responses of aboveground biomass and grain yield in barley increased by 38 and 50%, respectively, when plants were grown in growth chambers, than when they were grown in FACE or OTC, in accord with earlier studies (Long et al., 2006). Similar results have been reported in meta-analyses of rice (Wang et al., 2015) and wheat (Wang et al., 2013). However, another meta-analysis of wheat noted no significant difference between FACE and OTC experiments with respect to the response of grain yield to eCO₂ (Feng et al., 2008). Nonetheless, no study seems to have directly compared the response to different CO₂ exposure methods of the same genotype grown under identical soil, environmental condition, and cultivation practice.

The effect of rooting condition on the response of barley yield to eCO₂ varies with other factors, such as yield variable and crop species. The aboveground biomass (31.3%), grain number (30%), and grain yield (41.3%) significantly increased for barley grown in a pot rather than under field conditions. However, the responses of TGW and harvest index were insensitive to the rooting condition. In agreement with our findings, previous meta-analyses have also reported higher yield responses of pot-rooted wheat plants under eCO₂ compared to field-rooted plants (Taub and Wang, 2008; Wang et al., 2013). But other studies have also reported similar responses of grain yield to eCO₂ for field-grown and pot-grown wheat plants (Feng et al., 2008). Lastly, a significant decline in TGW in wheat of up to 3% in OTC experiments can occur due to limited rooting volume (Ainsworth, 2008).

Conclusions

This meta-analysis quantified the effect of eCO₂ as a single factor and its interaction with N and temperature on barley production. A strong positive effect of eCO₂ was observed for aboveground biomass, grain yield, and grain number. However, the responses of aboveground biomass, grain number, and grain yield to eCO₂ were lower under limited N fertilizer (<50 kg ha⁻¹) or higher temperature (>25 °C). In general, the magnitude of the CO₂-induced effect on barley grain yield will depend on the future atmospheric CO₂ concentration and agronomic practices such as genotype choice, and growing conditions. Modern barley genotypes "*Anakin*" had higher aboveground biomass under eCO₂ than older ones, with "*Genebank accessions*" having the highest grain yield and grain number response to eCO₂. Uncertainties remain, however, regarding the responses to environmental stress of yield variables, mainly aboveground biomass, grain number, and grain yield. Field experiments that better characterize the

responses of barley and its interaction with additional stressors to eCO₂ can help reduce uncertainties due to climate change in estimating future food production.

Declarations

Funding:

This publication is an output of a Ph.D. scholarship at the University of Hohenheim in the framework of the project “German-Ethiopian SDG Graduate School: Climate Change Effects on Food Security (CLIFOOD)” between the University of Hohenheim (Germany) and the Hawassa University (Ethiopia), supported by the DAAD with funds from the Federal Ministry for Economic Cooperation and Development (BMZ).

Conflict of interest:

The authors have no conflict of interest.

Availability of data and material:

Data obtained for the current study is available.

Code availability:

Software application code is available.

Author Contributions:

M.G. and P.H. conceived and designed the study; data collection was made by M.G. M.G and W.A.M participated in the analysis of the data; M.G. wrote the paper with substantial input from P.H. B.H and W.A.M.

Acknowledgments:

The authors are grateful to Andrea Giovanna Nino Savala, Ireen Drebestedt, and Philipp Neher for their critical reading and valuable comments. Also, a special thank goes to CLIFOOD team members and coordinators. We also like to acknowledge DFG grant PI 377/24-1.

References

1. Ainsworth, E.A., 2008. Rice production in a changing climate: a meta-analysis of responses to elevated carbon dioxide and elevated ozone concentration. *Glob. Chang. Biol.* 14, 1642–1650. <https://doi.org/10.1111/j.1365-2486.2008.01594.x>
2. Ainsworth, E.A., Davey, P.A., Bernacchi, C.J., Dermody, O.C., Heaton, E.A., Moore, D.J., Morgan, P.B., Naidu, S.L., Ra, H.S.Y., Zhu, X.G., Curtis, P.S., Long, S.P., 2002. A meta-analysis of elevated [CO₂] effects on soybean (*Glycine max*) physiology, growth, and yield. *Glob. Chang. Biol.* 8, 695–709. <https://doi.org/10.1046/j.1365-2486.2002.00498.x>
3. Ainsworth, E.A., McGrath, J.M., 2010. Direct Effects of Rising Atmospheric Carbon Dioxide and Ozone on Crop Yields, in: *Advances in Global Change Research*. <https://doi.org/10.1007/978-90-481-2953-9-7>
4. Alemayehu, F.R., Frenck, G., van der Linden, L., Mikkelsen, T.N., Jørgensen, R.B., 2014. Can barley (*Hordeum vulgare* L.) adapt to fast climate changes? A controlled selection experiment. *Genet. Resour. Crop Evol.* 61, 151–161. <https://doi.org/10.1007/s10722-013-0021-1>
5. Amthor, J.S., 2001. Effects of atmospheric CO₂ concentration on wheat yield: a review of results from experiments using various approaches to control CO₂ concentration. *Field Crops Research* 73, 1–34
6. Barnabás, B., Jäger, K., Fehér, A., 2008. The effect of drought and heat stress on reproductive processes in cereals. *Plant, Cell Environ.* 31, 11–38. <https://doi.org/10.1111/j.1365-3040.2007.01727.x>
7. Broberg, M.C., Högy, P., Feng, Z., Pleijel, H., 2019. Effects of elevated CO₂ on wheat yield: Non-linear response and relation to site productivity. *Agronomy*. 9, 1–18. <https://doi.org/10.3390/agronomy9050243>
8. Clausen, S.K., Frenck, G., Linden, L.G., Mikkelsen, T.N., Lunde, C., Jørgensen, R.B., 2011. Effects of Single and Multifactor Treatments with Elevated Temperature, CO₂ and Ozone on Oilseed Rape and Barley. *J. Agron. Crop Sci.* 197, 442–453. <https://doi.org/10.1111/j.1439-037X.2011.00478.x>
9. Connor, D., 2002. Climate Change and Global Crop Productivity. *Crop Sci.* 42, 978. <https://doi.org/10.2135/cropsci2002.9780>
10. Conroyac, J.P., Seneweera, S., Basra, A.S., Rogers, G., Wooller, B.N., 1994. Influence of Rising Atmospheric CO₂ Concentrations and Temperature on Growth, Yield and Grain Quality of Cereal Crops. *Aust. J. Plant Physiol.* 21(6), 741 – 758. <https://doi.org/10.1071/PP9940741>
11. Cox, P.M., Betts, R.A., Jones, C.D., Spall, S.A., Totterdell, I.J., 2000. Acceleration of global warming due to carbon-cycle feedbacks in a coupled climate model. *Nature*. 408, 184–187(2000). <https://doi.org/10.1038/35041539>
12. Curtis, P.S., Wang, X., 1998. A meta-analysis of elevated CO₂ effects on woody plant mass, form, and physiology. *Oecologia*. 113(3), 299-313. <https://doi.org/10.1007/s004420050381>
13. Drake, B., Gonzalez-Meler, M., Long, S., 1997. More efficient plants: A consequence of rising atmospheric CO₂. *Plant Mol. Biol.* 48, 609–639. <https://doi.org/10.1146/annurev.arplant.48.1.609>

14. Fangmeier, A., Chrost, B., Högy, P., Krupinska, K., 2000. CO₂ enrichment enhances flag leaf senescence in barley due to greater grain nitrogen sink capacity. *Environ. Exp. Bot.* 44(2), 151-164. [https://doi.org/10.1016/S0098-8472\(00\)00067-8](https://doi.org/10.1016/S0098-8472(00)00067-8)
15. Fangmeier, A., Gruters, U., Vermehren, B., Jager, H.J., 1996. Responses of some cereal cultivars to CO₂ enrichment and tropospheric ozone at different levels of nitrogen supply. *J. Appl. Bot. Bot.*
16. Food and Agriculture Organization of the United Nations (2020). FAOSTAT database is available at <http://www.fao.org/faostat/en/#data/QC>. Accessed on 19th December 2020.
17. Feng, Z., Kobayashi, K., Ainsworth, E.A., 2008. Impact of elevated ozone concentration on growth, physiology, and yield of wheat (*Triticum aestivum* L.): A meta-analysis. *Glob. Chang. Biol.* 14, 2696–2708. <https://doi.org/10.1111/j.1365-2486.2008.01673.x>
18. Franzaring, J., Holz, I., Fangmeier, A., 2013. Responses of old and modern cereals to CO₂-fertilisation. *Crop Pasture Sci.* 64, 943–956. <https://doi.org/10.1071/CP13311>
19. Fróna, D., Szenderák, J., and Harangi-Rákos., M., 2019. The challenge of Feeding the World. *Sustainability.* 11, 5816. <https://doi:10.3390/su11205816>
20. Gurevitch, J., and Hedges, L. V., 1999. Statistical issues in ecological meta-analyses. *Ecology.* 80 (4), 1142-1149. [https://doi.org/10.1890/0012-9658\(1999\)080\[1142:SIEMA\]2.0](https://doi.org/10.1890/0012-9658(1999)080[1142:SIEMA]2.0)
21. Hansen, J., Sato, M., Ruedy, R., Lacis, A., Oinas, V., 2000. Global warming in the twenty-first century: An alternative scenario. *Proc. Natl. Acad. Sci. U. S. A.* 97(18), 9875-9880. <https://doi.org/10.1073/pnas.170278997>
22. Hatfield, J.L., Boote, K.J., Kimball, B.A., Ziska, L.H., Izaurralde, R.C., Ort, D., Thomson, A.M., Wolfe, D., 2011. Climate impacts on agriculture: Implications for crop production. *Agron. J.* 103, 351–370. <https://doi.org/10.2134/agronj2010.0303>
23. Hay, R.K.M., 1995. Harvest index: a review of its use in plant breeding and crop physiology. *Ann. Appl. Biol.* 126, 197–216. <https://doi.org/10.1111/j.1744-7348.1995.tb05015.x>
24. Hay, R.K.M., Gilbert, R.A., 2001. Variation in the harvest index of tropical maize: Evaluation of recent evidence from Mexico and Malawi. *Ann. Appl. Biol.* 138, 103–109. <https://doi.org/10.1111/j.1744-7348.2001.tb00090.x>
25. Högy, P., Fangmeier, A., 2008. Effects of elevated atmospheric CO₂ on grain quality of wheat. *J. Cereal Sci.* 48, 580–591. <https://doi.org/10.1016/j.jcs.2008.01.006>
26. Högy, P., Franzaring, J., Schwadorf, K., Breuer, J., Schütze, W., Fangmeier, A., 2010a. Effects of free-air CO₂ enrichment on energy traits and seed quality of oilseed rape. *Agric. Ecosyst. Environ.* 139, 239–244. <https://doi.org/10.1016/j.agee.2010.08.009>
27. Högy, P., Keck, M., Niehaus, K., Franzaring, J., Fangmeier, A., 2010b. Effects of atmospheric CO₂ enrichment on biomass, yield, and low molecular weight metabolites in wheat grain. *J. Cereal Sci.* 52(2), 215-220. <https://doi.org/10.1016/j.jcs.2010.05.009>
28. Högy, P., Kottmann, L., Schmid, I., Fangmeier, A., 2019. Heat, wheat, and CO₂: The relevance of timing and the mode of temperature stress on biomass and yield. *J. Agron. Crop Sci.* 205, 608–615.

<https://doi.org/10.1111/jac.12345>

29. Högy, P., Poll, C., Marhan, S., Kandeler, E., Fangmeier, A., 2013. Impacts of temperature increase and change in precipitation pattern on crop yield and yield quality of barley. *Food Chem.* 136, 1470–1477. <https://doi.org/10.1016/j.foodchem.2012.09.056>
30. Högy, P., Wieser, H., Köhler, P., Schwadorf, K., Breuer, J., Franzaring, J., 2009. Effects of elevated CO₂ on grain yield and quality of wheat: results from a three-year FACE experiment. *Plant Biol.* 11, 60–69. <https://doi.org/10.1111/j.1438-8677.2009.00230.x>
31. IPCC, 2018. Proposed outline of the special report in 2018 on the impacts of global warming of 1.5 °C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate cha. *IPCC - Sr15 2*, 17–20.
32. Jablonski, L.M., Wang, X., Curtis, P.S., 2002. Plant reproduction under elevated CO₂ conditions: A meta-analysis of reports on 79 crop and wild species. *New Phytol.* 156, 9-26. <https://doi.org/10.1046/j.1469-8137.2002.00494.x>
33. Jaggard, K.W., Qi, A., Ober, S., 2010. Possible changes to arable crop yields by 2050. *Philos. Trans. R. Soc. B Biol. Sci.* 365, 2835–2851. <https://doi.org/10.1098/rstb.2010.0153>
34. Johannessen, M.M., Mikkelsen, T.N., Nersting, L.G., Gullord, M., Von Bothmer, R., Jørgensen, R.B., 2005. Effects of increased atmospheric CO₂ on varieties of oat. *Plant Breed.* 124, 253–256. <https://doi.org/10.1111/j.1439-0523.2005.01096.x>
35. Kim, H.Y., Lieffering, M., Kobayashi, K., Okada, M., Miura, S., 2003. Seasonal changes in the effects of elevated CO₂ on rice at three levels of nitrogen supply: A free-air CO₂ enrichment (FACE) experiment. *Glob. Chang. Biol.* 9, 826–837. <https://doi.org/10.1046/j.1365-2486.2003.00641.x>
36. Kimball, B.A., Kobayashi, K., Bindi, M., 2002. Responses of agricultural crops to free-air CO₂ enrichment. *Adv. Agron.* 77, 293–368. [https://doi.org/10.1016/S0065-2113\(02\)77017-X](https://doi.org/10.1016/S0065-2113(02)77017-X)
37. Long, S.P., Ainsworth, E.A., Leakey, A.D.B., Nösbsrger, J., Ort, D.R., 2006. Food for thought: Lower-than-expected crop yield stimulation with rising CO₂ concentrations. *Science* 312(5782), 1918–1921. <https://doi.org/10.1126/science.1114722>
38. Manderscheid, R., Weigel, J., 2006. Responses of cereals and sugar beet in a crop rotation to free air carbon dioxide enrichment. *Bibliotheca Fragm Agron* 11:309–310.
39. Manderscheid, R., Erbs, M., Weigel, H.J., 2014. Interactive effects of free-air CO₂ enrichment and drought stress on maize growth. *Eur. J. Agron.* 52, 11–21. <https://doi.org/10.1016/j.eja.2011.12.007>
40. Manderscheid, R., Pacholski, A., Frühauf, C., Weigel, H.J., 2009. Effects of free air carbon dioxide enrichment and nitrogen supply on growth and yield of winter barley cultivated in a crop rotation. *F. Crop. Res.* 110, 185–196. <https://doi.org/10.1016/j.fcr.2008.08.002>
41. Mangelsen, E., Kilian, J., Harter, K., Jansson, C., Wanke, D., Sundberg, E., 2011. Transcriptome analysis of high-temperature stress in developing barley caryopses: Early stress responses and effects on storage compound biosynthesis. *Mol. Plant* 4, 97–115. <https://doi.org/10.1093/mp/ssq058>

42. Mitterbauer, E., Enders, M., Bender, J., Erbs, M., Habekuß, A., Kilian, B., Ordon, F., Weigel, H.J., 2017. Growth response of 98 barley (*Hordeum vulgare* L.) genotypes to elevated CO₂ and identification of related quantitative trait loci using genome-wide association studies. *Plant Breed.* 136, 483–497. <https://doi.org/10.1111/pbr.12501>
43. National Oceanic and Atmospheric Administration (NOAA). Science Report 2019. 2020, 84.
44. Palta, J.A., Ludwig, C., 2000. Elevated CO₂ during pod filling increased seed yield but not harvest index in indeterminate narrow-leaved lupin. *Aust. J. Agric. Res.* <https://doi.org/10.1071/AR99099>
45. Plessl, M., Heller, W., Payer, H.D., Elstner, E.F., Habermeyer, J., Heiser, I., 2005. Growth parameters and resistance against *Drechslera teres* of spring barley (*Hordeum vulgare* L. cv. Scarlett) grown at elevated ozone and carbon dioxide concentrations. *Plant Biol.* 7, 694–705. <https://doi.org/10.1055/s-2005-873002>
46. Poorter, H., Navas, M.L., 2003. Plant growth and competition at elevated CO₂: On winners, losers, and functional groups. *New Phytol.* 157(2), 175-198. <https://doi.org/10.1046/j.1469-8137.2003.00680.x>
47. R Core Team (2019). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
48. Schapendonk, A.H.C.M., Van Oijen, M., Dijkstra, P., Pot, C.S., Jordi, W.J.R.M., Stoop, G.M., 2000. Effects of elevated CO₂ concentration on photosynthetic acclimation and productivity of two potato cultivars grown in open-top chambers. *Aust. J. Plant Physiol.* 27, 1119–1130. <https://doi.org/10.1071/PP99205>
49. Schmid, I., Franzaring, J., Müller, M., Brohon, N., Calvo, O.C., Högy, P., Fangmeier, A., 2016. Effects of CO₂ Enrichment and Drought on Photosynthesis, Growth, and Yield of an Old and a Modern Barley Cultivar. *J. Agron. Crop Sci.* 202, 81–95. <https://doi.org/10.1111/jac.12127>
50. Stone, P.J., 2001. The effects of heat stress on cereal yield and quality., in: *Crop Responses and Adaptations to Temperature Stress*.
51. Taub, D.R., Wang, X., 2008. Why are nitrogen concentrations in plant tissues lower under elevated CO₂? A critical examination of the hypotheses. *J. Integr. Plant Biol.* 50(11), 1365-1374. <https://doi.org/10.1111/j.1744-7909.2008.00754.x>
52. Thompson, G.B., Woodward, F.I., 2007. Some influences of CO₂ enrichment, nitrogen nutrition, and competition on grain yield and quality in spring wheat and barley. *J. Exp. Bot.* <https://doi.org/10.1093/jxb/45.7.937>
53. Tubiello, F., Soussana, J.-F., Howden, S., 2007. Climate change and food security special feature: in: *Crop and Pasture Response to Climate Change*. *Proc Natl Acad Sci USA.* 104(50), 19686–19690. <https://doi.org/10.1073/pnas.0701728104>
54. United Nations, 2011. *World Population Prospects: The 2010, Volume 1: Comprehensive Tables I*, 169.
55. Usui, Y., Sakai, H., Tokida, T., Nakamura, H., Nakagawa, H., Hasegawa, T., 2014. Heat-tolerant rice cultivars retain grain appearance quality under free-air CO₂ enrichment. *Rice* 7(1), 6

<https://doi.org/10.1186/s12284-014-0006-5>

56. Vara Prasad, P. V., Boote, K.J., Hartwell Allen, L., Thomas, J.M.G., 2002. Effects of elevated temperature and carbon dioxide on seed-set and yield of kidney bean (*Phaseolus vulgaris* L.). *Glob. Chang. Biol.* 8, 710–721. <https://doi.org/10.1046/j.1365-2486.2002.00508.x>
57. Wang, J., Wang, C., Chen, N., Xiong, Z., Wolfe, D., Zou, J., 2015. Response of rice production to elevated [CO₂] and its interaction with rising temperature or nitrogen supply: a meta-analysis. *Clim. Change* 130, 529–543. <https://doi.org/10.1007/s10584-015-1374-6>
58. Wang, L., Feng, Z., Schjoerring, J.K., 2013. Effects of elevated atmospheric CO₂ on physiology and yield of wheat (*Triticum aestivum* L.): A meta-analytic test of current hypotheses. *Agric. Ecosyst. Environ.* 178, 57–63. <https://doi.org/10.1016/j.agee.2013.06.013>
59. Weigel, H.J., Manderscheid, R., 2012. Crop growth responses to free-air CO₂ enrichment and nitrogen fertilization: Rotating barley, ryegrass, sugar beet, and wheat. *Eur. J. Agron.* 43, 97–107. <https://doi.org/10.1016/j.eja.2012.05.011>
60. Weigel, H.J., Manderscheid, R., Jäger, H.J., Mejer, G.J., 1994. Effects of season-long CO₂ enrichment on cereals. I. Growth performance and yield. *Agric. Ecosyst. Environ.* 48, 231–240. [https://doi.org/10.1016/0167-8809\(94\)90105-8](https://doi.org/10.1016/0167-8809(94)90105-8)
61. Wheeler, T.R., Hong, T.D., Ellis, R.H., Batts, G.R., Morison, J.I.L., Hadley, P., 1996. The duration and rate of grain growth, and harvest index, of wheat (*Triticum aestivum* L.) in response to temperature and CO₂. *J. Exp. Bot.* 47(298), 623-630. <https://doi.org/10.1093/jxb/47.5.623>

Figures




Figure 1

Relative percentage change in barley yield response to eCO₂. The symbols represent the percentage change (\pm 95% CI) in response relative to the corresponding control. The sample size is given in parenthesis. AGB, aboveground biomass; GN, grain number; GY, grain yield; HI, harvest index; TGW, thousand-grain weight.




Figure 2

Relative percentage change in barley yield response to three levels of CO₂ treatment (451-550, 551-650, 651-750 ppm). The symbols represent the percentage change (± 95% CI) in response relative to the corresponding control. The numbers of observations are given in parentheses. AGB, aboveground biomass; GN, grain number; GY, grain yield; HI, harvest index; TGW, thousand-grain weight.




Figure 3

Relative percentage change in barley yield response to eCO₂ under four different N fertilization treatments. The four levels of N were 0-50, 51-100, 101-150, 151-200 kg ha⁻¹. The symbols represent the percentage change (\pm 95% CI) in response relative to the corresponding control. The sample size is given in parenthesis. AGB, aboveground biomass; HI, harvest index; TGW, thousand-grain weight.




Figure 4

Relative percentage change in barley yield response to eCO₂ under four different temperature treatments. The four temperature levels were <15, 15-20, 21-25, >25 °C. The symbols represent the percentage change (\pm 95% CI) in response relative to the corresponding control. The sample size given in parenthesis. AGB, aboveground biomass; GN, grain number; GY, grain yield; TGW, thousand-grain weight.




Figure 5

Relative percentage change in yield response to eCO₂ for nine different barley genotypes. The symbols represent the percentage change (\pm 95% CI) in response relative to the corresponding control. The sample size is given in parenthesis. AGB, aboveground biomass; GN, grain number; GY, grain yield; HI, harvest index; TGW, thousand-grain weight.




Figure 6

Relative percentage change in barley yield response to eCO₂ under four different CO₂ exposure methods. The four exposure methods were free-air CO₂ enrichment (FACE), growth chambers (GC), greenhouses (GH), open-top chambers (OTC). The symbols represent the percentage change (\pm 95% CI) in response relative to the corresponding control. The sample size is given in parentheses. AGB, aboveground biomass; GN, grain number; GY, grain yield; HI, harvest index; TGW, thousand-grain weight.




Figure 7

Relative percentage change in barley yield response to eCO₂ under two different rooting conditions (Field rooted and Pot rooted). The symbols represent the percentage changes (\pm 95% CI) in response relative to the corresponding controls. The sample size is given in parenthesis. AGB, aboveground biomass; GN, grain number; GY, grain yield; HI, harvest index; TGW, thousand-grain weight.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementaryMekidesGardi.xlsx](#)