

Long Non-Coding RNA FAM83H-AS1 Acts as a Potential Oncogenic Driver in Human Ovarian Cancer

Xiaolei Yuan

Second Affiliated Hospital of Harbin Medical University

Ying Huang

Harbin Medical University

Man Guo

Second Affiliated Hospital of Harbin Medical University

Xiaowei Hu

Tumor Hospital of Harbin Medical University

Peiling Li (✉ samanly@126.com)

Second Affiliated Hospital of Harbin Medical University

Research

Keywords: Ovarian cancer, lncRNA FAM83H-AS1, network, survival, immune

Posted Date: July 22nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-46762/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published on January 7th, 2021. See the published version at <https://doi.org/10.1186/s13048-020-00756-y>.

Abstract

Objective Ovarian cancer (OC) is one of the most aggressive women cancers with increasing incidence and mortality rates worldwide. Long non-coding RNAs (lncRNAs) could be major players in OC process. Although FAM83H antisense RNA1 (FAM83H-AS1) is demonstrated to play an important role in many cancers, the detailed function and mechanism has not been reported in OC.

Results We integrated multiple kinds of bioinformatics approaches and experiments validated method to evaluate functions of FAM83H-AS1 in OC. Some differentially expressed lncRNAs were identified between OC and normal control tissues. FAM83H-AS1 was one of most differentially expressed lncRNAs and up-regulated in multiple cancer types. Specially, expression of FAM83H-AS1 was higher in OC and showed difference in diverse stages. High FAM83H-AS1 expression is associated with worse pan-cancer and OC outcomes. FAM83H-AS1-centric network including lncRNA-miRNA, lncRNA-protein and lncRNA-mRNA ceRNA network were constructed to infer the function and mechanism of FAM83H-AS1. There were two methylation sites including cg01399317 and cg20519035 located at FAM83H-AS1. The methylation level of cg01399317 was correlated with gene expression of FAM83H-AS1. The expression level of FAM83H-AS1 was correlated with infiltration level of immune cell including macrophage, neutrophil and dendritic cell in OC patients. Lastly, qRT-PCR showed that the expression of FAM83H-AS1 was higher in OC tissues than normal control tissues.

Conclusion Collectively, these results indicated that FAM83H-AS1 may act as an oncogenic driver and it may be a potential therapy target in OC.

Background

In the female reproductive system, ovarian cancer (OC) is one of the most malignant tumors (1). As cancer statistics in China and United States suggested that the mortality rate of OC has been rising for the past few years (2, 3), there will be approximately 22,240 new cases of ovarian cancer diagnosed and 14,070 ovarian cancer deaths in the United States (4). OC accounts for 2.5% of all female malignant tumors and part of deaths for cancer patients due to poor survival which largely driven by late stage diagnoses (5). Surgical resection combined with platinum-based and taxane-based chemotherapy is the major and standard approach for advanced-stage ovarian treatment (6). Although advancing insight about mechanism and treatment of OC has been evolved rapidly in recent years, survival rates have improved only slightly over the past 3 decades. Thus, improving prevention and early detection based on identifying molecular biomarkers are effective ways to enhance survival for OC patients.

Long non-coding RNAs (lncRNAs), which are defined as RNA transcripts of > 200 nucleotides that are not translated into protein (7). lncRNAs are a highly versatile class of transcripts that have sparked new lines of research in nearly all fields of the life sciences (8). More recently, emerging studies have identified lncRNAs as major players in many kinds of cancer processes (9). Many studies revealed that lncRNAs play essential roles in proliferation, migration, and invasion of cancers including OC (10, 11). For example, lncRNA LINC00176 was highly expressed in OC tissues as well as in OC cell lines, respectively. Knockdown of lncRNA LINC00176 suppresses OC progression by BCL-mediated down-regulation of ceruloplasmin (12). Li et al. suggested that lncRNA UCA1 was upregulated in cisplatin-resistant patient tissues and cell lines. Knockdown of UCA1 inhibited cell proliferation and promoted the cisplatin-induced cell apoptosis in OC cells (13). The expression of lncRNA ABHD11-AS1 in OC tissues was higher compared to normal ovarian tissue. Overexpression of ABHD11-AS1 promoted OC cell proliferation, invasion and migration, and inhibited apoptosis (14).

Recently, a novel lncRNA, FAM83H antisense RNA1 (FAM83H-AS1), demonstrated important roles in many cancers. Zhang et al. reported that FAM83H-AS1 is associated with clinical progression and modulates cell proliferation, migration, and invasion in bladder cancer (15). FAM83H-AS1 was found overexpressed in HPV-16 positive cervical cancer cell lines in an HPV-16 E6-dependent manner but independently of p53 regulation (16). Xu et al. indicated that the expression of FAM83H-AS1 was higher in glioma tissues and cell lines and overexpression of FAM83H-AS1 was associated with poor prognosis of glioma. Specially, FAM83H-AS1 was upregulated in OC (17). However, underlying mechanisms of FAM83H-AS1 regulating functions in OC have yet to be elucidated.

In present study, differentially expressed lncRNAs were identified between OC and normal ovarian tissues. FAM83H-AS1 was differentially expressed in many types of cancers. Specially, the expression of FAM83H-AS1 was higher in OC and fluctuated among diverse stages of OC. FAM83H-AS1 was also associated with survival in pan-cancer and OC. FAM83H-AS1-centric network including lncRNA-miRNA, lncRNA-protein and lncRNA-mRNA ceRNA network were constructed. qRT-PCR showed that FAM83H-AS1 was up-regulated in OC tissues. Overall, our findings indicated that FAM83H-AS1 may mediate the oncogenesis process and can be regarded as a prognostic biomarker in OC.

Results

Some dysregulated genes and lncRNAs were identified in OC

We identified differentially expressed coding genes and lncRNAs between OV tissues and normal control tissues to screen candidate genes. 2419 differentially expressed lncRNAs and coding genes were identified based on t-test ($P < 0.001$) for OV patients (Figure 1A). There were 920 (38.03%) up-regulated and 1499 (61.97%) down-regulated for all the differentially expressed lncRNAs and coding genes (Figure 1B). We further focused on dysregulated lncRNAs in OC. There were 64 (22.70%) and 218 (77.30%) up and down-regulated lncRNAs in OC tissues compared with normal ovarian tissues (Figure 1C). The results indicated that some dysregulated lncRNAs may play essential roles in OC patients. Specially, differential expression of lncRNA FAM83H-AS1 was extremely significant ($P = 1.72 \times 10^{-101}$, fold change value = 3.781). Thus lncRNA FAM83H-AS1 was considered as a candidate biomarker for OC to perform follow-up analyses.

FAM83H-AS1 expression is dysregulated in human pan-cancer and OC

In order to characterize FAM83H-AS1 in OC, we first explore the expression of FAM83H-AS1 in other cancers. We downloaded and analyzed TCGA data to assess the expression of FAM83H-AS1 in pan-cancer. The result indicated that expression of FAM83H-AS1 in all the cancer types including bladder urothelial carcinoma (BLCA); breast invasive carcinoma (BRCA); cervical squamous cell carcinoma (CESC); cholangiocarcinoma (CHOL); colon adenocarcinoma (COAD); esophageal carcinoma (ESCA); head and neck squamous cell carcinoma (HNSC); lung squamous cell carcinoma (LUSC); liver hepatocellular carcinoma (LIHC); lung adenocarcinoma (LUAD); kidney renal papillary cell carcinoma (KIPR); prostate adenocarcinoma (PRAD); rectum adenocarcinoma (READ); skin cutaneous melanoma (SKCM); stomach adenocarcinoma (STAD); pancreatic adenocarcinoma (PAAD); thymoma, (THYM) and ovarian serous cystadenocarcinoma (OV) was significantly differential compared with their noncancer counterparts (Fig. 2A). And the expression of FAM83H-AS1 was up-regulated in any cancer type. FAM83H-AS1 was also dysregulated in integrated pan-cancer patients (Fig. 2B). The abnormal expression of FAM83H-AS1 was also observed in OC (Fig. 2C). In addition, we also found that expression level of FAM83H-AS1 changed in diverse stages of OC (Fig. 2D).

FAM83H-AS1 was associated with prognosis in pan-cancer and OC

Kaplan–Meier analysis and the log-rank test were used to explore the association between FAM83H-AS1 expression and survival status in OC patients in present study. We found higher expression of FAM83H-AS1 was associated with worse overall survival in pan-cancer ($P < 0.0001$, Fig. 3A). As except, higher expression of FAM83H-AS1 was also associated with worse overall survival in OC patients ($P = 0.044$, Fig. 3B). The results indicated that lncRNA FAM83H-AS1 maybe could as a unfavorable prognostic biomarker for OC patients.

FAM83H-AS1-centric lncRNA-miRNA, lncRNA-protein and lncRNA-mRNA ceRNA networks could show the special function of FAM83H-AS1

We try to explore the function of FAM83H-AS1 based on studying its interacted miRNA and coding genes using multiple networks. First, a lncRNA-mRNA network was constructed, which contained 22 interacted miRNAs (Fig. 4A). The co-expression of FAM83H-AS1 and any interacted miRNA were also calculated. The expression of miR-211 was significantly positive correlated with FAM83H-AS1 in OC patients (Fig. 4B). Present study demonstrated that the expression of miR-211 is differential expressed in OC tissues and cell lines compared to normal epithelial ovarian tissue and human ovarian surface epithelial cells, respectively. miR-211 was also found to arrest cells in the G0/G1-phase, inhibit proliferation and induce apoptosis (18). Second, a lncRNA-protein (or coding gene) network was constructed (Fig. 4C). There were three coding genes could interacted with FAM83H-AS1. These three coding genes all play essential roles in cancers (19–21). At last, we also constructed a lncRNA-mRNA ceRNA network (Fig. 4D). TP53 was a ceRNA of FAM83H-AS1 and mutations in the TP53 gene are still by far the most frequent genomic event in cancer genomes (22). All the results indicated that FAM83H-AS1 maybe play its role by interacting with some OC-related coding genes and miRNAs.

Expression level of FAM83H-AS1 was associated with methylation and immune

We found there were two methylation sites including cg01399317 and cg20519035 located at FAM83H-AS1 (Fig. 5A). The methylation level of cg01399317 was correlated with gene expression of FAM83H-AS1 ($P = 0.011$, Fig. 5B). Recent years, immunotherapy have resulted in clinical success in treating late-stage cancers (23). Some immune signature genes have also been identified and could be applied to characterize immune infiltrates and predict clinical outcome for cancers. We also explored the associations between immune and FAM83H-AS1. The expression level of FAM83H-AS1 was associated with infiltration level of immune cell including macrophage ($P = 1.34e-9$), neutrophil ($P = 7.93e-03$) and dendritic cell ($P = 2.33e-03$) in OC patients (Fig. 5C). The results indicated that FAM83H-AS1 maybe could be used to characterize immune infiltrates for OC patients.

FAM83H-AS1 was up-regulated in OC tissues by qRT-PCR validation

FAM83H-AS1 levels was significantly differentially expressed in OC tissues compared with corresponding adjacent non-tumor tissues ($P = 0.045$, Fig. 6A). The fold-change value of FAM83H-AS1 was 2.9 in OC and control tissues. The expression level of FAM83H-AS1 was significantly higher in OC tissues than control tissues (Fig. 6B).

Discussion

Multiple lines of evidence suggested that non-coding RNAs are majority of genome transcripts, however, only a small part of them have been characterized to be biologically functional (24). In recent years, more and more studies focused on implicating the role of lncRNAs in diseases including cancers (25–27). Some studies had reported that lncRNA FAM83H-AS1 was associated with many types of cancers such as breast cancer (28), colorectal cancer (29) and colon cancer (30). However, the function and mechanism of FAM83H-AS1 in OC has not been systemically studied.

In present study, we found FAM83H-AS1 as a novel and essential lncRNA involved in OC. We showed that FAM83H-AS1 was significantly upregulated in many kinds of cancer tissues compared with control normal tissues. In OC patients, FAM83H-AS1 was one of top differential expressed lncRNAs. Specially, expression of FAM83H-AS1 also showed difference in diverse stage of OC. Higher expression of FAM83H-AS1 was associated with worse overall survival in pan-cancer and OC. All above results indicated that FAM83H-AS1 may act as an oncogenic driver in OC.

In order to further explore the biological function and mechanism of FAM83H-AS1, some regulatory networks were constructed. Previous studies suggested that interacted or co-expressed coding genes could represented the function of lncRNAs (31). Thus, we try to use the function of coding genes and miRNAs to estimate the role of FAM83H-AS1 in OC. In present study, the expression of miR-211 was significantly positive correlated with FAM83H-AS1 in OC patients. The

role of miR-211 in diagnosis, prognosis and treatment in OC had been reported in many studies. For example, wang et al. found that miR-211 inhibited most of DNA damage response-related genes, and proposed that miR-211 might affect the sensitivity of OC cells to platinum by targeting multiple DNA damage response-related genes and thereby determine the prognosis of OC (32). In addition, miR-211 could sponge lncRNA MALAT1 to suppress tumor growth and progression through inhibiting PHF19 in OC (33). We could infer that FAM83H-AS1 maybe also play a key role in OC based on above results.

Many evidences taken together suggested that OC patients could potentially benefit from immunotherapy (34). Identifying immune-related targets could provide assistance for immunotherapy of OC patients. In our study, we also explore the relationships between immune and FAM83H-AS1. The expression level of FAM83H-AS1 was associated with infiltration level of immune cell including macrophage, neutrophil and dendritic cell in OC patients. The result indicated that FAM83H-AS1 had potential to become a immune-related signature in OC.

Conclusions

The present study clarified the role of lncRNA FAM83H-AS1 in OC patients. It also demonstrated that the associations between immune and FAM83H-AS1 in OC. The analyses are helpful to identify specific biomarker and drug repurposing candidates for OC. In summary, the present study expands the oncogenic lncRNA landscape of OC and reveals that lncRNA FAM83H-AS1 may act as a potential therapy target for OC.

Materials And Methods

Obtain and procession of public pan-cancer and OC data

We obtained lncRNA, miRNA, gene expression and methylation (level 3) data, as well as clinical data of all cancer types including OC, from The Cancer Genome Atlas (TCGA, Release: 2017-09-08, <https://portal.gdc.cancer.gov/>). The gene expression data of normal tissues for ovarian was obtained from GTEX portal (<https://www.gtexportal.org/home/index.html>). Patients in all the cancer types were integrated as pan-cancer patients. The sample numbers in each cancer type were shown in **Table S1**. To filter gene, miRNA, and lncRNA not expressed across all samples, the items with expression values of 0 in all of the samples were excluded. Any remaining expression values of 0 were set to the minimum value of all samples, and all values were log2-transformed. Expression of FAM83H-AS1 was dichotomized using median expression as the cutoff to define “high value” at or above the median versus “low value” below the median.

Construction of lncRNA-miRNA, lncRNA-protein and lncRNA-mRNA ceRNA networks for FAM83H-AS1

In order to describe the functions of FAM83H-AS1, some interacted regulatory networks were constructed. lncRNA-miRNA network was constructed based on experimentally verified associations between miRNAs and lncRNAs were identified in starBase v3.0 (<http://starbase.sysu.edu.cn/>) (35) and DIANA-LncBase 3.0 (www.microna.gr/LncBase) (36). lncRNA-protein data was download from starBase v3.0 and NPInter v2.0 (37) supported by AGO CLIP-seq data. In order to build the lncRNA-mRNA ceRNA network, a hypergeometric test was used to evaluate whether the two lncRNAs have a potential ceRNA relationship by considering their shared interactive miRNAs. All the networks were constructed by Cytoscape 3.3.0 (<http://www.cytoscape.org/>).

Differential expressed and co-expressed analyses

T-test was used to calculate the differential expression of all genes and lncRNAs between cancer and normal control samples. Pearson's correlation coefficients (PCCs) were calculated between FAM83H-AS1 and its interacted miRNAs in lncRNA-miRNA network. In addition, co-expression of methylation level and FAM-83H-AS1 expression was also calculated.

Survival analysis for the FAM83H-AS1 in pan-cancer and OC

The patients were divided into two groups based on median value of FAM83H-AS1 expression. Kaplan-Meier method and log-rank test were used to evaluate the survival difference in patients with high and low expression of FER1L4. $P < .05$ was regarded as statistically significant.

Tumor-immune infiltrating cells associated with FAM83H-AS1 in OC

The associations between all tumor-immune infiltrating cells and the FAM83H-AS1 were analyzed via the Tumor Immune Estimation Resource (TIMER) platform (<https://cistrome.shinyapps.io/timer/>), a web tool for studying tumor-infiltrating immune cells and their interactions with cancer cells (38). B-cells, CD4+T-cells, CD8+T-cells, dendritic cells, macrophages and neutrophils were included in the correlated analyses.

Patients and tissue samples

OC patients with a histological diagnosis who had undergone surgical resection and had not received chemotherapy or radiotherapy were extracted in our study. Lastly, eight OC tissues and corresponding adjacent normal ovarian tissues were obtained from The Third Affiliated Hospital of Harbin Medical University. The tissue samples were frozen in liquid nitrogen and stored at -80°C until experiment. The pathological diagnosis were confirmed by three independent senior pathologists. The study was approved by the Research Ethics Committee of The Third Affiliated Hospital of Harbin Medical University. All

patients received written informed consent and disposed of specimens in accordance with accepted ethical standards. The clinicopathological features of all patients are indicated in Table 1.

Table 1:
The clinicopathological features of all patients.

| No. | Age (year) | Tumor size (cm) | Lymph node metastasis | Stage | Pathological type | Treatment | ER positive cells (%) | PR positive cells (%) | WT-1 | P53 | KI67 (%) | MDR |
|-----|------------|-----------------------------|-----------------------|-------|---|-----------|-----------------------|-----------------------|------|-----|----------|-----|
| 1 | 53 | R:13x13x10 | No | IA | Advanced serous adenocarcinoma of right ovary | No | 70 | < 5% | + | 3+ | 70 | + |
| 2 | 48 | L:4.5x3.5x2.0,R:7.5x4.5x3.0 | No | IIB | Advanced serous adenocarcinoma of bilateral ovary | No | 90 | 40 | — | 3+ | 60 | — |
| 3 | 43 | L:7x6x5,R:8x6x5 | Yes | IIB | Advanced serous adenocarcinoma of bilateral ovary | No | 70 | 30 | + | 2+ | 70 | - |
| 4 | 56 | L:8x7x7 | Yes | IIC | Advanced serous adenocarcinoma of left ovary | No | 90 | 5 | + | 3+ | 80 | - |
| 5 | 56 | L:8x7x6 | No | IA | Advanced serous adenocarcinoma of left ovary | No | 70 | — | + | 3+ | — | — |
| 6 | 53 | L:20x16x8 | No | IC | Advanced serous adenocarcinoma of left ovary | No | 50 | 10 | + | + | 30 | — |

Note: R:right ovary; L:left ovary

Quantitative real-time reverse transcription PCR (qRT-PCR)

Total RNA was extracted from fresh frozen samples and cells using Trizol Reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. Total RNA (2 µg) was reverse-transcribed into cDNA using Transcriptor First Strand cDNA Synthesis Kit (Roche, Vilvoord, Brussel, Belgium). The relative levels of EGOT to glyceraldehyde 3-phosphate dehydrogenase (GAPDH) control transcripts were determined by qPCR using the ABI 7500 Fast Real-Time PCR System (Invitrogen). The primer sequences were as follows. FAM83H-AS1: forward 5'-ACTACAGGCACCCACCACCAC-3', reverse 5'-TGAGACGGGCGGGATCACAAGG-3'; GAPDH: forward 5'-ACCACAGTCCATGCCATCAC-3', reverse 5'-TCCACCCTGTTGCTGTA-3'. The qRT-PCR amplification was performed in triplicate reactions starting at 95°C for 10 min, followed by 40 cycles at 95°C for 10 s, and 60°C for 60 s. Quantitative normalization of EGOT cDNA was performed in each sample using GAPDH expression as an internal control. The relative level of EGOT transcripts to control GAPDH was determined by the 2^{-ΔΔCT} method. Each sample was examined in triplicate.

Declarations

Funding

This study was supported by the National Natural Science Foundation of China (grant nos. 81660448 and 81360401), the special Health Technical Personnel Training program of Yunnan, China (grant no.L-201612) the Natural Science Foundation of Yunnan, China (grant no.2017FE468-006).

Disclosure of interest

The authors declare that they have no competing interest.

Acknowledgments

Not applicable.

Availability of data and material

The data that support the findings of this study are available from the corresponding author upon reasonable request.

Ethics approval and consent to participate

Not applicable.

Patient consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable.

Authors' contributions

LPL conceived and designed the experiments, YXL, HY, GM and HXW analyzed the data, and YXL wrote the manuscript.

References

1. J. Krzystyniak, L. Ceppi, D. S. Dizon, M. J. Birrer, Epithelial ovarian cancer: the molecular genetics of epithelial ovarian cancer. *Ann Oncol* **27 Suppl 1**, i4-i10 (2016); published online EpubApr (10.1093/annonc/mdw083).
2. W. Chen, R. Zheng, P. D. Baade, S. Zhang, H. Zeng, F. Bray, A. Jemal, X. Q. Yu, J. He, Cancer statistics in China, 2015. *CA Cancer J Clin* **66**, 115-132 (2016); published online EpubMar-Apr (10.3322/caac.21338).
3. R. L. Siegel, K. D. Miller, A. Jemal, Cancer statistics, 2015. *CA Cancer J Clin* **65**, 5-29 (2015); published online EpubJan-Feb (10.3322/caac.21254).
4. L. A. Torre, B. Trabert, C. E. DeSantis, K. D. Miller, G. Samimi, C. D. Runowicz, M. M. Gaudet, A. Jemal, R. L. Siegel, Ovarian cancer statistics, 2018. *CA Cancer J Clin* **68**, 284-296 (2018); published online EpubJul (10.3322/caac.21456).
5. J. Ubachs, J. Ziemons, I. J. G. Minis-Rutten, R. Kruitwagen, J. Kleijnen, S. Lambrechts, S. W. M. Olde Damink, S. S. Rensen, T. Van Gorp, Sarcopenia and ovarian cancer survival: a systematic review and meta-analysis. *J Cachexia Sarcopenia Muscle* **10**, 1165-1174 (2019); published online EpubDec (10.1002/jcsm.12468).
6. A. M. Nick, R. L. Coleman, P. T. Ramirez, A. K. Sood, A framework for a personalized surgical approach to ovarian cancer. *Nat Rev Clin Oncol* **12**, 239-245 (2015); published online EpubApr (10.1038/nrclinonc.2015.26).
7. K. C. Wang, H. Y. Chang, Molecular mechanisms of long noncoding RNAs. *Mol Cell* **43**, 904-914 (2011); published online EpubSep 16 (10.1016/j.molcel.2011.08.018).
8. P. Grote, R. A. Boon, LncRNAs Coming of Age. *Circ Res* **123**, 535-537 (2018); published online EpubAug 17 (10.1161/CIRCRESAHA.118.313447).
9. D. H. Bach, S. K. Lee, Long noncoding RNAs in cancer cells. *Cancer Lett* **419**, 152-166 (2018); published online EpubApr 10 (10.1016/j.canlet.2018.01.053).
10. T. Worku, D. Bhattarai, D. Ayers, K. Wang, C. Wang, Z. U. Rehman, H. S. Talpur, L. Yang, Long Non-Coding RNAs: the New Horizon of Gene Regulation in Ovarian Cancer. *Cell Physiol Biochem* **44**, 948-966 (2017)10.1159/000485395).
11. L. L. Fu, C. J. Li, Y. Xu, L. Y. Li, X. Zhou, D. D. Li, S. X. Chen, F. G. Wang, X. Y. Zhang, L. W. Zheng, Role of lncRNAs as Novel Biomarkers and Therapeutic Targets in Ovarian Cancer. *Crit Rev Eukaryot Gene Expr* **27**, 183-195 (2017)10.1615/CritRevEukaryotGeneExpr.2017019244).
12. L. Dai, J. Niu, Y. Feng, Knockdown of long non-coding RNA LINC00176 suppresses ovarian cancer progression by BCL3-mediated down-regulation of ceruloplasmin. *J Cell Mol Med*, (2019); published online EpubOct 30 (10.1111/jcmm.14701).
13. Z. Li, H. Niu, Q. Qin, S. Yang, Q. Wang, C. Yu, Z. Wei, Z. Jin, X. Wang, A. Yang, X. Chen, lncRNA UCA1 Mediates Resistance to Cisplatin by Regulating the miR-143/FOSL2-Signaling Pathway in Ovarian Cancer. *Mol Ther Nucleic Acids* **17**, 92-101 (2019); published online EpubSep 6 (10.1016/j.omtn.2019.05.007).
14. D. D. Wu, X. Chen, K. X. Sun, L. L. Wang, S. Chen, Y. Zhao, Role of the lncRNA ABHD11-AS1 in the tumorigenesis and progression of epithelial ovarian cancer through targeted regulation of RhoC. *Mol Cancer* **16**, 138 (2017); published online EpubAug 17 (10.1186/s12943-017-0709-5).
15. H. Shan, Y. Yang, X. Zhu, X. Han, P. Zhang, X. Zhang, FAM83H-AS1 is associated with clinical progression and modulates cell proliferation, migration, and invasion in bladder cancer. *J Cell Biochem* **120**, 4687-4693 (2019); published online EpubMar (10.1002/jcb.27758).
16. J. A. Barr, K. E. Hayes, T. Brownmiller, A. D. Harold, R. Jagannathan, P. R. Lockman, S. Khan, I. Martinez, Long non-coding RNA FAM83H-AS1 is regulated by human papillomavirus 16 E6 independently of p53 in cervical cancer cells. *Sci Rep* **9**, 3662 (2019); published online EpubMar 6 (10.1038/s41598-019-

- 40094-8).
17. Q. Dou, Y. Xu, Y. Zhu, Y. Hu, Y. Yan, H. Yan, LncRNA FAM83H-AS1 contributes to the radioresistance, proliferation, and metastasis in ovarian cancer through stabilizing HuR protein. *Eur J Pharmacol* **852**, 134-141 (2019); published online EpubJun 5 (10.1016/j.ejphar.2019.03.002).
 18. B. Xia, S. Yang, T. Liu, G. Lou, miR-211 suppresses epithelial ovarian cancer proliferation and cell-cycle progression by targeting Cyclin D1 and CDK6. *Mol Cancer* **14**, 57 (2015); published online EpubMar 11 (10.1186/s12943-015-0322-4).
 19. T. Murakami, A. S. Singh, T. Kiyuna, S. M. Dry, Y. Li, A. W. James, K. Igarashi, K. Kawaguchi, J. C. DeLong, Y. Zhang, Y. Hiroshima, T. Russell, M. A. Eckardt, J. Yanagawa, N. Federman, R. Matsuyama, T. Chishima, K. Tanaka, M. Bouvet, I. Endo, F. C. Eilber, R. M. Hoffman, Effective molecular targeting of CDK4/6 and IGF-1R in a rare FUS-ERG fusion CDKN2A-deletion doxorubicin-resistant Ewing's sarcoma patient-derived orthotopic xenograft (PDOX) nude-mouse model. *Oncotarget* **7**, 47556-47564 (2016); published online EpubJul 26 (10.18632/oncotarget.9879).
 20. L. Schonemann, U. Kuhn, G. Martin, P. Schafer, A. R. Gruber, W. Keller, M. Zavolan, E. Wahle, Reconstitution of CPSF active in polyadenylation: recognition of the polyadenylation signal by WDR33. *Genes Dev* **28**, 2381-2393 (2014); published online EpubNov 1 (10.1101/gad.250985.114).
 21. M. Santarosa, L. Del Col, A. Viel, N. Bivi, C. D'Ambrosio, A. Scaloni, G. Tell, R. Maestro, BRCA1 modulates the expression of hnRNP2B1 and KHSRP. *Cell Cycle* **9**, 4666-4673 (2010); published online EpubDec 1 (10.4161/cc.9.23.14022).
 22. L. Silwal-Pandit, A. Langerod, A. L. Borresen-Dale, TP53 Mutations in Breast and Ovarian Cancer. *Cold Spring Harb Perspect Med* **7**, (2017); published online EpubJan 3 (10.1101/cshperspect.a026252).
 23. A. Ribas, J. D. Wolchok, Cancer immunotherapy using checkpoint blockade. *Science* **359**, 1350-1355 (2018); published online EpubMar 23 (10.1126/science.aar4060).
 24. An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57-74 (2012); published online EpubSep 6 (10.1038/nature11247).
 25. K. Fang, C. Hu, X. Zhang, Y. Hou, D. Gao, Z. Guo, L. Li, LncRNA ST8SIA6-AS1 promotes proliferation, migration and invasion in breast cancer through the p38 MAPK signalling pathway. *Carcinogenesis*, (2019); published online EpubNov 30 (10.1093/carcin/bgz197).
 26. X. Wang, Y. Wang, L. Li, X. Xue, H. Xie, H. Shi, Y. Hu, A lncRNA coordinates with Ezh2 to inhibit HIF-1alpha transcription and suppress cancer cell adaption to hypoxia. *Oncogene*, (2019); published online EpubNov 29 (10.1038/s41388-019-1123-9).
 27. J. Yang, Q. Qiu, X. Qian, J. Yi, Y. Jiao, M. Yu, X. Li, J. Li, C. Mi, J. Zhang, B. Lu, E. Chen, P. Liu, Y. Lu, Long noncoding RNA LCAT1 functions as a ceRNA to regulate RAC1 function by sponging miR-4715-5p in lung cancer. *Mol Cancer* **18**, 171 (2019); published online EpubNov 29 (10.1186/s12943-019-1107-y).
 28. F. Yang, S. X. Lv, L. Lv, Y. H. Liu, S. Y. Dong, Z. H. Yao, X. X. Dai, X. H. Zhang, O. C. Wang, Identification of lncRNA FAM83H-AS1 as a novel prognostic marker in luminal subtype breast cancer. *Onco Targets Ther* **9**, 7039-7045 (2016)10.2147/OTT.S110055).
 29. S. Lu, W. Dong, P. Zhao, Z. Liu, lncRNA FAM83H-AS1 is associated with the prognosis of colorectal carcinoma and promotes cell proliferation by targeting the Notch signaling pathway. *Oncol Lett* **15**, 1861-1868 (2018); published online EpubFeb (10.3892/ol.2017.7520).
 30. L. Yang, J. Cui, Y. Wang, J. Tan, FAM83H-AS1 is upregulated and predicts poor prognosis in colon cancer. *Biomed Pharmacother* **118**, 109342 (2019); published online EpubOct (10.1016/j.biopha.2019.109342).
 31. Z. Zhao, J. Bai, A. Wu, Y. Wang, J. Zhang, Z. Wang, Y. Li, J. Xu, X. Li, Co-LncRNA: investigating the lncRNA combinatorial effects in GO annotations and KEGG pathways based on human RNA-Seq data. *Database (Oxford)* **2015**, (2015)10.1093/database/bav082).
 32. T. Wang, D. Hao, S. Yang, J. Ma, W. Yang, Y. Zhu, M. Weng, X. An, X. Wang, Y. Li, D. Wu, J. Tang, C. Yang, Y. He, L. Zhang, X. Jin, G. Wang, Z. Li, T. Zheng, H. Meng, Y. Feng, X. Li, miR-211 facilitates platinum chemosensitivity by blocking the DNA damage response (DDR) in ovarian cancer. *Cell Death Dis* **10**, 495 (2019); published online EpubJun 24 (10.1038/s41419-019-1715-x).
 33. F. Tao, X. Tian, S. Ruan, M. Shen, Z. Zhang, miR-211 sponges lncRNA MALAT1 to suppress tumor growth and progression through inhibiting PHF19 in ovarian carcinoma. *FASEB J*, fj201800495RR (2018); published online EpubJun 6 (10.1096/fj.201800495RR).
 34. E. Ghisoni, M. Imbimbo, S. Zimmermann, G. Valabrega, Ovarian Cancer Immunotherapy: Turning up the Heat. *Int J Mol Sci* **20**, (2019); published online EpubJun 15 (10.3390/ijms20122927).
 35. J. H. Li, S. Liu, H. Zhou, L. H. Qu, J. H. Yang, starBase v2.0: decoding miRNA-ceRNA, miRNA-ncRNA and protein-RNA interaction networks from large-scale CLIP-Seq data. *Nucleic acids research* **42**, D92-97 (2014); published online EpubJan (10.1093/nar/gkt1248).
 36. D. Karagkouni, M. D. Paraskevopoulou, S. Tastsoglou, G. Skoufos, A. Karavangeli, V. Pierros, E. Zacharopoulou, A. G. Hatzigeorgiou, DIANA-LncBase v3: indexing experimentally supported miRNA targets on non-coding transcripts. *Nucleic Acids Res*, (2019); published online EpubNov 16 (10.1093/nar/gkz1036).
 37. Y. Hao, W. Wu, H. Li, J. Yuan, J. Luo, Y. Zhao, R. Chen, NPInter v3.0: an upgraded database of noncoding RNA-associated interactions. *Database (Oxford)* **2016**, (2016)10.1093/database/baw057).
 38. B. Li, E. Severson, J. C. Pignon, H. Zhao, T. Li, J. Novak, P. Jiang, H. Shen, J. C. Aster, S. Rodig, S. Signoretti, J. S. Liu, X. S. Liu, Comprehensive analyses of tumor immunity: implications for cancer immunotherapy. *Genome Biol* **17**, 174 (2016); published online EpubAug 22 (10.1186/s13059-016-1028-7).

Figures




Figure 1
 Some dysregulated genes and lncRNAs were identified in OC. (A) Heatmap of significantly differentially expressed lncRNAs and coding genes in OC patients. Red indicates genes that had higher expression level and green indicates genes had lower expression. (B) The pie chart shows percent of up- and down-regulated lncRNAs and coding genes. (C) The pie chart shows percent of up- and down-regulated lncRNAs.




Figure 2
 FAM83H-AS1 expression is upregulated in human pan-cancer and OC. (A) The box plots shows the expression of FAM84H-AS1 in multiple cancer types. Red and blue represent tumor and normal samples. (B) The expression of FAM84H-AS1 in pan-cancer. (C) The expression of FAM84H-AS1 in OC. (D) The expression of FAM84H-AS1 in diverse stages of OC.




Figure 3

Kaplan-Meier survival curves for FAM83H-AS1 associated with overall survival. (A) High expression of FAM83H-AS1 correlates with worse OS in pan-cancer patients and (B) OC patients. X- and Y-axis represent survival time and overall survival. Red and blue lines represent high and low expression of FAM83H-AS1.




Figure 4
 FAM83H-AS1-centric IncRNA-miRNA, IncRNA-protein and IncRNA-mRNA ceRNA networks. (A) FAM83H-AS1-centric IncRNA-miRNA network. (B) The point plot shows expression of miR-211 (X-axis) and FAM83H-AS1 (Y-axis) in OC. (C) FAM83H-AS1-centric IncRNA-protein network. (D) FAM83H-AS1-centric IncRNA-mRNA ceRNA networks.




Figure 5

Expression level of FAM83H-AS1 was associated with methylation and immune. (A) There are two methylation sites locate at FAM83H-AS1. (B) The point plot shows expression of FAM83H-AS1 (X-axis) and methylation level of cg01399317 (Y-axis) in OC. (C) The point plots show the correlations between FAM83H-AS1 and immune cells including B cell, CD8+ T cell, CD4+ cell, Macrophage, Neutrophil and Dendritic cell. X- and Y-axis represent infiltration level and FAM83H-AS1 expression.




Figure 6

Expression levels of FAM83H-AS1 in OC by qRT-PCR. (A) Relative expression levels of FAM83H-AS1 in OC tissue and adjacent noncancerous tissues. (B) The heatmap shows the expression of FAM83H-AS1. Red and green represent high and low expression level.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [TableS1.xlsx](#)