On-the-job vocational training of nonprofessional ethnic health workers of a primary health care team improves their sense of coherence

Cintia Katona (✉ katona.cintia@med.unideb.hu)

University of Debrecen Faculty of Medicine: Debreceni Egyetem Általános Orvostudományi Kar
<https://orcid.org/0000-0002-5977-4313>

Éva Bíró

University of Debrecen Faculty of Medicine: Debreceni Egyetem Általános Orvostudományi Kar

Szilvia Vincze

Faculty of Economics and Business University of Debrecen

Karolina Kósa

University of Debrecen Faculty of Medicine: Debreceni Egyetem Általános Orvostudományi Kar

Research

Keywords: Roma population, primary health care, monitoring, health status

Posted Date: May 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-468414/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Human Resources for Health on February 7th, 2022. See the published version at <https://doi.org/10.1186/s12960-021-00690-0>.

Abstract

Objectives

A Primary Care Model Programme had been implemented in Hungary between 2013–2017 in which group practices were established that employed – among others – nonprofessional health workers (health mediators, similar to community health workers) to facilitate access for the most disadvantaged population groups. The health of mediators, themselves mostly disadvantaged ethnic Roma, was monitored every odd year of the Programme.

Methods

A repeated cross-sectional health interview survey had been implemented inviting all health mediators who were employed at the time of the survey. The same questionnaire was used in all 3 surveys with items from the European Health Interview Survey 2009 and validated versions of other scales.

Results

Positive changes occurred in the health status of mediators during 5 years of follow-up, especially among those who completed vocational training in the Programme. By 2017, significant increase in sense of coherence was observed among those who obtained vocational qualification as opposed to those who did not. The proportion of highly stressed mediators showed a significant increase among those with no vocational training. Improvement was detected in all mediators in health awareness, dysfunctional attitudes, psychological stress and smoking prevalence.

Conclusions

Positive changes among those who obtained on-the-job vocational qualification were observed during follow-up of ethnic Roma health mediators in the programme in which they were equal members of the primary health care team. Employment of health mediators in primary care teams not only contributed to improving access to care for disadvantaged groups but also improved the mental health of mediators themselves.

Introduction

Many international reports and publications detailed the manifold disadvantages experienced by the Roma, the largest ethnic group in Europe [1,2,3] and in Hungary [4,5] that led in 2005 to the launching of the international initiative „Decade of Roma Inclusion” [6] and in 2011 to the development of national Roma integration strategies in EU member states [7] including Hungary [8].

Reduction of economic and social disparities within Hungary, including the Roma, was also one of the main objectives of the Swiss-Hungarian bilateral framework agreement concluded in 2007 [9] that specified four

priority areas for funding, human resource and social development as one of them. A large-scale community oriented Programme in primary health care (titled „Primary Care Development Model Programme”) had been implemented in this priority area between 2012 and 2017 in Hungary that gives the framework for the present paper.

The Model Programme was designed to introduce group practices in primary care to create a new operational model in Hungary (where all general practices were single-handed until then) which would enable general practitioners to employ nonmedical professionals in order to widen primary care services and improve access to and use of these services for all population groups, including disadvantaged, among them Roma, people. Summary [10] and details [11,12,13] of the Model Programme have been published elsewhere.

Briefly, four group practices (called GP clusters, each formed by six general practitioners) were established in two economically disadvantaged regions of the country. GP clusters funded the employment of nonmedical health professionals such as public health specialists, dietitian, physiotherapist, health psychologist, and as a novelty, GP clusters also employed non-professional workers called health mediators specifically for the purpose of facilitating communication and access to primary services for the most disadvantaged population groups.

Health mediators, similarly to community health workers in the healthcare systems of developing countries [14], had worked in various arrangements in primary care in European countries but for a different reason compared to CHWs. The major task of mediators is to facilitate access of Roma minority groups to healthcare [15] rather than substitute for professional workers. The employment of mediators in the Hungarian Model Programme was supported by positive international experiences [16] including the ROMED initiative of the Council of Europe that was established to train mediators for various settings (healthcare, schools, local authorities, etc.) in order to facilitate communication and cooperation between the Roma and public institutions [17].

Employment of 48 health mediators (12 per GP cluster) was planned. Mediators (called „segéd-egészségőrök” in Hungarian) were recruited from the local communities with no requirement for professional or vocational training, and were employed half-time (20 hours per week) by the GP clusters. Preference was given to those applicants who self-identified as Roma or had experience working with Roma population groups in the serviced areas. They were supervised by the so-called public health coordinator who supervised all nonmedical workers of the team [18]. It is worth noting that the Hungarian Model Programme was Europe-wide the first in which health mediators were employed as equal members of the primary care team, being funded from the same budget and covered by the same obligations and rules as all other members of the GP cluster.

In order to prepare health mediators for their jobs, mandatory 30-hour training in health mediation was provided in 2013 (within one month of the start of employment) and in 2015 by the Hungarian partner organization of the ROMED initiative [19]. To increase their capacity and long-term employment prospects, vocational training in assistant nursing and assistant social services was offered in the first year of employment to all those mediators who did not have any health-related qualification, had the educational requirements to start this training, and volunteered to do so. Altogether 20 persons in assistant nursing and 2

persons in assistant social services obtained this qualification by 2014. Mandatory one-day trainings were provided to all mediators approximately every half year throughout the Model Programme.

All trainings, including vocational training, were delivered during work hours, took place in Debrecen, geographically away from the workplaces and co-workers of health mediators, and participation was completely free of charge for them. Transfer, accommodation, meals, and all training tools and equipment were covered from the budget of the Model Programme.

Health mediators, being a group of rural and mostly Roma women with low education whose life substantially changed by employment in the Model Programme, presented unique opportunity to monitor what impact employment, higher income, work responsibilities and being part of a professional team would have on their health. Therefore, in accordance with international recommendations [20], the major aims of the Swiss Contribution, and the Hungarian National Social Integration Strategy, and also contributing to the evaluation of the Programme, a project to monitor the health status of mediators throughout the Model Programme was set up. The aim of this paper is to describe the results of this monitoring project.

Methods

Study design and participants

The questionnaire-based study had a repeated cross-sectional design. The Programme funded the employment of altogether 48 health mediators but this number fluctuated during the Programme (Figure 1). All health mediators employed at the given time of data collection were invited to participate voluntarily and anonymously (since individual identification was not granted by the ethical permission) except in 2013 as shown in Figure 1. Turnover among the mediators during 5 the years of the Programme and the obligation of anonymity excluded panel design.

Data collection

Data collection was carried out first in October 2013 (1st), repeated in August 2015 (2nd) and January 2017 (3rd) during trainings organized by the University of Debrecen. The first data collection occurred during the vocational training for assistant nursing described above, and those 20 mediators were invited who entered this particular training. The second and third data collection was completed during one-day trainings that were mandatory for all health mediators. Vocational training had already been completed at those surveys. The arrangement of these trainings (including the vocational training) and the timing of data collection was such that their willingness to participate be highest: data collection was carried out during work hours; geographically away from their workplaces, co-workers and supervisors; there were no monetary obligations whatsoever to participate; voluntary participation and anonymity was granted, and there was no feedback of the results to their workplaces.

Questionnaire

The same self-report questionnaire was used during all 3 data collections that included demographic data, items on health status and mental health as follows. Scales and items of the questionnaire were taken from the Hungarian version of the European Health Interview Survey 2009 [21] or they were validated Hungarian versions of widely used scales [22].

Demographic data

These included year of birth from which age was calculated for the year of data collection; gender; education (in three categories: primary, secondary, tertiary); marital status (in two categories: lives alone/lives with partner); number of persons in the household; number of children; subjective wealth (five categories collapsed into two categories: bad (very bad and bad) or at least acceptable (acceptable, good, very good). Ethnic identity was self-reported on two items taken from the Hungarian census of 2011 [23]. Primary ethnicity was asked by the question of „Which ethnicity do you feel you belong to?”; secondary ethnicity by the question “Do you belong to another ethnicity?”. The answer listed all recognized minorities in Hungary including Roma from which respondents could choose. A dummy variable was created for Roma ethnic identity (yes/no). One question (yes/no) was related to completion of the vocational training.

Health status and health awareness

Self-rated health was measured on a 5-step Likert scale ranging from „very poor” to „excellent” and collapsed into 3 categories (very poor and poor as „poor”, acceptable as such, and good and excellent as „good”). Health problems limiting daily functioning in the past 6 months were assessed by a 3-step frequency scale (severely limiting, mildly limiting, no limitation) and collapsed into a binary variable for regression (limitation vs no limitation).

Health awareness was approximated by the question of „How much one can do for their health” that was answerable on a 4-step Likert scale from „nothing” to „very much”. Smoking was assessed by a 6-step frequency scale that was collapsed into two categories: nonsmoker and former smoker vs. current smoker.

Mental health

The short (12-item) version of GHQ had been used to detect psychological distress in the past weeks. Its use above the age of 17 years has been well established [24] and had been validated in Hungarian [21]. Answers are given on a Likert scale from 1 to 4 yielding a total score between 12-48. Another, simple scoring assigns a score of 1 to each present symptom, while lack of symptom is 0. A score above 4 identifies persons with high distress, a score of 4 or below average stress. This scoring yields a binary variable of 0 or 1, and had been used in Hungary to identify persons who are highly stressed [25].

Sense of coherence as a pervasive, enduring and dynamic feeling of confidence was defined by Antonovsky who also developed tools to measure the construct [26]. Of those, the short version validated in Hungarian [21]

was used. Items can be answered on a 7-point Likert scale producing a total score between 13-91 where higher scores represent higher sense of coherence.

The 17-item version [27] of the Dysfunctional Attitudes Scale (DAS) [28] was used to measure the presence and intensity of perfectionism and dependence as dysfunctional attitudes. DAS-A-17 is a self-report scale containing 17 items 11 of which is related to perfectionism, 6 items to dependence answerable on a 7-step Likert scale. The total score for perfectionism ranges between 6-42, for dependence between 11-77, higher scores representing more intense dysfunctional attitudes.

Social support was measured by the Hungarian version of the Health and Lifestyle Survey and Health Survey for England [29]. Respondents answered seven questions on a 3-step frequency scale that yielded a total score between 7-21. Support from co-workers was estimated by one item answered on a 3-step scale („My co-workers fully support me – not as much as I would like – not at all”). A binary variable for social support was created for logistic regression modeling by dichotomizing responses to reflect optimal (full) support (score: 21) or less than optimal support (score <21).

Data analysis

Two-way tables of frequencies were used to analyse categorical variables across the study years, and the null hypothesis (no change across the years) was tested by Fisher's exact test.

Normally distributed interval variables were described by the mean and standard deviation, and were compared across the study years by analysis of variance. Non-normally distributed interval variables were described by the median and interquartile range, and compared across the years by Kruskal-Wallis test. Correlation between interval variables was tested by Pearson's; between categorical variables by Spearman's rank correlation test. Logistic regression was carried out defining psychological distress as a binary dependent variable described above. The value of alpha was set at 0.05. Stata 13.1 and MS Excel 2013 were used for statistical analysis.

Results

Demographic data of health mediators

All positions for mediators were filled in October 2013, and 78% of the mediators self-identified as Roma. The number of health mediators varied during the Programme because some left and vacancies were filled. However, only 4 persons joined the Programme more than 18 months after its start. Health mediators who completed vocational training in the Programme (n=22) had been employed 7.2 months longer than those who did not (35.9 ± 8.4 vs 28.7 ± 14 months, $p=0.017$).

The survey in 2013 had been restricted to those who were involved in vocational training in the Programme (n=20, response rate: 90%). The survey was extended to all mediators in 2015 (n= 41, response rate=100%) and 2017 (n=32, response rate=100%). Since the ethical permission of the survey did not allow for personal

identification, it can only be stated that all respondents in 2017 were also employed in 2015. There was no significant difference in any demographic variables of the respondents in the 3 years of the surveys. (Table 1)

Health mediators were dominantly young to middle-age women, approximately two-third of them with primary education, more than one-third in bad subjective wealth who raised on average 2 children, and lived in multiperson households (Table 1). The number of children varied between 0-5, and the number of persons living in the household ranged between at least 2 and no more than 7

Health status

Two-third of the respondents were in good or excellent health in 2013, and this did not significantly change by 2017. Similarly, there was no significant change in the proportion of those with mild or severe functional limitation.

Health awareness increased further, almost reaching significance ($p=0.079$) though its degree had already been quite high even in 2013 (Table 2). Spearman correlation between health awareness, subjective health and functional limitation was carried out separately for 2015 and 2017. No significant correlation was found either between health awareness and subjective health ($p=0.059$ in 2015, $p=0.310$ for 2017) or health awareness and functional limitation ($p=0.698$ for 2015, $p=0.059$ for 2017).

Improving though nonsignificant trends could be observed in dysfunctional attitudes: the median for dependence decreased by 5 points, perfectionism 3.5 points. In terms of social support from family and friends, at least half of all respondents felt fully supported in each study year (Table 2). Support from co-workers was estimated by one item as described in Methods; more than three-fourth of mediators felt fully supported in each study year, and not a single person during 3 rounds of data collection responded that they did not feel supported at all by their co-workers.

Smoking

44% percent of the mediators were daily smokers both in 2013 and 2015 but this decreased considerably, by 37% in 2017. Stratifying the group of nonsmokers to former smokers and never smokers, the prevalence of former smokers rose from 5.6% in 2013 to 12.5%. The proportion of never smokers was 38.9%, 36.6% and 37.5% in the subsequent study years (data not shown).

Mental health

Sense of coherence showed a mildly significant increase from 2013 to 2017 ($d=8.5$ points, $p=0.057$) (Table 2). However, when mediators were stratified by vocational training on-the-job, it was found that among those who did complete this training, a significant rise occurred in sense of coherence from 2015 to 2017 ($d= 9.6$, $p=0.033$) but it remained unchanged among those who did not participate in this training ($d=0.8$, $p=0.862$) (Table 3).

When psychological distress was analyzed by completion of on-the-job vocational training, it was found that the proportion of highly stressed mediators did not significantly change (though it showed declining trend) in

those who did ($d = -6.6\%$, $p = 0.546$) but significantly increased from 2015 to 2017 among those who did not participate in this training ($d = 14.2\%$; $p = 0.016$) (Table 3).

Sense of coherence of the mediators was also compared to sense of coherence of females obtained from population surveys representative of the Hungarian population [³⁰] (since the overwhelming majority of mediators were females). Sense of coherence among health mediators in 2013 was significantly higher not only compared to females with primary education in the general population ($d = 4.7$; $p < 0.001$) but also compared to females in the general population ($d = 1.4$, $p = 0.026$) to whom they were most similar in terms of education. This difference became even greater by 2017 (difference compared to women with primary education: $d = 11.7$, $p < 0.001$; difference compared to females: $d = 7.6$, $p < 0.001$) (Table 4).

The proportion of highly stressed health mediators decreased from 16.7% in 2013 to 12.5% in 2017 (Table 4) but this did not reach significance ($p = 0.683$). The proportion of highly distressed mediators was not significantly different from either compared to all females or females with primary education in the general population in 2013 (Table 4). However, the proportion of highly stressed mediators became significantly higher compared to all females (but was not different from their educational strata) in 2017 (Table 4).

Models for the determinants of psychological stress among health mediators

In order to identify determinants of psychological stress, data from all study years were pooled. Correlation was examined among psychological distress and its potential determinants such as sex, educational level, subjective wealth, Roma ethnicity, living with a partner, sense of coherence, social support and participation in vocational training as described in Methods. Significantly correlated variables were included in a logistic regression model defining psychological stress as a binary outcome (Model 1 in Table 5).

Additional models were tested by removing non-significant independent variables as shown in Table 5. Based on the significance of independent variables, the squared pseudo-regression coefficient (highest) and the Akaike information criterion (lowest), the best model proved to be Model 1. In this model, tertiary compared to primary education significantly increased ($p = 0.034$), sense of coherence significantly ($p = 0.008$) decreased the odds of becoming psychologically distressed as shown in Table 5.

Discussion

Our results showed improving trends in the health status of health mediators employed in the Primary Care Development Model Programme in Hungary between 2013-2017. Significant improvement in mental health was observed among those who obtained vocational qualification in the form of on-the-job training. By the end of the Model Programme, significant increase in sense of coherence was observed among those who completed vocational training as opposed to those who did not. The proportion of highly stressed mediators significantly increased among those who did not complete vocational training as opposed to those who did.

Changes in other health measures also reflected improvements such as decrease in dysfunctional attitudes, and decrease in smoking, in concert with a large body of research that has shown that employed individuals

are healthier compared to those who are not [^{31,32,33}]. A notable finding was the very high level of social support among health mediators, especially the high level of feeling completely supported by their co-workers. Roma people, though far from heterogeneous, tend to form closed groups especially in neighbourhoods where all members of the community know each other. Mediators in the same GP clusters lived in the area and most of them knew each other before joining the Programme. This likely contributed to the high level of support they experienced which was strengthened by working as a team within the healthcare team [³⁴]. Social support played an important role in facilitating access to health care among Roma living in settlements in Slovakia [³⁵]. Support from mediators can overcome lack of trust in healthcare and its providers among Roma people [³⁶].

An advantage of our study is its follow-up design – the first such study on the health status of nonprofessional support workers employed in primary health care – in which all health mediators employed in the Model Programme were invited in 2015 and 2017. Another advantage of the study is their full participation in 2015 and 2017 that was probably due to the good working relations and mutual trust between mediators and the authors who had been responsible for the professional support of the mediators including all trainings.

One of the limitations of the study is the relatively low number of participants that resulted in the underpowering of statistical tests. This likely led to most results being underestimated, making significant results even more meaningful. Another shortcoming of the study is the partial involvement of health mediators in 2013 that was due to organizational and logistical limitations at the time of data collection.

The positive changes described in our paper provide evidence for the multiple benefits of on-the-job training for health mediators. Healthier workers provide better services and make better role models which is greatly needed in Roma communities [^{37,38,39}]. In addition to less favourable morbidity and mortality, Roma struggle with other issues, for example Roma patients suffering from coronary heart disease had higher levels of anxiety and lower sense of coherence compared to non-Roma patients [⁴⁰].

The significantly increased distress of those who had no vocational training at the end of the Programme was probably due to the fact that there was uncertainty about their further employment, and those with vocational training had better job prospects.

Education, employment and health care constitute major areas for the development of human resources in all large-scale international frameworks aimed at disadvantaged populations. The Primary Care Model Programme of Hungary funded by the Swiss Contribution implemented activities in two of these areas (employment and health care) for the target population, and activities in all three areas for its Roma employees, the health mediators who became valued members of the primary care teams. Their contribution to the performance of the GP clusters such as the achievement of a very high participation rate in one of the new services (health status assessment) was detailed elsewhere [⁴¹].

The majority of mediators had been Roma women who tend to be in the most disadvantaged position in terms of education, employment, income and health status when compared to Roma men and either gender groups of majority populations [^{42,43}]. Positive changes in their health status, behaviour and health

awareness, in addition to their meaningful work contribution [40] provides justification for the employment of health mediators in primary health care in those areas where sizable disadvantaged Roma population groups need to be cared for.

Conclusions

On-the-job vocational training improved the mental health of nonprofessional community workers (health mediators) in the long term. The 5-year follow-up of mental health of mediators in the Primary Care Development Model Programme in Hungary revealed an improving trend in terms of all investigated indicators between 2013–2017. A notable result was the significant increase in sense of coherence – a measure of mental resilience – among those mediators who completed vocational training as opposed to those who did not. In terms of high psychological stress, the opposite was found. The proportion of highly stressed mediators significantly increased among those who did not complete vocational training as opposed to those who did.

Declarations

Ethics approval and consent to participate

The Model Programme had been implemented in the framework of the Swiss Contribution Programme SH/8/1 that specified the indicators for evaluation. Indicators monitored in primary health care are specified in the Hungarian Health Care Act of 1997 (1997. évi CLIV. törvény az egészségügyről). The Programme was created on the basis of the Framework Agreement between Switzerland and Hungary (declared by 348/2007. (XII. 20.)) and signed on 20 December 2007. Ethical approval for data collection for research purposes in the Programme was issued by the Scientific and Research Ethics Committee of the Medical Research Council of Hungary (ETT-TUKEB) (16676-3/2016/EKU (0361-16)).

The protocol of the health status assessment was approved by the Scientific and Research Ethics Committee of the Medical Research Council of Hungary (ETT-TUKEB) (52119–2/2015 EKU). All participants signed an informed consent form, and the anonymity of personal information was guaranteed during data analysis.

Consent for publication

Not applicable.

Availability of data and materials

The datasets generated and/or analysed during the current study are not publicly available due to containing personal and special data the availability and use of which is regulated by law (*2011. évi CXII. törvény az információs önrendelkezési jogról és az információszabadságról*). Data collected in the framework of the Programme is stored and managed by the National Public Health Institute of Hungary and available on reasonable request from the director of the Institute.

Competing interests

None of the authors have competing interests.

Funding

The study was carried out in the „Public Health Focused Model Programme for Organising Primary Care Services Backed by a Virtual Care Service Centre”. The Model Programme was implemented in the framework of the Swiss Contribution Programme SH/8/1 that had been supported by a grant from Switzerland through the Swiss Contribution to Hungary.

The authors were supported during the writing of the manuscript by the GINOP-2.3.2-15-2016-00005 project financed by the European Union under the European Social Fund and European Regional Development Fund. The funders had no influence on study design, data collection and analyses, interpretation of results, writing of the manuscript or the decision to submit it for publication.

Authors' contribution

CK contributed to study design, performed the literature review, collected and analysed data, and drafted the manuscript. ÉB contributed to study design, management of vocational training, data analysis, and writing of the manuscript. SZV contributed to data analysis and writing of the manuscript. KK conceptualized the research, finalized the study design, contributed to literature review, analysed data, and finalised the manuscript. The final manuscript was read and approved by all authors.

Acknowledgement

The authors express their gratitude to Róza Ádány and Magor Papp for their help in establishing the vocational training for health mediators. Drs. János Cséki, Gábor Benkő, István Völgyi, and János Szabó were Head GPs (directors) of the four new primary care teams or GP clusters. Éva Gombkötő, Edit Szabó, Ágnes Elek, and Adrienn Csilla Tóth as Public Health Coordinators had been supervisors of the health mediators. Their contribution to the integration of health mediators as equal members of the group practices is gratefully acknowledged.

The authors thank Éva Gutási, Zsuzsa Nagy-Belgyár, Anita Nagy Farkas and Zsanett Kiss for their help in data collection.

References

- 1 UCL Institute of Health Equity. "Health inequalities in the EU. Final report of a consortium led by M. Marmot." UCL Institute of Health Equity.
https://ec.europa.eu/health/sites/health/files/social_determinants/docs/healthinequalitiesineu_2013_en.pdf
Accessed 21 April 2021.
- 2 Hajioff, S., and M. McKee. 2000. "The health of the Roma people: a review of the published literature." *Journal of Epidemiology and Community Health* 54 (11):864-9. doi: 10.1136/jech.54.11.864.

- 3 McKee-Ryan, F., Z. Song, C. R. Wanberg, and A. J. Kinicki. 2005. "Psychological and physical well-being during unemployment: a meta-analytic study." *Journal of Applied Psychology* 90 (1):53-76. doi: 10.1037/0021-9010.90.1.53.
- 4 Kósa, Z., G. Szeles, L. Kardos, K. Kosa, R. Nemeth, S. Orszagh, G. Fesus, M. McKee, R. Adany, and Z. Voko. 2007. "A comparative health survey of the inhabitants of Roma settlements in Hungary." *American Journal of Public Health* 97 (5):853-9. doi: 10.2105/AJPH.2005.072173.
- 5 Masseria, C., P. Mladovsky, and C. Hernandez-Quevedo. 2010. "The socio-economic determinants of the health status of Roma in comparison with non-Roma in Bulgaria, Hungary and Romania." *Eur J Public Health* 20 (5):549-54. doi: 10.1093/eurpub/ckq102.
- 6 RI2020. "Roma Decade and the EU." Roma Integration 2020, Accessed 13 January 2020. <https://www.rcc.int/romaintegration2020/pages/4/roma-decade-and-the-eu>. Accessed 21 April 2021.
- 7 EUR-LEX. "An EU Framework for National Roma Integration Strategies up to 2020. Communication From The Commission To The European Parliament, The Council, The European Economic And Social Committee And The Committee Of The Regions." Eur-Lex. <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=COM:2011:173:FIN>. Accessed 21 April 2021.
- 8 EMMI. "Magyar Nemzeti Társadalmi Felzárkózási Stratégia (Hungarian National Social Integration Strategy) II 2011-2020." Emberi Erőforrások Minisztériuma http://romagov.hu/wp-content/uploads/2020/12/MNTFS2030_1201-tervezet.pdf Accessed 21 April 2021.
- 9 Programme, Swiss-Hungarian Cooperation. "Framework Agreement between the Swiss Federal Council and the Government of the Republic of Hungary." https://www.dfae.admin.ch/dam/erweiterungsbeitrag/en/documents/Laender/Rahmenabkommen-Ungarn_en.pdf Accessed 21 April 2021.
- 10 FDFA. "Pilot general practitioner teams in North-East Hungary. Swiss Agency for Development and Cooperation" Federal Department of Foreign Affairs. https://www.erweiterungsbeitrag.admin.ch/erweiterungsbeitrag/en/home/projects/focus-on-projects.par_projectfilter_page8.html/content/dezaprojects/SDC/en/2012/7F08179/phase1 Accessed 21 April 2021.
- 11 Adany, R., K. Kosa, J. Sandor, M. Papp, and G. Furjes. 2013. "General practitioners' cluster: a model to reorient primary health care to public health services." *Eur J Public Health* 23 (4):529-30. doi: 10.1093/eurpub/ckt095.
- 12 Kósa, K., A. Hegedüs-Erdős, R. Ádány, M. Papp, and J. Sándor. 2018. "Mental health assessment in one cluster of the Primary Care Development Model Programme in Hungary." *Central European Journal of Occupational and Environmental Medicine* 24 (3-4):174-89.
- 13 Rurik, I. 2014. "How can the Swiss Contribution improve Hungarian primary care?" *Primary Care* 14 (9):155-6.

- 14 Christopher, J.B., Le May, A., Lewin, S. et al. Thirty years after Alma-Ata: a systematic review of the impact of community health workers delivering curative interventions against malaria, pneumonia and diarrhoea on child mortality and morbidity in sub-Saharan Africa. *Hum Resour Health* 9, 27 (2011).
<https://doi.org/10.1186/1478-4491-9-27>
- 15 OSI. "Mediating Romani health: Policy and programme opportunities." Open Society Institute.
https://www.opensocietyfoundations.org/sites/default/files/roma_health_mediators.pdf Accessed 21 April 2021.
- 16 OSF. "Roma Health Mediators: successes and challenges " Open Society Foundations,
<https://www.opensocietyfoundations.org/sites/default/files/roma-health-mediators-20111022.pdf>. Accessed 21 April 2021.
- 17 Council of Europe. "About ROMED1." Council of Europe <http://www.coe-romed.org/romed1>. Accessed 21 April 2021.
- 18 Kósa, K., J. Sandor, E. Dobos, M. Papp, G. Furjes, and R. Adany. 2013. "Human resources development for the operation of general practitioners' cluster." *Eur J Public Health* 23 (4):532-3. doi: 10.1093/eurpub/ckt097.
- 19 PHA. "ROMED." Partners Hungary Alapítvány <https://partnershungary.hu/romed/> Accessed 21 April 2021.
- 20 McFadden A, Siebelt L, Gavine A, Atkin K, Bell K, Innes N, Jones H, Jackson C, Haggi H, MacGillivray S. 2018. Gypsy, Roma and Traveller access to and engagement with health services: a systematic review. *Eur J Public Health*. 28(1):74-81. doi: 10.1093/eurpub/ckx226.
- 21 KSH. "Európai Lakossági Egészségfelmérés kérdőív 2009." (European Health Interview Survey Questionnaire 2009) Központi Statisztikai Hivatal (Central Statistical Office), Hungary
http://www.ksh.gov.hu/elef/archiv/2009/pdf/elef_kerdoiv_alap.pdf Accessed 21 April 2021.
- 22 Balajti, I., Z. Vokó, R. Ádány, and K. Kósa. 2007. "A koherencia-érzés mérésére szolgáló rövidített kérdőív és a lelki egészség (GHQ-12) kérdőív magyar nyelvű változatainak validálása." *Mentálhigiéné és Pszichoszomatika* 2007 (2):147-61. doi: 10.1556/Mentál.8.2007.2.4.
- 23 KSH. "Személyi kérdőív. Népszámlálás 2011." (Personal questionnaire of the Census of 2011) Központi Statisztikai Hivatal (Central Statistical Office), Hungary
<https://www.ksh.hu/nepszamlalas/docs/kerdoivek/szemely.pdf> Accessed 21 April 2021.
- 24 Tait, R. J., G. K. Hulse, and S. I. Robertson. 2002. "A review of the validity of the General Health Questionnaire in adolescent populations." *Australian and New Zealand Journal of Psychiatry* 36 (4):550-7. doi: 10.1046/j.1440-1614.2002.01028.x.
- 25 Örkényi, Á. 2003. Mental health. Hungarian National Health Interview Survey 2003. Budapest: National Center of Epidemiology.
- 26 Antonovsky, A. . 1987. Unraveling the mystery of health. San Francisco: Jossey-Bass Publishers.

- 27 de Graaf, L. E., J. Roelofs, and M. J. Huibers. 2009. "Measuring Dysfunctional Attitudes in the General Population: The Dysfunctional Attitude Scale (form A) Revised." *Cognitive Therapy and Research* 33 (4):345-55. doi: 10.1007/s10608-009-9229-y.
- 28 Lovibond, S.H., and P.F. Lovibond. 1995. *Manual for the Depression Anxiety Stress Scales*. 2nd ed. Sydney: Psychology Foundation.
- 29 Albert, F., B. Dávid, and R. Németh. 2005. Társas támogatás, társadalmi kohézió. Országos Lakossági Egészségfelmérés (OLEF) 2003. (Social support, social cohesion. Hungarian National Health Interview Survey 2003). Budapest: Országos Epidemiológiai Központ.
- 30 Kósa K., Vitrai J., Vincze Sz., Bíró É.,. 2021. "Trends of the mental health of the Hungarian adult population between 2010-2019." Manuscript submitted.
- 31 McKee-Ryan, F., Z. Song, C. R. Wanberg, and A. J. Kinicki. 2005. "Psychological and physical well-being during unemployment: a meta-analytic study." *Journal of Applied Psychology* 90 (1):53-76. doi: 10.1037/0021-9010.90.1.53.
- 32 Emerson, E., C. Hatton, S. Baines, and J. Robertson. 2018. "The association between employment status and health among British adults with and without intellectual impairments: cross-sectional analyses of a cohort study." *Bmc Public Health* 18:ARTN 401. doi: 10.1186/s12889-018-5337-5.
- 33 Park, S. J., S. Y. Kim, E. S. Lee, and S. Park. 2020. "Associations among Employment Status, Health Behaviors, and Mental Health in a Representative Sample of South Koreans." *International Journal of Environmental Research and Public Health* 17 (7). doi: 10.3390/ijerph17072456.
- 34 Katona, C., E. Gutasi, M. Papp, O. Varga, and K. Kosa. 2020. "Facilitating equal access to primary care for all: work experiences of health mediators in a primary health care model programme in Hungary." *BMC Family Practice* 21 (1). doi: 10.1186/s12875-020-01281-z.
- 35 Bobakova, D., Z. D. Veselska, I. Babinska, D. Klein, A. M. Geckova, L. Cislakova, and HEPA-META Team. 2015. "Differences between Roma and non-Roma in how social support from family and friends helps to overcome health care accessibility problems." *International Journal for Equity in Health* 14:ARTN 37. doi: 10.1186/s12939-015-0165-z.
- 36 Hanssens, L. G. M., I. Devisch, J. Lobbestael, B. Cottenie, and S. Willems. 2016. "Accessible health care for Roma: a gypsy's tale a qualitative in-depth study of access to health care for Roma in Ghent." *International Journal for Equity in Health* 15. doi: 10.1186/s12939-016-0327-7.
- 37 Sandor, J., Z. Kosa, K. Boruzs, J. Boros, I. Tokaji, M. McKee, and R. Adany. 2017. "The decade of Roma Inclusion: did it make a difference to health and use of health care services?" *International Journal of Public Health* 62 (7):803-15. doi: 10.1007/s00038-017-0954-9.
- 38 Sprocha, B., and B. Bleha. 2021. "Mortality, Health Status and Self-Perception of Health in Slovak Roma Communities." *Social Indicators Research* 153 (3):1065-86. doi: 10.1007/s11205-020-02533-2.

39 Zeman, C. L., D. E. Depken, and D. S. Senchina. 2003. "Roma health issues: A review of the literature and discussion." *Ethnicity & Health* 8 (3):223-49. doi: 10.1080/1355785032000136434.

40 Silarova, B., I. Nagyova, J. P. Van Dijk, J. Rosenberger, and S. A. Reijneveld. 2014. "Anxiety and sense of coherence in Roma and non-Roma coronary heart disease patients (vol 19, pg 500, 2014)." *Ethnicity & Health* 19 (5):I-I.

41 Kósa, K., C. Katona, M. Papp, G. Fürjes, J. Sándor, K. Bíró, and R. Ádány. 2020. "Health mediators as members of multidisciplinary group practice: lessons learned from a primary health care model programme in Hungary." *BMC Family Practice* 21:1-9.

42 FRA. "Second European Union Minorities and Discrimination Survey. Roma women in nine EU Member States." European Union Agency for Fundamental Rights
https://fra.europa.eu/sites/default/files/fra_uploads/fra-2019-eu-minorities-survey-roma-women_en.pdf
Accessed 21 April 2021.

43 Cserti Csapó, T. 2019. "The social situation of Roma women in Hungary: A statistical analysis." *Orvosi Hetilap* 9 (1):67–78. doi: 10.1556/063.9.2019.1.5.

Tables

Table 1. Demographic features of health mediators at the time of data collection

Variables	2013	2015	2017	p
Invited participants (n)	20	41	32	
Response rate (%)	90%	100%	100%	
Gender (female, %)	100%	95%	91%	0.590
Age (mean±SD, years)	33.5 ± 7.6	37.3 ± 7.1	39.1 ± 8.4	0.061
Age range (min-max. years)	18-44	22-57	19-59	-
Education				
primary (%)	61	62	69	0.981
secondary (%)	33	30	25	
tertiary (%)	6	7	6	
Subjective wealth				
bad (%)	44	39	41	0.946
acceptable (%)	50	46	47	
good (%)	6	15	12	
Lives with partner (%)	39	59	63	0.283
Number of children (median; IQR)	2 (1;3)	2 (1;2)	2 (1.5;2.5)	0.982
Number of persons in household, median (IQR)	4 (3;5)	4 (3;4)	3 (3;4)	0.116

Table 2. Health variables of health mediators

Variables	2013	2015	2017	p
Self-rated health (%)				
-good/excellent	67	61	65	0.537
-acceptable	20	31	32	
-bad/very bad	13	8	3	
Health problem causing at least mild functional limitation (%)	17	23	31	0.928
Health awareness (%)				
-can do much or very much for health	87	92	97	0.079
-can do little or nothing for health	13	10	6	
Sense of coherence (mean $\pm$ SD)	61.7 ($\pm$ 14.6)	64.6 ($\pm$ 11.8)	70.2 ($\pm$ 12.8)	0.057
Highly stressed (%)	16.8	9.7	12.5	0.674
Dependence (median; IQR)	20 (11;22)	15 (9;21)	15 (9.5;22.5)	0.513
Perfectionism (median; IQR)	23.5 (17;38)	20 (17;29.5)	20 (15;25)	0.484
Social support, general				
-full (%)	50	60	73	0.333
-partial (%)	50	40	27	
Full social support from co-workers (%)	77.8	82.9	87.5	0.660
Smoking (current vs nonsmokers, %)	56 vs 44	54 vs 46	50 vs 50	0.688

Table 3. Indicators of mental health among health mediators by on-the-job vocational training

	2015	2017	p (2017 vs 2015)
Sense of coherence (mean ± SD)			
No vocational training	67.1 (±11.9)	67.9 (13.5)	0.862
On-the-job vocational training	62.4 (±12.1)	72 (±10.8)	0.033
p (with vs without vocational training)	0.249	0.369	
High psychological distress (%)			
No vocational training	4.6	18.8	0.016
On-the-job vocational training	13.3	6.7	0.546
p (with vs without vocational training)	0.341	0.315	

Table 4. Psychological distress and sense of coherence among health mediators compared to the general population

Variable	Group	2013	2017
Sense of coherence (SoC-13)	4. health mediators (mean±SD)	61.8 (±14.6)	70.2 (±12.3)
	5. females in the population (mean±95%CI)	60.4 (58.8;61.9)	62.6* (61.1;64.2)
	6. females with primary education in the population (mean±95%CI)	57.1 (55.1; 59.1)	58.5* (55.3;61.7)
p	4. vs 5.	0.026	<0.001
	4. vs 6.	<0.001	<0.001
High psychological distress (GHQ-12)	1. health mediators (%)	16.7	12.5
	2. females in the population (%±95%CI)	10.8 (8.2;14.1)	6.8* (4.7;9.6)
	3. females with primary education in the population (%±95%CI)	16.8 (11.8;23.4)	13.7* (8.07;22.51)
p	1. vs 2.	0.397	<0.001
	1. vs 3.	0.986	0.850

* Data are for 2019

Table 5. Comparison of the models of psychological stress⁺

	Model 1	Model 2	Model 3	Model 4	Model 5
secondary education (ref: primary)	2.488	1.990			1.440
tertiary education (ref: primary)	6.706*	5.356*			3.744*
subjective wealth acceptable (ref: poor)	-2.379		-1.734	-2.251	
subjective wealth good (ref: poor)	1.117		1.251	0.532	
sense of coherence	-0.34*	-0.29**	-0.17**	-0.16**	-0.22**
social support (ref: less than optimal)	-0.310	-0.663	-0.884		
Pseudo R-squared	0.594	0.507	0.463	0.410	0.394
AIC	37.22	38.18	40.81	47.07	48.00

+ Regression coefficients of the variables are shown in the table.

* p<0.05, ** p<0.01, *** p<0.001