

# Predicting the Effect of Environmental Humidity on COVID-19 Trend in Medan, Indonesia

Tri Bayu Purnama (✉ [tribayupurnama@uinsu.ac.id](mailto:tribayupurnama@uinsu.ac.id))

---

## Research

**Keywords:** COVID-19, Environment Humidity, Time Series

**Posted Date:** July 23rd, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-46871/v1>

**License:**  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

COVID-19 remained a public health problem and reportedly associated with the environment. However, this environment parameters have not determined yet the correlation to COVID-19. This study used ecological study to investigate time series of environment parameters and COVID-19 data. Spearman correlation test was applied to analyzed the correlation. Humidity has detected as positive correlation with COVID-19 and temperature has shown as negative correlation. Predicting trend of environment parameters and COVID-19 is needed to conduct public health preparedness of COVID-19 strategy.

## Introduction

COVID-19 (coronavirus disease) remains public health problem in Indonesia which is estimated confirmed and mortality cases respectively per May, 9 2020 around 13.645 cases and 959 cases (1). COVID-19 is not only occurring in Jakarta but also in all province in Indonesia (1). Medan as the most prominent capital city of North Sumatera Province and also the largest city in the western part of Indonesia has reported 132 confirmed cases with 12 death cases (2). Several risk factor associated with COVID-19 have been identified and reported in the studies elsewhere including weather indicators (3–5).

Weather effect as a environmental parameters consists of temperature, humidity, rainfall and sunlight exposure are associated with respiratory disease (6–8). Studies found elsewhere reported climatics indicator might exacerbated influenza, severe acute respiratory syndrom and COVID-19 incidence(3–5). Optimum seasonal parameter contributes to viral multiplication on virus (9) and it might be associated with higher COVID-19 incidence and mortality rate (3, 10–12). Elaborating surveillance data with weather data can be used to forecast seasonal trends for public health planning and to detect unusual incidence of the diseases (13).

As the capital of North Sumatera Province and the largest city in the western part of Indonesia, Medan is located close to equator line, which has a typical tropical, humid, rain dan dry season (14). The average annual temperature are 22 °C–29 °C. Besides, Medan owns an area of 265.1 km<sup>2</sup> and a population over 2.26 million (as of 2019) which is equal to 16% of total population in North Sumatera Province.(14). Limited information about potential COVID-19 spread and seasonality trends, this study aims to mapping the trend in both data and also to investigate correlation between COVID-19 and seasonal parameters in Medan, Indonesia.

## Methods

### COVID-19 case definition

COVID-19 case definition referred to Ministry of Health criteria with COVID-19 symptoms and laboratory results (15, 16). In this study, we used confirmed COVID-19 cases and recovery COVID-19 cases in Medan.

The data was collected from Medan COVID-19 task force that is available online access for public (2). This study obtained COVID-19 data on March, 25 to April, 30 2020.

## Environment Parameters

The environmental data were obtained from Indonesian Climatology and Meteorology Bureau Regional 1, Medan, North Sumatera Province, Indonesia (17). The environmental data in this study included minimum temperature, maximum temperature, average temperature, humidity and sunlight exposure data. We collected environmental data for two month (March to April 2020) to observe whether is there environmental variation before and after COVID-19 detected in Medan.

## Data Analysis

We run the environmental and COVID-19 data with trend-line graph to observe the variables per day. We compared environmental data before and after initial COVID-19 spread over in Medan with t-test. Correlation test was performed to analyzed environmental and new and recoveries cases of COVID-19. Due to COVID-19 data have not distributed normally, Spearman Correlation Test (non-parametric test) has been applied to investigating correlation statistically.

## Results

In this study result found that the temperature of Medan has been reported on range 23°C-29°C and tend to fluctuate in 2-month observation period due to rainfall pattern. Humidity and sunlight exposure also have similar pattern with temperature that observed usual climate on tropical countries. There is no significant different on environmental data before and after COVID-19 detected in Medan.

In the table 2 found that there is no correlation between new COVID-19 cases and environmental indicators. Most of environmental indicators shown negative correlation with low Spearman correlation score (r). However, recoveries COVID-19 cases present negative correlation on average temperature, which means higher number of recovery COVID-19 cases, are in line with lower average temperature. Humidity also found statistically significant with recovery cases with intermediate spearman correlation score.

Table 1  
Environmental correlation with new and recovery COVID-19 cases

Variable	New cases		Recovery cases	
	r	Pvalue	r	Pvalue
Min Temperature	-0.164	0.332	-0.310	0.062
Max temperature	-0.029	0.867	-0.191	0.256
Average Temperature	-0.198	0.240	-0.479	0.003
Humidity	0.094	0.579	0.498	0.003
Sunlight exposure	-0.046	0.786	0.058	0.731

## Discussion

In the end of April, 2020, 132 confirmed cases has been reported in Medan that detected in line with exponential trend cases on national/Indonesia data. More than 100 cases detected in a month after one cases detected in March 25, 2020 that equal to more than 80% of cases originally domicile from Medan in total confirmed of COVID-19 cases in North Sumatera Province. There is many study investigating the empirical plausability of environmental parameters in infectiousness of COVID-19 based on disease and weather time series. However, the impacts of environmental parameters has not empirically correlated yet. This is the first study to describe trend of COVID-19 and environment data and to evaluate the impact of environment parameters. Our results found that environment parameters might correlate COVID-19 transmission.

This study found that COVID-19 has spread in Medan with fluctuation environment parameters through study period. It might be complicated to interpreted the data due to some environment parameters shown correlation with COVID-19 data. Previous study reported that respiratory disease mainly on influenza is correlated with fluctuations in temperature rather than low temperature that affects the onset of virus activities (18–21).

Our result found that low temperature is correlated with positive impact of number of recovery COVID-19 cases with intermediate correlation. This is in line with the research of Tosepu et al (2020) that the average temperature ( $^{\circ}\text{C}$ ) is significantly correlated to the COVID-19 recovery cases ( $r = 0.392$ ;  $p < 0.001$ ) (4). Temperature becomes determinant to the transmission acceleration of the COVID-19 outbreak in China (10). At the SARS transmission, temperature, relative humidity and wind velocity become a factor of transmission acceleration (6). Seasonality of respiratory problem likely influenza is associated with temperature and humidity whereas more frequently found in humid-rainy conditions occur outbreaks in tropical and subtropical zones (22). It might be caused by salts and proteins in the droplets (23). The seasonality in infectiousness of COVID-19 would be plausible that COVID-19 shows higher infectivity low

temperature (24). This suggest that positive impacts on the infectivity of COVID-19 might be more suited on constant low temperature in normal tropical condition.

In this study found that positive correlation between humidity and COVID-19 cases. Average humidity determines influenza virus transmission and survival variability around 50% and 90% respectively (3,25). Several study also reported that humidity as most crucial parameter to predicts heat-related mortality among environment/weather parameters. (21,26–28). This study found that humidity has been intermediate correlation with COVID-19 and this finding is in line with study found that humidity as better indicator of health problem on environment parameters (29). Further research is needed to understand these relationships between environment parameter and COVID-19.

The results of this research are also in line with previous research, that humidity significantly distribute against seasons in viruses that attack the respiratory system (11) and has a strong correlation at the level below 1% (8). Thus, the correlation between weathering and COVID-19 transmissions in Wuhan becomes an important factor against the emphasis of cases when weather temperature and humidity rises (12). Humidity may correlate to mortality rates (7). In general, enveloped virus that is wrapped in lipid membranes will survive better at lower humidity, while non-enveloped will be more stable when in an environment with high humidity (30).

Rous sarcoma virus (RSV) and infectious bovine rhinotracheitis virus (IBRV) is a more stable enveloped virus in the environment with high humidity, while pigeon pox virus, also enveloped after researched not resistant to humidity. In areas that have temperate climates, the seasonal influenza virus outbreak turns out to have a strong correlation to low humidity (31). The viral transmission will increase in line with low ambient humidity, but the impact on humans (hosts) on influenza virus infections related to humidity remains unclear. In research Eriko et al (2019) showed that rat experiment on housing Mx1 congenic rats with relatively low humidity led to a significant impact so that rats would be more susceptible to the impact of breathing on an influenza A virus infection (32). In tropical climates, influenza virus outbreak and respiratory syncytial virus (RSV) are more common in the rainy season (high humidity rate), but the mechanism of the seasonal pattern is still unclear (33).

There are two physical properties of the environment that have a very close association, namely humidity and temperature. Humidity is a measure of the humidity content in the gas, relative to the gas capacity to withstand humidity that varies with temperature. Thus, humidity becomes the most relevant measurement parameter e.g. water will evaporate from respiratory droplet which indicates humidity become very complicated when describing the water content in air (18). The virus viability of the humidity effect review will be in mediation by the concentration of salts in droplet, wherein at high humidity, the virus will be relatively stable as a result of the defence of physiological concentrations. At intermediate humidity, there is an increase in salt concentrations as a result of evaporation process resulting in virus inactivation. While at low humidity (50%), the concentration of salts will be low and increase the stability of the virions as a result of salt that crystallizes out of the solution (34).

Influenza disease can be prevented at high relative humidity where the mechanism will increase the expression levels of anti-IAV ISGs (including BST2, Mx1, IFITM2, IFITM3, Viperin, ZAP, and ISG15). Inability of the host animal exposed to low humidity clears the virus may be caused by the declining MCC as a result of low relative humidity exposure and the elimination of the virus particles in the airway to the trachea (32). These results are in line with human observation, that low humidity and temperature result in a long decline of periciliary coating resulting in a decrease in cell movement of the cilia, and the slowing of MCC with the final result will be increased pathogenic spread (35). At low humidity, ISG will be induced in different cell types in the lungs, resulting in higher viral loads and caspase-1/11-dependent pathology. Together, in low humidity conditions, the disease will be seen from a combination of various factors (32).

Aerosol transmission can be influenced by humidity with a mechanism of respiratory droplets proportions that become aerosolised and the ability of the virus to survive in the aerosol (33). Respiratory droplets that enter the environment with low humidity produced by the respiratory tract with high humidity will disappear very quickly because it is affected by evaporation, while at high humidity, Respiratory droplets will evaporate slower and aerosols will remain on the environment (36,37). An environment with a high level of humidity implicates the increased durability of influenza viruses and high RSV on the droplet surface supported by the slowing of evaporation (however, high humidity will reduce the virus's survival if droplets becomes dry). In addition, the spread of viruses coming from the surface of the hand and droplets will increase in high humidity conditions (38).

When respiratory droplet is released from the respiratory tract to the ambient air (aerosol transition from higher humidity to lower humidity), evaporation will occur due to the vapor pressure gradient between the surface and the ambient air. As a result of evaporation, water will disappear along with evaporation while the solute (protein and salt) will remain, because the molar fraction of water in an aerosol is comparable to the humidity pressure on the surface (Raoult law). The cessation of evaporation occurs when the point at which the water activity in aerosols is equal to the ambient humidity (vapor pressure on the surface of the aerosol is reduced by ambient air) (39).

The ambient humidity has implications for physical and immune to viral infections that attack the respiratory system (32). At the correlation of epidemiology, the death of influenza virus infection in temperate regions with decreased humidity as a mechanism of impaired MCC and ISG induction (40). Influenza viruses can develop better in regions with tropical and subtropical climates, however, humidity does not affect the host against the durability of influenza virus infections in all situations (41,42). In the experiment of Yang & Marr (2012), lipid membranes in enveloped viruses will protect the capsid from threats of damage due to environmental humidity conditions. Viruses that attack the respiratory system will experience slow rehydration when inhalations. The humidity effect on non-enveloped viruses that attack the respiratory system such as rhinovirus needs to be taken into account in this effect. Inactivation of the non-enveloped virus is performed by fundamental restructuring during rehydration in the Impinger collection medium and slowing the level of rehydration (31).

In human populations, relatively low humidity (RH) is not associated with influenza epidemics. Therefore, further epidemiological studies are necessary to determine the correlation of the season during influenza outbreak with humidity, as well as indoor and outdoor temperatures. This is important because between temperature and humidity in the room is associated with outdoor conditions as well as an unimpossible influenza outbreak (18). Although there is a link between the two parameters, namely humidity and temperature with epidemic, humidity becomes the most dominant factor at the rate of viral spread (influenza) in terms of temperate climates (40). In conclusion, this study showed a plausibility of environment parameters and COVID-19 diseases.

This study has a number of limitations due to data availability to mapping risk of environment parameters to COVID-19. We were unable to obtain the environment data on all places in Medan due to testing limitation on the environment parameters. This result might be not generalized to COVID-19 patient due to study design limitation. Despite of these limitations, this study is primarily to investigate environment parameters in western part of Indonesia dominantly in Medan, Indonesia.

## Declarations

## Ethical Approval

Ethical approval is not applicable for this study due to this study used secondary data that open access for public and also does not contain personal information.

## Conflict of Interest

The authors declare no competing interest in this paper


## References

1. Indonesian COVID-19 Task Force. Situation Report of COVID-19 in Indonesia. 2020.
2. Medan City Communication and Information Office. General Data Related to COVID-19 in Medan City. 2020.
3. Liu J, Zhou J, Yao J, Zhang X, Li L, Xu X, et al. Impact of meteorological factors on the COVID-19 transmission: A multi-city study in China. *Sci Total Environ*. 2020.
4. Tosepu R, Gunawan J, Savitri D, Ode L, Imran A, Lestari H, et al. Correlation between weather and Covid-19 pandemic in Jakarta, Indonesia. *Sci Total Environ*. 2020;725:1–4.
5. Bashir MF, Ma B, Bilal, Komal B, Bashir MA, Tan D, et al. Correlation between climate indicators and COVID-19 pandemic in New York, USA. *Sci Total Environ*. 2020.
6. Yuan J, Yun H, Lan W, Wang W, Sullivan SG, Jia S, et al. A climatologic investigation of the SARS-CoV outbreak in Beijing, China. *Am J Infect Control*. 2006;34(4):234–6.

7. Ma Y, Zhao Y, Liu J, He X, Wang B, Fu S, et al. Effects of temperature variation and humidity on the death of COVID-19 in Wuhan, China. *Sci Total Environ*. 2020.
8. Wang J, Tang K, Feng K, Lv W. High temperature and high humidity reduce the transmission of COVID-19. *SSRN*. 2020;2020:1–26.
9. Scholtissek C, Rott R. Effect of temperature on the multiplication of an Influenza virus. *J Gen Virol*. 1969.
10. Shi P, Dong Y, Yan H, Li X, Zhao C, Liu W, et al. The impact of temperature and absolute humidity on the coronavirus disease 2019 (COVID-19) outbreak - evidence from China. *medRxiv*. 2020;(77).
11. Sajadi MM, Habibzadeh P, Vintzileos A, Miralles-wilhelm F, Amoroso A. Temperature, humidity, and latitude analysis to predict potential spread and seasonality for COVID-19. *SSRN*. 2020;2020:6–7.
12. Guo X-J, Zhang H, Zeng Y-P. Transmissibility of COVID-19 and its Association with Temperature and Humidity. *Res Sq Prepr*. 2020;1–10.
13. World Health Organization. Global Surveillance for human infection with coronavirus disease (COVID-19). *Interim Guid*. 2020;(February):27–9.
14. BPS Statistics Indonesia. Kota Medan dalam Angka (Medan Profile in Number). 2018.
15. MOH. HK 01.07/MENKES/182/2020 tentang Jejaring Laboratorium Pemeriksaan Coronavirus Disease 19 (Covid-19). 2020.
16. MOH. Pedoman Kesiapsiagaan Menghadapi Coronavirus Disease (COVID-19). Direktorat Jenderal Pencegah dan Pengendali Penyakit. 2020;1–88.
17. Indonesian Meteorology and Climatology. Data Online-Database Center, BMKG. 2020.
18. Lowen AC, Steel J. Roles of Humidity and Temperature in Shaping Influenza Seasonality. *J Virol*. 2014.
19. Jaakkola K, Saukkoriipi A, Jokelainen J, Juvonen R, Kauppila J, Vainio O, et al. Decline in temperature and humidity increases the occurrence of influenza in cold climate. *Environ Heal A Glob Access Sci Source*. 2014.
20. Shaw Stewart PD. Seasonality and selective trends in viral acute respiratory tract infections. *Med Hypotheses*. 2016.
21. Peci A, Winter AL, Li Y, Gnaneshan S, Liu J, Mubareka S, et al. Effects of absolute humidity, relative humidity, temperature, and wind speed on influenza activity in Toronto, Ontario, Canada. *Appl Environ Microbiol*. 2019.
22. Chew FT, Doraisingham S, Ling AE, Kumarasinghe G, Lee BW. Seasonal trends of viral respiratory tract infections in the tropics. *Epidemiol Infect*. 1998.
23. Sooryanarain H, Elankumaran S. Environmental Role in Influenza Virus Outbreaks. *Annu Rev Anim Biosci*. 2015.
24. Ujiie M, Tsuzuki S, Ohmagari N. Effect of temperature on the infectivity of COVID-19. *Int J Infect Dis* [Internet]. 2020 Apr 30;S1201-9712(20)30284-8. Available from: <https://pubmed.ncbi.nlm.nih.gov/32360939>.

25. Shaman J, Kohn M. Absolute humidity modulates influenza survival, transmission, and seasonality. *Proc Natl Acad Sci U S A*. 2009.
26. Quilodran CS, Currat M, Montoya-Burgos JI. Climatic factors influence COVID-19 outbreak as revealed by worldwide mortality. *medRxiv*. 2020.
27. Gupta S, Raghuwanshi GS, Chanda A. Effect of weather on COVID-19 spread in the US: A prediction model for India in 2020. *Sci Total Environ*. 2020.
28. Ahmadi M, Sharifi A, Dorosti S, Jafarzadeh Ghouschi S, Ghanbari N. Investigation of effective climatology parameters on COVID-19 outbreak in Iran. *Sci Total Environ*. 2020.
29. Zhang K, Li Y, Schwartz JD, O'Neill MS. What weather variables are important in predicting heat-related mortality? A new application of statistical learning methods. *Environ Res*. 2014.
30. Sobsey M, Meschke J. Virus survival in the environment with special attention to survival in sewage droplets and other environmental media of fecal or respiratory origin. Geneva; 2003.
31. Yang W, Marr LC. Mechanisms by Which Ambient Humidity May Affect Viruses in Aerosols AND VIABILITY. *Appl Environ Microbiol*. 2012;78(19):6781–8.
32. Kudo E, Song E, Yockey LJ, Rakib T, Wong PW, Homer RJ. Low ambient humidity impairs barrier function and innate resistance against influenza infection. *PNAS*. 2019;116(22):10905–10.
33. Paynter S. Humidity and respiratory virus transmission in tropical and temperate settings. *Epidemiol Infect*. 2020;143(May):1110–8.
34. Yang W, Elankumaran S, Marr L. Relationship between humidity and influenza viability in droplets and implications for influenza's seasonality. *PLoS One*. 2012;7(e46789).
35. Oozawa H, Kimura H, Noda T, Hamada K. Effect of prehydration on nasal mucociliary clearance in low relative humidity. *Auris Nasus Larynx*. 2012;39(1):48–52.
36. Xie X, Li Y, Chwang ATY, Ho PL, Seto WH. How far droplets can move in indoor environments—revisiting the Wells evaporation–falling curve. *Indoor Air*. 2007;17(3):211–25.
37. Yang W, Marr LC. Dynamics of airborne influenza A viruses indoors and dependence on humidity. *PLoS One*. 2011;6(1):e21481.
38. Lopez GU, Gerba CP, Tamimi AH, Kitajima M, Maxwell SL, Rose JB. Transfer efficiency of bacteria and viruses from porous and nonporous fomites to fingers under different relative humidity conditions. *Appl Environ Microbiol*. 2013;79(18):5728–34.
39. Posada J, Redrow J, Celik I. A mathematical model for predicting the viability of airborne viruses. *J Virol Methods*. 2010;164(1):88 – 95.
40. Shaman J, Pitzer V, Viboud C, Grenfell B. M L. Absolute humidity and the seasonal onset of influenza in the continental United States. *PLoS Biol*. 2010;8:e1000316.
41. Shek L, Lee B. Epidemiology and seasonality of respiratory tract virus infections in the tropics. *Paediatr Respir Rev*. 2003;4(1):105–11.
42. Moura F, Perdigão A, Siqueira M. Seasonality of influenza in the tropics: A distinct pattern in northeastern Brazil. *Am J Trop Med Hyg*. 2009;81(1):180–3.

# Figures



**Figure 1**

Environmental indicators and COVID-19 cases of Medan, North Sumatera from March to April. Figure 1a-c indicates minimum, maximum and average temperature of Medan respectively. Figure 1d presents average humidity and 1e also presents sunlight exposure per day. Figure 1f highlights number of new COVID-19 cases (black bar), cumulative (red line) and recovery case (blue line)