DNMT1 inhibits hepatocellular carcinoma cells by altering the epigenetic modification of Marvel1 gene

Song Yan Zhang

Harbin Medical University Cancer Hospital

Qian Luo

Harbin Medical University Cancer Hospital

Peng Wang

Harbin Medical University Cancer Hospital

Hao Ran Zhao

Harbin Finance University

Hai Tao Xu

Harbin Medical University Cancer Hospital

Liang Zhao

Harbin Medical University Cancer Hospital

Zhao Wei Qu

Harbin Medical University Cancer Hospital

Jian Min Sun

Harbin Medical University Cancer Hospital

Yu Bao Zhang (✉ aite_cherry@163.com)

Harbin Medical University Cancer Hospital

Biao Zheng

Shenzhen University Gender Hospital/Shenzhen University Clinical Medical Academy

Research article

Keywords: Hepatocellular Carcinoma, Epigenetic Modification, DNMT1 Gene, Marvel1 Gene, Methylation, Proliferation

Posted Date: September 5th, 2019

DOI: <https://doi.org/10.21203/rs.2.14010/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Objective To study the effect of DNA methyltransferase (DNMT1) on Marveld1 in hepatocellular Carcinoma (HCC), and explore its role in the molecular mechanism of HCC occurrence and development.

Methods The expression level of Dnmt1 and Marveld1 genes was detected by RT-PCR in the intraoperative pathological specimens. The human hepatoma Bel7402 and SMMC7721 cell lines down-regulated by DNMT1 gene expression were constructed by SiRNA transfection. The methylation level of Marveld1 promoter region was detected after RT-PCR and West-blot verification. The expression of Marveld1 gene, cell proliferation, invasion and change of cell cycle were detected by RT-PCR and West-blot. The expressions of P53, CyclinD1, P21 and P16 protein were detected by West-blot method.

Results The expression of DNMT1 gene in human hepatocarcinoma tissues was higher than that in adjacent tissues. The methylation level of the Marveld1 gene promoter region in human hepatoma cell line down-regulated by DNMT1 gene expression was decreased, the expression level of Marveld1 gene was increased, and the proliferation and invasion ability of cells were weakened. The cell cycle was showed as G1-S phase arrest; P53 and P16 protein expressions were up-regulated, and CyclinD1 and P21 protein expressions were down-regulated.

Conclusions Dnmt1 gene is highly expressed in human hepatocarcinoma tissues. Down-regulation of DNMT1 gene expression can decrease the methylation level of Marveld1 promoter region, up-regulate Marveld1 gene and human P53 and P16 proteins expressions, and down-regulate CyclinD1 and P21 proteins expressions, inhibit the proliferation and invasion of human hepatoma cells and can cause cell cycle G1-S arrest.

Introduction

Hepatocellular carcinoma (HCC) has become one of the most common malignant tumors worldwide. About 780,000 new cases of liver cancer occur each year, which accounting for 7.4% of all malignant tumors, and the mortality rate is about 9%. Statistics from IARC (International Agency for Research on Cancer) showed that HCC has become the third leading cause of cancer death in the world ^[1]. HCC has a high degree of malignancy, which is easy for recurrence after surgery, and has poor prognosis. Therefore, exploring the causes of HCC and the molecular mechanism of its pathogenesis is of great value for its early diagnosis, treatment and prognosis. Existing studies have found that the occurrence of HCC is closely related to viral hepatitis, alcoholism, diet and cirrhosis, but the molecular mechanism of the disease still involves two aspects of genetic mechanism and epigenetic modification mechanism ^[2]. More and more evidence also indicates that epigenetic modification plays a very important role in the occurrence and development of HCC. In the study of epigenetic modification, DNA methylation research is one of the important fields of current research ^[3]. Studies have shown that

hypermethylation inactivation of multiple tumor suppressor genes (Ecadherin, RASSF1A, GSTP, SOCS1, SFRP1, PTEN) in HCC is one of the important molecular mechanisms for the occurrence and development of HCC [4-6].

DNA methyltransferase (DNMT) plays a key role in DNA methylation. The methylation of genomic DNA mainly occurs on the cytosine of CpG island, and methyl group is added to 5' end of the pyrimidine ring by the action of DNMT. The DNMTs family has approximately five members: DNMT1, DNMT2, DNMT3a, DNMT3b, and DNMT3L, of which DNMT1 is the most important DNA methylase and plays an important role in the maintenance of methylation status [7-8]. DNMT inhibitors have been used in a few fields such as myeloid leukemia, so the new treatment with DNMT as the entry point is a valuable direction in the field of HCC treatment.

Marveld1 (Marvel domain containing-1, MD1) gene is located in the long arm 2 region 4 band 2 sub-band of human chromosome 10 (10q24.2). The mRNA is 3238 bp in length and has two exons. The coding region is located in the first exons, and the encoded protein product consists of 173 amino acid residues with a molecular weight of approximately 18.9 kDa. Li et al [9] first reported the biological function of Marveld1 gene, and confirmed that it plays an important role in the regulation of malignant biological behavior of tumors in malignant tumors such as liver cancer [10] and lung cancer [11].

In order to study the effect of DNMT1 on Marveld1 in HCC, and explore its role in the molecular mechanism of HCC occurrence and development, and then explore a new possible method for the diagnosis and treatment of HCC, we use the method of SiRNA transfection to down-regulate the expression of DNMT1 gene in human HCC cell lines, and its effect on HCC cells was studied.

Material And Methods

1.1 Search the TCGA database

The expression difference of DNMT1 and Marveld1 genes in liver cancer and adjacent tissues and the relationship between the expression difference and survival, the difference of methylation level of Marveld promoter region, and the related genes of DNMT1 and Marveld1 genes were examined respectively.

1.2 Cell culture

Human HCC Bel7402 and SMMC7721 cell lines were used, and DMEM medium (Gibco, Thermo Fisher Scientific Suzhou Instrument Co., Ltd.) supplemented with 10% serum (Gibco, Grand Island, NY, USA) was used at 37 °C - 5% CO2 incubator for cell culture.

1.3 Construction of transfected cell lines with down-regulated expression of Dnmt1 gene

SiRNA (Suzhou Gemma Gene) sequence:

Control group (Dnmt1-NC)

Positive-sense strand (5'-3'): UUCUCCGAACGUGUCACGUTT

Antisense strand (5'-3'): ACGUGACACGUUCGGAGAATT

Dnmt1 gene down-regulation group (Dnmt1-2505)

Positive-sense strand (5'-3'): GGGACUGUGUCUCUGUUAUTT

Antisense strand (5'-3'): AUAACAGAGACACAGUCCCTT.

Human HCC Bel7402 and SMMC7721 cells were evenly plated into 6-well plates and cultured for 24 hours. The cell density was about 50%. Premixed transfection solution was added to each well (containing 75 pmol of SiRNA, 7.5 uL of lipofectamin2000, and 200 uL of Opti-MEM medium). Further experiments were carried out after incubation for 48 hours at 37 °C - 5% CO₂.

1.4 RT-PCR

From January 2017 to December 2017, 22 pairs of HCC and adjacent cancer tissue samples were collected from the Affiliated Tumor Hospital of Harbin Medical University. All samples were preserved in liquid nitrogen immediately after collection. All the samples were confirmed to be HCC by postoperative pathology. The written informed consent of the patients for the study was obtained before surgery. The intraoperative samples used in this study were approved by the medical ethics of the Human Experimental Committee of Affiliated Cancer Hospital of Harbin Medical University.

1 µg of total RNA was extracted from tissue samples and adherent cultured cell samples with trizol, and then reverse-transcribed into cDNA by reverse transcription kit, and detected by RT-PCR using a ViiATM7 real-time PCR instrument (Applied Biosystems).

Dnmt1 gene primer sequence:

Upstream 5'-AGGCGGCTCAAAGATTTGGAA-3'

Downstream 5'-GCAGAAATTCGTGCAAGAGATTTC-3'

Marveld1 gene primer sequence:

Upstream 5'- ACTGAGAAGTCCCGCTGTTACAG -3'

Downstream 5'- GGGATGCTGGGAATCTTAAGG-3'

GAPDH Primer Sequence:

Upstream 5'-AGGCTGTGGGCAAGGTCATC-3'

Downstream 5'-TCAGGTCCACCACTGACACG-3'

1.5 Marveld1 gene promoter region methylation level sequencing detection

Marveld1 gene promoter region methylation level sequencing detection

The 1µg genomic DNA was extracted, and the bisulfite conversion and purification and collection were carried out according to the kit instructions (QIAGEN, No. 59824). The sequence of the Marveld1 gene promoter region was found in NCBI and designed the primers:

Upstream 5'-GCCAGGGCAAGCAGGAGGTGG-3'

Downstream 5'-CTGCTGAGACTAAGACGTGG-3'

After PCR amplification, the amplification products were sequenced to determine the methylation level of the Marveld1 gene promoter region, and the sequence and CpG island sites were as follows:

1CpG 2CpG 3CpG 4CpG 5CpG

CGAGTGGAAACGAGTGGCGGTGCCTCTGGTGGGTGATGGACTCTTACTCCGACATGGGTAAGTCGGACAG

6CpG 7CpG 8CpG

CTTTGGACAGGACGAGGCCGGGCCCTGGCAGGAGACTGCGGATGTCCAGGTGACAGTGGATGCCAGTTCA

 9CpG10CpG 11CpG 12CpG

AAACCAGCACTGCTCTCCTCCCG CGCCCAGGAAGAGCCCAGAGGCCCGGGGTGGGCGCTATGGCAGG

 13CpG 14CpG

AAAGAGGGAAAAAGGAGGAGGCAGGGCGGCAGGGGCG

1.6 Western-blot

Cell count: Taken the same number of cells and added with RIPA lysate (containing 5% Doc, 1% PMSF, 1% pepstain A, 1% ribonuclease) at a concentration of 1×10^6 cells/100 μ l, and after lysed at 4°C for 30 minutes, centrifuged at 12,000 rpm and then taken the supernatant protein solution. SDS-acrylamide gel electrophoresis was carried out and imprinted on a 0.2 μ m PVDF membrane. Washed with 0.1% PBST after sealed for 1 hour with 5% skim milk, and then immersed in 0.1% PBST diluted primary antibody overnight; washed with 0.1% PBST, and then immersed in secondary antibody for 1 hour, then washed with 0.1% PBST, and added with developer for development by ECL Western Blot Analysis System (Amersham Biosciences, Chalfont St. Giles, Buckinghamshire, UK), and grayscale analysis was conducted.

1.7 Measurement of cell proliferation ability by MTT assay

The cells were plated into the 96-well plates with 1500 cells/well, cultured and adherent with 200μl of DMEM medium, and then added 100μl of medium containing 10% MTT to each well at 0h, 24h, 48h, 72h and 96h, respectively, after cultured for 4 hours, 100μl of DMSO was added to each well, and the OD value was read at a wavelength of 450 nm by a microplate after shaking. The 0hOD value was used as the baseline, and the ratio of

the OD value at each time point to the OD value at 0h was the cell proliferation rate, and the proliferation curve of each group was plotted.

1.8 Changes of cell invasion ability after down-regulation of Dnmt1 gene expression was measured by Transwell method

The upper chamber of Transwell plate (3428, costar) was evenly spread with glue, and then spread with cells with 3×10^5 cells/well in each well, and 1.5 ml of DMEM medium containing 0.5% serum was added, and 2.6 ml of medium containing 10% serum was added to the lower chamber; After cultured at 37°C -5% for 24 hours, photographs were taken under a crystal violet staining microscope. Randomly selected 4 field of view from each group for counts.

1.9 Determination of Cell Cycle Changes after Down-regulation of Dnmt1 Gene Expression with flow cytometry

The cells were cultured in an adherent manner without EDTA trypsinization, and then the cells were collected after centrifugation, and after washed with PBS, the cells were fixed with 75% ethanol at 4 °C for 24 hours; then washed with PBS, PI staining was performed for 30 min, and the cell cycle was measured by flow cytometry after washing with PBS.

Results

2.1 TCGA database search results

The search results of TCGA database showed that the expression of DNMT1 gene in HCC was higher than that in adjacent tissues, the expression of Marvel1 gene was lower than that in adjacent tissues, the methylation level of promoter region was higher than that of adjacent tissues. The changes of expression of DNMT1 and Marvel1 gene was related with poor prognosis (Figure 1). There is currently no data on the association between DNMT1 gene and Marvel1 gene expression.

2.2 Differential expression of Dnmt1 and Marvel1 in HCC tissues and adjacent tissues in clinical samples

Paired analysis of RT-PCR results of 22 clinical samples showed that the expression level of Dnmt1 gene in HCC tissues was higher than that in adjacent tissues, and the expression of Marvel1 gene in HCC tissues was lower than that in adjacent tissues (Figure 2).

2.3 Construction of human HCC cell line with down-regulated Dnmt1 gene expression

After transfection with Si-RNA, the RNA and protein expression of Dnmt1 gene in the experimental group was significantly decreased (Figure 3, Figure 4).

2.4 Detection of Marvel1 gene expression level

2.4.1 Detection of methylation level of Marveld1 gene promoter region

The 14 CpG islands in the promoter region of Marveld1 gene were sequenced. The number of methylation in human HCC cells with down-regulated Dnmt1 expression was lower than that in the control group, and the methylation level of the Marveld1 gene promoter region was decreased (Figure 5, Figure 6).

2.4.2 Marveld1 gene expression detection

The RNA and protein expression of Marveld1 gene in human HCC cells with down-regulated Dnmt1 expression was up-regulated (Figure 7, Figure 8).

2.5 Changes of proliferative capacity of human HCC after down-regulation of Dnmt1 gene expression by MTT assay

After down-regulation of Dnmt1 gene expression, the proliferative capacity of human HCC was decreased significantly (Figure 9, Figure 10).

2.6 Detection of the change of invasive ability of HCC cells after down-regulation of Dnmt1 gene expression by Transwell assay

After down-regulation of Dnmt1 gene expression, the invasive ability of HCC cells was weakened (Figure 11, Figure 12).

2.7 Detection of change of cell cycle after down-regulation of Dnmt1 gene expression by flow cytometry

After down-regulation of Dnmt1 gene expression, HCC cells were increase in G1 phase, and decreased in S phase, suggesting that down-regulation of Dnmt1 gene expression may cause cell cycle G1-S arrest in HCC cells (Figure 13).

2.8 West-Blot detection of P53, CyclinD1, P21 and P16 protein expression

After transfection with Si-RNA, the protein expression of the Dnmt1 gene in the experimental group was significantly reduced. The protein expression of Marvel1 in human HCC cells down-regulated by Dnmt1 gene was up-regulated, the expression of P53, P21 and P16 proteins was up-regulated, and the expression of CyclinD1 protein was down-regulated (Figure 14).

Discussion

DNA methyltransferase (DNMT), as the key factor in the regulation of DNA methylation in tumor tissues, can affect the biological behavior of tumor cells by regulating the degree of methylation of gene promoters with different functional [12]. The DNMTs family has approximately five members: DNMT1, DNMT2, DNMT3a, DNMT3b, and DNMT3L, of which DNMT1 is the most important DNA methyltransferase and plays an important

role in methylation status maintenance; deletion of DNMT1 may lead to decreasing of DNA methylation, affect the transcription of genes, and then regulate cell growth and differentiation.

Proteins containing the MARVEL domain are evolutionarily conserved and widely expressed in normal adult tissues. They involve a variety of biological processes, including cell cycle progression, chemotactic activity, tight junctions, and clathrin-mediated endocytosis [9, 13-16]. As a member of this family, the research on the molecular function of Marveld-1 is limited so far. Studies have shown that the expression of Marveld-1 gene in liver cancer and lung cancer tissues is lower than that in adjacent tissues. Overexpression of Marveld-1 gene can inhibit the proliferation of tumor cells, up-regulate P53 and P16 genes, and can down-regulate the expression of CyclinD1 gene, Marveld-1 can affect the G1-S phase checkpoint-associated proteins of the cell cycle mentioned above, and then lead to G1-S phase arrest of Marveld-1 overexpressing cells, thus exert the function of tumor suppressor gene [10-11].

The data in the TCGA database showed that the DNMT1 gene was highly expressed in HCC tissues and low in Marveld1 gene, which was consistent with the clinical sample paired PCR results in this study. Marveld1 gene was hypermethylated in HCC tissues, and the survival of the samples with low expression of Dnmt1 gene was longer than that of the samples with high expression of Dnmt1 gene, suggesting that down-regulation of DNMT1 gene expression can up-regulate the expression of Marveld1 gene, and the results of inhibiting the malignant biological manifestations of HCC may have potential clinical application value. There is no data on the relationship between Marveld1 gene and DNMT1 gene in TCGA database. This study showed that the down-regulation of Dnmt1 gene can up-regulate the expression of Marveld1 gene by down-regulating the methylation level of Marveld1 gene promoter region, and there is certain correlation between the expressions of two genes. This study found that in human HCC tissues, the expression of Dnmt1 gene in HCC tissues was higher than that in adjacent tissues. Down-regulation of the expression of Dnmt1 gene in HCC cells could down-regulate the methylation level of the promoter region of Marveld1 gene. The expression of Marveld1 gene can be up-regulated by the epigenetic modification, and can inhibit the proliferation and invasion of HCC cells that consistent with the existed study on the overexpression of Marveld1 gene [10]. And then up-regulated P53, P21 and P16 genes, down-regulated the expression of CyclinD1 gene. Affecting the G1-S phase checkpoint-associated proteins in the cell cycle, leading to G1-S phase arrest in cells. The Dnmt1 gene can change pathway, cause the epigenetic modification of the Marveld1 gene through a promoter methylation level, thus resulting in up-regulation of its expression and enhancing the function of the tumor suppressor gene.

Dnmt inhibitors have been used in clinical applications, and the main therapeutic areas are bladder cancer and myeloid leukemia. The results of this study suggest that Dnmt inhibitors may have some application prospects in the field of HCC at the in vitro cytological level, and related cells, animals and clinical trials may be a valuable research direction.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All data generated or analysed during this study are included in this published article.

Competing interests

The authors declare that they have no competing interests.

Funding

This study is funded by Harbin Medical University Cancer Hospital HaiYan Science Foundation Key Program, No. JJZD2014-06.

Authors' contributions

Song-Yan Zhang and Qian Luo: conceived and designed the experiments, performed the experiments.

Yu-Bao Zhang and Biao Zheng: performed the experiments, analyzed the data, wrote the manuscript.

Peng Wang, Hao-Ran Zhao, Hai-Tao Xu, Liang Zhao, Zhao-Wei Qu and Jian Min Sun: analyzed the data and proofed the manuscript.

All authors have proofed the article.

Acknowledgements

Not applicable.

References

- [1] Jemal A, Bray F, Center MM. Global cancer statistics. *A Cancer J Clinicians* 2011; 61(2):69-90.
- [2] Eden A, Gaudet F, [Waghmare A](#), Jaenisch R. Chromosomal instability and tumors promoted by DNA hypomethylation. *Science* 2003; 300(5618):455.
- [3] Li XD, Wang YW, Hu J. Progress on epigenetic research of liver cancer. *Journal of Modern Oncology* 2016; 24(09):1501-1504.
- [4] Calvisi DF, [Ladu S](#), [Gorden A](#), [Farina M](#), [Lee JS](#), [Conner EA](#), [Schroeder I](#), [Factor VM](#), [Thorgeirsson SS](#). Mechanistic and prognostic significance of aberrant methylation in the molecular pathogenesis of human hepatocellular carcinoma. *J Clin Invest* 2007; 117(9):2713-22.
- [5] Tischoff I, Tannapfe A. DNA methylation in hepatocellular carcinoma. *World J Gastroenterol* 2008; 14(11):1741-8.
- [6] Lou C, Yang B, Gao YT, [Wang YJ](#), [Nie FH](#), [Yuan Q](#), [Zhang CL](#), [Du Z](#). Aberrant methylation of multiple genes and its clinical implication in hepatocellular carcinoma. *Chinese Journal of Oncology* 2008; 30(11):831-6.
- [7] Bird A. DNA methylation patterns and epigenetic memory. *Genes Dev* 2002; 16(1):6-21.
- [8] Liang G, [Chan MF](#), [Tomigahara Y](#), [Tsai YC](#), [Gonzales FA](#), [Li E](#), [Laird PW](#), [Jones PA](#). Cooperativity between DNA methyltransferases in the maintenance methylation of repetitive elements. *Mol Cell Biol* 2002; 22(2):480-91.
- [9] [S Wang](#), [Y Li](#), [F Han](#), et al. Identification and characterization of MARVELD1, a novel nuclear protein that is down-regulated in multiple cancers and silenced by DNA methylation. *Cancer Letters* 2009; 282 (1):77-86
- [10] Yu L, Zhang Y, Hu J, et al. MARVELD1 inhibited cell proliferation and enhance chemosensitivity via increasing expression of p53 and p16 in hepatocellular carcinoma. *Cancer Sci* 2012; 103(4):716-22.
- [11] [M Shi](#), [S Wang](#), [Y Yao](#), et al. Biological and clinical significance of epigenetic silencing of MARVELD1 gene in lung cancer. *Scientific Reports* 2013; 4:7545-7545.
- [12] Schutte M, Hruban RH, Geradts J, [Maynard R](#), [Hilgers W](#), [Rabindran SK](#), [Moskaluk CA](#), [Hahn SA](#), [Schwarte-Waldhoff I](#), [Schmiegel W](#), [Baylin SB](#), [Kern SE](#), [Herman JG](#). Abrogation of the Rb/p16 tumor-suppressive pathway in virtually all pancreatic carcinomas. *Cancer Res* 1997; 57(15):3126-30.
- [13] Mimori K, Shiraishi T, Mashino K et al. MAL gene expression in esophageal cancer suppresses motility, invasion and tumorigenicity and enhances apoptosis through the Fas pathway. *Oncogene* 2003; 22: 3463-71.

[14] Shi Y, Ding X, He ZH et al. Critical role of TRPC6 channels in G2 phasetransition and the development of human esophageal cancer. *Gut* 2009; 58:1443-50.

[15] Han W, Lou Y, Tang V et al; Molecular cloning and characterization of chemokine-like factor 1 (CKLF1), a novel human cytokine with uniquestructure and potential chemotactic activity. *Biochem J* 2001; 357: 127-3520.

[16] Martí'n-Belmonte F, Martí'nez-Mena'rguez JA, Aranda JF et al. MAL regulates clathrin-mediated endocytosis at the apical surface of Madin-Darby canine kidney cells. *J Cell Biol* 2003; 163: 155-64.

Figures

Figure 1

Figure 2
Paired RT-PCR detection of Dnmt1 and MarvelD1 clinical samples

Figure 3
Dnmt1 gene RT-PCR detection

Figure 4

Dnmt1 gene Western-blot detection

Figure 5

Marveld1 gene promoter region methylation sequencing

Figure 6

Methylation level of the Marvel1 gene promoter region A: Methylation CpGs rate: NC group $57.1 \pm 2.38\%$, SI group $43.6 \pm 6.61\%$, $p < 0.05$ B: Methylation CpGs rate: NC group $40.7 \pm 2.14\%$, SI group $32.9 \pm 4.42\%$, $p < 0.05$

Figure 7

RT-PCR detection of Marveld1 gene

Figure 8

Western-Blot detection of Marveld1 gene

Figure 9

Cell proliferation curve

Figure 10

The cell proliferation difference at 72 hours A: 72 h proliferation multiple: NC group 4.9670 ± 0.1597 ; SI group 2.9676 ± 0.3614 ; $P=0.0000$ B: 72 h proliferation multiple: NC group 3.2734 ± 0.1394 ; SI group 2.5646 ± 0.0789 ; $P=0.0000$

Figure 11

Invasive ability of cells

Figure 12

Counting of cell invasion A: Invading cell number: NC group 155.25 ± 15.97 ; SI group 65.00 ± 4.55 ; $P=0.0000$ B: Invading cell number: NC group 102.70 ± 12.90 ; SI group 41.00 ± 6.27 ; $P=0.0000$

Figure 13

Detection of cell cycle E: G1 stage: $P=0.0010$; S stage: $P=0.0480$; G2 stage: $P=0.0600$ F: G1 stage: $P=0.0000$; S stage: $P=0.0230$; G2 stage: $P=0.8440$

Figure 14

Western-blot detection of P53, CyclinD1, P21 and P16