

# Effect of luseogliflozin on bone microarchitecture in elderly patients with type 2 diabetes: Study protocol for a randomized controlled pilot trial using second-generation high-resolution peripheral quantitative computed tomography (HR-pQCT)

**Ai Haraguchi**

Nagasaki Daigaku <https://orcid.org/0000-0002-0671-2530>

**Riyoko Shigeno**

Nagasaki Daigaku

**Ichiro Horie** (✉ [horie@nagasaki-u.ac.jp](mailto:horie@nagasaki-u.ac.jp))

<https://orcid.org/0000-0003-3430-5796>

**Shimpei Morimoto**

Nagasaki Daigaku

**Ayako Ito**

Nagasaki Daigaku

**Ko Chiba**

Nagasaki Daigaku

**Yurika Kawazoe**

Nagasaki Daigaku

**Shigeki Tashiro**

Nagasaki Daigaku

**Junya Miyamoto**

Nagasaki Daigaku

**Shuntaro Sato**

Nagasaki Daigaku

**Hiroshi Yamamoto**

Nagasaki Daigaku

**Makoto Osaki**

Nagasaki Daigaku

**Atsushi Kawakami**

Nagasaki Daigaku

**Norio Abiru**

Nagasaki Daigaku

## Study protocol

**Keywords:** type 2 diabetes, luseogliflozin, SGLT2 inhibitor, HR-pQCT, fracture, bone, randomized controlled trial, pilot

**Posted Date:** March 23rd, 2020

**DOI:** <https://doi.org/10.21203/rs.2.14017/v2>

**License:**   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Trials on May 5th, 2020. See the published version at <https://doi.org/10.1186/s13063-020-04276-4>.

# Abstract

Background Elderly patients with type 2 diabetes mellitus (T2DM) have an increased risk of bone fracture independent of their bone mineral density (BMD), which is explained mainly by the deteriorated bone quality in T2DM compared to non-diabetic adults. Sodium-glucose co-transporter (SGLT) 2 inhibitors have been studied in several trials in T2DM, and the Canagliflozin Cardiovascular Assessment Study showed an increased fracture risk related to treatment with the SGLT2 inhibitor canagliflozin, although no evidence of increased fracture risk with treatment with other SGLT2 inhibitors has been reported. The mechanism of the difference in the fracture risk between the SGLT2 inhibitors is unknown, but the differences among the SGLT2 inhibitors in the selectivity of SGLT2 against SGLT1 may affect bone metabolism, since among the SGLT2 inhibitors the selectivity of canagliflozin is lowest. We will investigate whether the SGLT2 inhibitor luseogliflozin, which has higher SGLT2 selectivity, affects bone metabolism by using high-resolution peripheral quantitative computed tomography (HR-pQCT) which provides direct in vivo morphometric information about the bone microarchitecture. Methods This is a single-center, randomized, open-label, active-controlled, parallel pilot trial. Eligible participants are elderly (age  $\geq 60$  years) individuals with T2DM with HbA1c levels at 7.0%–8.9%. A total of 24 participants will be allocated to either the luseogliflozin group (taking luseogliflozin) or control group (taking metformin) in a 1:1 ratio to compare the groups' changes in bone microarchitecture of radius and tibia which are analyzed by HR-pQCT before and 48 weeks after the administration of each medication. The laboratory data associated with glycemic control and bone metabolism will be collected every 12 weeks during the study. Recruitment began in June 2019. Discussion The reason we use metformin as an active control is to avoid yielding differences in glycemic control between the luseogliflozin and control groups. Besides, metformin is considered to have neutral effect on bone. This trial will reveal the effect of luseogliflozin on bone metabolism in elderly patients with T2DM.

## Background

Type 2 diabetes mellitus (T2DM) is well known as a cause of secondary osteoporosis. Individuals with T2DM have shown a 40%–70% increase in the risk of hip fracture compared to age- and gender-matched healthy adults.(1, 2) However, the bone mineral density (BMD) of the spine and femoral bone in individuals with T2DM has been reported to be comparable to or somewhat higher than that in non-diabetic adults.<sup>1 (3)</sup> The cause of the elevated fracture risk in the individuals with T2DM is thus suspected to be deteriorated bone quality due to (1) the accumulation of advanced glycation end products in the bone collagen fibers, and (2) decreased bone turnover as a result of exposure to hyperglycemia(3, 4).

High-resolution peripheral quantitative computed tomography (HR-pQCT), first developed in 2004, is a three-dimensional imaging technology. HR-pQCT is effective for determining bone quality based on an evaluation of microarchitecture, volumetric BMD, and geometry separately for cortical and trabecular compartments of the distal radius and tibia. This technique can often detect a fracture risk that is independent of the BMD determined by dual-energy X-ray absorptiometry (DXA).(5-7) Several studies evaluated the bone microstructure in individuals with T2DM using first-generation HR-pQCT, and the

results regarding the deteriorated bone quality were controversial.(8-11) A second-generation HR-pQCT scanner was recently developed with an improved ability to analyze bone microarchitecture; this second-generation HR-pQCT scanner has not yet been used in a study of individuals with T2DM.

Sodium glucose co-transporter 2 (SGLT2) inhibitors have garnered attention as a breakthrough drug for patients with T2DM, as it was revealed that SGLT2 inhibitors have the potential to improve cardiovascular outcomes in T2DM.(12-14) However, there are several concerns about the SGLT2 inhibitors' effects on bone metabolism regarding the following:(15) (i) stimulation of parathyroid hormone and fibroblast growth factor 23 by the increased serum phosphate levels via tubular reabsorption in phosphate, (ii) weight loss, and (iii) volumetric depletion and fall risk (e.g., orthostatic hypotension and postural dizziness).

The U.S. Food and Drug Administration recently strengthened the warning for the SGLT2 inhibitor canagliflozin related to an increased risk of bone fracture, based on the outcomes in the Canagliflozin Cardiovascular Assessment Study (CANVAS).(16) In that study, the rate of fracture adverse events in the canagliflozin group was significantly increased compared to that of the placebo group (4.0% vs. 2.6%, respectively).(17) It was also reported that canagliflozin treatment was associated with a small but significant reduction in the BMD of the total hip and with increases in bone biomarkers of formation (osteocalcin) and resorption (collagen type 1  $\beta$ -carboxyteleopeptide) compared to placebo controls,(18) which might be explained due at least in part to canagliflozin-treated patients weight loss. In contrast, no evidence of an increased fracture risk has been reported in other cardiovascular outcome trials evaluating the effects of SGLT2 inhibitors in the patients with T2DM.(12, 19, 20)

There is no clear explanation for the difference in fracture risk between the CANVAS trial and other trials. It is possible that among the SGLT2 inhibitors, the selectivity of SGLT2 against SGLT1 may affect bone metabolism and the fracture risk, since the selectivity of canagliflozin is the lowest among the SGLT2 inhibitors under clinical development.(21) However, it is not yet known whether the CANVAS trial's finding of an increased rate of fracture events related to canagliflozin treatment would be reproduced by treatment with other SGLT2 inhibitors, especially those with higher SGLT2 selectivity.

## Methods/design

### Objective

This study will aim to determine whether treatment with the SGLT2 inhibitor luseogliflozin (which has high SGLT2 selectivity) affects bone metabolism by assessing the changes in bone microstructure using second-generation HR-pQCT in elderly patients with T2DM.

### Study design

This is a pilot trial to assess the effect of SGLT2 inhibitors on bone metabolism. We designed a single-center, randomized, open-label, parallel-group, active-controlled trial for patients with T2DM whose

conditions have never been complicated with osteoporosis. We will randomly assigned 24 patients into 1:1 ratio to the luseogliflozin group (receiving additional treatment with luseogliflozin) or the control group (receiving additional treatment with metformin). The study follow-up duration is 48 weeks. The study will be conducted at Nagasaki University Hospital, Japan. The study design is summarized in Figure 1.

## **Outcomes**

The primary outcome measures are changes in the predicted bone strength as determined by second-generation HR-pQCT, evaluated using the parameters of (1) bone stiffness, and (2) the estimated failure load of the radius and tibia of the nondominant body side. For each parameter, any change will be determined based on the difference between the measurement results at baseline (week 0) and week 48.

The study's secondary outcome measures are as follows. (1) The changes in the structures of cortical bone, trabecular bone, and the bone morphology measured by HR-pQCT as described below in the 'Image measurements' section from baseline (week 0) to week 48. (2) The changes in the laboratory data values, including the levels of glycated hemoglobin (HbA1c) from week 0 to weeks 12, 24, 36 and 48. (3) The changes in the areal BMD of the lumbar spine (L1–L4), femoral neck, and distal radius estimated by DXA from week 0 to week 48. (4) The incidence of vertebral fracture or femoral fracture from week 0 to week 48. (5) Changes in the bone metabolic markers from week 0 to week 48.

In addition to the primary and secondary outcomes, we will evaluate adverse effects from both luseogliflozin and metformin (control agent). We will also evaluate the recruitment rate and consent rate.

## **Sample size estimation**

This is a pilot trial to assess the changes in bone microstructure affected by luseogliflozin treatment compared with metformin treatment, evaluated by HR-pQCT. There is no prior similar study comparing the bone strength before and after intervention that can be used to estimate the precise optimal sample size. Julious et al. reported that 12 participants per group are needed for a pilot study (22). The justification for this sample size is based on the rationale concerning feasibility and precision regarding the mean and variance of the primary outcome measures.

## **Patients and public involvement statement**

There is no patient or public involved in this trial.

## **Participants and recruitment**

A total of 24 participants aged  $\geq 60$  years will be recruited into the study. The enrollment started in June 2019. All participants have been diagnosed with T2DM, and their cases have never been complicated with osteoporosis. Patients fulfilling the inclusion criteria described below will be invited to a screening for their eligibility. Principal investigator and co-investigators recruit the participants among their

outpatients, and obtain a written informed consent from the participants. There are no additional consent provisions for collection and use of participant data and biological specimens in ancillary studies. The recruitment rate and the consent rate will be evaluated at the end of the study.

### **Inclusion criteria**

As shown in Figure1, participants must fulfill the following criteria to be eligible for inclusion at their first visit to Nagasaki University Hospital (visit 1): (1) an individual aged  $\geq 60$  years; (2) diagnosed with T2DM; (3) an outpatient; (4) treated with diet therapy alone or with oral antidiabetic agents including metformin ( $\leq 1000$  mg per day), and/or an alpha-glucosidase inhibitor ( $\alpha$ -GI) and/or a dipeptidyl peptidase-4 (DPP-4) inhibitor, whose treatment has not been changed within 6 months before enrollment; and (6) providing written informed consent.

### **Run-in criteria**

Participants will be included when they also fulfill the following criteria at week 0 (visit2): (1) the patient's HbA1c (NGSP) level is  $\geq 7.0\%$  and  $< 9.0\%$ ; (2) the patient's T-score in both lumbar vertebrae and the femoral neck determined by DXA is  $> -2.5$  standard deviations (SD).

### **Exclusion criteria**

Eligible participants will be excluded if they meet any of the following criteria: (1) a previous history of treatment with an SGLT2 inhibitor; (2) complicated with a bone metabolic disorder (i.e., osteomalacia, thyroid dysfunction, hyperparathyroidism, or hypoparathyroidism which requires any medical treatments) (3) administered any treatment for osteoporosis within 12 months before enrollment (e.g., a bisphosphonate, vitamin D preparation, vitamin K preparation, calcium preparation, selective estrogen receptor modulator, antireceptor activator of nuclear factor kappa-B ligand antibody [denosumab], anti-sclerostin antibody [romosozumab], or recombinant human PTH [teriparatide]); (4) treated with estrogen or testosterone hormone replacement therapy; (5) complicated with diabetic retinopathy and/or diabetic neuropathy that required therapeutic intervention; (6) severe renal dysfunction defined as an estimated glomerular filtration rate (eGFR)  $< 30$  mL/min/1.73 m<sup>2</sup>; (7) severe anemia (Hb  $< 10$  g/dL); (8) complicated with malignancy; (9) hypersensitivity or allergy to metformin and/or luseogliflozin; (10) alcohol consumption  $> 20$  g/day; (11) a smoking habit, including smoking even one cigarette per day, within 12 months before enrollment; (12) participating in other clinical research within 6 months before enrollment; (13) complicated with chronic liver disease with  $\geq 6$  points as the Child-Pugh score; (14) body mass index (BMI)  $< 18.5$  kg/m<sup>2</sup>; (15) otherwise deemed inappropriate by the study's principal investigator.

### **Withdrawal criteria**

Participants will be withdrawn from this trial after randomization if they meet any of the following criteria: (1) exhibiting an increase in the HbA1c level that is  $\geq 0.5\%$  of the baseline (week 0) for any two successive visits; (2) an HbA1c level  $> 10.0\%$  at any visit; (3) the occurrence of any adverse event that is

≥grade 3 of the Common Terminology Criteria for Adverse Events (CTCAE) ver. 4.0; (4) showing any fracture(s) with the exception of an occult fracture; (5) has requested withdrawal from the study; (6) has requested a change or the discontinuation of the diabetes treatment; (7) it is determined that it is inappropriate for the patient to continue the study by the investigators because of the progression of an underlying disease or a complication, or the occurrence of unidentified disease; (8) it is judged by the investigators that it is inappropriate for the patient to continue participating in the study for any other reason.

## **Ethical considerations**

This protocol was approved by Clinical Research Review Board of Nagasaki University which is certified by Japan's Ministry of Health, Labour and Welfare (approval no. CRB18-0006). The study is registered on the University Hospital Medical Information Network Clinical Trials Registry (UMIN-CTR) as UMIN-CTR no. 000036202. The study is conducted in accordance with the 7th revision of the Declaration of Helsinki 2013 and the Clinical Trials Act enforced in April 2018 in Japan.

## **Setting and study timeline**

All participants are recruited at Nagasaki University Hospital. Each participant's information as listed below is collected at the prescreening for enrollment (visit 1); age, sex, height, comorbid complications, diabetes duration, history of diabetes treatment, family history of diabetes and osteoporosis within the second degree of kinship, and histories of smoking, drinking, and fracture. Body weight, blood pressure, and pulse rate in a sedentary position are measured at every visit of the study (at visits 1–6, and [if withdrawing from the study] at withdrawal). As shown in Table 1, the participants will undergo examinations by spine X-ray, DXA, and HR-pQCT at week 0 (visit 2) and week 48 (visit 6). When participants need to be withdrawn from the study for any reason, they will undergo HR-pQCT within 4 weeks after discontinuing the study drug. Blood specimens are obtained at every visit except the prescreening visit (visit 1).

## **Image measurements**

Frontal and lateral X-ray examinations are performed to detect compression fractures of thoracic and lumbar vertebrae using a semiquantitative technique.<sup>(23)</sup> DXA is performed for the assessment of the areal BMD at the lumbar spine (L1, L2, L3, L4, L1–4, L-min), total hip (right total, left total), femoral neck (rt. neck, lt. neck), and 1/3 distal of the radius (rt. radius, lt. radius) using DXA system (iDXA, GE Medical Systems, WI, USA). The second-generation HR-pQCT (Xtreme CT II, Scanco Medical, Brüttisellen, Switzerland) scans the radius and tibia of the nondominant body side to evaluate the bone microstructure.

We measure the following parameters by HR-pQCT: (1) bone density, i.e., total volumetric bone mineral density, trabecular volumetric bone mineral density, and cortical volumetric bone mineral density, (2) trabecular bone parameters, i.e., bone volume fraction, trabecular thickness, trabecular number, and

trabecular separation, (3) cortical bone parameters, i.e., cortical thickness, cortical porosity, and cortical pore diameter, (4) bone morphology, i.e., total area, trabecular area, cortical area, and periosteal perimeter, and (5) predicted bone strength, i.e., stiffness and failure load.

## Laboratory measurements

At visits 3–5, we measure the patients' complete blood count and plasma levels of sodium, potassium, chlorine, calcium, phosphate, magnesium, total protein, albumin, total cholesterol, high-density lipoprotein cholesterol, low-density lipoprotein cholesterol, triglyceride, blood urea nitrogen, creatinine, total bilirubin, aspartate aminotransferase, alanine aminotransferase, lactate dehydrogenase, alkaline phosphatase,  $\gamma$ -glutamyl transpeptidase, creatine kinase, uric acid, glucose, and HbA1c. At visit 2 and visit 6 (and at withdrawal if discontinuing participation), in addition to the above laboratory data, we measure the serum (plasma) levels of glycoalbumin, C-peptide, glucagon, glutamic acid decarboxylase antibody, intact parathyroid hormone (PTH), whole PTH, free thyroxine, thyroid-stimulating hormone, adrenocorticotrophic hormone, luteinizing hormone, follicle-stimulating hormone, total estradiol, total testosterone, sex hormone-binding globulin, total cortisol, growth hormone, insulin-like growth factor-1, homocysteine, glucagon, tartrate-resistant acid phosphatase 5b, total type I procollagen N-terminal propeptide, bone-specific alkaline phosphatase, undercarboxylated osteocalcin, osteocalcin, 25-hydroxyvitamin D, 1,25-dihydroxyvitamin D, sclerostin, dickkopf-1, and pentosidine. There are no plans for collection, laboratory evaluation, and storage of biological specimens for genetic or molecular analysis in the current trial and for future use in ancillary studies.

## Randomization and allocation concealment

A randomized, open-label, parallel pilot trial design has been set up. On the day on which the first administration of the study drug (visit 2) is planned after the eligibility assessment, each participant is sequentially allocated to either the luseogliflozin group or the control group in a 1:1 ratio, using a blocked randomization stratified by sex (female or male). The allocation is conducted by the participant-registration center, and the result is sent to the investigators via facsimile so that the block size. As the study design is open label with only outcome assessors being blinded, unblinding will not occur during the study.

## Interventions

(1) Luseogliflozin group: Participants allocated to the luseogliflozin group will take oral luseogliflozin (Taisho Pharma Co., Tokyo) at 2.5 mg/day from the beginning of the trial (visit 2) in addition to their previous treatment. The dose of luseogliflozin will be increased to 5 mg from visit 3 unless a participant achieves an HbA1c level <7% or a decrease in the HbA1c of  $\geq 0.5\%$  compared to the previous visit. The maximum dose of luseogliflozin permitted is up to 5 mg/day. If a participant cannot achieve the HbA1c levels described above with 5-mg luseogliflozin, an additional treatment with metformin (Sumitomo Dainippon Pharma Co., Osaka, Japan) will be prescribed at each visit by 500-mg increments up to 1500

mg. The daily dose of metformin will be increased by 250 mg at each visit if the participant already takes  $\geq 1500$  mg/day. The maximum dose of metformin permitted is 2250 mg/day.

(2) Control group (Metformin group): Participants allocated to the control group will be newly administered or added oral metformin 500 mg/day from the beginning of the trial (visit 2) in addition to the previous treatment. The additional metformin will be administered at each visit by 500-mg increments unless the participant achieves an HbA1c level  $<7\%$  or a decrease in HbA1c  $\geq 0.5\%$  compared to the previous visit. The daily dose of metformin will be increased by a 250-mg increment at each visit if the participant already takes  $\geq 1500$  mg/day. The maximum dose of metformin permitted is 2250 mg/day.

### **Adverse events (AEs)**

All adverse events (AEs) that occur during the trial will be recorded on a case report form and reviewed as part of the central data monitoring. Investigators explain AEs to the participants and offer appropriate care. If serious AEs such as death, a life-threatening condition, hospitalization, sequelae, disability, and congenital illness occur during the trial, the principal investigator will report the AE to the Minister of Health, Labour and Welfare (Japan) and the certified review board within 15 days.

### **Data collection, data management and monitoring**

All the data will be collected using the research electronic data capture (REDCap) system by the authorized investigators. Once the data are checked, they will be fixed by the trial steering committee (TSC). The TSC provides overall supervision for the trial on behalf of the sponsor and funder and to ensure that the trial is conducted to the rigorous standards set out in the guideline of the Clinical Trials Act. The meeting of the TSC will be held every month over the course of the trial to oversee conduct and progress. Clinical Porter Inc. (Tokyo), a site management organization (SMO) specializes in clinical trial management which is independent from the sponsor and competing interest, will monitor the overall conduct of the trial, safeguarding the interests of the trial participants and assessing the safety and efficacy of the intervention. The authorized person of the SMO visits before and every year after starting the trial to review protocol compliance, conduct source data verification, assess laboratory procedure, and ensure that the study is being conducted in accordance with protocol requirement. Because the risk of intervention in the trial is considered to be extremely low, a regular audit is not conducted in the trial unless serious concerns about the outcomes or/and serious AEs arise. Once a serious event occurred, the authorized persons of the Clinical Research Support Center of Ryukyu University Hospital, which is independent from the investigators and the sponsor; will make a site visit to audit the trial conduct. The sponsor is responsible for all aspects of local organization including identifying potential recruits and taking consent.

### **Statistical analyses**

As stated above in the Objectives section, the primary outcome of this study is the extent of luseogliflozin's effects on bone quality change. Hence, the primary data analysis will be conducted using

the set of participants with sufficient exposure to the allocated drugs. For this reason, we will use a per protocol set (PPS), the definition of which includes the adherence ratio, as the target population in the primary data analysis. The same analyses will be conducted on a full analysis set (FAS) modified FAS (mFAS) to provide supplemental information about the influence of selecting PPS subjects on the results.

As the primary and the secondary data analyses, the adjusted mean and the 95% confidence intervals (95%CI) of the bone deterioration effect of the treatment group will be estimated as regression coefficient and its standard error for the respective primary and secondary outcome measures, under a linear regression model with the following two covariates: the baseline measurement of the respective measures and the stratification factor in the randomization (i.e., sex). This study is set up as a pilot study, but we will test the null hypotheses that "the mean of the bone quality deterioration effect of the treatment group is 0" by determining whether the 95%CIs contain the null hypotheses.

The respective analysis sets are defined as follows. The intention to treat (ITT) population is defined as all participants registered for this trial; the safety analysis set (SAS) population is defined as the participants in the ITT population with at least one administration of luseogliflozin/metformin. The full analysis set (FAS) population is defined as the participants in the SAS for whom data about the predicted bone strength after the administration of luseogliflozin/metformin at one or more scheduled visits are available. The PPS is defined as the participants in the FAS with the primary outcome measure. Other criteria for the PPS will be determined before the database lock of this study. We defined the modified FAS (mFAS), the definition of which is a relaxed version of PPS by allowing missingness of the data of predicted bone strength at week 48 only if the data of the items at withdrawal from this study are obtained.

As a safety analysis, tabular summaries of AE incidence among subjects in a SAS will be created. All hypothesis testing will be conducted at the significance level of 0.05 (two-sided). The data collected will be summarized using the arithmetic mean, the standard deviation, and the quantiles. All statistical analyses are conducted under the computing environment R. (R Foundation for Statistical Computing, Vienna, Austria)

## Discussion

The main purpose of this study is to determine whether treatment with the SGLT2 inhibitor luseogliflozin affects bone metabolism in elderly patients with T2DM by using second-generation HR-pQCT.

For the evaluation of this effect of luseogliflozin, we use metformin as an active control drug for the following three reasons. (1) We should avoid yielding differences in glycemic control between the luseogliflozin and control groups. If we use a placebo control, the effect of luseogliflozin on bone may be confounded by the improvement of glycemic control. (2) Metformin is reported to have a neutral effect or at least a not-negative effect on bone.(24, 25) We can avoid overestimating the efficacy of luseogliflozin for bone metabolism by using metformin as a control drug. (3) Metformin is the most commonly used

antidiabetic agent worldwide and is recommended by most of the relevant guidelines as a first-line therapy for T2DM.(14, 26, 27)

Participants treated with an  $\alpha$ -GI and/or a DPP-4 inhibitor in addition to metformin can also be enrolled in the study. There is no report demonstrating an increased risk of fracture associated with  $\alpha$ -GIs and DPP-4 inhibitors.(25, 28) In addition, those drugs are used widely in Japan for elderly patients with T2DM.

It is recognized that bone strength is determined by bone mass, morphology, and microstructure. In a clinical setting, an individual's bone mass is generally represented as the BMD assessed by DXA.(7) We will identify the precise changes in bone strength brought about by 1-year treatment with luseogliflozin by using second-generation HR-pQCT, which can detect the changes of bone morphology and microstructure. Sundh et al. reported that increased cortical porosity determined by HR-pQCT is associated with hip fracture in postmenopausal women, independent of areal BMD in the femoral neck determined by DXA even after adjusting for other fracture risk factors.(29) Salmon et al. reported that the trabecular and cortical bone structure evaluated by HR-pQCT can predict a future risk of femoral fracture, which is independent of the estimated fracture risk provided by the fracture risk assessment tool (FRAX). (7)

The mega-trial CANVAS demonstrated that canagliflozin treatment increased fracture incidents in patients with T2DM compared to those administered a placebo.(17) The increased fracture AEs might to a certain degree be caused by an increased frequency of falls in that patient population, which may have developed secondary to volume contraction or a reduction in blood pressure.(17) The increase in bone fractures observed in the CANVAS trial was not observed in the other CANVAS-like trials regarding the efficacy of canagliflozin, i.e., the CANVAS-Renal (CANVAS-R) trial(13) and eight randomized controlled studies (non-CANVAS studies).(17) There is no clear explanation for the differences in fracture risk between CANVAS and the nine studies including the CANVAS-R and eight non-CANVAS studies. There is no evidence of an increased fracture risk associated with the other SGLT2 inhibitors with higher selectivity of SGLT2 against SGLT1 compared to that of canagliflozin.(12, 19, 20) It is unclear whether fracture risk is a class effect of the SGLT2 inhibitors.

There are some limitations in the trial. First, the sample size is small (n=24). Second, the follow-up period of the trial may be insufficient to lead to a definitive conclusion regarding the bone effect of luseogliflozin. Third, we cannot deny that the other antidiabetic agents concomitantly used during the trial will influence the participants' bone metabolism. Despite these limitations, our research also has strengths. There has been no study assessing the effect of antidiabetic drugs on the bone microstructure using HR-pQCT in patients with T2DM. We will be able to elucidate the effect of an SGLT2 inhibitor on bone metabolism by using second-generation HR-pQCT, which provides direct *in vivo* morphometric information about the bone microarchitecture with a three-dimensional morphological technique.

The main goal of treatment for adolescent or middle-aged patients with T2DM is to prevent cardiovascular disease and microvascular complications such as diabetic retinopathy. However, maintaining the patients' quality of life while preventing fractures is also required when we manage

hyperglycemia in fragile elderly patients with T2DM. It is an important task for clinicians to clarify whether adverse effects of SGLT2 inhibitors on bone develop in elderly T2DM patients with a higher fracture risk.

## **Trial Status**

The recruitment is conducted from June 2019 to December 2020. The current version of this protocol is version 5.0 which was updated on 19 July 2019.

## **Abbreviations**

T2DM; type 2 diabetes mellitus, BMD; bone mineral density, SGLT2; Sodium-glucose co-transporter 2, HR-pQCT; high-resolution peripheral quantitative computed tomography, DXA; dual-energy X-ray absorptiometry, CANVAS; the Canagliflozin Cardiovascular Assessment Study, CANVAS-R; CANVAS-Renal trial

## **Declarations**

### **Ethics approval and consent to participate**

The certified review board of Japan's Ministry of Health, Labour and Welfare (no. 071180061). Informed consent will be obtained from all study participants.

### **Consent for publication**

Not applicable

### **Availability of data and materials**

Not applicable.

### **Competing interests**

None declared.

### **Funding**

This study is funded by Taisho Pharma Co. (Tokyo, Japan).

The funder played no part in study design; collection, management, analysis, and interpretation of data; writing of the report; and the decision to submit the report for publication.

### **Author contributions**

AH, IH, and SM wrote this manuscript. IH, RS, NA, KC, MO, and AK designed the protocol. AH, IH, RS, AI, NA, KC, and YK are conducting the study. SM, ST, JM, SS, and HY will perform the statistical analyses. All authors have contributed to the study concept and design.

## Acknowledgements

Not applicable.

## References

1. Vestergaard P. Discrepancies in bone mineral density and fracture risk in patients with type 1 and type 2 diabetes—a meta-analysis. *Osteoporos Int.* 2007;18(4):427-44.
2. Janghorbani M, Van Dam RM, Willett WC, Hu FB. Systematic review of type 1 and type 2 diabetes mellitus and risk of fracture. *Am J Epidemiol.* 2007;166(5):495-505.
3. Napoli N, Chandran M, Pierroz DD, Abrahamsen B, Schwartz AV, Ferrari SL, et al. Mechanisms of diabetes mellitus-induced bone fragility. *Nat Rev Endocrinol.* 2017;13(4):208-19.
4. de Waard EA, van Geel TA, Savelberg HH, Koster A, Geusens PP, van den Bergh JP. Increased fracture risk in patients with type 2 diabetes mellitus: an overview of the underlying mechanisms and the usefulness of imaging modalities and fracture risk assessment tools. *Maturitas.* 2014;79(3):265-74.
5. Boutroy S, Khosla S, Sornay-Rendu E, Zanchetta MB, McMahon DJ, Zhang CA, et al. Microarchitecture and Peripheral BMD are Impaired in Postmenopausal White Women With Fracture Independently of Total Hip T-Score: An International Multicenter Study. *J Bone Miner Res.* 2016;31(6):1158-66.
6. Stein EM, Kepley A, Walker M, Nickolas TL, Nishiyama K, Zhou B, et al. Skeletal structure in postmenopausal women with osteopenia and fractures is characterized by abnormal trabecular plates and cortical thinning. *J Bone Miner Res.* 2014;29(5):1101-9.
7. Samelson EJ, Broe KE, Xu H, Yang L, Boyd S, Biver E, et al. Cortical and trabecular bone microarchitecture as an independent predictor of incident fracture risk in older women and men in the Bone Microarchitecture International Consortium (BoMIC): a prospective study. *Lancet Diabetes Endocrinol.* 2019;7(1):34-43.
8. Paccou J, Ward KA, Jameson KA, Dennison EM, Cooper C, Edwards MH. Bone Microarchitecture in Men and Women with Diabetes: The Importance of Cortical Porosity. *Calcif Tissue Int.* 2016;98(5):465-73.
9. Burghardt AJ, Issever AS, Schwartz AV, Davis KA, Masharani U, Majumdar S, et al. High-resolution peripheral quantitative computed tomographic imaging of cortical and trabecular bone microarchitecture in patients with type 2 diabetes mellitus. *J Clin Endocrinol Metab.* 2010;95(11):5045-55.
10. Yu EW, Putman MS, Derrico N, Abrishamian-Garcia G, Finkelstein JS, Bouxsein ML. Defects in cortical microarchitecture among African-American women with type 2 diabetes. *Osteoporos Int.*

- 2015;26(2):673-9.
11. Farr JN, Drake MT, Amin S, Melton LJ, 3rd, McCready LK, Khosla S. In vivo assessment of bone quality in postmenopausal women with type 2 diabetes. *J Bone Miner Res.* 2014;29(4):787-95.
  12. Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med.* 2015;373(22):2117-28.
  13. Neal B, Perkovic V, Mahaffey KW, de Zeeuw D, Fulcher G, Erondou N, et al. Canagliflozin and Cardiovascular and Renal Events in Type 2 Diabetes. *N Engl J Med.* 2017;377(7):644-57.
  14. Davies MJ, D'Alessio DA, Fradkin J, Kernan WN, Mathieu C, Mingrone G, et al. Management of Hyperglycemia in Type 2 Diabetes, 2018. A Consensus Report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetes Care.* 2018;41(12):2669-701.
  15. Taylor SI, Blau JE, Rother KI. Possible adverse effects of SGLT2 inhibitors on bone. *Lancet Diabetes Endocrinol.* 2015;3(1):8-10.
  16. Neal B, Perkovic V, de Zeeuw D, Mahaffey KW, Fulcher G, Stein P, et al. Rationale, design, and baseline characteristics of the Canagliflozin Cardiovascular Assessment Study (CANVAS)—a randomized placebo-controlled trial. *Am Heart J.* 2013;166(2):217-23 e11.
  17. Watts NB, Bilezikian JP, Usiskin K, Edwards R, Desai M, Law G, et al. Effects of Canagliflozin on Fracture Risk in Patients With Type 2 Diabetes Mellitus. *J Clin Endocrinol Metab.* 2016;101(1):157-66.
  18. Bilezikian JP, Watts NB, Usiskin K, Polidori D, Fung A, Sullivan D, et al. Evaluation of Bone Mineral Density and Bone Biomarkers in Patients With Type 2 Diabetes Treated With Canagliflozin. *J Clin Endocrinol Metab.* 2016;101(1):44-51.
  19. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med.* 2019;380(4):347-57.
  20. Toulis KA, Bilezikian JP, Thomas GN, Hanif W, Kotsa K, Thayakaran R, et al. Initiation of dapagliflozin and treatment-emergent fractures. *Diabetes Obes Metab.* 2018;20(4):1070-4.
  21. Suzuki M, Honda K, Fukazawa M, Ozawa K, Hagita H, Kawai T, et al. Tofogliflozin, a potent and highly specific sodium/glucose cotransporter 2 inhibitor, improves glycemic control in diabetic rats and mice. *J Pharmacol Exp Ther.* 2012;341(3):692-701.
  22. Julious SA. Sample size of 12 per group rule of thumb for a pilot study. *Pharmaceutical Statistics.* 2005;4:287-91.
  23. Genant HK, Wu CY, van Kuijk C, Nevitt MC. Vertebral fracture assessment using a semiquantitative technique. *J Bone Miner Res.* 1993;8(9):1137-48.
  24. Wolverton D, Blair MM. Fracture risk associated with common medications used in treating type 2 diabetes mellitus. *Am J Health Syst Pharm.* 2017;74(15):1143-51.
  25. Kalaitzoglou E, Fowlkes JL, Popescu I, Thrailkill KM. Diabetes pharmacotherapy and effects on the musculoskeletal system. *Diabetes Metab Res Rev.* 2019;35(2):e3100.

26. Davies MJ, D'Alessio DA, Fradkin J, Kernan WN, Mathieu C, Mingrone G, et al. Management of hyperglycaemia in type 2 diabetes, 2018. A consensus report by the American Diabetes Association (ADA) and the European Association for the Study of Diabetes (EASD). *Diabetologia*. 2018;61(12):2461-98.
27. International Diabetes Federation. Recommendations For Managing Type 2 Diabetes In Primary Care, 2017. [www.idf.org/managing-type2-diabetes](http://www.idf.org/managing-type2-diabetes). Accessed 2 Mar 2019.
28. Mamza J, Marlin C, Wang C, Chokkalingam K, Idris I. DPP-4 inhibitor therapy and bone fractures in people with Type 2 diabetes - A systematic review and meta-analysis. *Diabetes Res Clin Pract*. 2016;116:288-98.
29. Sundh D, Nilsson AG, Nilsson M, Johansson L, Mellstrom D, Lorentzon M. Increased cortical porosity in women with hip fracture. *J Intern Med*. 2017;281(5):496-506.

## Tables

**Table 1.** The schedule of pre-screening, interventions, and assessments

Assessment	Pre-screening	Baseline	Monitoring				Withdrawal
	Day 60~ -1	Week 0	Week 12	Week 24	Week 36	Week 48	withdrawal
Visit	Visit 1	Visit 2	Visit 3	Visit 4	Visit 5	Visit 6	
Informed consent	●	-	-	-	-	-	-
Baseline characterization check	●	-	-	-	-	-	-
Adherence check	-	-	.	.	.	.	.
Concomitant treatments							
Adverse events	-						
Vital sign	●	●	.	.	.	.	.
Laboratory data	-	●	.	.	.	.	.
X-ray	-	●	-	-	-	.	-
DXA (BMD measurement)	-	●	-	-	-	.	-
HR-pQCT	-	●	-	-	-	.	.


● Will be done before initiating the study drug. · Will be done after initiating the study drug.

A range of  $\pm 4$  weeks is allowed for each visit day.

The examination at withdrawal would be carried out within 4 weeks from the date of withdrawal.

BMD: bone mineral density, DXA: dual-energy X-ray absorptiometry, HR-pQCT: high-resolution peripheral quantitative computed tomography.

## Figures



**Figure 1**

The study design. The eligibility check is conducted and the patient's informed consent is obtained at Visit 1. Only the patients who satisfy the run-in criteria at Visit 2 are enrolled in the study; they are randomized at a 1:1 ratio to the Luseogliflozin group (administered luseogliflozin 2.5 mg) and the Control group (administered metformin 500 mg). The bone microarchitecture of the patients is analyzed before (Visit 2) and 48 weeks after the administration of each medication (Visit 6). BMD: bone mineral density, SD: standard deviation, T2DM: type 2 diabetes mellitus.

## Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SPIRITchecklist20200319.doc](#)