Particulate Matter of Different Sizes Associated With Acute Lower Respiratory Infection Outpatient Visits in Children: A Time-Series Analysis in Guangzhou, China

Zhenyu Liang

Guangdong Second Provincial General Hospital

Qiong Meng

Guangdong Second Provincial General Hospital

Qiaohuan Yang

Guangdong Second Provincial General Hospital

Na Chen

Guangdong Second Provincial General Hospital

Chuming You (✉ gd2hek@163.com)

Guangdong Second Provincial General Hospital <https://orcid.org/0000-0001-6262-3463>

Research

Keywords: particulate matter, particle, lower respiratory infection, China

Posted Date: May 6th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-474721/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

The burden of lower respiratory infection is primarily borne by developing countries. However, the association between particulate matter of different sizes and acute lower respiratory infection (ALRI) outpatient visits in developing countries is less studied.

Methods

We obtained data on ALRI outpatient visits ($N = 105,639$) from a tertiary hospital in Guangzhou, China between 2013 and 2019. Over-dispersed generalized additive Poisson models were employed to evaluate the excess risk (ER) associated with particulate matter [inhalable particulate matter (PM_{10}), coarse particulate matter (PM_c), and fine particulate matter ($PM_{2.5}$)]. Counterfactual analyses were used to examine the potential percent reduction of ALRI outpatient visits if the levels of air pollution were as low as those recommended by the World Health Organization (WHO).

Results

There were 35,310 pneumonia, 68,218 bronchiolitis, and 2,111 asthma outpatient visits included. Each $10 \mu g/m^3$ increase of three-day moving averages of particulate matter was associated with significant ER [95% confidence interval (CI)] of outpatient visits of pneumonia [$PM_{2.5}$: 3.71% (2.91%, 4.52%); PM_c : 9.19% (6.94%, 11.49%); PM_{10} : 4.36% (3.21%, 5.52%)], bronchiolitis [$PM_{2.5}$: 3.21% (2.49%, 3.93%); PM_c : 9.13% (7.09%, 11.21%); PM_{10} : 3.12% (2.10%, 4.15%)], and asthma [$PM_{2.5}$: 3.45% (1.18%, 5.78%); PM_c : 11.69% (4.45%, 19.43%); PM_{10} : 3.33% (0.26%, 6.49%)]. The association between particulate matter and pneumonia outpatient visits was stronger among male patients and in cold seasons. Counterfactual analyses suggested that $PM_{2.5}$ was associated with the largest potential decline of ALRI outpatient visits [pneumonia: 3.89%, 95% CI: (3.24%, 5.52%); bronchiolitis: 4.35% (3.06%, 4.82%); asthma: 5.98% (1.92%, 10.37%)] if the air pollutants were reduced to the level of the reference guidelines.

Conclusion

Short-term exposure to $PM_{2.5}$, PM_c , and PM_{10} is associated with significant risk of ALRI outpatient visits, among which $PM_{2.5}$ is associated with the highest potential decline in outpatient visits if it could be reduced to the WHO recommended level.

Introduction

Lower respiratory infections, also known as pneumonia or bronchiolitis, are the sixth leading cause of death for all ages, resulting in around 2.4 million deaths worldwide in 2016[1]. The burden of lower respiratory infections is unevenly distributed across the globe and primarily born in developing countries and socioeconomically disadvantaged communities, where adequate nutrition, clean fuel, sanitation, and clean air are unavailable[2, 3].

Exposure to ambient particulate matter (PM) has been widely reported to be associated with lower respiratory infection[4–6]. However, evidence on the association between different sizes of PM and lower respiratory infections, especially that from developing countries in which the level of air pollution is high, is relatively little[7–9]. Meanwhile, it is widely acknowledged that short-term exposure to particulate matter is associated with ALRI hospitalizations[10–14], but we have not found any study that investigated the association between exposure to particulate matter and ALRI outpatient visits.

In this time-series analyses of outpatient visits in a tertiary hospital in Guangzhou, China from 2013 to 2019, we investigate the association between PM of different sizes (PM_{10} , PM_c , and $PM_{2.5}$) and the outpatient visits of ALRI, potential mediators, and the potential percent reduction of ALRI outpatient visits if the levels of particulate matter were as low as those recommended by the WHO.

Methods

Acute lower respiratory infection data

Data on hospital outpatient visits for ALRI were retrieved from the Guangdong Second Provincial General Hospital located in the southwest of the city (Figure 1), which is one of the highest-level (tertiary) hospitals in Guangzhou[15]. According to the International Classification of Diseases, Tenth Revision (ICD-10), hospital outpatient visits with the primary diagnoses of pneumonia (J12-J18), bronchiolitis (J20-J21), and asthma (J45-J46)[16] were obtained between February 2013 and December 2019. We aggregated the three subtypes of ALRI to a series of daily time-series data[17-20].

Air pollution and meteorological data

Daily concentrations of air pollution during the study period were obtained from 11 air monitoring stations in Guangzhou (Figure 1), including inhalable particulate matter (PM_{10}), coarse particulate (PM_c), fine particulate matter ($PM_{2.5}$), nitrogen dioxide (NO_2), sulfur dioxide (SO_2) and ozone (O_3). Following previous study[19], the PM_c concentrations were calculated by subtracting $PM_{2.5}$ from PM_{10} , because PM_{10} was consisted of $PM_{2.5}$ and PM_c . Air pollution measurement details have been previously described[21]. Approximately 1% of observation days had missing data for the air pollution, the missing data were imputed using a linear interpolation approach (the “na.approx” function in “zoo” package in R).

Daily meteorological data (mean temperature and relative humidity) were obtained from the National Weather Data Sharing System (<http://data.cma.cn/>). Because there is potentially high correlation among

different air pollutants and meteorological factors, we examined the pairwise Pearson correlation coefficients among these variables.[22, 23]

Statistical models

The ALRI data, daily air pollution concentrations and meteorological data were linked by date. Following prior similar epidemiology studies[24], the association between PM pollution and hospital outpatient visits for acute lower respiratory infections diseases was examined by an over-dispersed generalized additive Poisson model (GAM). In the model, public holidays (PH) and day of the week (DOW) were adjusted for as categorical variables. Seasonal patterns, long-term trends, temperature, and relative humidity were controlled for as smoothing splines. In accordance with the approaches used in previous studies[25, 26], we selected six degrees of freedom (df) per year for temporal trends, a df of 6 for moving average temperature of the current day and previous three days (Temp03) and relative humidity (RH).

Considering the delayed health effects of air pollutants, we examined the lag effects for different lag structures. We begin with the same day (lag0) up to five days lag (lag5) in the single-lag day models. We also considered the accumulated effects of multi-day lags (moving averages for the current day and the previous 1, 2 and 3 days [lag01, lag02, and lag03]).

Stratified analyses

In order to evaluate the potential effect modifiers of the PM pollution-ALRI associations, we conducted stratified analyses by gender (male vs. female), age group (age <5 vs. age 5-14), and season (warm vs. cold). The warm season was defined as from April to September, and the cold season was from October to March. The 95% confidence interval (CI) of the difference between group was calculated by the formula below:

$$Q_1 - Q_2 \pm 1.96\sqrt{(SE_1)^2 + (SE_2)^2}$$

where Q represents the estimated coefficient in each stratum, and SE is the corresponding standard error[27]. The difference was considered as statistically significant if the 95% CI did not include unity.

Sensitivity analyses

To examine the robustness of the main models, we applied a series of sensitivity studies. The main findings were assessed by changing the df in the smooth functions for temporal trends and meteorological factors. Additionally, we adjusted for gaseous air pollutants (SO₂, NO₂, and O₃) in two-pollutant models. The models were regarded as robust if there were no significant changes after df-changed or further adjustment for gaseous air pollutants.

Counterfactual analyses on the burden of ALRI attributable to air pollution

We estimated the burden of ALRI attributable to $PM_{2.5}$, PM_{10} , and $PM_{2.5}$ by calculating the difference between the observed ALRI outpatient visits and the counterfactual visits predicted using well-recognized reference values of air pollution recommended by the World Health Organization (WHO) and our previously built over-dispersed generalized additive Poisson models. This difference between the observed and counterfactual ALRI outpatient visits represents the estimated burden of ALRI outpatient visits associated with particulate matter of different sizes. The counterfactual scenarios were set to be hypothetical values of $PM_{2.5}$, PM_{10} , and $PM_{2.5}$ set by the WHO Air Quality Guidelines (24 hours mean: $25 \mu g/m^3$ for $PM_{2.5}$, $25 \mu g/m^3$ for PM_{10} , and $50 \mu g/m^3$ for $PM_{2.5}$). The observed air pollution levels lower than the reference values were kept the same in the counterfactual scenario. The 95% CIs were constructed using 1,000 bootstrap replicates with replacement for each model.[28, 29]

In all statistical analyses, a P value ≤ 0.05 was considered statistically significant. All data cleaning, aggregation, and visualization, and statistical analyses were done using statistical computing environment R (version 4.0.5)[30].

Results

Figure 1 presents the geographical location of Guangzhou city and the sample hospital, as well as the geographical distribution of the air monitoring stations in Guangzhou. A total of 35,310 pneumonia, 68,218 bronchiolitis, and 2,111 asthma cases were included in our analyses. Table 1 shows the summary statistics of ALRI subtypes, particulate matter of different sizes (PM_{10} , PM_{10} , and $PM_{2.5}$), and gaseous pollutants (SO_2 , NO_2 , and O_3). The daily averages [standard deviations (SD)] of pneumonia, bronchiolitis, and asthma cases were 12.5 (9.1), 24.3 (11.5), and 0.8 (1.4). The mean concentrations of PM_{10} , PM_{10} , and $PM_{2.5}$ in our study were 58.3, 21.0, and $37.8 \mu g/m^3$. The mean (SD) of temperature and relative humidity was 22.8 (5.9) °C and 81.8% (10.2%).

Table 1
Summary statistics of acute lower respiratory infections outpatient visits, air pollutants, and meteorological variables.

	Mean	SD	Percentile				
			Min	25th	50th	75th	Max
Acute lower respiratory infections							
Pneumonia	12.5	9.1	0.0	6.0	11.0	16.0	73.0
Bronchiolitis	24.3	11.5	0.0	16.0	23.0	31.0	81.0
Asthma	0.8	1.4	0.0	0.0	0.0	1.0	12.0
Air pollution, µg/m³							
PM ₁₀	58.3	28.1	10.0	38.2	51.1	73.4	217.8
PM _c	21.0	9.9	0.8	14.7	18.8	25.3	77.7
PM _{2.5}	37.8	21.2	4.6	22.7	32.3	48.3	156.4
SO ₂	13.6	8.5	2.8	8.6	11.9	16.5	166.4
NO ₂	45.2	18.6	4.4	33.6	41.2	53.7	177.7
O ₃	51.6	30.2	3.5	30.0	47.2	67.1	294.6
Meteorological variables							
Temperature, °C	22.8	5.9	1.7	19.0	25.0	27.5	32.8
Relative humidity, %	81.8	10.2	30.5	77.0	83.1	89.3	100.0
SD: standard deviation							

Figure 2 shows the correlation plot of the air pollutants and meteorological variables in our sample. All the Pearson correlation coefficients were statistically significant except for the correlation between NO₂ and O₃. PM₁₀ was significantly and strongly correlated with PM_c and PM_{2.5} (Pearson correlation coefficients: 0.81 and 0.93); NO₂ was moderately correlated with particulate matters (Pearson correlation coefficients for PM₁₀, PM_c, and PM_{2.5}: 0.78, 0.66, and 0.70). Meteorological variables were negatively correlated with air pollutants except for the positive correlation between temperature and O₃.

Table 2 exhibits the ER of pneumonia, bronchiolitis, and asthma outpatient visits associated with per 10 $\mu\text{g}/\text{m}^3$ increase of PM_{2.5}, PM_c, and PM₁₀ in lag03. The results revealed that particulate matters of different sizes were significantly associated with pneumonia, bronchiolitis, and asthma, respectively in single-pollutant models, where the ER of PM_c is the largest, followed by those of PM_{2.5} and PM₁₀. The

results were consistent and robust in two-pollutant models with further adjustment for SO₂, NO₂, and O₃, except for those asthma models controlling for NO₂. The corresponding exposure-response nonlinear curves for daily particulate matter and log relative risk are provided in Supplemental Fig. 1.

Table 2

Excess risk and 95% confidence intervals of pneumonia, bronchiolitis, and asthma for each 10 µg/m³ increase in PM_{2.5}, PM_c, PM₁₀ using single- and two-pollutants models.

Pollutants	Models	Pneumonia	Bronchiolitis	Asthma
PM ₁₀				
	Single-pollutant model	3.71 (2.91, 4.52)	3.21 (2.49, 3.93)	3.45 (1.18, 5.78)
	Two-pollutant models			
	Control for SO ₂	3.81 (2.97, 4.66)	3.44 (2.69, 4.21)	3.46 (1.13, 5.85)
	Control for NO ₂	2.47 (1.47, 3.47)	1.48 (0.58, 2.37)	0.26 (-2.58, 3.19)
	Control for O ₃	4.06 (3.22, 4.91)	3.48 (2.72, 4.25)	3.72 (1.30, 6.20)
PM _c				
	Single-pollutant model	9.19 (6.94, 11.49)	9.13 (7.09, 11.21)	11.69 (4.45, 19.43)
	Two-pollutant models			
	Control for SO ₂	9.32 (6.98, 11.72)	9.72 (7.58, 11.91)	11.70 (4.29, 19.63)
	Control for NO ₂	5.58 (3.03, 8.19)	4.80 (2.45, 7.20)	3.26 (-4.88, 12.09)
	Control for O ₃	9.52 (7.21, 11.87)	9.47 (7.37, 11.61)	12.09 (4.56, 20.17)
PM _{2.5}				
	Single-pollutant model	4.36 (3.21, 5.52)	3.12 (2.10, 4.15)	3.33 (0.26, 6.49)
	Two-pollutant models			
	Control for SO ₂	4.61 (3.37, 5.87)	3.50 (2.39, 4.63)	3.45 (0.14, 6.87)
	Control for NO ₂	2.30 (0.96, 3.65)	0.39 (-0.78, 1.58)	-0.40 (-3.86, 3.18)
	Control for O ₃	4.85 (3.63, 6.09)	3.38 (2.29, 4.48)	3.54 (0.26, 6.91)

Similar patterns of ER of ALRI outpatient visits associated with per 10 µg/m³ increase in different size fractions could be observed in Fig. 3. Each 10 µg/m³ increase in PM_{2.5}, PM_c, PM₁₀ was associated with outpatient visits of pneumonia, bronchiolitis, and asthma in different lag days. In contrast, the effects of matter of different sizes on asthma are less robust: the moving average lags of PM_c and PM_{2.5} were still

significantly associated with ALRI outpatient visits, but the effects of lag0 to lag5 of PM_c and $PM_{2.5}$ and different lags of PM_{10} were nonsignificant or at borderline significant.

Table 3 presents the estimated ER with 95% CI of pneumonia, bronchiolitis, and asthma stratified by gender, age group, and season, where the bold numbers indicate significant differences across strata. We can observe that each $10 \mu g/m^3$ increases in PM_{10} , PM_c , and $PM_{2.5}$ were consistently associated with significantly different effects on pneumonia outpatient visits by gender and season groups. Similar differential effects can be observed for bronchiolitis associated with increases in PM_{10} and PM_c by different season strata, but not for $PM_{2.5}$. However, the differential effects across strata were much less significant for asthma outpatient visits: it was only significantly different between warm and cold seasons.

Table 3

Excess risk and 95% confidence intervals of pneumonia, bronchiolitis, and asthma for each 10 $\mu\text{g}/\text{m}^3$ increase in $\text{PM}_{2.5}$, PM_c , and PM_{10} stratified by gender, age group, and season.

Pollutants	Stratum	Pneumonia	Bronchiolitis	Asthma
PM_{10}				
	Gender			
	Male	4.49 (3.54, 5.45)	3.44 (2.68, 4.21)	4.46 (1.62, 7.39)
	Female	2.68 (1.61, 3.75)	2.76 (1.84, 3.69)	1.78 (-1.51, 5.18)
	Age			
	< 5	3.50 (2.66, 4.34)	3.09 (2.36, 3.82)	1.78 (-0.97, 4.60)
	5–14	4.50 (2.71, 6.33)	3.70 (2.49, 4.92)	6.01 (2.39, 9.75)
	Season			
	Warm	-0.06 (-1.24, 1.13)	2.13 (1.03, 3.23)	6.53 (2.52, 10.69)
	Cold	5.12 (4.00, 6.25)	3.76 (2.74, 4.79)	1.76 (-1.00, 4.61)
PM_c				
	Gender			
	Male	11.07 (8.36, 13.83)	10.02 (7.82, 12.25)	15.65 (6.36, 25.74)
	Female	6.70 (3.78, 9.71)	7.57 (4.99, 10.21)	5.94 (-3.96, 16.87)
	Age			
	< 5	8.69 (6.37, 11.06)	8.76 (6.69, 10.88)	6.96 (-1.69, 16.38)
	5–14	10.76 (5.65, 16.13)	10.75 (7.34, 14.27)	17.98 (6.46, 30.74)
	Season			
	Warm	-0.93 (-4.33, 2.60)	3.77 (0.40, 7.26)	22.14 (6.92, 39.53)
	Cold	13.57 (10.46, 16.77)	11.92 (9.07, 14.84)	9.39 (0.93, 18.56)
$\text{PM}_{2.5}$				
	Gender			
	Male	5.31 (3.95, 6.69)	3.48 (2.39, 4.58)	4.13 (0.30, 8.09)

The bold type represents the statistically significant differences ($p < 0.05$).

Warm season: April to September; cold season: October to March.

Pollutants	Stratum	Pneumonia	Bronchiolitis	Asthma
	Female	3.11 (1.58, 4.65)	2.45 (1.14, 3.78)	1.92 (-2.54, 6.58)
	Age			
	< 5	4.07 (2.87, 5.28)	2.88 (1.84, 3.93)	1.53 (-2.16, 5.35)
	5–14	5.59 (3.07, 8.17)	4.10 (2.39, 5.85)	6.13 (1.22, 11.28)
	Season			
	Warm	0.08 (-1.49, 1.67)	3.00 (1.56, 4.46)	7.73 (2.62, 13.08)
	Cold	6.08 (4.42, 7.77)	3.01 (1.51, 4.53)	-0.30 (-4.05, 3.60)
The bold type represents the statistically significant differences ($p < 0.05$).				
Warm season: April to September; cold season: October to March.				

Table 4 shows the proportion reduction of ALRI (pneumonia, bronchiolitis, and asthma) outpatient visits attributable to $PM_{2.5}$, PM_c , and PM_{10} in Guangzhou from 2013 to 2019 using counterfactual analysis that compares to the WHO recommended levels ($25 \mu\text{g}/\text{m}^3$ for $PM_{2.5}$, $25 \mu\text{g}/\text{m}^3$ for PM_c , and $50 \mu\text{g}/\text{m}^3$ for PM_{10}). We found that $PM_{2.5}$ was associated with the largest decline of ALRI outpatient visits [pneumonia: 3.89%, 95% CI: (3.24%, 5.52%); bronchiolitis: 4.35%, 95% CI: (3.06%, 4.82%); asthma: 5.98%, 95% CI: (1.92%, 10.37%)] if the levels of air pollution were reduced to the level of the reference guidelines specified by the WHO.

Table 4

Counterfactual analysis on the percent of decline (95% confidence intervals) in acute lower respiratory infection outpatient visits if the level of $PM_{2.5}$, PM_c , and PM_{10} were reduced to the reference levels in Guangzhou from 2013 to 2019.

	Pneumonia	Bronchiolitis	Asthma
PM_{10}	3.89% (3.24%, 5.52%)	4.35% (3.06%, 4.82%)	5.98% (1.92%, 10.37%)
PM_c	1.42% (1.10%, 1.82%)	1.91% (1.48%, 2.36%)	1.79% (0.60%, 3.09%)
$PM_{2.5}$	4.63% (3.47%, 5.98%)	4.38% (2.68%, 5.97%)	6.07% (0.08%, 12.37%)
The references of PM_{10} , $PM_{2.5}$, and PM_c concentration were based on the World Health Organization's Ambient Air Quality guidelines (24h mean: $50 \mu\text{g}/\text{m}^3$ for PM_{10} , $25 \mu\text{g}/\text{m}^3$ for PM_c , and $25 \mu\text{g}/\text{m}^3$ for $PM_{2.5}$, respectively).			

Discussion

In this seven-year time-series analyses of daily ALRI outpatient visits data ($N = 105,639$) in Guangzhou, China from 2013 to 2019, we observed statistically significant ERs and potential declination of ALRI

(including pneumonia, bronchiolitis, and asthma) outpatient visits associated with particulate matter of different sizes. The results were consistent in exposure assessment using different lags (lag 0 to 5 and moving averages of 1 to 3 days), two-pollutant models adjusting for SO₂, NO₂, and O₃, and various degrees of freedom. In counterfactual analyses that are of more public health significance, PM_{2.5} was associated with the largest decline of ALRI outpatient visits if the exposure were as low as the WHO guideline reference.

Different effect estimates were observed among different sized PM with the largest ER being observed in PM_c, followed by PM_{2.5} and PM₁₀. However, these results should be interpreted with caution that PM_{2.5}, PM_c, and PM₁₀ have different means and standard deviations: the mean of PM_c in our sample (21.0 µg/m³) was lower than the WHO recommended level (25 µg/m³); the SD of PM_c (9.9 µg/m³) was much smaller compared to those of PM_{2.5} (21.2 µg/m³) and PM₁₀ (28.1 µg/m³). Therefore, the ER of ALRI associated with PM_c appeared to be the largest, which likely results from its small standard deviation.

In view of the limitation that the calculation of ER largely depends on the statistical distribution of the exposures, we further examined the potential proportion declination that would occur if exposure to particulate matter of different sizes were reduced to the WHO recommended levels (25 µg/m³ for PM_{2.5}, 25 µg/m³ for PM_c, and 50 µg/m³ for PM₁₀). In contrast to the result that PM_c was associated with the highest ER, our counterfactual analysis suggested that reducing PM_{2.5} to the WHO reference was associated with the largest potential decline in ALRI outpatient visits, followed closely by the reduction of PM₁₀; while reducing PM_c to the WHO reference is associated with the lowest potential decline in ALRI outpatient visits, which is likely explained by the fact that the mean level of PM_c (21.0 µg/m³) in our sample is lower than that of WHO reference level (25 µg/m³).

Our counterfactual analysis results have much more practical public health meaning than those of ER. The implication that reducing the level of PM_{2.5} may be associated with the largest decline in ALRI outpatient visits is consistent with previous reports about the toxicity of smaller-sized particulate matter on lower respiratory infection hospitalizations[10–14]. For example, Wang et al. specifically focused on the association between particulate matter of different sizes and childhood pneumonia, and they reported a graded impact of particulate matter of different sizes on childhood pneumonia (PM₁ > PM_{2.5} > PM₁₀). Smaller-sized particulate matter is more likely to enter lower and deeper lobes of lung and cause more severe consequences of health.

Although the air quality has been substantially improved attributable to the effort of air quality management in China over the past decade[31, 32], the average level of particulate matter (especially PM_{2.5} and PM₁₀), are still above the WHO recommended level. Northern China cities with high population density can experience anomalously high levels of air pollution during the winter[33]. Our results highlight the importance of focusing on smaller-sized particulate matter due to its harmful effect on ALRI outpatient visits.

This study should be interpreted in view of several limitations. First, similar to previous time-series studies on air pollution associated health outcomes, we used daily aggregated data to evaluate the short-term effect of particulate matter on health outcomes, but this aggregated nature of data could be subject to ecological bias. Second, we used fixed-site measurement of particulate matter instead of personal exposure, which could lead to inaccurate exposure measurement. Third, we included relatively small number of asthma outpatient visits, which led to unstable point estimates and CIs. Fourth, since we used secondary data collected from the hospital administrative database, environmental and behavior variables that could serve as confounders were not available to us. Lastly, COVID-19 has greatly reshaped the behavior of patient inpatient and outpatient visits, but we were not able to investigate the potential effect of the COVID-19 pandemic on ALRI outpatient visits since the data in this period is not available [34, 35].

Nonetheless, this study has several strengths. First, this is the first study that investigates the association between particulate matter of different sizes and subtypes of ALRI outpatient visits, while previous studies either reported the exposure of $PM_{2.5}$ or hospitalization as the health outcome. Second, we used counterfactual analyses to estimate the potential percent reduction in ALRI outpatient visits compared to the WHO recommended levels. The results of counterfactual analyses have more substantial public health significance compared to ER, OR, and any other estimates associated with fixed amount of increase in particulate matter (such as $10 \mu g/m^3$ increase in $PM_{2.5}$).

Conclusions

In summary, this study suggests a larger potential percent of reduction in ALRI outpatient visits if $PM_{2.5}$ could be as low as the level recommended by the WHO. The association between particulate matter and pneumonia outpatient visits was stronger among male patients and in cold seasons. The results highlight the need for a consolidated effort to reduce the particulate matter pollution of smaller sizes and consequently improve the health outcomes of residents in China.

Abbreviations

ALRI: Acute lower respiratory infection; CI: Confidential intervals; RR: Relative risk; ER: Excess risk; WHO: World Health Organization; PM: Particulate matter; PM_{10} : Particulate matter with an aerodynamic diameter $\leq 10mm$; PM_c : Particulate matter between 2.5 and $10\mu m$ in aerodynamic diameter; $PM_{2.5}$: Particulate matter with an aerodynamic diameter $\leq 2.5mm$ SO_2 : Sulfur dioxide; NO_2 : Nitrogen dioxide; O_3 : Ozone; GAM: Generalized additive models; DOW: Day of the week; PH: Public holidays; df: Degrees of freedom; ICD-10: International Classification of Diseases, Tenth Revision

Declarations

Acknowledgements

The authors thank the Chinese Meteorological Data Sharing Service System for providing the meteorological data used in this study.

Availability of data and materials

Please contact author for data requests.

Funding

This study was supported by the Science Foundation of Guangdong Second Provincial General Hospital (2021BSGZ001).

Authors' contributions

Study concept and design were contributed by ZL; statistical analysis was contributed by QM and QY; drafting of the manuscript was contributed by ZL, NC, and CY; All authors contributed to the interpretation of results and manuscript editing; All authors read and approved the final manuscript.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing of interests

The authors declare that they have no competing interests.

References

1. G. B. D. Causes of Death Collaborators: **Global, regional, and national age-sex specific mortality for 264 causes of death, 1980-2016: a systematic analysis for the Global Burden of Disease Study 2016.** *Lancet* 2017, **390**(10100):1151-1210.
2. G. B. D. Lower Respiratory Infections Collaborators: **Estimates of the global, regional, and national morbidity, mortality, and aetiologies of lower respiratory infections in 195 countries, 1990-2016: a systematic analysis for the Global Burden of Disease Study 2016.** *Lancet Infect Dis* 2018, **18**(11):1191-1210.
3. World Health Organization: **End preventable deaths: Global action plan for prevention and control of pneumonia and diarrhea.** *France: UNICEF* 2013.
4. Mehta S, Shin H, Burnett R, North T, Cohen AJ: **Ambient particulate air pollution and acute lower respiratory infections: a systematic review and implications for estimating the global burden of disease.** *Air Qual Atmos Health* 2013, **6**(1):69-83.

5. Johnston FH, Salimi F, Williamson GJ, Henderson SB, Yao J, Dennekamp M, Smith K, Abramson MJ, Morgan GG: **Ambient Particulate Matter and Paramedic Assessments of Acute Diabetic, Cardiovascular, and Respiratory Conditions.** *Epidemiology* 2019, **30**(1):11-19.
6. Burnett RT, Pope CA, 3rd, Ezzati M, Olives C, Lim SS, Mehta S, Shin HH, Singh G, Hubbell B, Brauer M *et al*: **An integrated risk function for estimating the global burden of disease attributable to ambient fine particulate matter exposure.** *Environ Health Perspect* 2014, **122**(4):397-403.
7. Kurmi OP, Lam KB, Ayres JG: **Indoor air pollution and the lung in low- and medium-income countries.** *Eur Respir J* 2012, **40**(1):239-254.
8. Horne BD, Joy EA, Hofmann MG, Gesteland PH, Cannon JB, Lefler JS, Blagev DP, Korgenski EK, Torosyan N, Hansen GI *et al*: **Short-Term Elevation of Fine Particulate Matter Air Pollution and Acute Lower Respiratory Infection.** *Am J Respir Crit Care Med* 2018, **198**(6):759-766.
9. Gurley ES, Salje H, Homaira N, Ram PK, Haque R, Petri WA, Jr., Bresee J, Moss WJ, Luby SP, Breyse P *et al*: **Indoor exposure to particulate matter and age at first acute lower respiratory infection in a low-income urban community in Bangladesh.** *Am J Epidemiol* 2014, **179**(8):967-973.
10. Wang X, Xu Z, Su H, Ho HC, Song Y, Zheng H, Hossain MZ, Khan MA, Bogale D, Zhang H *et al*: **Ambient particulate matter (PM₁, PM_{2.5}, PM₁₀) and childhood pneumonia: The smaller particle, the greater short-term impact?** *Sci Total Environ* 2021, **772**:145509.
11. Pu X, Wang L, Chen L, Pan J, Tang L, Wen J, Qiu H: **Differential effects of size-specific particulate matter on lower respiratory infections in children: A multi-city time-series analysis in Sichuan, China.** *Environ Res* 2021, **193**:110581.
12. Tian Y, Liu H, Wu Y, Si Y, Li M, Wu Y, Wang X, Wang M, Chen L, Wei C *et al*: **Ambient particulate matter pollution and adult hospital admissions for pneumonia in urban China: A national time series analysis for 2014 through 2017.** *PLoS medicine* 2019, **16**(12):e1003010.
13. Li D, Wang JB, Zhang ZY, Shen P, Zheng PW, Jin MJ, Lu HC, Lin HB, Chen K: **Effects of air pollution on hospital visits for pneumonia in children: a two-year analysis from China.** *Environ Sci Pollut Res Int* 2018, **25**(10):10049-10057.
14. Cheng CY, Cheng SY, Chen CC, Pan HY, Wu KH, Cheng FJ: **Ambient air pollution is associated with pediatric pneumonia: a time-stratified case-crossover study in an urban area.** *Environ Health* 2019, **18**(1):77.
15. Cai M, Liu E, Tao H, Qian Z, Lin X, Cheng Z: **Does Level of Hospital Matter? A Study of Mortality of Acute Myocardial Infarction Patients in Shanxi, China.** *Am J Med Qual* 2018, **33**(2):185-192.
16. Cai M, Xie Y, Bowe B, Gibson AK, Zayed MA, Li T, Al-Aly Z: **Temporal Trends in Incidence Rates of Lower Extremity Amputation and Associated Risk Factors Among Patients Using Veterans Health Administration Services From 2008 to 2018.** *JAMA Netw Open* 2021, **4**(1):e2033953.
17. Wang X, Leng M, Liu Y, Qian ZM, Zhang J, Li Z, Sun L, Qin L, Wang C, Howard SW: **Different sized particles associated with all-cause and cause-specific emergency ambulance calls: A multicity time-series analysis in China.** *Science of The Total Environment* 2021, **783**:147060.

18. Wang X, Qian Z, Wang X, Hong H, Yang Y, Xu Y, Xu X, Yao Z, Zhang L, Rolling CA *et al*: **Estimating the acute effects of fine and coarse particle pollution on stroke mortality of in six Chinese subtropical cities.** *Environmental pollution (Barking, Essex : 1987)* 2018, **239**:812-817.
19. Wang X, Tian J, Li Z, Lai J, Huang X, He Y, Ye Z, Li G: **Relationship between different particle size fractions and all-cause and cause-specific emergency ambulance dispatches.** *Environmental health : a global access science source* 2020, **19**(1):69.
20. Zhao Y, Wang S, Lang L, Huang C, Ma W, Lin H: **Ambient fine and coarse particulate matter pollution and respiratory morbidity in Dongguan, China.** *Environmental pollution (Barking, Essex : 1987)* 2017, **222**:126-131.
21. Wang X, Zhang L, Yao Z, Ai S, Qian ZM, Wang H, BeLue R, Liu T, Xiao J, Li X *et al*: **Ambient coarse particulate pollution and mortality in three Chinese cities: Association and attributable mortality burden.** *The Science of the total environment* 2018, **628-629**:1037-1042.
22. Cai M, Yazdi MAA, Mehdizadeh A, Hu Q, Vinel A, Davis K, Xian H, Megahed FM, Rigdon SE: **The association between crashes and safety-critical events: Synthesized evidence from crash reports and naturalistic driving data among commercial truck drivers.** *Transportation Research Part C: Emerging Technologies* 2021, **126**:103016.
23. Revelle W, Condon DM, Wilt J, French JA, Brown A, Elleman LG: **Web and phone based data collection using planned missing designs.** *Sage handbook of online research methods (2nd ed, p 578-595)* Sage Publications, Inc 2016.
24. Chen R, Yang J, Chen D, Liu WJ, Zhang C, Wang H, Li B, Xiong P, Wang B, Wang Y *et al*: **Air pollution and hospital outpatient visits for conjunctivitis: a time-series analysis in Tai'an, China.** *Environ Sci Pollut Res Int* 2021, **28**(12):15453-15461.
25. Lin H, Liu T, Xiao J, Zeng W, Li X, Guo L, Zhang Y, Xu Y, Tao J, Xian H *et al*: **Mortality burden of ambient fine particulate air pollution in six Chinese cities: Results from the Pearl River Delta study.** *Environment international* 2016, **96**:91-97.
26. Nhung NTT, Schindler C, Dien TM, Probst-Hensch N, Perez L, Kunzli N: **Acute effects of ambient air pollution on lower respiratory infections in Hanoi children: An eight-year time series study.** *Environ Int* 2018, **110**:139-148.
27. Zeka A, Zanobetti A, Schwartz J: **Individual-level modifiers of the effects of particulate matter on daily mortality.** *Am J Epidemiol* 2006, **163**(9):849-859.
28. Davison AC, Hinkley DV: **Bootstrap methods and their application:** Cambridge university press; 1997.
29. Cai M, Liu E, Zhang R, Lin X, Rigdon SE, Qian Z, Belue R, Chang JJ: **Comparing the Performance of Charlson and Elixhauser Comorbidity Indices to Predict In-Hospital Mortality Among a Chinese Population.** *Clin Epidemiol* 2020, **12**:307-316.
30. R Core Team: **R: A language and environment for statistical computing.** 2013.
31. Zhang Q, Zheng Y, Tong D, Shao M, Wang S, Zhang Y, Xu X, Wang J, He H, Liu W *et al*: **Drivers of improved PM2.5 air quality in China from 2013 to 2017.** *Proc Natl Acad Sci U S A* 2019, **116**(49):24463-24469.

32. Lu X, Zhang S, Xing J, Wang Y, Chen W, Ding D, Wu Y, Wang S, Duan L, Hao J: **Progress of air pollution control in China and its challenges and opportunities in the ecological civilization era.** *Engineering* 2020.
33. Le T, Wang Y, Liu L, Yang J, Yung YL, Li G, Seinfeld JH: **Unexpected air pollution with marked emission reductions during the COVID-19 outbreak in China.** *Science* 2020, **369**(6504):702-706.
34. Bowe B, Xie Y, Gibson AK, Cai M, van Donkelaar A, Martin RV, Burnett R, Al-Aly Z: **Ambient Fine Particulate Matter Air Pollution and the Risk of Hospitalization among COVID-19 Positive Individuals: Cohort Study.** *Environment International* 2021:106564.
35. Wang Y, Yuan Y, Wang Q, Liu C, Zhi Q, Cao J: **Changes in air quality related to the control of coronavirus in China: Implications for traffic and industrial emissions.** *Sci Total Environ* 2020, **731**:139133.

Figures

Figure 1

Geographical distribution of the sample hospitals and air monitoring stations. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion

whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 2

Correlation plot of the air pollutants and meteorological variables. The white cells indicate insignificance correlation coefficient (NO2 and O3).

Figure 3

Excess risk (95% confidence intervals) of hospital outpatient visits associated with 10 µg/m³ increase in PM₁₀, PM_c, and PM_{2.5}.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementalmaterials.docx](#)