Genome- wide characterization, expression profile analysis of WRKY family genes in *Santalum album* and functional identification of their role in abiotic stress

Haifeng Yan (✉ gstsyh@163.com)

South China Botanical Garden

Mingzhi Li

Biodata biotechnology CO. Ltd

Yuping Xiong

South China Botanical Garden

Yueya Zhang

South China Botanical Garden

Hanzhi Liang

South China Botanical Garden

Yuan Li

South China Botanical Garden

Xinhua Zhang

South China Botanical Garden

Meiyun Niu

South China Botanical Garden

Beiyi Guo

South China Botanical Garden

Jaime A. Teixeira da Silva

P.O. Box 7 Michi-Cho Post office

Guohua Ma

South China Botanical Garden

Research article

Keywords: WRKY transcription factor, sandalwood, SA (salicylic acid), MeJA (methyl jasmonate), salt tolerance

Posted Date: September 7th, 2019

DOI: <https://doi.org/10.21203/rs.2.14060/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at International Journal of Molecular Sciences on November 13th, 2019. See the published version at <https://doi.org/10.3390/ijms20225676>.

Abstract

Background: WRKY proteins are a large superfamily of transcription factors that are involved in diverse biological processes including development, as well as biotic and abiotic stress responses in plants. WRKY family proteins have been extensively characterized and analyzed in many plant species, including *Arabidopsis*, rice and poplar. However, knowledge on WRKY transcription factors in *Santalum album* is scarce. **Results:** Based on *S. album* genome and transcriptome data, 64 SaWRKY genes were identified in this study. A phylogenetic analysis based on the structures of WRKY protein sequences divided these genes into three major groups (I, II, III) together with WRKY protein sequences from *Arabidopsis*. Tissue-specific expression patterns showed that 37 SaWRKY genes were expressed in at least one of five tissues (leaves, roots, heartwood, sapwood, or the transition zone) while the remaining four genes were weakly expressed in all of these tissues. Analysis of the expression profiles of the 42 SaWRKY genes after callus was stimulated by salicylic acid (SA) and methyl jasmonate (MeJA) revealed that 34 and 19 SaWRKY genes, respectively were significantly induced. The function of SaWRKY1, which was significantly up-regulated by SA and MeJA, was analyzed. SaWRKY1 was localized in the nucleus and its overexpression improved salt tolerance in transgenic *Arabidopsis*. **Conclusions:** Our study provides important information to further identify the functions of SaWRKY genes and to understand the roles of SaWRKY family genes involved in development and in SA- and MeJA-mediated stress responses.

Background

Transcription factors (TFs) are proteins that can bind to specific DNA target sites and activate and/or repress transcription of their target genes [1]. Among them, WRKY TFs are mainly found in plants while a few are found in the protist *Giardia lamblia* and the slime mold *Dictyostelium discoideum* [2, 3]. In green plants, the WRKY family is the eighth largest TF family following ethylene responsive factor (ERF), C₂H₂, basic helix-loop-helix (bHLH), myeloblastosis (MYB), NAC (NAM, ATAF1/2 and CUC2), basic leucine zipper (bZIP), and MYB-related families [4] (<http://planttfdb.cbi.pku.edu.cn/>). The first WRKY TF, SWEETPOTATO FACTOR1 (SPF1), was identified in sweet potato (*Ipomoea batatas*) in 1994 [5]. Since then, numerous WRKY genes have been identified after genome-wide analyses of plants over the past 24 years, including 74 in *Arabidopsis* (*Arabidopsis thaliana*) [6], 102 in rice (*Oryza sativa*) [7], 197 in soybean (*Glycine max*) [8], and 100 in poplar (*Populus trichocarpa* and *P. tomentosa*) [9]. The WRKY family is characterized by the presence of one or two conserved 60 amino acid WRKY domains (WDs), which include the highly conserved WRKYGQK sequence at its N-terminal end and a zinc-finger-like motif at the C-terminal end [6]. The C-terminal WRKY domain is mainly responsible for recognizing the cognate *cis*-acting W box elements (T)(T)TGAC(C/T) while the N-terminal end participates in increasing the affinity or specificity of their target sites [6]. Based on the number of WDs and the pattern of their zinc-finger-like motifs, all members of the WRKY family proteins have been classified into three groups I-III [6, 10]. WRKY proteins with two WDs fall into group I whereas proteins with only one WD are assigned to groups II or III. Members of groups I and II have the same type of finger motif, C-C-H-H, while members of group III

contain a unique C-C-H-C motif [6]. In addition, group II proteins were further split into five distinct subgroups (IIa, IIb, IIc, IId and IIe) based on a phylogenetic analysis [6, 10]. Apart from the WDs and zinc-finger-like motif, domains for transcobalamin (R) proteins also coexist with WRKY TFs forming R protein-WRKY genes in monocot genomes such as in switchgrass (*Panicum virgatum*) [11].

WRKY TFs play well-documented roles in regulating various biological processes in plants. The majority of reports demonstrated that numerous WRKY TFs are involved in transcriptional reprogramming associated with the plant's biotic stress response. *AtWRKY3* and *AtWRKY4* from *A. thaliana* were induced by pathogen infection and salicylic acid (SA), and the T-DNA insertion mutants and transgenic overexpression lines of *AtWRKY3* and *AtWRKY4* suggested that they all had enhanced plant tolerance to necrotrophic pathogens, while *AtWRKY4* negatively regulated plant tolerance to biotrophic pathogens [12]. *AtWRKY38* and *AtWRKY62* were induced by pathogen infection and SA treatment, serving as negative regulators of plant basal defense [13]. In rice, *OsWRKY03* (an *AtWRKY29/22* homolog) was involved in SA-dependent or jasmonic acid (JA)-dependent defense signaling cascades and its overexpression upregulated pathogenesis-related genes *OsNPR1*, *OsPR1b*, phenylalanine ammonia-lyase *ZB8* and peroxidase *POX22.3* [14]. Furthermore, accumulating data has shown that WRKY TFs are involved in abiotic stresses, including cold, wounding, heat, ultraviolet light, drought and salinity [15, 16]. In addition, WRKY TFs also play roles in regulating plant development, such as seed dormancy and germination [17, 18], root hair growth [19], and leaf senescence [20].

Santalum album L., commonly known as sandalwood, is a hemiparasitic tropical tree distributed in India, Indonesia, Malaysia, and Australia [21]. It is valued for its essential oil, which is extracted from aromatic heartwood and roots, and is used in aromatherapy, perfumes, cosmetics, medicine and sacred unguents [22, 23]. (*Z*)- α -Santalol, (*Z*)- β -santalol, (*Z*)-*epi*- β -santalol, (*Z*)- α -*exo*-bergamotol, and sesquiterpene alcohols are the major components (70%) of *S. album* essential oil [23]. Unsustainable demand for and exploitation of this slow-growing tree has threatened wild sandalwood populations in their native locations [24]. In an effort to meet the demand for its essential oil, *S. album* has been planted in Malaysia, Indonesia, the Philippines, Northern Australia, and South China [25]. *S. album* was introduced to South China Botanical Garden in 1970, and is now cultivated in several subtropical regions of China such as Guangdong, Guangxi, Hainan, and Fujian provinces. Various stresses, including low temperatures [26] or highly saline or alkaline soil, negatively impact the cultivation and growth of this species. The molecular biology of *S. album* has made considerable strides in recent years [27], including abiotic stress tolerance. WRKY TFs play important roles in plants against abiotic stress, so characterization and functional analysis of the WRKY family genes in *S. album* would shed light on the stress mechanisms underlying the involvement of WRKY family genes, and it could also serve to breed *S. album* varieties that are resistant to these stresses. As far as we are aware, there is scarce knowledge on WRKY TFs in *S. album*. Large-scale available RNA-seq data sets [26, 28, 29] and genomic data generated recently in *S. album* [30] might offer additional insight about WRKY family genes.

In this study, a total of 64 *SaWRKY* genes from *S. album* were identified, and their conserved motif and tissue expression patterns were analyzed. In order to understand their evolutionary relationship, a

phylogenetic tree that combined WRKY proteins from *A. thaliana* was constructed. Furthermore, the expression profiles of 42 *SaWRKY* genes under SA and methyl jasmonate (MeJA) stress were generated using RT-qPCR and the function of *SaWRKY1*, a gene that was induced both by SA and MeJA, was characterized in transgenic *A. thaliana*. The findings of our study will help to understand the roles of *SaWRKY* genes in SA- and MeJA-mediated pathways and to further identify the functions of this essential gene family in *S. album*.

Methods

Plant materials and hormone treatments

The leaves, roots and stems were collected from three seven-year-old *S. album* trees growing in South China Botanical Garden. Callus was induced from newly sprouted shoots following previously reported methods [25, 53], with a few modifications, as described next. Shoot segments were inoculated on solid Murashige and Skoog (MS) basal medium [54] to which 1.0 mg/L thidiazuron (TDZ) was added to induce callus. The resulting callus was proliferated on solid MS medium supplemented with 1.5 mg/L 2,4-dichlorophenoxyacetic acid (2,4-D) and 0.2 mg/L TDZ. After about 20 days, similarly good callus (Additional file 5: Figure S2) was proliferated after transferring into liquid MS basal medium for 20 days then shaken continuously at 100 rpm under the same culture conditions. After 24 h, SA and MeJA were added separately at a final concentration of 100 μ M each. Samples were collected at 0, 3 and 6 h with three biological triplicates per treatment. All samples were frozen immediately in liquid nitrogen (N₂) and stored at -80°C until use.

Identification of *WRKY* genes in *S. album* and prediction of subcellular localization

The *S. album* genome was downloaded from the National Center for Biotechnology Information (NCBI) (GCA_002925775.1) and then annotated using sandalwood transcriptome sequencing data and genome sequence data from maize, rice and Arabidopsis by a web service for training AUGUSTUS and predicting genes in eukaryotes (WebAUGUSTUS; <https://bio.tools/webaugustus>). *S. album* transcriptome sequencing data was downloaded from the NCBI Sequence Read Archive (SRA) (GenBank accessions: PRJNA297453, SRA150639, SRR3731808, and SRR3731809), and the *AtWRKY* gene sequences were searched and downloaded from The Arabidopsis Information Resource (TAIR) website (<http://www.arabidopsis.org/index.jsp>). All the *S. album* *WRKY* genes were identified following He et al. [55]. In brief, candidate *SaWRKY* genes were obtained by searching *S. album* protein sequences using the hidden Markov model (<http://hmmer.janelia.org/>). Putative *SaWRKY* genes were further verified for the presence of the PFAM domain (PF03106) in the pFAM database (<http://pfam.xfam.org/>) and then screened for the WRKY motif and zinc-finger domains, and finally confirmed as *SaWRKY* genes after removing redundant sequences by CD-HIT software (see Additional file 1: Table S1). The subcellular localization of *SaWRKY* genes was predicted using the PSORT program (<https://psort.hgc.jp/>).

Construction of phylogenetic tree and conserved motif analysis of SaWRKY proteins

An unrooted phylogenetic tree of *S. album* and Arabidopsis WRKY proteins was constructed using MEGA6.0 software with 1000 bootstrap replicates. The parameters used to construct trees were: gaps/missing data: complete deletion; substitution model: maximum composite likelihood; rates among sites: gamma distributed with invariant sites (G).

The online program MEME v 4.11.2 (<http://meme.nbcr.net/meme/>) was used to analyze conserved motifs for the 64 *S. album* WRKY protein sequences. MEME parameters were set as advised by He et al. [55]. The TBtool was used to draw a schematic diagram of the conserved motifs and gene structures.

Identification and functional enrichment analysis of potential SaWRKY target genes

The 2-k DNA sequence upstream of the ATG start codon of each gene assembled from the *S. album* genome was used to scan the WRKY TF binding site element with the sequence (C/T)TGAC(C/T). The *SaWRKY* genes with at least five WRKY binding sites were identified as potential *SaWRKY* target genes, and were used for further pathway enrichment analysis using the KEGG database. The top 20 enrichment KEGG pathways were drawn with R package ggplot2.

Gene expression analysis by quantitative real-time PCR (RT-qPCR)

Total RNA from *S. album* callus was isolated using the Eastep[®] Super Total RNA Extraction kit (Promega, Shanghai, China), and the RNA from leaves, stems and roots was extracted using a protocol reported for the isolation of RNA from woody plants [56]. The quality and quantity of DNA-free total RNA was assessed using a NanoDrop ND-1000 spectrophotometer (Nanodrop Technologies, Wilmington, NC, USA). RNA samples with an A_{260}/A_{280} ratio between 1.9 and 2.1 and an A_{260}/A_{230} ratio greater than 2.0 were used for subsequent analysis. RNA integrity was assessed by 1.5% (w/v) agarose gel electrophoresis. Total RNA (1 µg) was used to synthesize first-strand cDNAs using an equivalent amount of oligo-(dT)₁₅ and random primers in 20 µL volume with the GoScript[™] Reverse Transcriptase system (Promega, Madison, WI, USA) according to the manufacturer's protocols. Amplification was performed at 25°C for 5 min, 42°C for 60 min, and 70°C for 15 min. The successfully synthesized cDNA samples were diluted 1:10 with nuclease-free water and stored at -20°C until further use.

RT-qPCR was performed in 96-well plates in an ABI 7500 Real-time system (ABI, Alameda, CA, USA) using the SoAdvanced[™] Universal SYBR[®] Green Supermix detection system (Bio-Rad, Hercules, CA, USA). The RT-qPCR reaction in a total volume of 10 µL consisted of 5 µL SYBR[®] Green Supermix, 0.5 µL of forward primer (10 µM), 0.5 µL of reverse primer (10 µM), 1 µL of cDNA, and 3 µL of ddH₂O. The cycling

conditions were 95°C for 2 min, followed by 40 cycles at 95°C for 15 s and 60°C for 1 min. After 40 cycles, melting curve analysis was performed ranging from 60 to 95°C. *S. album* actin (Genbank accession: EF452617) was used as the internal reference gene. The relative gene expression level was calculated using the $2^{-\Delta\Delta C_q}$ method. All RT-qPCR experiments were carried out using three biological replicates of each sample. The gene-specific primers used for RT-qPCR are listed in Additional file 6: Table S4.

Cluster analysis of expression data

For the expression profiles of all putative *SaWRKY* genes in different tissues including leaves, roots and wood (containing heartwood, sapwood and the transition zone), the data that was obtained from the NCBI SRA database was mapped with *SaWRKY* gene nucleotide sequences using TopHat version 2.0.8. The level of gene expression was calculated by reads per kilobase of exon model per million mapped reads (RPKM), and the expression profiles via a heat map were calculated from the log₂ (FPKM+1) value, and shown as a green-black-red gradient by using the heat map package in R version 3.4.0. Expression (Cq) values of the 42 putative *SaWRKY* genes at each time point in SA and MeJA treatments were determined using RT-qPCR. The expression profiles were generated by a heat map using R language based on the log₂ ($2^{-\Delta\Delta C_q}$) value. Up- and down-regulated genes were defined as those showing more than a 1.5-fold change or lower than a 0.5-fold change, respectively ($P < 0.05$).

Cloning and subcellular localization of *SaWRKY1*

RACE was performed to obtain the full-length sequence of *SaWRKY1* using the SMARTer RACE cDNA Amplification Kit (Clontech Laboratories Inc., CA, USA; see Additional file 7: Method S1, Additional file 8: Figure S3). The amplification product was sequenced at the Beijing Genomics Institute (BGI, Shenzhen, China), and then submitted to NCBI (accession number: MG655189). The theoretical isoelectric point and protein molecular weight were determined using Expasy server (<https://web.expasy.org/cgi-bin/protparam/protparam>).

The full length coding sequence of *SaWRKY1* without the termination codon was amplified with gene-specific primers (SaWRKY1YFPF, 5'-CGAACGATAGCCATGG

CAATGGATTACTCTTCTTGGAC-3'; SaWRKY1YFPR, 5'-TGAGTCCGGACCAT

GGTAAATTTTTCATTTTGACTTT-3') and then fused to yellow fluorescent protein (YFP) behind the Cauliflower mosaic virus (CaMV) 35S promoter in the *Nco*I site of the pSAT6-EYFP-N1 vector following the In-Fusion HD Cloning Kit (Clontech) instructions. The 35S: *SaWRKY1*: pSAT6-EYFP-N1 construct was confirmed by sequencing and then co-transformed with a nuclear location protein mCherry into *A. thaliana* mesophyll protoplasts according to a protocol described by Yoo et al. [57]. After 12 h incubation

at 22°C under light conditions, YFP fluorescence was visualized with a Zeiss LSM 510 confocal microscope (Zeiss, Jena, Germany), and images were captured using the same software.

***Arabidopsis thaliana* transformation and salinity stress**

The full-length coding sequence of *SaWRKY1* was amplified with gene-specific primers (SaWRKY10XF, 5'-GGACTCTTGACCATGGTAATGGATTACTCTTCTT

GGACGGC-3'; SaWRKY10XR, 5'-GTCAGATCTACCATGGTAAATTTTTCATT

TTGACTTT-3') and then cloned into the pCAMBIA1302 vector behind the CaMV 35S promoter in the *NcoI* site using the In-Fusion HD Cloning Kit. The resulting *35S: SaWRKY1: pCAMBIA1302* construct, which was confirmed by sequencing and then transformed into *Agrobacterium tumefaciens* EHA105 using the freeze-thaw method described by Weigel and Glazebrook [58], was introduced into *A. thaliana* by the floral dip method [59]. Transformants were screened on half-strength MS medium containing 25 mg/L hygromycin. After two generations of selfing hygromycin-resistant lines, a homozygote was generated without segregation. RT-qPCR was performed using gene-specific primers (SaWRKY1qPCRf/R see Additional file 6: Table S4) to confirm the expression of *SaWRKY1* in transformed plants (Additional file 9: Figure S4). Two *SaWRKY1* overexpression transgenic lines from the homozygous T3 generation, which were about 4 weeks old, were irrigated with 300 mM NaCl at three-day intervals to observe their phenotypes and to determine chlorophyll content using aqueous acetone as the solvent, as described by Porra et al. [60]. On the same dates, *A. thaliana* ecotype Columbia (Col-0) was also watered with 300 mM NaCl and used as the control. Experiments were performed at least three times to observe resulting phenotypes and to determine chlorophyll content.

Statistical analyses

Statistical analysis was performed using SPSS 19.0 (IBM Corp., Armonk, NY, USA). Following one-way analysis of variance (ANOVA), significance between treatment means was assessed by Duncan's multiple range test at $P < 0.05$ and $P < 0.01$.

Results

Identification and subcellular localization of *SaWRKYs*

A total of 64 *SaWRKY* genes named *SaWRKY1* to *SaWRKY64* were identified (Table 1) by searching the *S. album* genome and transcriptome datasets using total *A. thaliana AtWRKY* genes as queries. All of the identified *SaWRKY* proteins contained at least one highly conserved heptapeptide WRKYGQK domain, while 13 out of 64 *SaWRKY* proteins contained two WRKYGQK domains. The length of the amino acid

sequence in the 66 SaWRKY proteins (Table 1, Additional file 1, Table S1) ranged from 121 (SaWRKY45) to 764 (SaWRKY12) amino acids, with an average of 384 amino acids. A C-C-H-H type zinc-finger motif was found in 56 SaWRKY proteins whereas SaWRKY7, 9, 28, 47 and 57 had a C-C-H-C type zinc-finger motif, and other variants of zinc-finger motifs such as C-C-H-T (SaWRKY12, 53), C-C-H-L (SaWRKY25, 51), C-C-H-V (SaWRKY42), C-C-H-Y (SaWRKY45), and C-C-H-S (SaWRKY58), were also found (Table 1).

Based on a prediction by the PSORT program, the subcellular localization of most of the 64 SaWRKY proteins was in the nucleus, except for SaWRKY8, which was found in the Golgi apparatus, SaWRKY45 in the cytoplasm, SaWRKY46 in the peroxisome, and SaWRKY49 in the chloroplast (Table 1).

Phylogenetic analysis of SaWRKY proteins

To understand the evolutionary relationship between SaWRKY proteins, the 64 identified SaWRKY proteins was examined based on AtWRKY proteins from the three groups, and an unrooted tree was built by MEGA6.0 software using the NJ method (Fig. 1). All 64 SaWRKY proteins were classified into three major groups (I, II and III) (Table 1, Fig. 1). Among the 15 SaWRKY proteins in group I, 13 contained two WRKY domains and the remaining two SaWRKY proteins (SaWRKY8 and 25) contained only one WRKY domain (Table 1). There were 43 SaWRKY proteins with only one WRKY domain in group II, and these could be further divided into an additional five subgroups, i.e., IIa, IIb, IIc, IId, and IIe. SaWRKY7, 9, 28, 47, 53 and 57, which contained only one WRKY domain, formed group III (Table 1).

Exon-intron organization of *SaWRKY* genes

To further understand the pivotal role that exon-intron structural features play in the evolution of *S. album* gene families, the structure of *SaWRKY* genes was obtained through exon-intron organization analysis. Among the 64 *SaWRKY* genes, three had one intron and two exons, 35 had two introns and three exons, six had three introns and four exons, nine had four introns and five exons, two had five introns and six exons, three had five introns and five exons, while the remaining *SaWRKY49* had six introns and seven exons, and *SaWRKY54* had 13 introns and 14 exons (Fig. 2b). It is noteworthy that *SaWRKY* genes in the same subgroup had a similar intron and exon composition, such as two introns and three exons in group II, subgroups IId, IIe and group III, or four introns and five exons in subgroup IIb. *SaWRKY54* had 14 exons, *SaWRKY49* had seven exons, and *SaWRKY2* and *SaWRKY16* had only two exons. This indicates an occurrence of both exon gain and loss during evolution of the *SaWRKY* gene family, thus leading to functional diversity among *SaWRKY* genes.

Motif composition of SaWRKY proteins

To gain insight into the functional regions of SaWRKY proteins, the MEME program was used to predict the composition of the 64 SaWRKY protein motifs. A total of 20 conserved motifs were detected (Fig. 2c). Among them, motifs 1 and 3 contained the heptapeptide stretch WRKYGQK while all 64 SaWRKY proteins contained one or two WRKYGQK motifs. Motif 2 was the conserved zinc-finger structure at the C-terminal end and was found in 61 SaWRKY proteins, but not in SaWRKY8, 11 and 51 (Fig. 2). Motif 9 was unique to all members of subgroup IId, motif 16 was unique to SaWRKY42, 43 and 52, and motif 17 was unique to SaWRKY30 and 36. Similar motif compositions were found in the same groups, especially in the same subgroups, such as in subgroups IIa or IIb.

Prediction and functional enrichment analysis of potential SaWRKY target genes

A total of 13,306 genes that contained at least one W-box in their putative promoters were identified in the assembled *S. album* genome. The number of genes decreased as more W-boxes were identified. Among all 13,306 genes, 2563 genes contained one W-box, 3007 genes contained two W-boxes, while 1328 genes contained at least five W-boxes and were used for further pathway enrichment analysis using the Kyoto Encyclopedia of Genes and Genomes (KEGG) database. As shown in Fig. 3, the top enriched KEGG pathways included plant-pathogen interactions, environmental adaption, metabolism and organismal systems. These results indicate that *SaWRKY* genes are closely involved in biotic and abiotic stress responses, as well as in other biological pathways.

Expression patterns of *SaWRKY* genes in different tissues

A gene expression pattern may reflect its biological function. To explore the possible functions of *SaWRKY* genes in *S. album* development, the expression patterns of 41 *SaWRKY* genes in various tissues (leaves, roots, heartwood, sapwood, and transition zone) were obtained from the transcriptome data (Fig. 4). Five *SaWRKY* genes (*SaWRKY16*, 20, 22, 3 and 44) showed a higher expression level in wood tissue (heartwood, sapwood, and transition zone) than in leaves and roots. The expression of *SaWRKY2*, 15 and 27 occurred preferentially in heartwood. Higher levels of mRNA were observed in roots for *SaWRKY4*, 10, 26 and 34 while 12 *SaWRKY* genes (*SaWRKY3*, 5, 8, 9, 17, 21, 28, 29, 33, 36, 38, and 41) showed a higher level of expression in leaves than in other tissues. *SaWRKY6*, 25, 35 and 37 had a low level of expression in all the tissues examined, while *SaWRKY12*, 30, 31, 39 and 42 had consistently high expression levels in all tissues (Fig. 4). It was not possible to assess the expression level of 23 *SaWRKY* genes in all tissues from the transcriptome data.

Expression profiles of *SaWRKY* genes in response to SA and MeJA

To detect whether the *SaWRKY* genes were induced by different hormones, RT-qPCR was performed to determine the expression levels of the 42 *SaWRKY* genes when stimulated by SA and MeJA in callus. The data shows that nine out of 42 *SaWRKY* genes were up-regulated by SA, namely *SaWRKY1*, 3, 9, 11, 25, 28, 37, 38 and 40 (Fig. 5A, Additional file 2: Table S2). In contrast, 25 *SaWRKY* genes were down-regulated by SA, and were mainly dispersed in group I and subgroup IIc as well as in a few members of subgroups IIb, IIc and IIe. Several *SaWRKY* genes were regulated by MeJA. Among them, *SaWRKY1*, 3, 4, 7, 8, 11, 16, 26, 29, 36, 38 and 42 were up-regulated whereas *SaWRKY10*, 13, 24, 25, 27, 28 and 38 were down-regulated (Fig. 5B, Additional file 3: Table S3). Interestingly, four *SaWRKY* genes (*SaWRKY1*, *SaWRKY3*, *SaWRKY11* and *SaWRKY38*) were up-regulated both by SA and MeJA.

Characterization, tissue expression patterns and subcellular localization of *SaWRKY1*

As indicated above, the expression level of *SaWRKY1* was not detected in five tissues from the transcriptome data. More interestingly, since *SaWRKY1* is one of the genes that was up-regulated both by SA and MeJA, it was selected for further analysis. The full length of the coding nucleotide sequence of *SaWRKY1* was 999 bp, encoding a protein of 332 amino acid residues with a predicted theoretical isoelectric point and protein molecular weight of 7.55 and 37.01 kD, respectively.

Tissue expression patterns, which were examined by RT-qPCR, showed that transcription of *SaWRKY1* took place mainly in callus, leaves and roots (Fig. 6). The abundance of *SaWRKY1* mRNA accumulated preferentially in callus (Fig. 6).

The subcellular localization of the *SaWRKY1* protein is most likely in the nucleus. To verify its subcellular location, a C-terminal YFP fusion construct for the *SaWRKY1* protein and a nuclear location protein mCherry was co-transformed into *Arabidopsis* mesophyll protoplasts. The *SaWRKY1*-YFP fusion protein was localized in the nucleus and co-localized with mCherry (Fig. 7), demonstrating that the *SaWRKY1* protein is located in the nucleus, in accordance with its putative function as a TF.

Overexpression of *SaWRKY1* enhances salinity tolerance in transgenic *Arabidopsis* plants

SaWRKY1 was significantly induced both by SA and MeJA, indicating that it might be a node of convergence in signal transduction pathways mediated by SA and MeJA. To further explore the roles of *SaWRKY1* in abiotic stress responses, the 35S: *SaWRKY1*: *pCambia1302* construct was transformed into *A. thaliana*, and two independent T3 homozygous progeny lines were used for further analysis.

In order to examine the role of *SaWRKY1* in abiotic stress, the two *A. thaliana* transgenic lines, which showed the same level of germination, growth properties, and chlorophyll content (Fig. 8A, 9A), were randomly selected and irrigated with 300 mM NaCl. After three days, both transgenic lines grew well but the wild type control showed obvious inhibited growth (Fig. 8A, B; Fig. 9A, B). The content of chlorophyll a,

b, and *a + b* in the wild type declined by 34.6%, 36.1% and 35.5%, respectively compared with the content of each component averaged over the two transgenic lines (Fig. 8B, 9B). When irrigated once again with 300 mM NaCl, unlike the wild type plants which withered and nearly died, the *35S:WRKY1* transgenic lines showed little withering, had greener leaves and slightly inhibited growth after 5 days (Fig. 8A, B, C). Accordingly, either the content of chlorophyll *a* and *b*, or the total chlorophyll content was reduced by more than 80% in wild type *A. thaliana* compared to the content of each component in *35S:WRKY1* transgenic lines (Fig. 8C, 9C).

Discussion

As one of the largest TF families in flowering plants, the *WRKY* gene family participates in plant development as well as in response to abiotic and biotic stresses. Based on the *S. album* genome and transcriptome data, 64 *SaWRKY* genes were identified in our work, accounting for 100% of the *S. album* *WRKY* family genes that were determined by Mahesh et al. [30] using the sandalwood genome. A total of 5 *SaWRKY* genes (*SaWRKY1*, 19, 43, 45, and 64) were only found in the genome data but lacked transcriptome data sets, either because these genes might have a lower level of expression in some tissues while others might be pseudogenes. Three *SaWRKY* genes (*SaWRKY8*, 29 and 41) was only found in transcriptome data, which may be caused by incomplete genome assembling. The 64 *SaWRKY* proteins identified could be divided into three groups based on a phylogenetic analysis for *A. thaliana* and *S. album*. Group II, which contained 43 *SaWRKY* proteins, was further divided into five subgroups (IIa, IIb, IIc, IId, and IIe), accounting for the largest proportion (67%) among all the groups. Other plant species showed a comparable proportion, namely 58% in *A. thaliana* [6], 64% in *Caragana intermedia* [31], and 66% in cassava (*Manihot esculenta*) [32]. Chimeric proteins such as R protein-WRKY (RW) genes are a feature of some *WRKY* family genes in switchgrass (*Panicum virgatum*), sorghum (*Sorghum bicolor*) and rice (*Oryza sativa*) [11]. We searched for 64 *SaWRKY* proteins, and no other domains except for WDs and zinc-finger motifs were found. This suggest that RW genes may be a characteristic of some monocot genomes.

Publicly available transcriptome datasets were processed to obtain the RPKM expression values for the 41 *SaWRKY* genes in five different tissues of *S. album*. The heat map (Fig. 4) generated using these expression values exhibited different expression patterns for all the identified *SaWRKY* genes except for *SaWRKY1* in the five tissues, indicating that they may perform diversified functions in plant development or in response to stress. *SaWRKY12*, 30, 31, 39, and 42 shared similarly high expression levels in all the tested tissues (Fig. 4), indicating that these genes may play essential roles in *S. album* tissue development. In contrast, four *SaWRKY* genes (*SaWRKY6*, 25, 35, and 37) showed a low level of expression in all five tissues and may thus have a limited function in their development. Several *SaWRKY* genes that were identified showed higher tissue-specific expression patterns, i.e., *SaWRKY2*, 15, 20, 22, 27, 32 and 44, which were preferentially expressed in wood tissue (sapwood, transition zone and heartwood) (Fig. 4). In particular, four of these genes (*SaWRKY16*, 20, 22, and 32) accumulated obviously higher levels of mRNA in the transition zone, which is the main location of the synthesis of valuable sandal essential oil [21]. For now, it can be speculated that several of these genes may be involved in essential

oil biosynthesis, although this will require extensive additional analyses. In addition, RT-qPCR was performed to detect the levels of *SaWRKY1* expression in roots, stems (wood tissue), leaves and callus (Fig. 6). *SaWRKY1* mRNA mainly accumulated in the leaves, roots and callus, showing the highest expression level in fast-growing tissue (i.e., callus), similar to apple [34].

Due to their sessile lifestyle, plants must adjust to various biotic and abiotic stresses [33]. Phytohormones play a central role in a plant's ability to integrate various environmental stimuli or cues with the genetic program in their life cycles [34]. Among several phytohormones, JA and SA play crucial roles in orchestrating different stress responses within a plant, such as in pathogen defense [35], or protection against abiotic stresses (drought, low temperature, and salinity) [36]. Increasing evidence shows that the expression levels of plant *WRKY* genes are regulated by different plant hormones, such as 37 *VvWRKY* genes that were induced by SA, and 48 *VvWRKY* genes that were regulated following JA treatment in grape (*Vitis vinifera*) [37], or 55 *PtrWRKY* genes that were induced by SA and MeJA treatment in poplar (*Populus trichocarpa* and *P. tomentosa*) [9]. It is difficult to conduct basic research in woody plants because of their slow growth, so callus is a useful and efficient system to conduct gene function research on plants such as oil palm (*Elaeis guineensis*) [38] and apple (*Malus x domestica*) [39]. In order to detect the expression levels of the *SaWRKY* genes in response to SA and MeJA, RT-qPCR was performed in callus generated from newly sprouted *S. album* shoots. RT-qPCR results showed nine up-regulated and 25 down-regulated *SaWRKY* genes after SA treatment (Fig. 5A, Additional file 2: Table S2), and 12 up-regulated and seven down-regulated *SaWRKY* genes after JA treatment (Fig. 5B, Additional file 3: Table S3). This indicates that these genes may be involved in biotic and abiotic stress responses mediated by SA- and/or JA-dependent defense signaling pathways. *AtWRKY53*, a paralogous gene of *SaWRKY7* and *SaWRKY28* in *S. album*, was significantly induced in five- and seven-week-old *A. thaliana* plants sprayed with 2 mM SA [40]. *AtWRKY53* expression was up-regulated after pathogen attack [41], and this gene participates in the regulation of senescence, being positively regulated by SA signaling and negatively regulated by JA signaling [40, 42], and negatively regulates drought tolerance via controlled stomatal movement independent of abscisic acid levels [43]. It can thus be inferred that paralogous genes *SaWRKY7* and *SaWRKY28* may perform a similar function in *S. album*. A member of subgroup IIc, *SaWRKY16*, which showed the highest level of expression in the transition zone of wood tissue, was significantly induced by JA (Fig. 5B, Additional file 3: Table S3). Its paralogous gene, *AtWRKY75*, played a positive role in the defense of *A. thaliana* against pathogens (including *Pectobacterium carotovorum* ssp. *carotovorum* and *Sclerotinia sclerotiorum*) and oxalic acid stress, and was mainly dependent on the JA/ethylene pathway [44, 45]. Three *SaWRKY* genes (*SaWRKY25*, 28 and 37) were up-regulated by SA but down-regulated by JA (Fig. 5A, B; Additional file 2, 3: Table S2, S3), consistent with some reports showing that SA and JA function antagonistically [46-48]. Conversely, four *SaWRKY* genes, including *SaWRKY1*, 3, 11 and 38, were up-regulated when stimulated by SA and JA. *SaWRKY1* was one of the genes induced both by SA and JA treatment, and its overexpression strengthened salt tolerance and increased the content of chlorophyll *a*, *b*, and *a + b* in transgenic *A. thaliana* plants compared to the wild type. Plants often enhance the content of non-enzymatic compounds such as chlorophyll to improve tolerance to salt stress, and a rich literature on this topic illustrates that SA and JA perform important roles in these

processes. For example, strawberry (*Fragaria × ananassa*) plants treated with 1.0 mM SA had a higher chlorophyll content and showed better growth under salt stress than unstressed plants [49]. The application of 0.01 mM SA to tomato (*Solanum lycopersicum*) plants caused the accumulation of photosynthetic pigments and soluble sugars, and enhanced plant growth and development under salt stress [50]. Foliar application of 2.0 mM JA to wheat seedlings enhanced their salt stress tolerance and growth by increasing the activity of antioxidant enzymes and the content of glutathione, chlorophyll *b* and carotenoids [51]. Cumulatively, these findings coincide with our results and suggest that *SaWRKY1* improves plant salt tolerance by increasing the content of [photosynthetic](#) pigments, probably mediated by SA and/or JA. However, the accurate regulatory mechanism still needs to be studied in further detail. In *A. thaliana*, the paralogous genes of *SaWRKY1*, *SaWRKY3* and *SaWRKY49*, namely *AtWRKY18*, *AtWRKY40* and *AtWRKY60*, are evolutionarily closely related WRKY domain TFs. Based on multiple alignments of amino acid sequences (Additional file 4: Figure S1), there was 36.02%, 40.82% and 33.87% identity between *SaWRKY1* and *AtWRKY18*, *40*, and *60*, respectively. *SaWRKY1* has the closest evolutionary relationship and highest amino acid sequence identity with *AtWRKY40*, which is also involved in salt stress [52], suggesting that these paralogous genes play similar roles in plants against abiotic stress and confirming that *SaWRKY1* plays a regulatory role in salt resistance. Thus, *SaWRKY1* may be useful in engineering salinity tolerance of *S. album* trees via *Agrobacterium*- or gene gun-mediated transformation of callus. As two members of subgroup IIa, *SaWRKY3* and *SaWRKY49* might also be involved in SA- and JA-mediated pathogen and salt stress responses, but the putative function needs to be further investigated.

Conclusions

In this study, we identified 64 *SaWRKY* genes in Indian sandalwood based on genome and transcriptome data. The phylogenetic relationships, composition of conserved motifs and tissue expression patterns of these genes were analyzed. Callus was used to efficiently detect the expression profiles of 42 *SaWRKY* genes under different phytohormone treatments, suggesting that 34 and 19 *SaWRKY* genes are involved in SA- and MeJA-mediated abiotic or biotic response pathways, respectively. In addition, functional analysis of *SaWRKY1* under salt stress increased our understanding of the roles that *SaWRKY1* play in abiotic stress.

Abbreviations

2,4-D: 2,4-dichlorophenoxyacetic acid; chlo: chloroplast; cyto: cytoplasm; ER: [endoplasmic reticulum](#); golg: golgi apparatus; MeJA: methyl jasmonate; mito: mitochondrion; nucl: nucleus; plas: plastid; RACE: rapid amplification of cDNA ends; RPKM: reads per kilobase of exon model per million mapped reads; RT-qPCR: reverse transcription quantitative polymerase chain reaction; SA: salicylic acid; TDZ: thidiazuron; TF: transcription factor.

Declarations

Ethics approval and consent to participate

The *S. album* plants used in this study were from the *S. album* plantation base of South China Botanical Garden cultivated by Guohua Ma, in Guangzhou, Guangdong, China, and no permissions were needed to obtain the material. Our study fully complies with institutional regulations.

Consent for publication

Not applicable.

Availability of data and materials

All data generated and analyzed during this study were included in this published article and the additional files.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was financially supported by the National Natural Science Foundation of China (grant numbers 31470685, 31270720, and 31100498), the Natural Science Foundation of Guangdong Province (S2012010009025), and the Guangdong Science and Technology project (2015B020231008). The funding agencies had no role in the design, analysis, and interpretation of the data, or writing of the manuscript.

Authors' contributions

GM conceived and designed the experiment. HY, YZ, and YX performed the experiments. HL, MN, and BG contributed materials. ML performed bioinformatics analyses. HY, YL, XZ and JATS wrote the manuscript. JATS provided critical analysis of the design and data. All authors read and approved the final manuscript.

References

1. Riechmann JL HJ, Martin G, Reuber L, Jiang C-Z, Keddie J, Adam L, Pineda O, Ratcliffe OJ, Samaha RR, Creelman R, Pilgrim M, Broun P, Zhang JZ, Ghandehari D, Sherman BK, Yu G-L. *Arabidopsis* transcription factors: genome-wide comparative analysis among eukaryotes. *Science*. 2000; 290:2105-2110.
2. Ulker B, Somssich IE. WRKY transcription factors: from DNA binding towards biological function. *Curr Opin Plant Biol*. 2004; 7(5):491-498.
3. Pan YJ, Cho CC, Kao YY, Sun CH. A novel WRKY-like protein involved in transcriptional activation of cyst wall protein genes in *Giardia lamblia*. *J Biol Chem*. 2009; 284(27):17975-17988.
4. Jin J, Tian F, Yang DC, Meng YQ, Kong L, Luo J, Gao G. PlantTFDB 4.0: toward a central hub for transcription factors and regulatory interactions in plants. *Nucleic Acids Res*. 2017; 45(D1):D1040-D1045.
5. Ishiguro S, Nakamura K. Characterization of a cDNA encoding a novel DNA-binding protein, SPF1, that recognizes SP8 sequences in the 5' upstream regions of genes coding for sporamin and β -amylase from sweet potato. *Mol Gen Genet*. 1994; 244(6):563-571.
6. Eulgem T RP, Robatzek S, Somssich IE. The WRKY superfamily of plant transcription factors. *Trends Plant Sci*. 2000; 5(5).
7. Ross CA, Liu Y, Shen QXJ. The WRKY gene family in rice (*Oryza sativa*). *J Integr Plant Biol*. 2007; 49(6):827-842.
8. Schmutz J, Cannon SB, Schlueter J, Ma J, Mitros T, Nelson W, Hyten DL, Song Q, Thelen JJ, Cheng J et al. Genome sequence of the palaeopolyploid soybean. *Nature*. 2010; 463(7278):178-183.
9. Jiang Y, Duan Y, Yin J, Ye S, Zhu J, Zhang F, Lu W, Fan D, Luo K. Genome-wide identification and characterization of the *Populus* WRKY transcription factor family and analysis of their expression in response to biotic and abiotic stresses. *J Exp Bot*. 2014; 65(22):6629-6644.
10. Rushton PJ, Somssich IE, Ringler P, Shen QJ. WRKY transcription factors. *Trends Plant Sci*. 2010; 15(5):247-258.
11. Rinerson CI, Scully ED, Palmer NA, Donze-Reiner T, Rabara RC, Tripathi P, Shen QJ, Sattler SE, Rohila JS, Sarath G et al. The WRKY transcription factor family and senescence in switchgrass. *BMC Genomics*. 2015; 16:912.
12. Lai Z, Vinod K, Zheng Z, Fan B, Chen Z. Roles of *Arabidopsis* WRKY3 and WRKY4 transcription factors in plant responses to pathogens. *BMC Plant Biol*. 2008; 8:68.
13. Kim KC, Lai Z, Fan B, Chen Z. *Arabidopsis* WRKY38 and WRKY62 transcription factors interact with histone deacetylase 19 in basal defense. *Plant Cell*. 2008; 20(9):2357-2371.
14. Liu XQ, Bai XQ, Qian Q, Wang XJ, Chen MS, and Chu CC. OsWRKY03, a rice transcriptional activator that functions in defense signaling pathway upstream of OsNPR1. *Cell Res*. 15(8):593-603.
15. Ramamoorthy R, Jiang SY, Kumar N, Venkatesh PN, Ramachandran S. A comprehensive transcriptional profiling of the *WRKY* gene family in rice under various abiotic and phytohormone treatments. *Plant Cell Physiol*. 2008; 49(6):865-879.

16. Chen L, Song Y, Li S, Zhang L, Zou C, Yu D. The role of WRKY transcription factors in plant abiotic stresses. *Biochim Biophys Acta*. 2012; 1819(2):120-128.
17. Ding ZJ, Yan JY, Li GX, Wu ZC, Zhang SQ, Zheng SJ. WRKY41 controls *Arabidopsis* seed dormancy via direct regulation of *ABI3* transcript levels not downstream of ABA. *Plant J*. 2014; 79(5):810-823.
18. Guan Y, Meng X, Khanna R, LaMontagne E, Liu Y, Zhang S. Phosphorylation of a WRKY transcription factor by MAPKs is required for pollen development and function in *Arabidopsis*. *PLoS Genet*. 2014; 10(5):e1004384.
19. Devaiah BN, Karthikeyan AS, Raghothama KG. WRKY75 transcription factor is a modulator of phosphate acquisition and root development in *Arabidopsis*. *Plant Physiol*. 2007; 143(4):1789-1801.
20. Jiang Y, Liang G, Yang S, Yu D. *Arabidopsis* WRKY57 functions as a node of convergence for jasmonic acid- and auxin-mediated signaling in jasmonic acid-induced leaf senescence. *Plant Cell*. 2014; 26(1):230-245.
21. Jones CG, Keeling CI, Ghisalberti EL, Barbour EL, Plummer JA, Bohlmann J. Isolation of cDNAs and functional characterisation of two multi-product terpene synthase enzymes from sandalwood, *Santalum album* L. *Arch Biochem Biophys*. 2008; 477(1):121-130.
22. Kim TH, Ito H, Hayashi K, Hasegawa T, Machiguchi T, Yoshida T. Aromatic constituents from the heartwood of *Santalum album* L. *Chem Pharm Bull*. 2005; 53(6):641-644.
23. Baldovini N, Delasalle C, Joulain D. Phytochemistry of the heartwood from fragrant *Santalum* species: a review. *Flavour Fragr J*. 2011; 26(1):7-26.
24. Rashkow ED. Perfumed the axe that laid it low: The endangerment of sandalwood in southern India. *Indian Econ. Soc. Hist. Rev*. 2014; 51(1):41-70.
25. Teixeira da Silva JA, Kher MM, Soner D, Page T, Zhang X, Nataraj M, Ma G. Sandalwood: basic biology, tissue culture, and genetic transformation. *Planta*. 2016; 243(4):847-887.
26. Zhang X, Teixeira da Silva JA, Niu M, Li M, He C, Zhao J, Zeng S, Duan J, Ma G. Physiological and transcriptomic analyses reveal a response mechanism to cold stress in *Santalum album* L. leaves. *Sci Rep*. 2017; 7:42165.
27. Teixeira da Silva JA, Kher MM, Soner D, Nataraj M, Dobránszki J, Millar MA. *Santalum* molecular biology: molecular markers for genetic diversity, phylogenetics and taxonomy, and genetic transformation. *Agroforest Syst*. 2017; 92(5):1301-1315.
28. Zhang X, Berkowitz O, Teixeira da Silva JA, Zhang M, Ma G, Whelan J, Duan J. RNA-Seq analysis identifies key genes associated with haustorial development in the root hemiparasite *Santalum album*. *Front Plant Sci*. 2015; 6:661.
29. Celedon JM, Chiang A, Yuen MM, Diaz-Chavez ML, Madilao LL, Finnegan PM, Barbour EL, Bohlmann J. Heartwood-specific transcriptome and metabolite signatures of tropical sandalwood (*Santalum album*) reveal the final step of (Z)-santalol fragrance biosynthesis. *Plant J*. 2016; 86(4):289-299.
30. Mahesh HB, Subba P, Advani J, Shirke MD, Loganathan RM, Chandana SL, Shilpa S, Chatterjee O, Pinto SM, Prasad TSK et al. Multi-omics driven assembly and annotation of the sandalwood (*Santalum album*) genome. *Plant Physiol*. 2018; 176(4):2772-2788.

31. Wan Y, Mao M, Wan D, Yang Q, Yang F, Mandlaa, Li G, Wang R. Identification of the *WRKY* gene family and functional analysis of two genes in *Caragana intermedia*. BMC Plant Biol. 2018; 18(1):31.
32. Wei Y, Shi H, Xia Z, Tie W, Ding Z, Yan Y, Wang W, Hu W, Li K. Genome-wide identification and expression analysis of the *WRKY* gene family in cassava. Front Plant Sci. 2016; 7:25.
33. Ahuja I, de Vos RC, Bones AM, Hall RD. Plant molecular stress responses face climate change. Trends Plant Sci. 2010; 15(12):664-674.
34. Santner A, Estelle M. Recent advances and emerging trends in plant hormone signalling. Nature. 2009; 459(7250):1071-1078.
35. Halim VA, Vess A, Scheel D, Rosahl S. The role of salicylic acid and jasmonic acid in pathogen defence. Plant Biol (Stuttg). 2006; 8(3):307-313.
36. Miura K, Tada Y. Regulation of water, salinity, and cold stress responses by salicylic acid. Front Plant Sci. 2014; 5:4.
37. Guo C, Guo R, Xu X, Gao M, Li X, Song J, Zheng Y, Wang X. Evolution and expression analysis of the grape (*Vitis vinifera* L.) *WRKY* gene family. J Exp Bot. 2014; 65(6):1513-1528.
38. Low ET, Alias H, Boon SH, Shariff EM, Tan CY, Ooi LC, Cheah SC, Raha AR, Wan KL, Singh R. Oil palm (*Elaeis guineensis* Jacq.) tissue culture ESTs: identifying genes associated with callogenesis and embryogenesis. BMC Plant Biol. 2008; 8:62.
39. Dong QL, Liu DD, An XH, Hu DG, Yao YX, Hao YJ. *MdVHP1* encodes an apple vacuolar H⁺-PPase and enhances stress tolerance in transgenic apple callus and tomato. J Plant Physiol. 2011; 168(17):2124-2133.
40. Miao Y, Zentgraf U. The antagonist function of *Arabidopsis* *WRKY53* and *ESR/ESP* in leaf senescence is modulated by the jasmonic and salicylic acid equilibrium. Plant Cell. 2007; 19(3):819-830.
41. Murray SL, Ingle RA, Petersen LN, Denby KJ. Basal resistance against *pseudomonas syringae* in *Arabidopsis* involves *WRKY53* and a protein with homology to a nematode resistance protein. Mol Plant-Microbe Interact. 2007; 20(11):1431-1438.
42. Zentgraf U, Laun T, Miao Y. The complex regulation of *WRKY53* during leaf senescence of *Arabidopsis thaliana*. Eur J Cell Biol. 2010; 89(2-3):133-137.
43. Sun Y, Yu D. Activated expression of *AtWRKY53* negatively regulates drought tolerance by mediating stomatal movement. Plant Cell Rep. 2015; 34(8):1295-1306.
44. Chen X, Liu J, Lin G, Wang A, Wang Z, Lu G. Overexpression of *AtWRKY28* and *AtWRKY75* in *Arabidopsis* enhances resistance to oxalic acid and *Sclerotinia sclerotiorum*. Plant Cell Rep. 2013; 32(10):1589-1599.
45. Choi C, Park YH, Kwon SI, Yun C, Ahn I, Park SR, Hwang D-J. Identification of *AtWRKY75* as a transcriptional regulator in the defense response to *Pcc* through the screening of *Arabidopsis* activation-tagged lines. Plant Biotechnol Rep. 2013; 8(2):183-192.

46. Kunkel BN, Brooks DM. Cross talk between signaling pathways in pathogen defense. *Curr Opin Plant Biol.* 2002; 5(4):325-331.
47. Mur LA, Kenton P, Atzorn R, Miersch O, Wasternack C. The outcomes of concentration-specific interactions between salicylate and jasmonate signaling include synergy, antagonism, and oxidative stress leading to cell death. *Plant Physiol.* 2006; 140(1):249-262.
48. Mur LA, Prats E, Pierre S, Hall MA, Hebelstrup KH. Integrating nitric oxide into salicylic acid and jasmonic acid/ ethylene plant defense pathways. *Front Plant Sci.* 2013; 4:215.
49. Karlidag H, Yildirim E, and Turan M. Salicylic acid ameliorates the adverse effect of salt stress on strawberry. *Sci Agric.* 2009; 66(2):180-187.
50. Wasti S, Mimouni H, Smiti S, Zid E, Ben Ahmed H. Enhanced salt tolerance of tomatoes by exogenous salicylic acid applied through rooting medium. *OMICS.* 2012; 16(4):200-207.
51. Qiu Z, Guo J, Zhu A, Zhang L, Zhang M. Exogenous jasmonic acid can enhance tolerance of wheat seedlings to salt stress. *Ecotoxicol Environ Saf.* 2014; 104:202-208.
52. Chen H, Lai Z, Shi J, Xiao Y, Chen Z, and Xu X. Roles of arabidopsis WRKY18, WRKY40 and WRKY60 transcription factors in plant responses to abscisic acid and abiotic stress. *BMC Plant Biol.* 2010; 10(1):281.
53. Singh CK, Raj SR, Jaiswal PS, Patil VR, Punwar BS, Chavda JC, Subhash N. Effect of plant growth regulators on *in vitro* plant regeneration of sandalwood (*Santalum album* L.) via organogenesis. *Agroforest Syst.* 2015; 90(2):281-288.
54. Murashige T, Skoog F. A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol Plant.* 1962; 15:473-497.
55. He C, Teixeira da Silva JA, Tan J, Zhang J, Pan X, Li M, Luo J, Duan J. A genome-wide identification of the WRKY family genes and a survey of potential WRKY target genes in *Dendrobium officinale*. *Sci Rep.* 2017; 7(1):9200.
56. Kolosova N, Miller B, Ralph S, Ellis BE, Douglas C, Ritland K, Bohlmann J. Isolation of high-quality RNA from gymnosperm and angiosperm trees. *BioTechniques.* 2004; 36(5):821-824.
57. Yoo SD, Cho YH, Sheen J. *Arabidopsis* mesophyll protoplasts: a versatile cell system for transient gene expression analysis. *Nat Protoc.* 2007; 2(7):1565-1572.
58. Weigel D, Glazebrook J. Transformation of *Agrobacterium* using the freeze-thaw method. *CSH Protoc.* 2006; 7.
59. Zhang X, Henriques R, Lin SS, Niu QW, Chua NH. *Agrobacterium*-mediated transformation of *Arabidopsis thaliana* using the floral dip method. *Nat Protoc.* 2006; 1(2):641-646.
60. Porra RJ, Thompson WA, Kriedemann PE. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls *a* and *b* extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. *Biochim Biophys Acta.* 1989; 975(3):384-394.

Tables

Table 1. List of *SaWRKY* genes identified in *Santalum album*.

Gene name	ORF (aa)	Conserved motif	Zinc-finger type	Subcellular location	Group
SaWRKY5	549	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH/	Nucleus	I
SaWRKY8	273	WRKYGQK	C-X ₄ -C-X ₂₂ -HXH	Golgi	I
SaWRKY12	764	WRKYGQK/WRKYGQK	C-X ₅ -C-X ₂₃ -HXT/C-X ₄ -C-X ₂₂ -HXH/ C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY18	700	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY20	507	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY25	342	WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH/ C-X ₇ -C-X ₂₃ -HXL	Nucleus	I
SaWRKY30	427	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY31	443	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY34	580	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY36	440	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY38	511	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY39	521	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY41	621	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₅ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY44	627	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY55	577	WRKYGQK/WRKYGQK	C-X ₄ -C-X ₂₂ -HXH/C-X ₄ -C-X ₂₃ -HXH	Nucleus	I
SaWRKY1	333	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIa
SaWRKY3	320	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIa
SaWRKY49	369	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Chloroplast	IIa
SaWRKY10	555	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIb
SaWRKY11	331	WRKYGQK	C-X ₆ -CX ₂₃ -HXH	Nucleus	IIb
SaWRKY17	639	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIb
SaWRKY24	545	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIb
SaWRKY35	435	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIb
SaWRKY40	631	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIb
SaWRKY2	179	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY4	329	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY13	295	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY15	306	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY16	199	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY19	296	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY23	234	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY27	266	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY29	181	WRKYGKK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY32	311	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY33	343	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY37	294	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc

Table 1. Continued

Gene name	ORF (aa)	Conserved motif	Zinc-finger type	Subcellular location	Group
SaWRKY45	121	WRKYGHK	C-X ₄ -C-X ₂₃ -HXY	cytoplasm	IIc
SaWRKY46	177	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	peroxisome	IIc
SaWRKY54	589	WRKYGKK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY62	341	WRKYGQK	C-X ₄ -C-X ₂₃ -HXH	Nucleus	IIc
SaWRKY22	356	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY42	335	WRKYGQK	C-X ₄ -C-X ₂₂ -HXV/C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY43	310	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY50	375	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY52	347	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY59	314	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY60	314	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY61	315	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY64	336	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IId
SaWRKY6	394	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY14	282	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY21	316	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY26	300	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY48	242	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY51	369	WRKYGQK	C-X ₃ -C-X ₂₂ -HXL/C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY56	422	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY58	319	WRKYGQK	C-X ₅ -C-X ₂₂ -HXS	Nucleus	IIe
SaWRKY63	334	WRKYGQK	C-X ₅ -C-X ₂₃ -HXH	Nucleus	IIe
SaWRKY7	367	WRKYGQK	C-X ₇ -C-X ₂₃ -HXC	Nucleus	III
SaWRKY9	357	WRKYGQK	C-X ₇ -C-X ₂₃ -HXC	Nucleus	III
SaWRKY28	366	WRKYGQK	C-X ₇ -C-X ₂₃ -HXC	Nucleus	III
SaWRKY47	290	WRKYGQK	C-X ₇ -C-X ₂₃ -HXC	Nucleus	III
SaWRKY53	362	WRKYGQK	C-X ₇ -C-X ₂₂ -HXT	Nucleus	III
SaWRKY57	330	WRKYGQK	C-X ₇ -C-X ₂₃ -HXC	Nucleus	III

Figures

Top20 of pathway enrichment

Figure 3

The 20 top Kyoto Encyclopedia of Genes and Genomes (KEGG) enrichment pathways of potential SaWRKY target genes with at least five W-boxes

Figure 4

The expression patterns of 41 SaWRKY genes in five tissues of *Santalum album*. The heat map was generated based on the number of reads per kilobase of exon modelled per million mapped reads (RPKM) of transcriptome data

Figure 5

Expression profiles of SaWRKY genes following SA and MeJA treatments. The expression of 42 SaWRKY genes was determined under SA (A) and MeJA (B) stimuli at three time points (0, 3, and 6 h) in callus. The heat map was generated based on log2-transformed count value from RT-qPCR data using R language. RT-qPCR data were collected from three biological replicates

Figure 6

Expression of *SaWRKY1* in different tissues of *Santalum album*. Relative expression levels of *SaWRKY1* in four tissues were determined by RT-qPCR with three biological replicates. Values shown are means $\pm$ SE. Asterisk (*) indicates significance at $P < 0.05$ using Duncan's multiple range test

Figure 7

Subcellular localization assay of the SaWRKY1 protein

Figure 8

Phenotypes of two lines of overexpression SaWRKY1 transgenic plants under salt stress. (A) Day 0. (B) Day 3 after irrigation with 300 mM NaCl. (C) Day 5 after second irrigation with 300 mM NaCl

Figure 9

Chlorophyll content of two lines of overexpression SaWRKY1 transgenic plants under salt stress. All values are means $\pm$ SE from at least three independent biological replicates (8 seedlings per replicate). Double asterisk (**) indicates significance at $P < 0.01$ for each component between wild type and transgenic lines using Duncan's multiple range test. (A) Day 0. (B) Day 3 after irrigation with 300 mM NaCl. (C) Day 5 after second irrigation with 300 mM NaCl

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1S.albumWRKYTFsfinal2.docx](#)