

Bacterial but no SARS-CoV-2 contamination after terminal disinfection of tertiary care intensive care units treating COVID-19 patients

Daniel A Hofmänner

UniversitätsSpital Zurich

Pedro D Wendel Garcia

UniversitätsSpital Zurich

Branko Duvnjak

UniversitätsSpital Zurich

Bhavya Chakrakodi

UniversitätsSpital Zurich

Julian D Maier

UniversitätsSpital Zurich

Michael Huber

Universität Zurich

Jon Huder

Universität Zurich

Aline Wolfensberger

UniversitätsSpital Zurich

Peter W Schreiber

UniversitätsSpital Zurich

Reto A. Schüpbach

UniversitätsSpital Zurich

Annelies S Zinkernagel

UniversitätsSpital Zurich

Philipp K Bühler

UniversitätsSpital Zurich

Silvio D Brugger (✉ silvio.brugger@usz.ch)

UniversitätsSpital Zurich <https://orcid.org/0000-0001-9492-9088>

Research

Keywords: SARS-CoV-2, disinfection, hydrogen peroxide nebulisation, ICU, hospital epidemiology, nosocomial infection

Posted Date: July 28th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-47872/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on January 12th, 2021. See the published version at <https://doi.org/10.1186/s13756-021-00885-z>.

Abstract

Background: In intensive care units (ICUs) treating patients with Coronavirus disease 2019 (COVID-19) invasive ventilation poses a high risk for aerosol and droplet formation. Surface contamination of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2) or bacteria can result in nosocomial transmission.

Methods: Two tertiary care COVID-19 intensive care units treating 53 patients for 870 patient days were sampled after terminal cleaning and preparation for regular use to treat non-COVID-19 patients.

Results: A total of 176 swabs were sampled of defined locations covering both ICUs. No SARS-CoV-2 ribonucleic acid (RNA) was detected. Gram-negative bacterial contamination was mainly linked to sinks and siphons. Skin flora was isolated from most swabbed areas and *Enterococcus faecium* was detected on two keyboards.

Conclusions: After basic cleaning with standard disinfection measures no remaining SARS-CoV-2 RNA was detected. Bacterial contamination was low and mainly localised in sinks and siphons.

Background

During the last months, the novel Coronavirus disease (COVID-19) pandemic has challenged many health-care systems worldwide. In line with the growing knowledge about specific pathophysiologic mechanisms, COVID-19 has increasingly been associated with bacterial co-infections, exhibiting high rates in intensive care units (ICUs) (1–4). It is estimated that around 50% of fatalities associated with COVID-19 had secondary infections contributing to the disease course with a relevant impact on patient outcomes (5, 6). Treating critically ill COVID-19 patients often involves mechanical ventilation, which poses a high risk for droplet and aerosol transmission. In addition to the risk of severe acute respiratory syndrome Coronavirus 2 (SARS-CoV-2) contamination of surfaces and medical equipment, bacterial pathogens complicating the disease course can be transmitted to the environment surrounding the patient. Equipment and surfaces (e.g. respirators, beds) are well known for contributing to the transmission of bacterial pathogens promoting nosocomial infections (7, 8).

With the decline of the current COVID-19 outbreak in some countries, affected ICUs now returning to treating non-COVID-19 patients are being faced with the challenge of disinfecting their equipment and facilities in order to prevent hospital-acquired infections. Moreover, it is currently not well characterised whether or how long SARS-CoV-2 ribonucleic acid (RNA) persists on surfaces after complete disinfection of ICUs.

The aim of this study was to investigate the burden of pathogenic bacteria and SARS-CoV-2 RNA on different devices and surfaces after terminal cleaning measures in two specialised ICUs of a tertiary care hospital exclusively treating COVID-19 patients.

Methods

Study setting. This single-blinded study was conducted at the University Hospital Zurich, Switzerland, a tertiary care hospital with around 980 beds. During the COVID-19 pandemic, two defined subunits of the ICU were designated to exclusively treat COVID-19 patients. The first (ICU 1) had the capacity to treat 16 patients, the second (ICU 2) 12 patients. In total, ICU 1 treated 39 COVID-19 patients (661 patient days). ICU 2 treated 14 COVID-19 patients (209 patient days). No institutional review board (IRB) approval was required as this study did not involve any patients.

ICU cleaning/disinfection. After the first COVID-19 wave, both subunits were cleaned in order to provide care for non-COVID-19 patients. All surfaces, devices and floors on ICU 1 were disinfected by nurses and cleaning staff with aldehyde-based Kohrsolin® 1% (Wilbert Hygiene, Germany). In addition, a dry mist hydrogen peroxide (H₂O₂) nebulisation technique (Dosymist XL, SolidFog Technologies®, H₂O₂ à 12% Fisher Scientific®) was used on ICU 1 in the ICU pharmacy room. With the aid of the hydrogen peroxide technology, a dry aerosol is generated, with the oxidative action of hydroxyl radicals leading to antimicrobial properties. Quality control of the H₂O₂ disinfection was performed with chemical indicators (3M Healthcare®, successful decontamination indicated by color change on test kit) and biological indicators (*Geobacillus stearothermophilus*, Crosstex®, successful decontamination indicated by 6-log reduction between positive and negative control). ICU 2 was disinfected with Kohrsolin® 1% (Wilbert Hygiene, Germany) only. Cleaning and disinfection were not supervised by the study team.

Virus sampling and analysis. After the cleaning procedures, 86 viral swabs (FLOQSwabs, UTM-RT transport medium, Copan®) were taken from different locations on both ICUs (representing high-frequency touch surfaces, patient-surrounding areas, healthcare-worker changing areas and sinks including siphons) with a minimal swabbing surface dimension of 10 × 10 cm. Sampling objects were 30 monitors/keyboards (bedside), 8 tabletops to prepare drugs for application including adjacent sinks, 17 individual sinks including siphons in patient rooms, 6 floors of patient compartments and 1 main exit gate (floor), 13 respirator touch screens and handling buttons, 2 physician's work desks, 2 non-personalised telephones, 2 perfusors, 4 rolling trolleys/tables, 2 fridge doors in the ICU pharmacy, 1 chair and 2 surfaces in kitchens. Due to the different configurations and designs of the two ICU subunits, it was not possible to take an equal amount of swabs from both ICU subunits. Figure 1 demonstrates both ICU subunits and sampling points are indicated.

The swabbed area was individualised according to the swabbed object size and shape. Real-time reverse transcriptase polymerase chain reaction (rtPCR) was performed to detect SARS-CoV-2 RNA with inclusion of positive and negative control. PCR conditions were as previously described by Corman et al (9).

Bacterial sampling, culture, identification and resistance testing. In parallel to the viral sampling, a total of 90 moistured swabs (eSwabs™ comprising 1 millilitre of Amies medium) were taken from the same locations on both ICUs, analogous to the viral sampling (Fig. 1).

Dependent on the object size and shape, the swabbed area was individualised. In general, the largest possible area to swab was chosen.

All swabs were plated on 5% sheep blood agar and MacConkey agar plates in three fractions. Subsequently, the samples were incubated for 48 hours at 37 °C without carbon dioxide atmosphere. After incubation, colony forming units (CFU) were analysed semiquantitatively. Growth in the first fraction of the agar plate corresponded to 10^3 - 10^4 CFU/ml, in the first and second fraction to 10^4 - 10^5 CFU/ml and in all three fractions to 10^5 - 10^6 CFU/ml.

Subcultures were plated on 5% sheep blood agar or MacConkey agar and identification was done as previously described (10) with the exception that identification was done using both a BD Phoenix™ automated identification and susceptibility testing system as well as a Bruker MALDI-TOF-MS biotyper. Susceptibility testing was done using the BD Phoenix system.

Results

All of the 86 viral swabs sampled tested negative for SARS-CoV-2 RNA. Two samples from ICU 1 (sink in the pharmacy and one monitor/keyboard) were initially weakly positive but inconclusive. The quantification cycle values for those two swabs were 43.17 and 38.94 respectively. However, this finding could not be reproduced in a subsequent confirmatory test.

All chemical and biological indicators demonstrated successful (bacterial) decontamination of the ICU 1 pharmacy room. Tables 1 and 2 summarise cultivated bacteria on different surfaces of both subunits of the ICU after disinfection. In ICU 1, most bacteria were found in the sinks and associated siphons – in the individual sinks in patient rooms and at tabletops employed for drug preparation by health care staffing (Table 1). *Pseudomonas aeruginosa*, *Pseudomonas putida*, *Stenotrophomonas maltophilia*, *Klebsiella oxytoca* and *Enterobacter cloacae* were cultured from these areas. In line with ICU 1, *Pseudomonas aeruginosa*, *Pseudomonas putida*, *Stenotrophomonas maltophilia* and *Achromobacter* spp. were found in ICU 2 in the sinks (Table 2). No multidrug-resistant (MDR) *P. aeruginosa* was detected.

Table 1.

Bacterial contamination on ICU 1 according to swabbed objects and swab count. Numbers in () indicating the fraction of positive samples compared to the total number of swabs per object group.

	number of swabs from distinct objects (n=67)	<i>Pseudomonas</i> spp.	other detected bacteria
Bacterial contamination			
ICU 1			
Monitors/keyboards	18		coagulase negative staphylococci (10/18)
Tabletops, sinks and siphons in drug preparation areas	5	<i>Pseudomonas aeruginosa</i> 10 ⁵ -10 ⁶ /ml (2/5), <i>Pseudomonas putida</i> 10 ⁵ -10 ⁶ /ml (1/5)	<i>Stenotrophomonas maltophilia</i> 10 ⁵ -10 ⁶ /ml (3/5), <i>Enterobacter cloacae</i> 10 ⁵ -10 ⁶ /ml (1/5), coagulase negative staphylococci (1/5)
Individual sinks in patient area	12	<i>Pseudomonas aeruginosa</i> 10 ⁵ -10 ⁶ /ml (2/12), <i>Pseudomonas</i> spp. 10 ³ -10 ⁴ /ml (1/12)	<i>Klebsiella oxytoca</i> 10 ⁴ -10 ⁵ /ml (1/12), <i>Stenotrophomonas maltophilia</i> 10 ⁵ -10 ⁶ /ml (1/12), coagulase negative staphylococci (1/12)
Floors in patient compartment	6		coagulase negative staphylococci (3/6)
Respirators	13		coagulase negative staphylococci (1/13)
Work desk physicians	2		coagulase negative staphylococci (2/2)
Telephones non-personalized	2		coagulase negative staphylococci (1/2)
Perfusors	2		
Rolling trolleys/tables	4		coagulase negative staphylococci (2/4)
Fridges pharmacy	2		
Chair	1		

Table 2

Bacterial contamination on ICU 2 according to swabbed objects and swab count. Numbers in () indicating

	number of swabs from distinct objects (n = 67)	<i>Pseudomonas</i> spp.	other detected bacteria
Bacterial contamination			
ICU 1			
Monitors/keyboards	18		coagulase negative staphylococci (10/18)
Tabletops, sinks and siphons in drug preparation areas	5	<i>Pseudomonas aeruginosa</i> 10 ⁵ -10 ⁶ /ml (2/5), <i>Pseudomonas putida</i> 10 ⁵ -10 ⁶ /ml (1/5)	<i>Stenotrophomonas maltophilia</i> 10 ⁵ -10 ⁶ /ml (3/5), <i>Enterobacter cloacae</i> 10 ⁵ -10 ⁶ /ml (1/5), coagulase negative staphylococci (1/5)
Individual sinks in patient area	12	<i>Pseudomonas aeruginosa</i> 10 ⁵ -10 ⁶ /ml (2/12), <i>Pseudomonas</i> spp. 10 ³ -10 ⁴ /ml (1/12)	<i>Klebsiella oxytoca</i> 10 ⁴ -10 ⁵ /ml (1/12), <i>Stenotrophomonas maltophilia</i> 10 ⁵ -10 ⁶ /ml (1/12), coagulase negative staphylococci (1/12)
Floors in patient compartment	6		coagulase negative staphylococci (3/6)
Respirators	13		coagulase negative staphylococci (1/13)
Work desk physicians	2		coagulase negative staphylococci (2/2)
Telephones non-personalized	2		coagulase negative staphylococci (1/2)
Perfusors	2		
Rolling trolleys/tables	4		coagulase negative staphylococci (2/4)
Fridges pharmacy	2		
Chair	1		

the fraction of positive samples compared to the total number of swabs per object group.

	number of swabs from distinct objects (n = 67)	<i>Pseudomonas</i> spp.	other detected bacteria
	number of swabs from distinct objects (n = 23)	<i>Pseudomonas</i> spp.	other detected bacteria
Bacterial contamination			
ICU 2			
Monitors/keyboards	12		enterococcus faecium 10 ³ -10 ⁴ /ml (2/12), coagulase negative staphylococci (8/12)
Tabletops, sinks and siphons in drug preparation areas	3		coagulase negative staphylococci (3/3)
Individual sinks in patient area	5	<i>Pseudomonas aeruginosa</i> 10 ⁴ -10 ⁵ /ml (1/5), <i>Pseudomonas putida</i> 10 ⁴ -10 ⁵ /ml (1/5), <i>Pseudomonas</i> spp. 10 ⁴ -10 ⁵ /ml (2/5)	<i>Achromobacter</i> 10 ³ -10 ⁴ /ml (1/5), <i>Stenotrophomonas maltophilia</i> 10 ⁴ -10 ⁵ /ml (1/5), coagulase negative staphylococci (2/5)
Floor main exit gate	1		coagulase negative staphylococci (1/1)
Surface patient kitchen	2		coagulase negative staphylococci (1/2)
the fraction of positive samples compared to the total number of swabs per object group.			

Enterococcus faecium was cultivated in two out of 12 swabs from monitors/keyboards. On both ICUs, *coagulase-negative* Staphylococci were found in most swabbed areas but with low CFU numbers (< 10³ per ml corresponding to less than 5 CFU/cm² (Tables 1 and 2).

Discussion

The aim of study was to investigate the persistence of SARS-CoV-2 and bacteria on different devices and surfaces after disinfecting measures in two tertiary care ICU subunits that had exclusively treated COVID-19 patients. The main finding of this study is that no SARS-CoV-2 RNA was detected by rRT-PCR. However, pathogenic bacteria were abounded mainly in sinks and corresponding siphons of the ICUs and

enterococci were isolated from two monitors. Gram-positive bacteria were mainly isolated from surfaces (e.g. monitors/keyboards) whereas Gram-negative bacteria were mainly detected in plumbing units.

Up to date, it is not clearly established how long and to what extent SARS-CoV-2 RNA contaminates environmental surfaces and devices in hospitals. A recent study has described a low percentage of positive samples in an emergency room and the sub-intensive care ward (11). In another study, almost 57% of rooms treating COVID-19 patients had at least one environmental surface contaminated (12).

In our study, viral RNA was not detectable on any of the swabbed locations of the two ICU subunits, which suggests that conventional disinfection measures are likely sufficient to prevent further spread of the virus. Especially, since viruses likely lose infectivity before their RNA cannot be amplified by PCR. This finding thus might contribute to infection and prevention policies.

Already back in 1974, The Lancet published an article describing isolation of *Pseudomonas aeruginosa* in hospital sinks (13). Since then, various studies have investigated hospital water supply systems as potential reservoirs for the transmission of pathogenic bacteria in hospitals (14–18).

In line with previous studies (15, 16, 18), we found *Pseudomonas aeruginosa* in sinks/siphons in both ICUs. Specifically regarding the ICU setting, Zhou et al demonstrated that sink traps may act as an important source of *Pseudomonas* spp. leading to colonisation or infection of critically ill patients (18). Additionally, stagnant water in general bears a risk for being a reservoir and source of outbreaks with several pathogens (19). This finding highlights the need for constant awareness of possible bacterial reservoirs and the need for further refining cleaning methods in the ICU.

In this line of research it has been suggested that devices applying heat and electromechanical vibration to siphons lead to lower colonisation of patients with multidrug-resistant (MDR) *Pseudomonas* spp. (16) and that self-disinfecting sinks reduced *Pseudomonas* spp. bioburden in a pediatric ICU (20), offering possibilities to lower transmission of MDR pathogens. Whether such measures prove feasible and effective in other ICU settings remains to be elucidated in further studies and is beyond the scope of this paper. However, no MDR *Pseudomonas aeruginosa* was detected in our ICUs despite the fact that several patients on the ICU had been colonised with MDR *Pseudomonas aeruginosa*.

Further cultured bacteria associated with moist environments were *Stenotrophomonas maltophilia*, *Klebsiella oxytoca* and *Achromobacter* spp., all Gram-negative, potentially harmful bacteria. *Stenotrophomonas maltophilia* is known to cause hospital-acquired infections and is associated with multidrug resistance (21) (22). *Klebsiella* spp. can be responsible for infectious complications, especially hospital-acquired pneumonias, urinary tract infections or septicemias (23). Further, *Achromobacter* spp. have been linked to increased risk of morbidity and mortality in critically ill patients in a study reporting a 18 month lasting epidemic in an ICU (24). Additionally, *Enterobacter cloacae* was cultured at sinks located next to tabletops employed for drug preparation in ICU 1. *Enterobacteriaceae* (especially carbapenem-resistant strains) are an important cause of healthcare-associated infections and transmission by means of environmental reservoirs, such as therapeutic beds, has been described (25)

(26). The colonisation of critically ill patients with carbapenemase-producing *Enterobacteriaceae* has been associated with an increased length of stay, as well as with a 1.8 higher death hazard as opposed to non-colonised ICU patients (27).

Such findings mandate thorough disinfection of ICUs and surveillance of sinks and siphons as a source for further outbreaks. Water-free ICUs might reduce the risk of nosocomial transmission from bacteria residing in sinks and siphons.

Monitors and keyboards are frequently used devices in the everyday care on the ICU. We found *Enterococcus faecium* on two out of 12 monitors/keyboards in ICU 2, a pathogen known to cause hospital-acquired infections and possibly exhibiting complex resistance patterns, e.g. vancomycin-resistant strains. The failure of disinfection in this case is most likely because of the applied manual disinfection, which is prone to failure. However, we were not able to assess the individual cleaning performance. In a large multicenter, randomised trial, a cleaning bundle (REACH) showed promising results in reducing especially enterococcal infections (28). The implementation of such bundles in ICUs might probably lead to less nosocomial infections in critically ill patients and warrants investigation in further trials.

Coagulase-negative staphylococci were detected in the majority of the swabbed locations. These bacteria are known to be associated with human skin and mucosa and can be a source for infections of foreign bodies or indwelling catheters (29). It remains speculative, whether the finding of these bacteria on the disinfected ICUs has a clinical relevance with regard to the transmission of potentially harmful infections.

The use of the hydrogen peroxide nebulisation technique has been shown to reduce bacterial burden on medical devices (30). In the present study, we did not analyse the effectiveness of this technique as compared to control locations without its use. However, we were not able to culture bacteria in the areas where hydrogen peroxide nebulisation was used. This supports the possible role of hydrogen peroxide as a useful adjunct for ICU cleanings. Another advantage of this technique is the operator-independence.

Strengths of this study are the structured and comprehensive viral/bacterial sampling of different objects and its pragmatic design during COVID-19 pandemics reflecting everyday situations on ICUs with possibly relevant impacts on patient outcomes. Furthermore, the single-blinded study design ensured the investigation of regular cleaning quality.

Our study has several limitations. First, the ICUs were cleaned and disinfected by numerous employees of the ICU department and facility management. As a consequence, different parts of the facilities were probably not cleaned uniformly or in the same intensity. However, this fact reflects daily real-life work processes in hospitals. Second, the investigators could not verify if the objects actually had been cleaned or had been omitted owing to human factors such as overlooking or ignoring.

Third, the analysed objects differed in shape and size, which might have influenced the swabbing technique. Fourth, no cluster-randomisation could be performed between the two ICU subunits due to

time-constraints and urgent demand for ICU beds.

Conclusion

In conclusion, several bacteria possibly leading to colonisation or infections of critically ill patients were cultured from swabs taken from disinfected ICUs after the COVID-19 pandemic. SARS-CoV-2 RNA was not detected on any of the swabbed objects further supporting that conventional disinfection is sufficient to safely repurpose COVID-19 ICUs to routine ICUs despite the contagious nature of the disease.

List Of Abbreviations

ICU - intensive care unit

SARS-CoV-2 - severe acute respiratory syndrome Coronavirus 2

RNA - ribonucleic acid

COVID-19 - Coronavirus disease 2019

IRB - institutional review board

H₂O₂ - hydrogen peroxide

rtPCR - Real-time reverse transcriptase polymerase chain reaction

CFU - colony forming units

MDR - multidrug-resistant

Declarations

Ethics approval and consent to participate

Not applicable. No institutional review board (IRB) approval was required as this study did not involve any patients

Consent for publication

Not applicable

Availability of data and materials

The datasets used and analysed during the current study are available from the corresponding author on request

Competing interests

The authors declare that they have no competing interests. PWS has received travel grants from Gilead and Pfizer, and speakers honorary from Pfizer, outside of the submitted work.

Funding

AW and PWS are supported by the academic career program “Filling the gap” of the Medical Faculty of the University of Zurich and SDB is supported by a Fellowship from the Promedica Foundation (1449/M).

Authors' contributions

DAS, PWG, MH, RAS, ASZ, PKB, SDB conceptualised the study

DAH, BD, BC, JDM, MH, JH, ASZ, PKB, SDB were responsible for the methodology

DAH, PWG, BD, BC, JDM, MH, JH, PKB, SDB conducted the investigation

DAH, BD, BC, JDM, MH, JH, PKB, SDB did the formal analysis

DAH, PKB, SDB wrote the original draft

DAH, PWG, BD, BC, JDM, MH, JH, AW, PWS, RAS, ASZ, PKB, SDB reviewed and edited the manuscript

All authors read and approved the final manuscript

Acknowledgments

Not applicable

The COVID-19 ICU-Research Group Zurich. University Hospital of Zurich, University Zurich, Switzerland.

Institute for Intensive Care: *Bartussek, Jan; Buehler, Phillip; Heuberger, Dorothea Monika; Hilty, Matthias Peter; Hofmänner, Daniel Andrea; Maibach, Martina; Schuepbach Reto Andreas; Wendel Garcia, Pedro David.*

Department of Infectious Diseases and Hospital Epidemiology: *Acevedo, Claudio; Chang, Chun-Chi, Brugger, Silvio Daniel; Gomez-Mejia, Alejandro; Mairpady Shambat, Srikanth; Zinkernagel, Annelies*

References

1. Rawson TM, Moore LSP, Zhu N, Ranganathan N, Skolimowska K, Gilchrist M, et al. Bacterial and fungal co-infection in individuals with coronavirus: A rapid review to support COVID-19 antimicrobial prescribing. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America.* 2020.

2. Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of Co-infection Between SARS-CoV-2 and Other Respiratory Pathogens. *Jama*. 2020;323(20):2085-6.
3. Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *The Lancet Respiratory medicine*. 2020;8(5):475-81.
4. Ruan Q, Yang K, Wang W, Jiang L, Song J. Clinical predictors of mortality due to COVID-19 based on an analysis of data of 150 patients from Wuhan, China. *Intensive care medicine*. 2020;46(5):846-8.
5. Wendel Garcia PD, Fumeaux T, Guerci P, Heuberger DM, Montomoli J, Roche-Campo F, et al. Prognostic factors associated with mortality risk and disease progression in 639 critically ill patients with COVID-19 in Europe: Initial report of the international RISC-19-ICU prospective observational cohort. *EClinicalMedicine*. 2020:100449.
6. Zhou F, Yu T, Du R, Fan G, Liu Y, Liu Z, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet (London, England)*. 2020;395(10229):1054-62.
7. Russotto V, Cortegiani A, Raineri SM, Giarratano A. Bacterial contamination of inanimate surfaces and equipment in the intensive care unit. *Journal of intensive care*. 2015;3:54.
8. Schabrun S, Chipchase L. Healthcare equipment as a source of nosocomial infection: a systematic review. *The Journal of hospital infection*. 2006;63(3):239-45.
9. Corman VM, Landt O, Kaiser M, Molenkamp R, Meijer A, Chu DK, et al. Detection of 2019 novel coronavirus (2019-nCoV) by real-time RT-PCR. *Euro surveillance : bulletin Europeen sur les maladies transmissibles = European communicable disease bulletin*. 2020;25(3).
10. Frey PM, Marti GR, Droz S, de Roche von Arx M, Suter-Riniker F, Aujesky D, et al. Bacterial colonization of handheld devices in a tertiary care setting: a hygiene intervention study. *Antimicrobial resistance and infection control*. 2019;8:97.
11. Colaneri M, Seminari E, Novati S, Asperges E, Biscarini S, Piralla A, et al. Severe acute respiratory syndrome coronavirus 2 RNA contamination of inanimate surfaces and virus viability in a health care emergency unit. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2020:S1198-743X(20)30286-X.
12. Chia PY, Coleman KK, Tan YK, Ong SWX, Gum M, Lau SK, et al. Detection of air and surface contamination by SARS-CoV-2 in hospital rooms of infected patients. *Nature communications*. 2020;11(1):2800.
13. Ayliffe GA, Babb JR, Collins BJ, Lowbury EJ, Newsom SW. *Pseudomonas aeruginosa* in hospital sinks. *Lancet (London, England)*. 1974;2(7880):578-81.
14. Aranega-Bou P, George RP, Verlander NQ, Paton S, Bennett A, Moore G. Carbapenem-resistant Enterobacteriaceae dispersal from sinks is linked to drain position and drainage rates in a laboratory model system. *The Journal of hospital infection*. 2019;102(1):63-9.
15. Breathnach AS, Cubbon MD, Karunaharan RN, Pope CF, Planche TD. Multidrug-resistant *Pseudomonas aeruginosa* outbreaks in two hospitals: association with contaminated hospital

- waste-water systems. *The Journal of hospital infection*. 2012;82(1):19-24.
16. de Jonge E, de Boer MGJ, van Essen EHR, Dogterom-Ballering HCM, Veldkamp KE. Effects of a disinfection device on colonization of sink drains and patients during a prolonged outbreak of multidrug-resistant *Pseudomonas aeruginosa* in an intensive care unit. *The Journal of hospital infection*. 2019;102(1):70-4.
 17. Shaw E, Gavaldà L, Càmara J, Gasull R, Gallego S, Tubau F, et al. Control of endemic multidrug-resistant Gram-negative bacteria after removal of sinks and implementing a new water-safe policy in an intensive care unit. *The Journal of hospital infection*. 2018;98(3):275-81.
 18. Zhou Z, Hu B, Gao X, Bao R, Chen M, Li H. Sources of sporadic *Pseudomonas aeruginosa* colonizations/infections in surgical ICUs: Association with contaminated sink trap. *Journal of infection and chemotherapy : official journal of the Japan Society of Chemotherapy*. 2016;22(7):450-5.
 19. Kanamori H, Weber DJ, Rutala WA. Healthcare Outbreaks Associated With a Water Reservoir and Infection Prevention Strategies. *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America*. 2016;62(11):1423-35.
 20. Fusch C, Pogorzelski D, Main C, Meyer CL, El Helou S, Mertz D. Self-disinfecting sink drains reduce the *Pseudomonas aeruginosa* bioburden in a neonatal intensive care unit. *Acta paediatrica (Oslo, Norway : 1992)*. 2015;104(8):e344-9.
 21. Looney WJ. Role of *Stenotrophomonas maltophilia* in hospital-acquired infection. *British journal of biomedical science*. 2005;62(3):145-54; quiz 1 p following 54.
 22. Looney WJ, Narita M, Mühlemann K. *Stenotrophomonas maltophilia*: an emerging opportunist human pathogen. *The Lancet Infectious diseases*. 2009;9(5):312-23.
 23. Podschun R, Ullmann U. *Klebsiella* spp. as nosocomial pathogens: epidemiology, taxonomy, typing methods, and pathogenicity factors. *Clinical microbiology reviews*. 1998;11(4):589-603.
 24. Reverdy ME, Freney J, Fleurette J, Coulet M, Surgot M, Marmet D, et al. Nosocomial colonization and infection by *Achromobacter xylosoxidans*. *Journal of clinical microbiology*. 1984;19(2):140-3.
 25. Logan LK, Weinstein RA. The Epidemiology of Carbapenem-Resistant Enterobacteriaceae: The Impact and Evolution of a Global Menace. *The Journal of infectious diseases*. 2017;215(suppl_1):S28-s36.
 26. van der Mee-Marquet N, Girard S, Lagarrigue F, Leroux I, Voyer I, Bloc D, et al. Multiresistant *Enterobacter cloacae* outbreak in an intensive care unit associated with therapeutic beds. *Critical care (London, England)*. 2006;10(1):405.
 27. Dautzenberg MJ, Wekesa AN, Gniadkowski M, Antoniadou A, Giamarellou H, Petrikos GL, et al. The association between colonization with carbapenemase-producing enterobacteriaceae and overall ICU mortality: an observational cohort study. *Critical care medicine*. 2015;43(6):1170-7.
 28. Mitchell BG, Hall L, White N, Barnett AG, Halton K, Paterson DL, et al. An environmental cleaning bundle and health-care-associated infections in hospitals (REACH): a multicentre, randomised trial. *The Lancet Infectious diseases*. 2019;19(4):410-8.

29. Becker K, Heilmann C, Peters G. Coagulase-Negative Staphylococci. 2014;27(4):870-926.
30. Amodio E, Kuster SP, Garzoni C, Zinkernagel AS, Sax H, Wolfensberger A. Disinfecting noncritical medical equipment-Effectiveness of hydrogen peroxide dry mist as an adjunctive method. American journal of infection control. 2020.

Figures




Figure 1

Both ICU subunits with indicated sampling points. Part A represents ICU 1, Part B represents ICU 2