Effects of Quran Recitation on the Reduction of Preoperative Anxiety in Elective Surgery: A Systematic Review and Meta-Analysis of Randomized Controlled Trials

Vahideh Zarea Gavgani

Tabriz University of Medical Sciences

Mortaza Ghojzadeh

Tabriz University of Medical Sciences

Fatemeh Sadeghi-Ghyassi (✉ ghyassi2722@yahoo.com)

Tabriz University of Medical Sciences <https://orcid.org/0000-0001-9446-9763>

Tahmineh Khodapanah

Tabriz University of Medical Sciences

Research article

Keywords: Elective Surgical Procedures, Anxiety, Meta-analysis, Systematic Review

Posted Date: December 11th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-48044/v2>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Anxiety is a common reaction among patients undergoing surgery. This study aim to assess effectiveness of Quran recitation on reducing the preoperative anxiety.

Methods: A systematic review of the citations in the Medline, EMBASE, Cochrane Library, PsycInfo, Arab World Research Source and other relevant databases was performed to collect the data. Randomized clinical trials about the effect of Quran recitation on the reduction of preoperative anxiety in elective surgery were included without any Language and date restriction. Interventions with self-reading/self-recitation were excluded. The Cochran's Q statistic and the (I^2) with 50 percent threshold was used for calculation of the heterogeneity and inconsistency index. Subgroup analysis was conducted based on the type of surgery. The funnel plot has been used to evaluate the possibility of the publication bias.

Results: Twelve studies were included in the qualitative synthesis and nine studies included in the quantitative synthesis. Our meta-analysis showed a significant reduction in the anxiety level with Quran recitation. The heterogeneity of the included studies was statistically significant ($Q=23.05$, $I^2=65.29$, $P=0.003$). The pooled effect size of the anxiety was $d=-8.893$; 95% CI=[-10.763 to -7.022] ($P < 0.001$). The subgroup analysis showed that Quran recitation has a beneficial effect on reducing anxiety in both major and minor surgeries. There was no publication bias ($t=0.907$, $P=0.39$) in the studies.

Conclusion: Quran recitation can be considered as a non-invasive and peaceful intervention to reduce preoperative anxiety in elective surgery.

Background

There is an increasing interest in non-pharmacological treatments, such as psychological therapy [1]. Music therapy is known as one of the most effective interventions for reducing stress and anxiety. In surgery, music and melodious sounds are used as safe, non-invasive, low-cost and easy to access therapeutic approaches [2, 3]. This plays an outstanding role in controlling the anxiety and stress of patients before and after the surgery [4, 5]. Listening to Quran recitation has a great value among Muslims as a religious melodious sound and extraordinary arrangement of words [6]. It can be considered as an effective psychological therapy with a substantive effect on creating peace of mind and eliminating anxiety. It also proved to be effective in reducing the heart rate and improving calmness in patient in intensive care unit [7].

Anxiety is a common reaction among patients who are admitted to a hospital for surgery [9], that can be described as an unpleasant state of unease or tension caused by the patients' uncertainty about the results of the operation. Anxiety could also be a consequence of a higher analgesic requirement, a prolonged hospital stay, and postoperative pain [10, 11]. It is primarily a cognitive process and the more a person is able to control his or her mind, the better s/he can cope with the anxiety [12]. Various non-pharmacological methods, such as sports, nutrition, diet, relaxation techniques, bibliotherapy [13, 14] and calming music [15] are used for the reduction of sympathetic reactions [16]. Quran recitation itself and listening to Quran recitation are regarded as non-pharmacological approaches like the psychological methods and cognitive therapies that are used for the management of anxiety [17].

The positive effect of Quran recitation has already been reported in terms of the significant reduction of stress (anxiety) and pain in addition to palliative effects [18-20]. A literature review has also revealed that there is a significant relationship between Quran recitation and health and well-being [24]. A recent systematic review assessed the effect of listening to the Quran on anxiety [25]. The review assessed the effect of Quran recitation in various fields in health care systematically but no meta-analysis was performed.

The scope of this study was limited to elective surgery because the non-emergency nature of elective surgeries provides enough time to allow for listening to Quran recitation for patients before the operation. Elective surgery is defined as "a surgery that is a planned, nonemergency surgery that can be scheduled at the convenience of the patient or the physician" [26]. The present study aims to systematically review and synthesize the original articles on the effectiveness of listening to Quran recitation for the management of anxiety in elective surgeries.

Methods

This systematic review was performed based on the statements of the Preferred Reporting Items for Systematic Review and Meta-Analyses (PRISMA) [27].

Review Question

The PICO schema was used, comprising: Patients: All patients with elective surgeries; Intervention/Exposure: Quran recitation; Comparison: None; Outcome: Reduction of anxiety (based on State-Trait Anxiety Inventory (STAI) checklist).

The research question was: What is the effect of Quran recitation on the reduction of anxiety before elective surgeries?

Search strategy

A comprehensive search strategy was conducted in Medline (via Ovid), EMBASE, Cochrane Library, PsycInfo, Scopus, Web of Science, Arab World Research Source, Irandoc (Iranian Research Institute for Information Science and Technology), Magiran, Barakat knowledge network system and Islamic Science Citation (ISC) and Scisearch. Dissertations and theses were searched via Proquest. For citation tracking, Scopus and Web of Science were also searched. ClinicalTrials.gov and IRCT.ir were searched for gray literature. Google Scholar was searched for any further articles. We contacted authors of unpublished and ongoing studies, abstracts found in gray literatures and conference proceedings. The references of the selected studies were manually checked for additional articles which might be useful for the study. The search keywords were Quran, Koran, Kuran, recitation, voice, listening, anxiety, surgery and tune. A combination of controlled vocabulary, synonyms and different spellings were used in the search strategy. Since the publications in this field are mostly from

Muslim countries, Arabic and Persian equivalents of search keywords were also searched. The first round of the search was completed in December 2018 and was updated in January 2020 for any new eligible studies.

Inclusion Criteria

All possible randomized control trials (RCTs) on the effect of Quran recitation on the reduction of anxiety in elective surgery were included in this study. No language restriction was applied. Only studies in which the anxiety level was examined using the State-Trait Anxiety Inventory (STAI) checklist were included in this study.

Exclusion Criteria

Studies without a control group, without a pre-/post-test design, studies of poor quality according to quality assessment, studies using anxiety inventory tools other than the STAI, studies with different outcomes (such as depression) and studies involving the self-recitation of Quran by the patients themselves were excluded. Also studies involving Quran recitation along with its translation were excluded because the main aim of our study was assessment of the effect of listening to Quran voice in its original language.

Study selection and quality assessment

All studies retrieved from databases were imported to EndNote X8 and duplicates eliminated. According to the inclusion and exclusion criteria, two authors (VZG & TKh) independently screened the studies for eligibility based on title and abstract. In the next stage, the remaining articles were examined by full text review. The quality of selected studies was assessed using the Joanna Briggs Institute (JBI) Critical Appraisal Checklist for Randomized Controlled Trials (Table 1) [28]. Any disagreements were resolved through discussion. Two authors extracted data from studies using a data extraction form in Microsoft Excel 2013.

Table 1. JBI Critical Appraisal Checklist for Randomized Controlled Trials

Statistical analysis

The difference between end points with standard deviation (SD) was used for meta-analysis. To assess the heterogeneity, the Cochran’s Q test and the inconsistency index (I^2) were applied. The I^2 statistical threshold value was indicated 50 percent. If I^2 is less than 50 percent, the Mantel-Haenszel fixed effect size model should be used, and if I^2 is more than 50 percent or the p-value is less than 0.05, the random effect model should be used in order to calculate the effect size. Subgroup analysis carried out based on the type of surgery. The funnel plot and the Egger regression test were used to evaluate publication bias ($p < 0.05$ was considered as statistically significant publication bias). A Comprehensive Meta-Analysis (version 2) was employed for data analysis. A p-value of <0.05 was considered statistically significant.

Results

We identified 2381 articles from our systematic search for related literature. After removing duplicates, 986 articles were screened based on title and abstract. Finally, 30 articles were assessed for eligibility. Some related articles were available with two different versions [29-32]. We excluded the version containing the less comprehensive data [29, 32]. Some studies were also excluded because of the study methodology, for example because of assessing the effect of Quran recitation only after surgery [34], or the intervention was a combined model and it was not possible to separate the data for Quran recitation for example Quran recitation with aromatherapy [35] or Quran recitation with its translation [33]. Reasons for exclusion of the other studies from the review were: implementation of different assessment tools, different outcome measures, different publication types and not studying elective surgeries. After reviewing the full text of eligible studies, twelve RCTs were critically appraised. Finally, a total of twelve studies were included in the qualitative synthesis and nine studies were included for the quantitative synthesis [30, 31, 36-42]. The flow diagram of the study is shown in the Figure 1.

The range of patient ages was 25 to 63 years old. All studies investigated the effect of Quran recitation on the reduction of anxiety in the elective surgeries. In eight studies (66.7%), the experimental group received only Quran recitation, whereas in three studies (25%), there were two experimental groups receiving Quran and music [30, 38, 41] and one study receiving Quran recitation and its translation [31]. All the studies were conducted in Iran, as all studies from other Arabic-speaking countries were eliminated in the study selection process as none of them studied elective surgeries. The characteristics of the sample and the level of anxiety before and after intervention are shown in Table 2.

Table 2. State anxiety of included Studies

Quality of studies

Two authors (FSG & MG) independently appraised the quality of studies. The quality of the majority of studies (83.3%) was moderate to high, with less than 17% of studies having low quality (Table 3). Allocation blinding was unclear in most studies (75%). Blinding of participants was not possible due to the nature of study, as participants were aware of the intervention at the time, but in some studies, the researchers wore headphone when assessing patients in the intervention and control groups during the intervention period in order to reduce the risk of bias. In two of the studies, the outcome assessors were blinded to treatment assignment. In three studies, the outcome measure was described in a reliable way.

Table 3. Results of critical appraisal of included studies

Qualitative synthesis

Instrument: In all of the studies the validated Persian version of the STAI was utilized as the outcome measure.

Quran recitation: In all of the studies except three [36, 42, 43], the Quran recitation was played for a duration of 20 minutes. The chapters (Surahs) Yusuf (Joseph) and Ar-Rahman (The Compassionate) were the most commonly used sections, and these were used in three studies. Most of the studies had two patient groups (intervention and control) but three studies included two interventions (one Quran recitation and the other music) with one control group. In these studies, 20 minutes of relaxing music was played to the second patient group.

Anxiety: In all of the studies, Quran recitation significantly reduced the level of anxiety in the intervention group. In the three studies that measured the effect of both music and Quran recitation as interventions, it was revealed that the impact of Quran recitation was larger than that of the music [30, 38, 41]. Four studies evaluated the levels of trait and state anxiety separately [36, 39, 42, 43]. These studies showed a significant decrease in both the trait anxiety and state anxiety in intervention group. One study reported just the total anxiety level [44].

Vital signs and pain: Four studies evaluated the impact of Quran recitation and music on patients' vital signs [30, 31, 42, 44]. The results of Mirbagher et al. and Sharafi show that Quran recitation had a significant effect on patients' breathing, blood pressure and pulse rate [30, 44]. The study of Tayebi et al. showed a significant reduction in breathing and blood pressure but no significant reduction in pulse rate [42]. The results of Shafiei's study were in contradiction with those of Tayebi et al. in that breathing and blood pressure exhibited no significant changes while pulse rate reduced significantly after intervention [31]. Conversely, Asgari's study showed a significant pain reduction in patients undergoing abdominal surgeries [37].

Quantitative synthesis

In this review 9 studies were included comprising a total of 652 patients with elective surgeries who were pooled in the meta-analysis. Three studies were excluded from meta-analysis because the SD was not reported for the mean anxiety score [43, 45] or only the total state and trait anxiety score was reported [44]. The intervention group was formed of 336 patients who listened to between 15 and 20 minutes of Quran recitation.

The mean anxiety score before and after intervention was used to calculate the pooled effect size. According to the results of the meta-analysis, Quran recitation had a positive effect on reducing anxiety in elective surgeries. Since the heterogeneity among included studies was statistically significant ($Q=23.05$, $I^2=65.29$, $P=0.003$), a random effects model was used to calculate the pooled effect size. The pooled difference in means calculated using a random effects model for anxiety showed a statistically significant decrease in anxiety for the Quran recitation group (difference in means: -8.893 ; 95% CI: -10.763 to -7.022 ; $P < 0.001$). Figure 2 shows the forest plot of the combined effect size. Subgroup analysis was performed based on the type of surgery which was categorized into minor and major surgical procedures. The definition of minor and major surgical procedures was based on prior literature [46]. Major surgeries included abdominal surgeries, elective general surgeries and cesarean section. Minor surgeries included endoscopy, angiography and cardiac catheterization. The subgroup analysis showed that Quran recitation has a beneficial effect on reducing anxiety in both major surgical procedures (difference in means: -8.327 ; 95% CI: -11.005 to -5.649 ; $P < 0.001$) and minor surgical procedures (difference in means: -9.877 ; 95% CI: -12.808 to -6.946 ; $P < 0.001$) (Figure 3).

According to the funnel plot and the Egger regression test (Figure 4), there was no publication bias (intercept= -1.64 , 95% CI: -5.94 to 2.64 , $t=0.907$, $df=7$, $P=0.39$).

Discussion

This meta-analysis was conducted to understand if listening to Quran recitation is effective in the management of anxiety before elective surgery. All possible RCTs on the effect of listening to the Quran recitation before the elective surgeries on reducing the anxiety were included in the analysis. The results revealed that listening to the Quran recitation before surgery leads to a significant reduction in anxiety, as measured using the STAI. This finding is consistent with the results of a previous meta-analyses finding a reduction of anxiety from listening to music [47, 48]. This also fulfills the Quranic verses as stated: The Quran is healing for those who perceive its meaning (Quran: 10:57; 17:82). The Quran is known as the ultimate source of guidance for Muslims, a book that contains the divine knowledge, and its instructions are sufficient for guidance of the men [49].

In our study, the quality of 58% of included studies was moderate. Nevertheless, less than half of studies reported the details about the randomization procedure used for assignment of participants and in the rest of studies it was unclear. In half of the studies, both true randomization and allocation concealment were unclear. Only one of the studies declared that those delivering treatment were blind to treatment assignment. Blindness declaration for outcomes assessors was reported by two studies. However, in all of the studies an appropriate statistical analysis was used. There was no selection bias in any of the studies.

According to a study by El-shahat and colleagues, the level of preoperative anxiety varies between minor and major surgical procedures and this rate is higher in the major ones [50]. They recommend psychological counseling at hospitals to help patients for anxiety reduction. The results of our study showed that Quran recitation has a significant effect on reducing preoperative anxiety in both major and minor elective surgeries. Other studies have confirmed the positive effect of various interventions such as music, information therapy and education on reducing anxiety before major and minor surgeries [51-53].

It should be emphasized that although both interventions (Quran and music) have a positive effect on anxiety reduction, the studies indicate that the effect of Quran recitation on anxiety reduction is higher than that of music [30, 38, 41]. But to reach the certainty in this regard, more studies need to be conducted.

The evidence reveals that Quran recitation has been used as a non-medical method for different purposes, especially for psychological issues in researches. An investigation on the effect of reading the Quran on controlling stress has shown that reading the Quran leads to a significant decrease in the stress level among 16-20 year olds [54]. In addition, Quran recitation itself or even listening to Quran recitation has a positive effect on the mental health by decreasing the risk of depression and anxiety-related disorders among caregivers [55], as well as hospitalized patients. It was also found that learning life skills through a

Quranic approach is more effective in the reduction of students' depression [24]. This means that the intervention of the Quran in any form of reading, perception and listening may be effective in improving health and wellbeing. However, the outcomes of a study about the relationship between Quran recitation and post-operative abdominal pain showed no significant relationship [56].

As emphasized by Kaleem and Ahmed, there is a relationship between religious belief and anxiety [57]. Therefore, it is possible that presence of religious belief modulates the effect of listening to Quran recitation. From this, the clinical recommendation may need to take culture into account, and as such the use of Quran recitation before elective surgery can be recommended in Islamic countries to reduce patient anxiety.

In this review, although the included studies were from an Islamic country, their findings did not address the likely relationship between religious beliefs and the positive effect of Quran recitation on anxiety. Additional studies are needed to evaluate the interaction between the differing religious beliefs and the effect of Quran recitation on the anxiety.

The main limitation of the current study was the lack of access to some Arabic Databases to search local language studies. Nevertheless, the current study utilized Google Scholar and Arab World Research Source with regard to Arabic language studies, in addition to international databases. The small sample size may also be considered a limitation of this meta-analysis. The source of heterogeneity could not be identified due to insufficient studies.

One of the strengths of this review was that a validated questionnaire was used to assess the effectiveness of the intervention of all studies. Homogeneity was observed in majority of the studies in terms of the duration of Quran recitation, representing another strength of this study.

Conclusion

Quran recitation has a significant effect on decreasing the anxiety among Muslim patients undergoing elective surgery. Sound of Quran as a non-invasive and peaceful intervention can be supported and advocated by the findings of this study. However, due to the importance of evidence-based decision making in medical science, further studies in different populations especially in non-believers and non-Muslim communities are still needed to achieve certainty.

List Of Abbreviations

PRISMA: Preferred Reporting Items for Systematic Review and Meta-Analyses

STAI: State-Trait Anxiety Inventory

JB: Joanna Briggs Institute

Declarations

Ethics approval and consent to participate: Ethical approval was obtained from the Ethics Committee of the Tabriz University of Medical Sciences under Grant [TBZMED.REC.1394.975].

Consent for publication: Not applicable.

Competing Interest: The authors declare no competing interest.

Availability of data and material: Data available on request from the authors

Competing interests: Not applicable.

Funding: This work is supported by Tabriz University of Medical Sciences under Grant [TBZMED.REC.1394.975]. The funding body played no role in the design of the study and collection, analysis, and interpretation of data and in writing the manuscript.

Authors' contributions: All authors contributed to the design of the study. VZ and TK were in charge of data collection. FSG, MG and VZ conducted the quality evaluation of the included studies. MG and FSG performed the data analysis. All authors drafted the manuscript and approved the final manuscript.

Acknowledgments: We expand our appreciation to Diala Al Samarani Moussa, Head of the Medical Library of Holy Spirit University of Kaslik in Lebanon for introducing the AlManhal database for searching Arabic theses; and Nursing and Midwifery librarians Maryam Razmkhah (Tabriz University of Medical Sciences) and Anahita Dashti (Shahid Beheshti University of Medical Sciences) for their assistance in providing us with access to citations. Also, we would extend our gratitude to Rebecca Dewey, sir Peter Mansfield Imaging Centre, The University of Nottingham for her valuable recommendations on English editing.

We are also grateful to the Iranian Center for Evidence Based Medicine and the Research Development & Cooperation Center (RDCC) of Tabriz University of Medical Sciences for their invaluable support in conducting this meta-analysis. This work is supported by Tabriz University of Medical Sciences under Grant [TBZMED.REC.1394.975].

References

1. McHugh RK, Whitton SW, Peckham AD, Welge JA, Otto MW: **Patient preference for psychological vs. pharmacological treatment of psychiatric disorders: a meta-analytic review.** *The Journal of clinical psychiatry* 2013, **74**(6):595.

2. Micozzi MS: **Fundamentals of complementary and alternative medicine**: Elsevier Health Sciences; 2014.
3. Forooghy M, Tabrizi EM, Hajizadeh E, Pishgoo B: **Effect of music therapy on patients' anxiety and hemodynamic parameters during coronary angioplasty: a randomized controlled trial**. *Nursing and midwifery studies* 2015, **4**(2).
4. Center EI, Center CR, Center PF: **Music therapy**. 2013.
5. Kühlmann A, De Rooij A, Kroese L, van Dijk M, Hunink M, Jeekel J: **Meta-analysis evaluating music interventions for anxiety and pain in surgery**. *British Journal of Surgery* 2018, **105**(7):773-783.
6. Nakhavali F, Seyedi SH: **A Research on " Rhythm & Music" in the Qur'an**. *International Journal of Linguistics* 2013, **5**(3):21.
7. Daud N, Sharif Z: **Effect of Listening to the Al-Quran on Heart Sound**. In: *IOP Conference Series: Materials Science and Engineering: 2018*. IOP Publishing; 2018: 012023.
8. Mahjoob M, Nejati J, Hosseini A, Bakhshani NM: **The effect of Holy Quran voice on mental health**. *Journal of religion and health* 2016, **55**(1):38-42.
9. Spielberger CD: **Anxiety: Current trends in theory and research**: Elsevier; 2013.
10. Eysenck MW: **Anxiety: The cognitive perspective**: Psychology Press; 2013.
11. Eysenck M: **Anxiety and cognition: A unified theory**: Psychology Press; 2014.
12. Chaplin TM, Aldao A: **Gender differences in emotion expression in children: A meta-analytic review**. In.: American Psychological Association; 2013.
13. Reeves T, Stace J: **Improving patient access and choice: assisted bibliotherapy for mild to moderate stress/anxiety in primary care**. *Journal of psychiatric and mental health nursing* 2005, **12**(3):341-346.
14. Jeffcoat T, Hayes SC: **A randomized trial of ACT bibliotherapy on the mental health of K-12 teachers and staff**. *Behaviour research and therapy* 2012, **50**(9):571-579.
15. Jeong GS, Kang Y: **The Effects of Music Therapy on Anxiety, Sedation, and Stress Responses of Patients Undergoing Surgery with Spinal Anesthesia**. *Korean Journal of Adult Nursing* 2016, **28**(5):525-535.
16. Bellocchio L, Soria-Gómez E, Quarta C, Metna-Laurent M, Cardinal P, Binder E, Cannich A, Delamarre A, Häring M, Martín-Fontecha M: **Activation of the sympathetic nervous system mediates hypophagic and anxiety-like effects of CB1 receptor blockade**. *Proceedings of the National Academy of Sciences* 2013, **110**(12):4786-4791.
17. Shafiei N, Salari S, Sharifi M: **Camparison between hearing the quran arabic voice and arabic voice with persian meaning on decreasing the anxiety and vital sighns stabilization of patients befor induction of anesthesia**. *Quran and Medicine* 2011, **2011**(1, Summer [En]):11-15.
18. Alhouseini AMA, Al-Shaikhli IF, bin Abdul Rahman AW, Alarabi K, Dzulkifli MA: **Stress Assessment While Listening to Quran Recitation**. In: *Computer Assisted System in Health (CASH), 2014 International Conference on: 2014*. IEEE; 2014: 67-72.
19. Al-Zaben A, Hamad H, Alfahoum A, Saefan W: **Heart Rate Variability While Listening to Quran Recitation**. *Arabian Journal for Science & Engineering (Springer Science & Business Media BV)* 2014, **39**(2).
20. Jafari H, Bagheri-Nesami M, Abdoli-Nejad MR: **The Effect of Quran Recitation and Religious Music on Mental and Physical Health: A Review Article**. *Journal of Clinical Excellence* 2016, **4**(2):14-11.
21. Heidari M, Shahbazi S: **An Assessment of the Effect of Recitation of the Quran on Exam Anxiety in Nursing and Emergency Medicine Students**. 2015.
22. Elhadj YOM, Alghamdi M, Alkanhal M: **Phoneme-based recognizer to assist reading the Holy Quran**. In: *Recent Advances in Intelligent Informatics*. edn.: Springer; 2014: 141-152.
23. Hematti S, Baradaran-Ghahfarokhi M, Khajooei-Fard R, Mohammadi-Bertiani Z: **Spiritual Well-Being for Increasing Life Expectancy in Palliative Radiotherapy Patients: A Questionnaire-Based Study**. *Journal of religion and health* 2015, **54**(5):1563-1572.
24. Sadeghi H: **Voice of Quran and health: A review of performed studies in Iran**. *Quran and Medicine* 2011, **2011**(1, Summer [En]):33-37.
25. Ghiasi A, Keramat A: **The effect of listening to holy quran recitation on anxiety: A systematic review**. *Iranian journal of nursing and midwifery research* 2018, **23**(6):411.
26. Furterer SL: **Leaning the Surgical Outpatient Screening Process in an Acute Care Hospital**. In: *Lean Six Sigma Case Studies in the Healthcare Enterprise*. edn.: Springer; 2014: 169-201.
27. Moher D, Liberati A, Tetzlaff J, Altman DG: **Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement**. *Int J Surg* 2010, **8**(5):336-341.
28. Tufanaru C, Munn Z, Aromataris E, Campbell J, Hopp L: **Chapter 3: Systematic reviews of effectiveness**. *Joanna Briggs Institute Reviewer's Manual The Joanna Briggs Institute* 2017.
29. Aghajani M, Mirbagher N: **Comparing the Effect of Holy Quran Recitation and Music on Patient Anxiety and Vital Signs before Abdominal Surgeries**. *Islamic Life Style* 2012, **1**(1):66-84.
30. Mirbagher Ajorpaz N, Aghajani M: **The effects of music and Holy Quran on patient's anxiety and vital signs before abdominal surgery**. *Evidence Based Care* 2011, **1**(1):63-76.
31. Shafiei N: **Comparison of the Effect of Two Methods of Quran and Audio Records with the Meaning of Verses on Reducing Anxiety and Stabilizing Signs of Patients Before Anesthesia induction in Imam Khomeini Hospital in Jiroft**. In: *Regional conference on health research in the Holy Qur'an*. Yasuj: Yasuj University of Medical Sciences; 2011: 104-117.
32. Shafiei N, Salari S, Sharifi M: **Camparison between hearing the Quran Arabic Voice and Arabic voice with Persian Meaning on Decreasing the anxiety and Vital Sighns Stabilization of Patients Befor induction of Anesthesia**. *Quran and Medicine* 2017, **2**(1):11-15.

33. Mirsane SA, Kheirkhah D, Shafagh S, Mirbagher Ajorpaz N, Aminpour J: **The effect of listening to Vaghe'a Surah and its translation on the state and trait anxiety before general surgeries: A Randomized Controlled Clinical Trial.** *Health, Spirituality and Medical Ethics* 2016, **3**(3):12-17.
34. Attari M, Sajedi P, Haydari S: **Effect on reducing anxiety and reciting the Quran proved vital signs of patients before induction of anesthesia.** *Teb Tazkiye J* 1998, **1**(37):91-94.
35. Babaii A, Abasnia M, Dehghani F: **Comparison the Effect of Listening to the Holy Quran and Aromatherapy on Anxiety before Cardiac Catheterization.** *Quarterly Journal of Health Breeze* 2015, **3**(3):14-21.
36. Babaii A, Abbasinia M, Hejazi SF, Tabaei S, Reza S, Dehghani F: **The Effect of Listening to the Voice of Quran on Anxiety before Cardiac Catheterization: A Randomized Controlled Trial.** *Health, Spirituality and Medical Ethics* 2015, **2**(2):2-7.
37. Asgari GH: **Effect of the Holy Quran Voice on the Anxiety and Pain in Patients Undergoing Abdominal Surgeries with an Emphasis on the Islamic Culture.** In: *The First international conference on culture, psychopathology and education: 2017; Alzahra University*, 2017.
38. Heidari M, Shahbazi S: **Effect of Quran and music on anxiety in patients during endoscopy.** *Knowledge & Health* 2013, **8**(2):Page: 67-70.
39. Majidi S: **Recitation effect of holy Quran on anxiety of patients before undergoing coronary artery angiography.** *Journal of Guilan University of Medical Sciences* 2004, **13**(49):61-67.
40. Mirbagher AjorPaz N, Ranjbar N: **Effects of recitation of holy Quran on anxiety of women before cesarean section: a randomize clinical trial.** *Qom University of Medical Sciences Journal* 2010, **4**(1):15-19.
41. Sharifi A, Alipour A, Baharloe S: **Comparison of the effect of instrumental music and voices of holy Quran on anxiety of woman before cesarean.** *Journal of Nursing and Midwifery Urmia University of Medical Sciences* 2013, **10**(6):841-846.
42. Tayebi V, Asadi R, Taherpour M, Mousavi Jajarmi SM: **The effect of listening to prayers on anxiety and vital signs before an elective surgery.** *Journal of Medicine and Cultivation* 2015, **23**(3):45-54.
43. Tajvidi M, Mohammadi E, Memarian R: **The effect of Quran sound on anxiety patients before of open heart surgery.** *Danshvar Journal* 2001, **8**(32):55-60.
44. Sharafi E: **Quran sound impact on patients anxiety before lithotripsy surgery in the Baqiyatallah hospital.** School of Nursing and Midwifery, Shahid Beheshti University of medical sciences; 2000.
45. Nikbakht A, Jafari G, Naji H: **Evaluate the effect of reciting the Quran on reducing anxiety before and after diagnostic and therapeutic.** *Teb Tazkiye J* 1998, **1**(29):78-84.
46. Ding L, Hua H, Zhu H, Zhu S, Lu J, Zhao K, Xu Q: **Effects of virtual reality on relieving postoperative pain in surgical patients: A systematic review and meta-analysis.** *International Journal of Surgery* 2020.
47. Loomba RS, Arora R, Shah PH, Chandrasekar S, Molnar J: **Effects of music on systolic blood pressure, diastolic blood pressure, and heart rate: a meta-analysis.** *Indian heart journal* 2012, **64**(3):309-313.
48. Tsai HF, Chen YR, Chung MH, Liao YM, Chi MJ, Chang CC, Chou KR: **Effectiveness of music intervention in ameliorating cancer patients' anxiety, depression, pain, and fatigue: A meta-analysis.** *Cancer Nursing* 2014, **37**(6):E35-E50.
49. Baloch HU: **THE HOLISTIC APPROACH TO HEALTH AND HEALING.** *Mystic Thoughts-Research Journal of Sufism and Peace* 2015, **1**(1).
50. El-shahat M, Taha NM, Ali HH, Metwaly S: **Comparative Study Between Major and Minor Pre-operative Anxiety and Fear Regarding Patients Undergoing Surgery.** *Zagazig Nursing Journal* 2017, **13**(2):190-206.
51. Gupta A, Ahmed B: **Experience of listening to music on patient anxiety during minor oral surgery procedures: a pilot study.** *British Dental Journal* 2020, **228**(2):89-92.
52. Isher SK: **Does preoperative education reduce anxiety in patients undergoing coronary artery bypass surgery?** 2010.
53. Tulgar S, Boga I, Piroglu MD, Ates NG, Bombaci E, Can T, Selvi O, Tas Z, Kose HC: **Preoperative anxiety before spinal anesthesia: Does internet-based visual information/multimedia research decrease anxiety and information desire? A prospective multicentered study.** *Anesthesia, essays and researches* 2017, **11**(2):390.
54. Taghiloo S: **The effect of the holy Quran reading instruction on stress reduction in the young and adolescents.** *Journal of Guilan University of Medical Sciences* 2009, **18**(71):72-81.
55. Kazemi M, Ansari A, Allah Tavakoli M, Karimi S: **The effect of the recitation of holy Quran on mental health in nursing students of Rafsanjan university of medical sciences.** *Journal of Rafsanjan University of Medical Sciences* 2004, **3**(1):52-57.
56. Nikbakht A: **Investigation the effectiveness of Holy Quran sounds on reduction of postoperative pain after abdominal surgery.** *Daneshvar* 1996, **4**(13-14):31-36.
57. Kaleem A, Ahmed S: **The Quran and poverty alleviation: A theoretical model for charity-based Islamic microfinance institutions (MFIs).** *Nonprofit and Voluntary Sector Quarterly* 2010, **39**(3):409-428.

Tables

Table1. JBI Critical Appraisal Checklist for Randomized Controlled Trials

No.	Question	Answers
Q1	Was true randomization used for assignment of participants to treatment groups?	Yes (Y)/No (N)/ Unclear (U)/NA
Q2	Was allocation to treatment groups concealed?	Yes (Y)/No (N)/ Unclear (U)/NA
Q3	Were treatment groups similar at the baseline?	Yes (Y)/No (N)/ Unclear (U)/NA
Q4	Were participants blind to treatment assignment?	Yes (Y)/No (N)/ Unclear (U)/NA
Q5	Were those delivering treatment blind to treatment assignment?	Yes (Y)/No (N)/ Unclear (U)/NA
Q6	Were outcomes assessors blind to treatment assignment?	Yes (Y)/No (N)/ Unclear (U)/NA
Q7	Were treatment groups treated identically other than the intervention of interest?	Yes (Y)/No (N)/ Unclear (U)/NA
Q8	Was follow up complete and if not, were differences between groups in terms of their follow up adequately described and analyzed?	Yes (Y)/No (N)/ Unclear (U)/NA
Q9	Were participants analyzed in the groups to which they were randomized?	Yes (Y)/No (N)/ Unclear (U)/NA
Q10	Were outcomes measured in the same way for treatment groups?	Yes (Y)/No (N)/ Unclear (U)/NA
Q11	Were outcomes measured in a reliable way?	Yes (Y)/No (N)/ Unclear (U)/NA
Q12	Was appropriate statistical analysis used?	Yes (Y)/No (N)/ Unclear (U)/NA
Q13	Was the trial design appropriate, and any deviations from the standard RCT design (individual randomization,	Yes

	parallel groups) accounted for in the conduct and analysis of the trial?	(Y)/No (N)/ Unclear (U)/NA
--	--	-------------------------------------

Table 2. State anxiety of included Studies

Authors	Publication Year	Mean age (Intervention, Control)	Elective surgery type	Quran recitation Time period	Sample Size		State Anxiety							
					Control	Intervention	Control				Intervention			
							Before		After		Before		After	
							Mean	Standard Deviation	Mean	Standard Deviation	Mean	Standard Deviation	Mean	Standard Deviation
Asgari	2017	NA	Abdominal surgeries	20 min	30	30	45.83	10.96	43.66	11.57	52.03	9.24	36.53	
Babaii, A., et al	2015	53.63±9.99, 56.96±7.89	Cardiac catheterization	18 min	30	30	49.8	5.87	50.43	5.63	50.1	5.8	41.2	
Tayebi et al	2014	32.44±11.77, 40.35±14.19	Elective general surgeries	15 min	32	32	42.69	8.6	42.78	8.7	43.83	7.9	41.14	
Heydari, M. & Shahbazi, S.	2013	40±2.1, 39±2.5	Endoscopy	20 min	20	20	43.3	2.61	43.3	2.61	42.1	3.75	34.24	
Sharifi et al	2013	28.9±4.76	Cesarean section	20 min	15	15	46.13	9.61	48.06	8.58	45.33	10.15	35.48	
Shafiei	2011	29.6±7.7, 30.5±8.5	Abdominal surgeries	20 min	60	60	35.23	8.1	37.93	8.6	38.73	8.6	35.37	
Mirbagher Ajorpaz, N., et al	2011	30±1.2, 30±2.4	Abdominal surgery	20 min	45	45	41.38	6.19	43.13	5.32	42.1	7.06	35.21	
Sharafi	2000	30-39	Shock wave lithotripsy	20 min	30	30	102.83**	22.71	101.6**	21.41	108.1**	23.76	78.27**	
Mirbagher AjourPaz, N. & Ranjbar, N.	2010	25-35	Cesarean section	20 min	30	50	38.58	8.69	41.43	8.34	42.1	7.06	35.21	
Majidi.S.	2004	NA*	Angiography	20 min	54	54	48.12	9.56	50.48	10	48.11	7.09	37.31	
Tajvidi et al	2001	NA	Open heart surgery	30 min	30	50	38.43	--	39.56	--	36.26	--	36.3	
Nikbakht et al	1998	38.41	Diagnostic & therapeutic operations	20 min	40	40	47.72	--	44.75	--	44.95	--	37.47	

* Not Available

** State Anxiety + Trait Anxiety

Table 3. Results of critical appraisal of included studies

Studies	Q1	Q2	Q3	Q4	Q5	Q6	Q7	Q8	Q9	Q10	Q11	Q12	Q13	Risk of Bias (%)
Asgari (2017)	Y	Y	Y	U	U	U	Y	Y	Y	Y	Y	Y	Y	Low (83%)
Babaii et al (2015)	Y	U	Y	U	N	U	Y	Y	Y	Y	U	Y	Y	Moderate (61%)
Tayebi et al (2014)	U	U	Y	Y	U	U	Y	Y	Y	U	Y	Y	Y	Moderate (61%)
Heydari, M. & Shahbazi, S. (2013)	U	U	U	U	U	U	Y	Y	Y	Y	U	Y	Y	High (46%)
Sharifi et al (2013)	U	U	Y	Y	U	U	Y	Y	Y	Y	U	Y	Y	Moderate (61%)
Shafiei (2011)	U	Y	Y	Y	U	Y	Y	Y	Y	Y	Y	Y	Y	Low (84%)
Mirbagher Ajorpaz, N., et al (2011)	Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	U	Y	Y	Low (92%)
Sharafi (2000)	Y	U	Y	U	U	U	Y	Y	Y	Y	U	Y	Y	Moderate (61%)
Mirbagher AjorPaz, N. & Ranjbar, N. (2010)	U	U	Y	Y	U	U	Y	Y	Y	Y	U	Y	Y	Moderate (61%)
Majidi.S. (2004)	Y	U	Y	U	U	U	Y	Y	Y	Y	U	Y	Y	Moderate (61%)
Tajvidi et al (2001)	U	U	Y	U	U	U	Y	Y	Y	Y	U	N	Y	High (46%)
Nikbakht et al (1998)	U	U	Y	U	U	U	Y	Y	Y	Y	U	Y	Y	Moderate (53%)

* High: up to 50% of "yes" scores, Moderate: 51% to 80% of "yes" scores, Low: more than 81% of "yes" scores.

Figures

Figure 1

Flow diagram of the present study investigating the effects of Quran recitation on anxiety before surgery

Figure 2

Forest plot of studies investigating the effect of Quran recitation on anxiety before surgery

Figure 3

Forest plot of subgroup analyses based on the type of surgery

Figure 4

Funnel plot and Egger regression of studies showing the effects of Quran recitation on anxiety before surgery ($t=0.907$, $p=0.39$)