

Transcriptional alteration of genes linked to gastritis concerning *Helicobacter pylori* infection status and its virulence factors

Seyedeh Zohre Mirbagheri

Tehran University of Medical Sciences

Ronak Bakhtiari

Tehran University of Medical Sciences

Hashem Fakhre Yaseri

Iran University of Medical Sciences

Abbas Rahimi Foroushani

Tehran University of Medical Sciences

Seyyed Saeed Eshraghi

Tehran University of Medical Sciences

Masoud Alebouyeh (✉ masoud.alebouyeh@gmail.com)

Shahid Beheshti University of Medical Sciences School of Medicine <https://orcid.org/0000-0001-7474-2515>

Research Article

Keywords: *Helicobacter pylori*, Gastritis, Inflammation, virulence genotype, NF- κ B pathway

Posted Date: May 14th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-484619/v1>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background

Helicobacter pylori infection and heterogeneity in its pathogenesis could describe diversity in the expression of inflammatory genes in the gastric tissue. We aimed to investigate transcriptional alteration of genes linked to gastritis concerning the *H. pylori* infection status and its virulence factors.

Methods and Results

Biopsy samples of 12 infected and 12 non-infected patients with *H. pylori* that showed moderate chronic gastritis were selected for transcriptional analysis. Genotyping of *H. pylori* strains was done using PCR and relative expression of inflammatory genes was compared between the infected and non-infected patients using relative quantitative real-time PCR. Positive correlations between transcriptional changes of *IL8* with *TNF- α* and *Noxo1* in the infected and *TNF- α* with *Noxo1*, *MMP7*, and *Atp4A* in the non-infected patients were detected. Six distinct genotypes of *H. pylori* were detected that showed no correlation with gender, ethnicity, age, endoscopic findings, and transcriptional levels of host genes. Irrespective of the characterized genotypes, our results showed overexpression of *TNF- α* , *MMP7*, *Noxo1*, and *ATP4A* in the infected and *IL-8*, *Noxo1*, and *ATP4A* in the non-infected patients.

Conclusions

A complexity in transcription of genes respective to the characterized *H. pylori* genotypes in the infected patients was detected in our study. The observed difference in co-regulation of genes linked to gastritis in the infected and non-infected patients proposed involvement of different regulatory pathways in the inflammation of the gastric tissue in the studied groups.

Introduction

Gastritis is a common disease among humans in different populations. Although this disease can control by therapeutic regimens, its chronic form could lead to cancer. Gastritis is a multifactorial and complex disease associated with both the environment, such as lifestyle, eating habits, and infections and genetic factors (1). *Helicobacter pylori* (*H. pylori*), which is known as the first carcinogenic bacteria, can cause gastritis and chronic inflammation through different mechanisms, including histopathological changes of the gastric tissue that are mediated by its virulence factors, induction of the immune system leading to the infiltration of neutrophils and lymphocytes as well as the production of proinflammatory cytokines, production of substances, such as ammonia, phospholipases, and cytotoxins, which lead to increased risk of malignant alterations of the gastric stem cells (1, 2). The infection occurs during infancy and may remain silent for decades in the gastric environment throughout life. The infection and pathophysiological abnormalities of chronic gastritis (CG) could lead to the loss of functional glands (atrophy) and replacement of the normal gland and foveolar epithelium with intestinal-type cells, intestinal metaplasia, and progress to dysplasia, and carcinoma (3). *H. pylori* also can cause other

diseases, such as peptic and duodenal ulcers, mucosa-associated lymphoid tissue lymphoma, or other conditions, such as recurrent aphthous stomatitis, anemia, altered serum levels of lipoproteins, and coronary atherosclerosis in some patients (3-7).

Different virulence factors are attributed to *H. pylori* that are linked to gastric disorders. Cytotoxic-associated gene A (CagA), vacuolating cytotoxin A (VacA), and several other virulence factors such as outer membrane proteins (OMPs), are involved in *H. pylori*-induced gastric inflammation via the activation of gene transcription (1, 8). Interaction of these virulence factors with host cell receptors on gastric epithelial cells has been described in both in vitro and in vivo studies (1). These interactions trigger intracellular signaling events that result in the release of proinflammatory cytokines and promote bacterial evasion from the acidic environment of the stomach, local mechanical stress, and native immune response (2, 4). Although infiltration of plasma cells, neutrophils, monocytes, differentiation of the gastric cells to the intestinal type cells, and depletion of the parietal cells in the gastric tissue are among common histological changes occurring following the infection, no congruency exist to link some of these virulence factors with these changes (8).

Many research works have been carried out to find gene expression profiles of gastric cancer in tumor tissues; however, there are very few studies that provide information about gene expression analysis in pre-cancerous tissue, especially in patients with chronic gastritis, and its link with different genotypes of *H. pylori* virulence factors. The induction of nuclear factor κ B /tumor necrosis factor-alpha (NF- κ B/TNF- α) inflammatory pathway is known to be the major route of immune-dependent carcinogenesis in the stomach (9). Activation of NF- κ B by *H. pylori* induces nuclear translocation, which causes an increase in transcription of NF- κ B responsive genes, like interleukin (*IL*)-8, and up or down-regulation of inflammatory genes (2). In this pathway, TNF- α , IL-8, nicotinamide dinucleotide phosphate (NADPH) oxidase 1 (NOX1), Matrix metalloproteinase 7 (MMP-7), and ATPase H⁺/K⁺ Transporting Subunit Alpha (ATP4A) cooperate to promote inflammation and histological changes in the gastric tissue (10-12).

Heterogeneity in the pathogenicity of the infection among patients seems to depend on the genotype of *H. pylori* strains carrying different virulence factors, and the extent of induction of genes mediating the activation of this pathway. Understanding this correlation possibly could illustrate observed diversity in the severity of the inflammation and its progress toward cancer. In this study, to describe this relationship, changes in transcription levels of key genes of the *NF- κ B* inflammatory pathway, including *TNF- α* , *IL-8*, *Nox1*, *Atp4aA*, and *MMP-7* were explored and their link with common virulence genotypes of related isolates was analyzed on gastric biopsy samples of patients with gastric disorders.

Materials And Methods

Patients and samples

The study was performed on outpatients who were referred to the endoscopic ward of a general hospital in Tehran, Iran from January to August of 2019. This study was approved by the ethical committee of the Research Center in Tehran University of Medical Science (accepted Number, IR.TUMS.SPH.REC.1398.167

1398/7/3) and an informed consent form was obtained from all the patients. Demographic and clinical information of patients recorded in a questionnaire form. Patients who had received non-steroidal anti-inflammatory drugs or antibiotics within the last six weeks were excluded from this study. Three biopsy specimens of patients with gastric complaints were obtained during endoscopy from the antrum (the distal region of the stomach), and were used for histological examination, *H. pylori* culture and rapid urease test, and RNA extraction.

Histological examination

Biopsy specimens in the pathology department were histologically examined by hematoxylin-eosin staining method and the grade of gastritis was described based on histological parameters and updated Sydney System. Patients samples with chronic gastritis were selected for further analysis.

Isolation and identification

Each biopsy specimen for culture were kept in a transport medium consisting of thioglycollate with 1.3 g/L Agar (Merck, Germany) with 3% yeast extract (Oxoid, UK) and were transferred to the laboratory in less than 2 hours, then were homogenized and cultured on Brucella Agar supplemented with 7% sheep blood, Campylobacter selective supplement (vancomycin 2.0 mg, polymyxin 0.05 mg, trimethoprim 1.0 mg), 10% fetal calf serum, and amphotericin B (2.5 mg/L). Incubation was performed in microaerophilic conditions at 37 °C for 5-7 days. Identification of *H. pylori* isolates was performed by analyzing colony morphology, Gram staining, positive reactions of oxidase and catalase, and urease activities. Confirmation of the identity was done using specific primers for *H. pylori* (*glmM*) by polymerase chain reaction (PCR) as described before (13). The isolates were preserved in BHI broth containing 20% glycerol and 10% fetal calf serum and stored at -70 °C.

Genotyping of *H. pylori* isolates

DNA extraction

Genomic DNA of harvested colonies was extracted as described by Douraghi M (14). Briefly, harvested colonies of the *H. pylori* isolates were suspended in 1ml PBS. After centrifugation (6,000 rpm, 5 min), the pellets were resuspended in 50 mM/L NaOH and heated at 100° C for 20 minutes. After a quick spin and addition of 1 M Tris-HCl, pH 7.5, centrifuged for 5 min at 3,000 rpm. The supernatants containing genomic DNA samples were stored at -20° C until used for molecular studies.

Genotyping

ollowed by a 10-min incuba-

tion in 1 mol / L Tris / HCl (pH 7.5)

ollowed by a 10-min incuba-

tion in 1 mol / L Tris / HCl (pH 7.5)

In this study, genotyping of the isolates was done by PCR. Primers for detection of *cagA*, *cagL*, *cagY*, *vacA* (s and m alleles), and *iceA1* and *iceA2* genes were used in separate reactions as described before (13, 15, 16). Each genotype was defined according to a distinct pattern of the virulence genes detected as follows, *cagA*^{+/-}/*cagL*^{+/-}/*cagY*^{+/-}/*vacA* (s_{1/2}/m_{1/2})/*iceA1*^{+/-}/*A2*^{+/-}.

Gene expression analysis

RNA extraction and cDNA synthesis

To analyze the extent of alteration in the transcription of inflammatory genes in patients with chronic gastritis, RNA extraction of the gastric biopsy samples was done using TRIzol reagent (BlueZol, Iran) following the manufacturer's guidelines with some modifications. Briefly, the tissue samples were cut into small pieces and mixed thoroughly with 600 µl of the extraction solution (BlueZol, Iran). After the addition of 150 µl chloroform and vigorous vortex for 15 seconds, the resulting mixture was centrifuged at 12000 rpm for 15 min at 4 ° C. The colorless upper aqueous phase was transferred to a new clean tube and mixed with 400 µl of isopropanol. The mixture was frozen at -70 ° C for 30 minutes, and the obtained RNA pellet (12,000 rpm, 15min at 4° C) was mixed with 1 ml 80% ethanol. A gentle vortex was applied to suspended the white plate at the bottom of the microtube. The extracted RNA in the pellet of the mixture was obtained after centrifuged at 7500 rpm for 5 min at 4° C. Each pellet was resuspended in 20-30 µl DEPC treated water, and stored at - 70 ° C after heat treatment in 60 ° C for 5 minutes. The extracted RNA concentration was measured by quantitative method (**Nano Drop™ One Microvolume UV-Vis Spectrophotometers**). cDNA synthesis was done after adjustment of RNA concentrations using the easy cDNA synthesis kit (Parstous, Iran), according to the manufacturer's instruction.

Detection of primer efficiency

The efficiency of primers targeting *TNF-α*, *IL-8*, *Noxo1*, *MMP-7*, *Atp4A*, *B2M*, and *ACTB* genes was measured before each analysis. The nucleotide sequence of these primers is shown in Table 1.

Relative quantitative real-time PCR

The expression level of *IL-8*, *TNF-α*, *Noxo1*, *MMP-7*, *ATP4A* genes in the *H. pylori*-infected compared with *H. pylori* non-infected patients with chronic gastritis was measured using SYBR green quantitative real-time PCR. *B2M* and *ACTB* genes were used as endogenous genes as described before (17). The reaction mixture consisted of 0.5 µL of each primer, 12.5 µL RealQ Plus Master Mix Green (Ampliqon, Denmark), 2 µL of cDNA, and distilled water up to the final volume of 25 µL. The thermal cycling conditions in Rotor Gene 6000 Corbett Sequence Detection System have comprised an initial denaturation step at 95 °C for 10 min, 40 cycles of denaturation at 95 °C for 30 seconds, annealing at 60 °C for 60 seconds, elongation at 72 °C for 60 seconds, and a final extension step at 72 °C for 3 min. All the reactions were tested in duplicate. To show the accuracy of the amplification for each gene, primer efficiency, melting curve

analysis, and gel electrophoresis were done. Relative gene expressions for all samples were determined from the obtained Ct (Crossing Threshold) values and using the $2^{-\Delta\Delta ct}$ ($2^{\Delta ct - \text{Average control group } \Delta ct}$) method. Up- and down-regulation were defined based on RQ values ≥ 2 and ≤ 0.5 , respectively (18).

Statistical analysis

Statistical analyses and graphical representation of results were performed using SPSS (25 version) and GraphPad Prism7 softwares. The correlation between the relative expression values of *IL-8*, *TNF- α* , *Noxo1*, *MMP-7*, *ATP4A* genes in the infected group was evaluated by Spearman's correlation nonparametric test. As well, the correlation between *H. pylori* genotypes and relative expression values of these genes in the *H. pylori*-infected patients with chronic gastritis was evaluated by Kruskal-Wallis nonparametric test. *p*-value ≤ 0.05 was considered statistically significant.

Results

Phenotypic and histological results

A total of 168 volunteer patients with gastric disorders (Male, 41.1% and Female, 58.9%), and a mean age of 46 years ranging from 17-86 years, were included in this study. *H. pylori* infection was detected in 27.4% (46/168) of the biopsy specimens. Histopathologic analysis showed that 87.5% (147/168) had chronic gastritis, where 41.49% (61/147) of them presented moderate chronic gastritis in their histological tests. Atrophy, metaplasia, and dysplasia were not detected in these samples. Among the patients with moderate chronic gastritis, biopsy samples of the *H. pylori*-infected and non-infected patients were selected for further analysis. Demographic and pathological information for the patients is presented in supplementary (Table 1). All of the *H. pylori* isolates in the infected patients showed positive results for common biochemical, enzymatic, and PCR (*glmM* gene) tests.

Characteristics of *H. pylori* genotypes

Genotypic characteristics of *H. pylori* isolates showed six distinct genotypes (supplementary, Table 1). Statistical analysis showed no correlation between these genotypes and gender, ethnicity, age, and endoscopic findings.

Expression and correlation of *IL-8*, *TNF- α* , *MMP7*, *Noxo1* and *ATP4A* genes in two groups of patients

The efficiency of primers targeting *TNF- α* , *IL-8*, *Noxo1*, *MMP-7*, *Atp4A* genes was between 0 and 1. Alteration in transcriptional levels of key inflammatory and carcinogenic genes was measured in the gastric tissue samples. The relative expression levels (mean \pm SD) of these genes revealed are presented in Table 2. Accordingly, overexpression of *TNF- α* , *MMP7*, *Noxo1* and *ATP4A* genes in the infected group and *IL-8*, *Noxo1*, and *ATP4A* genes in the non-infected group was detected (Figure 1). Given that the distribution of data was not normal, Spearman's correlation nonparametric test was used to compare coregulation of different genes linked to the inflammatory pathway in *H. pylori*-positive and -negative

groups. There were positive correlations between the expression of *TNF-α* and *IL-8* genes ($r=0.636$, $P=0.026$), *TNF-α* and *Noxo1* genes ($r=0.573$, $P=0.05$), *IL-8* and *Noxo1* genes ($r=0.601$, $P=0.039$) in *H. pylori*-positive group and positive correlations between the expression of *TNF-α* and *MMP-7* genes ($r=0.657$, $P=0.020$), *TNF-α* and *Noxo1* genes ($r=0.629$, $P=0.028$), *TNF-α* and *ATP4A* genes ($r=0.573$, $P=0.051$) in *H. pylori*-negative group that were statistically significant.

Comparison between *H. pylori* genotypes and relative expression values of genes

As was shown in Figure 1, relative expression levels of *IL-8*, *TNF α*, *MMP7*, *Noxo1*, and *ATP4A* genes in *H. pylori*-infected patients with defined genotypes were compared. Statistical analysis doesn't show a correlation between genotypes and the extent of expression for each gene. A link between *H. pylori* genotype A (*cagA*+/*vacA* s1m1) and increased levels of expression for *IL-8*, *TNF-α*, and *MMP-7* was shown, which were inversely accompanied with the lower level of *Noxo1* and *ATP4A* expression compared with the other genotypes. The lowest level of changes in the transcription of the studied genes compared with the non-infected patients was detected in a patient infected with *H. pylori* genotype F (*cagA*-/*vacA* s2m2). Patients infected with genotype A and C showed the highest level of *MMP7* and *TNF-α* expression compared with other genotypes (1.87 and 3.35-fold vs 0.77 ± 0.34 and 2.52 and 2.55-fold vs 0.47 ± 0.28 , respectively). While expression of *IL-8*, *TNF-α*, and *MMP-7* was higher in biopsy of patients infected with genotype A strains, the expression of *ATP4A* and *Noxo1* genes was higher in the patients infected with a more related genotype (genotype B, 16.91 and 3.76 folds, respectively). The patients infected with genotype C of *H. pylori* presented the highest transcription level of *TNF-α*, *MMP-7*, and *Noxo1* genes (2.55, 3.35 and 1.69 folds, respectively) compared to other genotypes (0.88 ± 0.94 , 0.96 ± 0.51 , and 0.62 ± 0.52).

Discussion

Gastritis is a common disease among humans in different populations. This is estimated that more than half of the world population experience chronic gastritis in different degree and extent. The role of chronic gastritis as a serious and insidious illness in the path of gastric carcinogenesis remains largely unknown. *H. pylori* infection is one of the most important contributors of gastritis that shows great diversity in its genomic content in different geographic locations (1, 15).

Host inflammatory responses are mediated through several mechanisms, including induction of oxidative-reductive stress response pathway, activation of cytotoxic immune cells, B cell activation, and permanent secretion of inflammatory cytokines and chemokines (2). Although real mechanisms of gastritis and its progression toward gastric cancer after *H. pylori* infection are poorly known, in vitro studies showed that induction of the nuclear factor- κ B (NF- κ B) inflammatory pathway is a critical regulator in this relationship. Activation of this pathway could promote immune-dependent inflammation and histological changes in the stomach (9).

A total of 1840 genomics types of *H. pylori* have been described according to the GenBank database, which shows great diversity in this bacterium that is due to its impaired DNA repair system and

evolutionary events. Pathogenicity of this bacterium in strains with intact type 4 secretion system is commonly linked to the expression of CagA, VacA, IceA, Urease, together with adhesions and outer membrane proteins. Finding a link between the genotypes of this bacterium and the extent of inflammatory response in the gastric tissue is hard, since there are several variants of each virulence gene. In this study, our results confirmed this complexity in the studied patients.

Results of our study showed a complexity between the extent of transcription in genes related to inflammation and gastric carcinogenesis and genotypes of the *H. pylori* isolates. Nearly half of the patients were infected with wild-type genes, while the absence of some virulence genes was detected in other samples. Totally, six different genotypes of *H. pylori* were detected in biopsy samples of patients with CG, which represented similar diversity as were reported from previous studies in Iran (15). Genotype A (*Cag A*⁺/*Cag Y*⁺/*Cag L*⁺/*Vac A m*₁*s*₁/*ice A1*⁺), which seems to present the highest pathogenicity based on *in vitro* studies, was detected in 25% of the patients. Histological analysis in the patients infected with this genotype showed similar pathogenicity compared with the patients infected with genotype C, a genotype with low virulence capacity (*Cag A*⁻/*Cag Y*/*Cag L*⁻/*Vac A m*₂*s*₂/*ice A1*⁺/*A2*⁺). Similar to this finding, in a study by Chiurillo *et al*, no congruency was detected between genotypes of *H. pylori* isolates and histological findings (19).

Besides variation in ethnicity of the participants, the observed diversity in genotypes could be explained through impaired DNA repair mechanisms during the replication that delivers several deletions and mutations (19). In our study, this diversity showed to have no impact on the transcription level of *TNF-α* in the infected patients compared with non-infected ones. Accordingly, overexpression of *TNF-α* was detected just for patients infected with the strains belonged to genotype A and C, which show no similarity in their genotypes. In general, a *H. pylori* strain with genotype A is considered virulent, where expression and secretion of CagA and VacA s₁m₁ in the gastric tissue is associated with the increased level of *TNF-α* expression and inflammation (20). The induction of *TNF-α* in three infected patients with *cagA* negative strain carrying an inactive variant of *vacA* s₂m₂ allele proposed the involvement of some other virulence factors, host factors, or microbes other than *Helicobacter* in this interplay. This discrepancy was previously reported by Zabaglia *et al* (21). Although carriage of *iceA1/A2* allele in *H. pylori* isolates with genotype C could explain the induction of *TNF-α* in the gastric tissue; lack of this induction in a patient infected with *H. pylori* genotype E has challenged this link. Similar to our finding, in a study by Yamaoka *et al*. which was done on the gastric tissue, no significant difference in expression of *TNF-α* was shown between the infected patients with *cagA*⁺ and *cagA*⁻ strains (22). A positive correlation between the expression of *TNF-α* gene with *IL-8* genes and *Noxo1* genes in the *H. pylori*-positive group and with *MMP-7*, *Noxo1*, and *ATP4A* genes in the *H. pylori*-negative group confirms its role as a main inflammatory factor that acts as a master switch in establishing between inflammation and cancer (23).

Overexpression of IL-8, as the second important cytokine mediating a role in inflammation, as shown by these authors both in the corpus and antrum of infected patients with *cagA*⁺ genotype compared with *cagA*⁻ ones. This finding is inconsistent with our results since the overexpression was just detected in the

infected patients with genotype A. These results are in agreement with Iqbal Siddique' (2014) and Celin Audibert' (2000) findings which showed that the presence of *cagA* is not associated with increased expression of *IL-8* gene (24, 25). Regardless of defined genotypes, although 2.5-fold higher level of IL-8 transcription was measured in *cagA*⁺ compared with *cagA*⁻ strains in our study, carriage of *vacA* s1m1 allele was a hallmark for explaining IL-8 overexpression in our samples since this allele was unique in the strains with genotype A, where nearly 8-fold higher level of the expression was detected compared with other genotypes.

One unanticipated finding was that the relative expression of the *IL-8* gene was lower in the infected patients with *H. pylori* compared to non-infected patients. This result differs from some published studies (26-30), which suggested that *H. pylori* infection is not the only factor associated with IL-8 induction in the gastric tissue. Gene polymorphisms in the mediators of NF-κB pathway, induction of TGF-β, IL-10, and Muc-1, and interaction of non-Helicobacter bacteria are among other factors that can influence the expression of IL-8 in patients with gastritis (31). In the case of studied patients with *H. pylori* infection, these results suggested that there is a link between genotypes of *H. pylori* and the level of *IL-8* gene expression. According to our knowledge, while no study examined the effect of complete *H. pylori* genotype on the extent of IL-8 transcription, similar to our results the higher level of IL-8 expression was previously established for *H. pylori* strains with *cagA*⁺/*vacA* s1m1 variant (20, 25, 29).

MMP-7 is a member of the MMP family and a key player in the inflammation process and carcinogenesis, which increase in *H. pylori* gastritis and early gastric carcinoma (32, 33). Based on our results, a higher level of the MMP-7 mRNA was detected in the *H. pylori*-positive group than in the *H. pylori*-negative group, but this difference was not significant. These results match those observed in earlier studies (32-36). In the *H. pylori*-positive group, the patients infected with genotype A and C of *H. pylori* had higher expression of the *MMP-7* gene compared to other genotypes and unlike previous studies, it was found that there was no relationship between the presence of *cagA*⁺ *H. pylori* strains and the increase in *MMP-7* gene in patients (36, 37). The ice A1 and ice A1/A2 seem to be involved in *MMP-7* expression that is suggested due to the presence of inactive *vacA* allele and *cagPAI* in the stains with genotype C. However, lack of MMP-7 induction in a sample with *H. pylori* genotype E, which carry this virulence gene, again showed the complexity for the description of this correlation.

H⁺, K⁺-adenosine triphosphatase (H⁺, K⁺-ATPase) as a proton pump and a marker of parietal cell function is the key pathway mediating the secretion of gastric acid and is affected by *H. pylori*. *H. pylori* or its products inhibit the activity of the promoter of the alpha-subunit of H⁺, K⁺ ATPase and suppress the expression of H⁺, K⁺-ATPase (38, 39). A few studies have examined the changes in the expression level of the *ATP4A* gene in human gastric. Our results are almost in line with Nayoung Kim and Ok Jae Lee's (2020) findings which showed there were no significant differences in ATP4A mRNA level between *H. pylori*-negative and *H. pylori*-positive groups (38). In our study, downregulation of *ATP4A* was detected for all genotypes, except genotypes B and E, where overexpression was detected in only one patient. This finding showed that transcription of this gene could be mediated by different pathways, independent of genotypes of *H. pylori* strains. In previous studies, the effect of existence *cagA* and [type IV secretory](#)

system (T4SS), *cagL*, for repression of *ATP4A* expression was confirmed in studies of *H. pylori* *cagPAI* strains (40-42), but in our study, such a relationship was not seen.

NADPH oxidase organizer 1 (NOXO 1) is one of the components forming the NOX1 complex. It is known as a TNF- α -dependent tumor-promoting factor for gastric inflammation (43). Several reports have shown that *Noxo1* expression is significantly upregulated in gastritis as well as the intestinal-type or diffuse-type gastric cancer (43, 44). In our study, there was no difference in the relative expression of the *Noxo1* gene between the two groups of patients. This result may be explained by the fact that all the samples used in our study were from the gastric tissue of patients with chronic gastritis. In the *H. pylori*-positive group, the patients infected with genotypes B and C showed higher levels of *Noxo1* expression compared with the other genotypes; however, no virulence factor could describe this increased level of the expression. Alteration in the expression level of *Noxo1* accompany with *TNF- α* and *IL-8* in the *H. pylori*-positive group and its link with *TNF- α* in the *H. pylori*-negative group represents its role in gastritis irrespective of the type of infections in the gastric tissue. In future studies, determining the role of infection with other microbes could shed light on this interaction.

Conclusion

In general, in this study, we investigated the impact of *H. pylori* infection and its characterized combined virulence genotypes on transcriptional changes of genes linked to the inflammatory pathway in the gastric tissue of the infected compared with non-infected patients with chronic gastritis. Our results showed complexity in the transcription of genes link to the inflammatory pathways in the gastric tissue respective to the characterized *H. pylori* genotypes. A direct relationship with overexpression of *IL-8*, *TNF- α* , and *MMP-7* genes and downregulation of *Noxo-1* and *ATP4A* was detected in the samples of patients infected with hypervirulent strains with genotype A compared with other genotypes. Irrespective of genotypes of *H. pylori*, our results showed a significant positive correlation between transcriptional changes of *IL8* in conjunction with *Noxo1* and/or *TNF- α* in *H. pylori*-infected patients that was different from the characterized correlation in the transcription of *TNF- α* and *ATP4A/Noxo1/MMP-7* genes in *H. pylori*-negative patients with gastritis. This finding proposed the involvement of some other host factors, such as gene polymorphisms in the mediators of the NF- κ B pathway, the interplay of the anti-inflammatory pathway, and the interaction of non-*Helicobacter* bacteria in this regard. More studies on a larger number of samples, especially those with characterized microbiota and related host immunogenetics and transcriptional data could provide valuable documents about this interaction.

Declarations

Acknowledgments

The authors would like to thank all colleagues of Pediatric Infections Research Center (PIRC), Research Institute for Children's Health, Mofid Children's Hospital, Shahid Beheshti University of Medical Sciences, Tehran, Iran for their cooperation in this study. The authors of this study also thank the kindly support of

gastroenterology and pathology units of Firoozgar hospital, Iran university of medical sciences, Tehran, Iran.

Author contributions

We declare that all the authors fulfilled the authorship criteria and All authors read and approved the final version of the manuscript.

Funding

This study was funded by a Ph.D. grant from the Department of Pathobiology, School of Public Health, Tehran University of Medical Sciences, Tehran, Iran.

Declarations

Conflict of interest

The authors declare that there is no conflict of interest.

Ethical approval

This study was approved by the ethical committee of the Research Center in Tehran University of Medical Science (accepted Number, IR.TUMS.SPH.REC.1398.167 1398/7/3) and an informed consent form was obtained from all the patients.

Informed consent

The authors included in the study consent to this manuscript to participate and for publication.

Consent for publication

The authors declare that they consent for publication of this study.

References

1. Varbanova M, Frauenschläger K, Malfertheiner PJP, Gastroenterology RC. Chronic gastritis–An update. 2014;28(6):1031-42.
2. Qadri Q, Rasool R, Gulzar G, Naqash S, Shah ZAJJogc. H. pylori infection, inflammation and gastric cancer. 2014;45(2):126-32.
3. Carrasco G, Corvalan AHJGr, practice. Helicobacter pylori-induced chronic gastritis and assessing risks for gastric cancer. 2013;2013.
4. Ricci V, Romano M, Boquet PJWjogW. Molecular cross-talk between Helicobacter pylori and human gastric mucosa. 2011;17(11):1383.

5. Riggio MP, Lennon A, Wray DJ, Joop, medicine. Detection of *Helicobacter pylori* DNA in recurrent aphthous stomatitis tissue by PCR. 2000;29(10):507-13.
6. Jia E-Z, Zhao F-J, Hao B, Zhu T-B, Wang L-S, Chen B, et al. *Helicobacter pylori* infection is associated with decreased serum levels of high density lipoprotein, but not with the severity of coronary atherosclerosis. 2009;8(1):1-7.
7. Farsak B, Yildirim A, Akyön Y, Pinar A, Öç M, Böke E, et al. Detection of *Chlamydia pneumoniae* and *Helicobacter pylori* DNA in Human Atherosclerotic Plaques by PCR. 2000;38(12):4408-11.
8. Correa P, Piazuelo M, Jodt. The gastric precancerous cascade. 2012;13(1):2-9.
9. Isomoto H, Mizuta Y, Miyazaki M, Takeshima F, Omagari K, Murase K, et al. Implication of NF- κ B in *Helicobacter pylori*-associated gastritis. 2000;95(10):2768-76.
10. Liu T, Zhang L, Joo D, Sun S-C. NF- κ B signaling in inflammation. 2017;2(1):1-9.
11. Williams RA, Timmis J, Qwarnstrom E. Computational models of the NF- κ B signalling pathway. 2014;2(4):131-58.
12. Hammond CE, Beeson C, Suarez G, Peek Jr RM, Backert S, Smolka A, JoP-G, et al. *Helicobacter pylori* virulence factors affecting gastric proton pump expression and acid secretion. 2015;309(3):G193-G201.
13. Yadegar A, Mobarez AM, Alebouyeh M, Mirzaei T, Kwok T, Zali MR, JoM, et al. Clinical relevance of *cagL* gene and virulence genotypes with disease outcomes in a *Helicobacter pylori* infected population from Iran. 2014;30(9):2481-90.
14. Saberi S, Douraghi M, Azadmanesh K, Shokrgozar MA, Zeraati H, Hosseini ME, et al. A potential association between *Helicobacter pylori* CagA EPIYA and multimerization motifs with cytokeratin 18 cleavage rate during early apoptosis. 2012;17(5):350-7.
15. Vaziri F, Peerayeh SN, Alebouyeh M, Mirzaei T, Yamaoka Y, Molaei M, et al. Diversity of *Helicobacter pylori* genotypes in Iranian patients with different gastroduodenal disorders. 2013;19(34):5685.
16. Ta LH, Hansen LM, Sause WE, Shiva O, Millstein A, Ottemann KM, et al. Conserved transcriptional unit organization of the *cag* pathogenicity island among *Helicobacter pylori* strains. 2012;2:46.
17. Wisnieski F, Calcagno DQ, Leal MF, dos Santos LC, de Oliveira Gigek C, Chen ES, et al. Reference genes for quantitative RT-PCR data in gastric tissues and cell lines. 2013;19(41):7121.
18. Festuccia C, Gravina GL, Giorgio C, Mancini A, Pellegrini C, Colapietro A, et al. UniPR1331, a small molecule targeting Eph/ephrin interaction, prolongs survival in glioblastoma and potentiates the effect of antiangiogenic therapy in mice. 2018;9(36):24347.
19. Chiurillo MA, Moran Y, Cañas M, Valderrama E, Granda N, Sayegh M, et al. Genotyping of *Helicobacter pylori* virulence-associated genes shows high diversity of strains infecting patients in western Venezuela. 2013;17(9):e750-e6.
20. Augusto AC, Miguel F, Mendonça S, Pedrazzoli Jr J, Gurgueira SA, JCb. Oxidative stress expression status associated to *Helicobacter pylori* virulence in gastric diseases. 2007;40(9-10):615-22.

21. Zabaglia LM, Ferraz MA, Pereira WN, Orcini WA, de Labio RW, Neto AC, et al. Lack of association among TNF- α gene expression,-308 polymorphism (G> A) and virulence markers of *Helicobacter pylori*. 2015;21(1):1-7.
22. Yamaoka Y, Kita M, Kodama T, Sawai N, Imanishi JJG. *Helicobacter pylori* cagA gene and expression of cytokine messenger RNA in gastric mucosa. 1996;110(6):1744-52.
23. Wu Y-d, Zhou BJBjoc. TNF- α /NF- κ B/Snail pathway in cancer cell migration and invasion. 2010;102(4):639-44.
24. Siddique I, Al-Qabandi A, Al-Ali J, Alazmi W, Memon A, Mustafa AS, et al. Association between *Helicobacter pylori* genotypes and severity of chronic gastritis, peptic ulcer disease and gastric mucosal interleukin-8 levels: Evidence from a study in the Middle East. 2014;6(1):1-10.
25. Audibert C, Janvier B, Grignon B, Salaüna L, Burucoa C, Lecron J-C, et al. Correlation between IL-8 induction, cagA status and vacA genotypes in 153 French *Helicobacter pylori* isolates. 2000;151(3):191-200.
26. Crabtree J, Wyatt J, Trejdosiewicz L, Peichl P, Nichols P, Ramsay N, et al. Interleukin-8 expression in *Helicobacter pylori* infected, normal, and neoplastic gastroduodenal mucosa. 1994;47(1):61-6.
27. Moss S, Legon S, Davies J, Calam JJG. Cytokine gene expression in *Helicobacter pylori* associated antral gastritis. 1994;35(11):1567-70.
28. Sharma SA, Tummuru MK, Blaser MJ, Kerr LDJTjoi. Activation of IL-8 gene expression by *Helicobacter pylori* is regulated by transcription factor nuclear factor- κ B in gastric epithelial cells. 1998;160(5):2401-7.
29. Bartchewsky Jr W, Martini MR, Masiero M, Squassoni AC, Alvarez MC, Ladeira MS, et al. Effect of *Helicobacter pylori* infection on IL-8, IL-1 β and COX-2 expression in patients with chronic gastritis and gastric cancer. 2009;44(2):153-61.
30. Outlioua A, Badre W, Desterke C, Echarke Z, El Hammani N, Rabhi M, et al. Gastric IL-1 β , IL-8, and IL-17A expression in Moroccan patients infected with *Helicobacter pylori* may be a predictive signature of severe pathological stages. 2020;126:154893.
31. Bornschein J, Kandulski A, Selgrad M, Malfertheiner PJDD. From gastric inflammation to gastric cancer. 2010;28(4-5):609-14.
32. Lu L, Ma G, Liu X, Sun R, Wang Q, Liu M, et al. Correlation between GDF15, MMP7 and gastric cancer and its prognosis. 2017;21(3):535-41.
33. Gontar Siregar SH, Sitepu RJOaMjoms. Serum IL-10, MMP-7, MMP-9 levels in *Helicobacter pylori* infection and correlation with degree of gastritis. 2016;4(3):359.
34. Lu P, Takai K, Weaver VM, Werb ZJCSHpib. Extracellular matrix degradation and remodeling in development and disease. 2011;3(12):a005058.
35. Wroblewski LE, Noble P-J, Pagliocca A, Pritchard DM, Hart CA, Campbell F, et al. Stimulation of MMP-7 (matrilysin) by *Helicobacter pylori* in human gastric epithelial cells: role in epithelial cell migration. 2003;116(14):3017-26.

36. Sadeghiani M, Bagheri N, Shahi H, Reisi S, Rahimian G, Rashidi R, et al. cag Pathogenicity island-dependent upregulation of matrix metalloproteinase-7 in infected patients with *Helicobacter pylori*. 2017;38(6):595-607.
37. Crawford HC, Krishna US, Israel DA, Matrisian LM, Washington MK, Peek Jr RMJG. *Helicobacter pylori* strain-selective induction of matrix metalloproteinase-7 in vitro and within gastric mucosa. 2003;125(4):1125-36.
38. Kim HJ, Kim N, Park JH, Choi S, Shin CM, Lee OJJG, et al. *Helicobacter pylori* eradication induced constant decrease in interleukin-1B expression over more than 5 years in patients with gastric cancer and dysplasia. 2020;14(6):735.
39. Yao X, Smolka AJJG. Gastric parietal cell physiology and *Helicobacter pylori*-induced disease. 2019;156(8):2158-73.
40. Saha A, Backert S, Hammond CE, Gooz M, Smolka AJJG. *Helicobacter pylori* CagL activates ADAM17 to induce repression of the gastric H, K-ATPase α subunit. 2010;139(1):239-48.
41. Chang W-L, Yeh Y-C, Sheu B-SJJobs. The impacts of *H. pylori* virulence factors on the development of gastroduodenal diseases. 2018;25(1):1-9.
42. Saha A, Hammond CE, Trojanowska M, Smolka AJJAJoP-G, Physiology L. *Helicobacter pylori*-induced H, K-ATPase α -subunit gene repression is mediated by NF- κ B p50 homodimer promoter binding. 2008;294(3):G795-G807.
43. Oshima H, Ishikawa T, Yoshida G, Naoi K, Maeda Y, Naka K, et al. TNF- α /TNFR1 signaling promotes gastric tumorigenesis through induction of Noxo1 and Gna14 in tumor cells. 2014;33(29):3820-9.
44. Echizen K, Horiuchi K, Aoki Y, Yamada Y, Minamoto T, Oshima H, et al. NF- κ B-induced NOX1 activation promotes gastric tumorigenesis through the expansion of SOX2-positive epithelial cells. 2019;38(22):4250-63.
45. Lou X, Zhu H, Ning L, Li C, Li S, Du H, et al. EZH2 regulates intestinal inflammation and necroptosis through the JNK signaling pathway in intestinal epithelial cells. 2019;64(12):3518-27.
46. Ohki R, Yamamoto K, Mano H, Lee RT, Ikeda U, Shimada KJJoh. Identification of mechanically induced genes in human monocytic cells by DNA microarrays. 2002;20(4):685-91.
47. Binato R, Santos EC, Boroni M, Demachki S, Assumpção P, Abdelhay EJO. A common molecular signature of intestinal-type gastric carcinoma indicates processes related to gastric carcinogenesis. 2018;9(7):7359.
48. Rubach M, Lang R, Hofmann T, Somoza VJAotNYAoS. Time-dependent component-specific regulation of gastric acid secretion-related proteins by roasted coffee constituents. 2008;1126(1):310-4.
49. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, et al. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. 2002;3(7):1-12.
50. Steinau M, Rajeevan MS, Unger ERJTJoMD. DNA and RNA references for qRT-PCR assays in exfoliated cervical cells. 2006;8(1):113-8.

Tables

Table 1 Primer sequences for the RT-PCR analysis

Name	Sequence (5' 3')	Length (bp)	Size of Product (bp)	Annealing tm	References
<i>TNF-α</i> ¹	F: GAGGCCAAGCCCTGGTATG	19	91	60	(45)
	R: CGGGCCGATTGATCTCAGC	19			
<i>IL-8</i> ²	F: GAACTGAGAGTGATTGAGAGTGGA	24	134	62	(46)
	R: CTCTTCAAAAACCTTCTCCACAACC	24			
<i>Noxo1</i> ³	F: AGATCAAGAGGCTCCAAACG	20	120	60	(44)
	R: GGAAGGTCTCCTTGAGGGTCT	21			
<i>MMP-7</i> ⁴	F: TGCAGAAGCCCAGATGTGGAGTG	23	96	62	(47)
	R: CGATCCTGTAGGTGACCACTTTGG	24			
<i>ATP4A</i> ⁵	F: CGGCCAGGAGTGGACATTCG	20	176	60	(48)
	R: ACACGATGGCGATCACCAGG	20			
<i>β2M</i> ⁶	F: TGCTGTCTCCATGTTTGATGTATCT	25	86	60	(49)
	R: TCTCTGCTCCCCACCTCTAAGT	22			
<i>β-actin</i> ⁷	F: ATGTGGCCGAGGACTTTGATT	21	107	60	(50)
	R: AGTGGGGTGGCTTTTAGGATG	21			

1. Tumor necrosis factor-alpha
2. Interleukin (*IL*)-8
3. Nicotinamide dinucleotide phosphate (NADPH) oxidase 1
4. Matrix metalloproteinase 7
5. ATPase H⁺/K⁺ Transporting Subunit Alpha
6. Beta-2 microglobulin
7. Beta-actin

Table 2 Relative expression (Mean±SD) of genes in two groups of patients, moderate chronic gastritis/ *H. pylori* positive and moderate chronic gastritis/ *H. pylori* negative.¹

Groups	<i>IL-8</i>	<i>TNF-α</i>	<i>MMP-7</i>	<i>NOXO1</i>	<i>ATP4A</i>
	Relative expression Mean±SD				
Moderate chronic gastritis/ <i>H. pylori</i> positive	1.203 ± 2.805	2.084 ± 2.684	3.242 ± 3.727	3.641 ± 4.805	4.299 ± 9.051
Moderate chronic gastritis/ <i>H. pylori</i> negative	2.587 ± 2.28	1.342 ± 1.068	1.939 ± 1.398	3.645 ± 8.613	2.779 ± 3.962

Gene expressions for all samples were determined from the obtained Ct (Crossing point) values and using the $2^{-\Delta\Delta Ct}$ ($2^{-(\text{Sample } \Delta Ct - \text{Average control group } \Delta Ct)}$) method. Up- and down-regulation was defined based on RQ values ≥ 2 and ≤ 0.5 , respectively.

Figures




Figure 1

A comparison of genes expression between the two groups of patients in all samples (moderate chronic gastritis/ *Helicobacter pylori* (*H. pylori*) positive, moderate chronic gastritis/ *H. pylori* negative), B Comparison between *H. pylori* genotypes and relative expression values of genes

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supplementary.docx](#)