

Maternal mortality following thromboembolism; Incidences and prophylaxis strategies

Mahboobeh Shirazi

Tehran University of Medical Sciences

Behrokh Sahebdel

Tehran University of Medical Sciences

Mahnoosh Torkzaban

Thomas Jefferson University

Elham Feyzabad

Tehran University of Medical Sciences

marjan ghaemi (✉ marjan_ghaemi@yahoo.com)

Tehran University of Medical Sciences

Research

Keywords: Maternal Mortality, Pregnancy, Thromboembolism, Heparin

Posted Date: August 5th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-49298/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on November 30th, 2020. See the published version at <https://doi.org/10.1186/s12959-020-00251-w>.

Abstract

Background: Thromboembolism is one of the main causes of maternal mortality, which can be prevented in many cases. The present study was designed to investigate the incidence and prophylaxis strategies for maternal mortality following thromboembolism.

Methods: In this case series study, the data of the mortality cases were extracted according to the ethical and security standards of the Ministry of Health of the country. The thromboembolism risk factors measured and scored using a questionnaire entitled "the evaluation of risk factors for maternal mortality following thromboembolism during pregnancy, labor, or post-partum".

Results: The maternal mortality rate was 16 per 100,000 live births. Among 297 mortality cases, 27 (9%) death were due to thromboembolism. The mean gestational age was 32.5 weeks. Dyspnea (88.8%) and tachycardia (18.5%) were found as common clinical manifestations in these patients. Sixteen cases (59.3%) did not get heparin, 6 (22.2%) received single dose and 5 (18.5%) received two doses and more. In these 11 cases, 5 (45%) patients received heparin before surgery, 1 after surgery, and 5 before and after surgery. Twenty cases deceased in the first hours after delivery and the rest after 2 to 12 days. The average score of risk for thromboembolism based on Royal College of Obstetricians & Gynecologist (RCOG) guideline was 4.6.

Conclusion: It seems that one of the most important cause of maternal mortality in this study was the lack of recognition of high-risk patients and the lack of prescription for prophylaxis with heparin and this clearly explains the need for accurate screening of high-risk mothers, designing a standard form and the care and treatment of these patients.

Introduction

Maternal mortality is one of the greatest tragedies with wide ranging consequences for the family and society (1). It is an important public health indicator that reflects both the quality of health care services and the women's status (2, 3). Reduction of maternal mortality serves as one of the most important international development goals (4). Several factors such as gravidity, type of delivery, socioeconomic status was found as the most important determinants of maternal mortalities in the region (3).

Primary causes of maternal mortality in U.S include infection, cardiac disease, hemorrhage, thromboembolism and hypertensive disorders (5). Iran has also experienced a transition pattern in its major causes of maternal mortality and morbidity from those characteristics of developing nations such as hemorrhage and infection to causes more commonly seen in developed nations including thromboembolism, cardiac diseases (6–8) and in not rare condition, post-operative complications (9). Accurate recognition and attention on maternal mortality is critical to have an effective prevention (10).

Pregnancy is a prothrombotic and temporary condition that increase the risk of venous thromboembolism (VTE) four to five times as compared to non-pregnant women (11, 12). The risk of

venous thromboembolism is higher in women ages 35; Thrombophilia, lupus, cardiac disease, sickle cell disease, obesity, fluid and electrolyte imbalance, postpartum infection and transfusion are the other significant risk factors (11). Many cases of VTE occur during the first trimester of pregnancy and the risk increases with progression of gestational age and reaches the maximum after delivery (13).

Thromboembolic events related to pregnancy are reported by 13 people per 10,000 cesarean and also suggests that cesarean delivery is associated with a five-fold increase in the risk of VTE incidence (14). Women with preeclampsia are at risk of treating thromboembolic events, three times more than healthy women, in the postpartum period. There is also a correlation between smoking and the increased risk of VTE (15).

The most common signs were dyspnea at first and tachycardia in the second grade. Although these symptoms can be completely similar in other illnesses such as heart disease or bleeding, but it indicates that these symptoms should consider in pregnant women. On the other hand, these two signs seem to be the most endearing signs of the disease, and in order to prevent maternal mortality, it should be determined by accurate screening of high-risk mothers at an earlier stage. An X-ray of chest helps to exclude other causes that lead to dyspnea and tachycardia (16).

Therefore, thromboembolism risk assessment should be performed at any time during pregnancy; especially after delivery, or in cases with medical or surgical conditions (12). Identification of these factors is important and may reduce the incidence and improve its timely diagnosis. In the current study we estimate the incidence, risk factors, and mortality from pregnancy-related venous thromboembolism.

Materials And Methods

In this retrospective study cases with thromboembolism coded as the underlying or contributory cause of death in women aged 15–44 from March 2017 to February 2018 were evaluated. We included all women with fatal VTE occurred during pregnancy or within 6 weeks post-partum, according to the definition of maternal death. Medical records, autopsy reports and data from the Ministry of Health were examined in order to detect contributing factors for VTE.

Obstetrical information such as parity, pregnancy complications and delivery complications. was retrieved either. The study protocol was approved by the local ethics committee of Tehran University of Medical Sciences. Patient records extracted in accordance with the ethical and security guidelines of the Ministry of Health and the risk factors of thromboembolism measured using a questionnaire entitled "Risk factors for thromboembolism during pregnancy and puerperium" that is summarized with the scores based on Royal College of Obstetricians & Gynecologist (RCOG) guideline (17) in Table 1. The scores will gather to reach a total value.

Table 1

The evaluation of risk factors for thromboembolism during pregnancy and puerperium

Risk Factors	Score
A) Risk factors related to general conditions	
Body mass index (BMI) equal to or greater than 40 (before or during early pregnancy)	2
Age older than 35 years	1
BMI between 35–40 (before or early pregnancy / weight greater than 80 kg)	1
Parity ≥ 3 regardless of the current pregnancy	1
Smoking	1
Gross varicose veins (symptomatic or above the knee or with phlebitis, edema, skin changes)	1
B) Risk factors related to medical conditions	
History of previous VTE (except for the reason of major surgery)	4
Acquired thrombophilia (antiphospholipid antibody syndrome): at least one laboratory test and at least a clinical criterion	4
History of previous VTE due to major surgery	3
Any medical conditions: cancer, cardiac disease, active lupus, inflammatory poly-arthritis or inflammatory bowel disease, nephrotic syndrome, type 1 diabetes mellitus with nephropathy, tuberculosis, current intravenous drug addiction	3
High risk inherited thrombophilia (deficiency of anti-thrombin, protein S or C deficiency, homozygote low risk thrombophilia or more than one heterozygote low risk thrombophilia)	3
Low risk inherited thrombophilia (heterozygote factor 5 Leiden, G20210A prothrombin gene mutation)	1
Family history of VTE in first degree relatives (spontaneous or estrogen dependent)	1
Anti-phospholipid antibodies (only laboratory criteria, without a clinical criteria)	1
C) Obstetrical Risk Factors	
Cesarean section in labor	2
Current preeclampsia	1
Pregnancy with fertility IVF / ART methods (only considered during pregnancy)	1
Multiple pregnancy	1
Non-emergency cesarean (elective)	1
Instrumental delivery	1
Prolonged labor (more than 24 hours)	1

Risk Factors	Score
Postpartum hemorrhage (greater than 1 liter / blood transfusion)	1
Preterm labor (less than 37 weeks) in the current pregnancy	1
Intrauterine fetal demise (IUFD) in the current pregnancy	1
D) Transient Risk Factors	
Ovarian hyper stimulation syndrome (only in the first trimester)	4
Surgery during pregnancy or postpartum (curettage, tubal closure, appendectomy..., except for perineal immediate repair)	3
Severe vomiting of pregnancy (hyperemesis)	3
Systemic infections (antibiotics or admission needed) for example: pneumonia, pyelonephritis, postpartum wound infection	1
Hospitalization or immobilization (≥ 3 days rest in bed), dehydration	1

If total score is 4 or more, anticoagulant administration with prophylaxis dose start from the beginning of the pregnancy. In score 3, it is recommended to start anticoagulant with prophylactic dosage from the 28th week of pregnancy. If total score is 2 or more postnatally or admitted to hospital antenatally, consider thromboprophylaxis for at least 10 days. In prolonged hospitalization (≥ 3 days) or readmission to hospital within the puerperium, consider thromboprophylaxis either (17).

Statistical Analysis

Mean (SD) (standard deviation) and percentage of frequency were calculated for descriptive objectives. The data were analyzed by the SPSS version 18.

Results

The maternal mortality rate was 16 per 100,000 live births. Among 297 mortality cases in 1 year, 27 death were due to thromboembolism. The mean (SD) age was 33.2 (3.3) years and the mean of gestational age was 33 weeks + 6 days.

Dyspnea (88.8%) and tachycardia (18.5%) were the common manifestations in these patients. The diagnosis of all cases was confirmed by autopsy. Demographic and clinical data of the cases are listed in Table 2.

Table 2
Demographic and clinical data of the cases deceased due to thromboembolism

variable		N (%)	
Type of hospital	Educational	21 (77.8)	
	Private	5 (18.5)	
	Charity	1 (3.7)	
BMI*	< 30	23 (85.2)	
	30–40	4 (14.8)	
	≥ 40	0	
Blood Group	A ⁺	11 (40.7)	
	O ⁺	8 (29.6)	
	B ⁺	5 (18.5)	
	AB ⁺	2 (7.4)	
	O ⁻	1 (3.7)	
	A ⁻	0	
	B ⁻	0	
	AB ⁻	0	
Delivery Type	NVD **	6 (22.2)	
	CS ***	Emergency	7 (25.9)
		Elective	14 (51.9)
Parity	≥ 3	10 (37)	
	< 3	17 (63)	
Infant gender	Male	14 (51.9)	
	Female	13 (48.1)	
1st min. Apgar Score	0	6 (22.2)	
	5	1 (3.7)	
	6	1 (3.7)	

* body mass index, **normal vaginal delivery, ** cesarean section, **** intra uterine fetal demise, ***** in vitro fertilization/ assisted reproductive technology

variable		N (%)
	7	4 (14.8)
	8	3 (11.1)
	9	12 (44.4)
Death time	Morning shift	18 (66.7)
	Evening shift	9 (33.3)
Delivery to death (days)	1	20 (74.1)
	2	3 (11.1)
	3	1 (3.7)
	4	1 (3.7)
	7	1 (3.7)
	12	1 (3.7)
Blood transfusion	Yes	11 (40.7)
	No	16 (59.3)
Heparin prescription	Yes	Single dose multiple doses
		6 (22.2) 5 (18.5)
	No	16 (59.3)
Time of Heparin prescription	Before surgery	5 (18.5)
	After surgery	1 (3.7)
	Before & after surgery	5 (18.5)
	None	16 (59.3)
Preterm labor	Yes	13 (48.1)
	No	14 (51.9)
IUFD****	Yes	7 (25.9)
	No	20 (74.1)
Postpartum hemorrhage	Yes	7 (25.9)
	No	20 (74.1)

* body mass index, **normal vaginal delivery, ** cesarean section, **** intra uterine fetal demise, ***** in vitro fertilization/ assisted reproductive technology

variable		N (%)
Prolonged labor	Yes	1 (3.7)
	No	26 (96.3)
Instrumental delivery	Yes	2 (7.4)
	No	25 (92.6)
Sepsis	Yes	3 (11.1)
	No	24 (88.9)
Multiple pregnancy	Yes	3 (11.1)
	No	24 (88.9)
IVF/ART****	Yes	4 (14.8)
	No	23 (85.2)
Antiphospholipid syndrome	Yes	1 (3.7)
	No	26 (96.3)
Cardiac Disease	Yes	8 (29.6)
	No	19 (70.4)
Hypothyroidism	Yes	1 (3.7)
	No	26 (96.3)
Cancer	Yes	1 (3.7)
	No	26 (96.3)
Diabetes	Yes	4 (14.8)
	No	23 (85.2)
Rheumatoid disease	Yes	1 (3.7)
	No	26 (96.3)
Hereditary thrombophilia	Yes	1 (3.7)
	No	26 (96.3)
Preeclampsia	Yes	1 (3.7)
	No	26 (96.3)

* body mass index, **normal vaginal delivery, ** cesarean section, **** intra uterine fetal demise, ***** in vitro fertilization/ assisted reproductive technology

variable		N (%)
Chronic hypertension	Yes	3 (11.1)
	No	24 (88.9)
Thromboembolism history	Yes	1 (3.7)
	No	26 (96.3)
* body mass index, **normal vaginal delivery, ** cesarean section, **** intra uterine fetal demise, ***** in vitro fertilization/ assisted reproductive technology		

There were no any case of family history for thromboembolism, hospitalized or immobilized (≥ 3 days rest in bed), dehydration, ovarian hyperstimulation syndrome, surgery during pregnancy or postpartum, hyperemesis gravidarum, nephropathic disorders, smoking, intravenous drug addiction, inflammatory bowel disease and sickle cell disease among these patients.

The mean (SD) of thromboembolism score was 4.6 (1.9). Nine patients had score < 4 and 18 patients ≥ 4 . Heparin prescription frequencies based on different reasons are listed in Table 3.

Table 3
Heparin prescription frequencies based on different reasons

variable	Heparin ⁺		Heparin ⁻	
		N (%)		N (%)
Total score < 4		2 (22.2)		7 (77.8)
Total score ≥ 4		9 (50.0)		9 (50.0)
Death in morning shift		2 (22.2)		7 (77.8)
Death in night shift		9 (50.0)		9 (50.0)
Normal Vaginal Delivery		3 (50.0)		3 (50.0)
Cesarean Section		8 (38.1)		13 (61.9)
Body Mass Index < 30		11 (40.7)		12 (59.3)
Body Mass Index ≥ 30		0		4 (100)
Parity ≥ 3		4 (40.0)		6 (60.0)
Parity < 3		7 (42.2)		10 (58.8)
Cardiac Disease	Yes	4 (50.0)		4 (50.0)
	No	7 (36.8)		12 (63.2)
Preterm Labor	Yes	6 (46.2)		7 (53.8)
	No	5 (36.8)		9 (64.3)

The rate of mortality with the history of VTE, anti-phospholipid syndrome and inherited thrombophilia was one in each group and all 3 received heparin as prophylaxis. There were 3 cases with sepsis and none of them got heparin.

Discussion

In this study, the risk factors for thromboembolism was investigated in 27 mortality due to mentioned reason. The mean gestational age was 33 weeks + 6 days and nearly half of the cases experienced premature births. This is the same reason as Simpson et al. study that reported delivery at gestational age < 36 as a risk factor for VTE (18).

The use of risk score calculator may effectively and safely determine the duration of heparin therapy (17). Darguad et al. indicated that in patients with moderate risk of VTE (score = 3–5), prescription of heparin prophylaxis in the third trimester prevent thrombosis (19). In our study, the mean score of the

thromboembolism was 4.6 suggesting that these cases were a high-risk population for thromboembolic events; however only half of them with score more than 4 received heparin.

Although the age of 35 years in pregnant women is associated with an increased risk of thromboembolic events (17), in this study the average age of the cases was 33.2 years that may reveal the impact of other variables to increase the risk of thromboembolism. Although Danilenko-Dixon et al. also reported no correlation between age and VTE in pregnancy and puerperium period (20).

More than 70% of cases have embolized in first day after delivery and about one half of them deceases intrapartum. Since more than half of them did not even receive a single dose of heparin, the need to quickly and accurately identify the indications of prophylactic treatment with heparin or other preventive methods to reduce mortality in these mothers is obvious.

Available data suggests that the risk of VTE is higher after cesarean section than vaginal delivery (21). The indications for cesarean and the conditions during/after the surgery could be involved in the occurrence of thromboembolism and possibly mortality. The puerperium is the time of maximal risk of pregnancy-associated VTE and several observational studies have assessed the risk of VTE after cesarean Sect. (22–24).

Blondon et al. accounted maternal Body mass index (BMI) as an important risk factor for postpartum VTE, grading from weak in overweight women to very strong in women with class III obesity (25). However, in the other study, BMI was not related to VTE. This is similar to our result that BMI was in few patients above or equal to 30 and in none of the mortalities more than 40.

Women with inherited thrombophilia or a positive family history even with no previous episode of VTE, have a risk of developing a first VTE episode in pregnancy or postpartum period (26). The highest risks were associated with homozygosity for factor V Leiden or the prothrombin G20210A variant (27). In the present survey, one case had a history of VTE, one with an anti-phospholipid antibody syndrome, and the other with hereditary thrombophilia that all received heparin. Thromboembolism happened may be due to not receiving proper dosage of heparin.

Low molecular heparin is the preferred option for prophylaxis in most patients because of its better bioavailability, longer plasma half-life, more predictable dose response, and improved maternal safety (28, 29). Also heparin is more effective than vitamin K antagonists in preventing recurrent VTE and post-thrombotic syndrome without increasing the risk of major bleeding events (30).

In our study, the most frequent blood group of mothers was A +. The study of Mirza Aghazadeh et al. also had the highest reports of VTE associated with the A + blood group (31); Although this blood group generally has a high prevalence in Iran (32). On the other population-based study blood groups A and AB were associated with increased risk estimates for VTE in pregnancy and the puerperium either (33).

37% of mothers had more than 3 times delivery, 30% had heart disease, 21% had high blood pressure and 25% had stillbirth, it seems that these factors have a significant effect on the risk of maternal conditions.

Frequency of multiple pregnancy, severe bleeding, instrumental delivery, pregnancy by assisted reproductive techniques, prolonged delivery, sepsis, preeclampsia, and underlying illness were observed in less than 30 percent of these 27 mothers and showed that although these factors are predisposed to the occurrence of thrombotic events, but they do not contribute much to our study.

There are studies on the prevalence of risk factors and etiology of maternal mortality that explain rupture of membrane and severe bleeding are important risk factors in thromboembolism; therefore, precise follow-up and, if necessary, preventive and interventional proceedings may be effective in preventing maternal deaths.

Conclusion

The study concludes that a high proportion of maternal mortality following thromboembolism are potentially preventable. It is suggested that pregnant women should be carefully evaluated for these risk factors, and that health care personnel and pregnant mothers should be trained to diagnose the early symptoms, as well as prevention and treatment of thromboembolic events.

The use of more accurate diagnostic methods for rapid diagnosis and control of the disease, as well as postpartum care, particularly in high-risk individuals, including women over 35 years of age, a history of more than three times pregnancy, a history of VTE, and the administration of prophylactic heparin with a sufficient dose and duration, decrease maternal mortality.

Abbreviations

VTE

Venous Thromboembolism

BMI

Body Mass Index

NVD

Normal Vaginal Delivery

CS

Cesarean Section

IUFD

Intra Uterine Fetal Demise

IVF/ART

In Vitro Fertilization/ Assisted Reproductive Technology

SD

Standard Deviation

Declarations

Ethics approval and consent to participate:

This manuscript was performed in accordance with Helsinki declaration. All patient's data were kept confidential. This study was approved by Tehran University of Medical Science's ethical committee.

Consent to publish:

The identity of the participants had not transpired.

Availability of data and materials:

The datasets used during the current study are available from the corresponding author on reasonable request. They are divided in two group. The data of the patients that declared in the article and are available with more detail by corresponding author and can be sent by her.

Competing interests: Not Applicable

Funding: Not Applicable

Authors' contributions:

M.S: Design of the work

M.G: Design of the work, Drafting the manuscript

B.S: Manuscript editing, Interpretation of data

M.T: Manuscript editing, Interpretation of data

E.L: Interpretation of data

All authors approved the submitted version (and any substantially modified version that involves the author's contribution to the study.

All authors are agreed both to be personally accountable for the author's own contributions and to ensure that questions related to the accuracy or integrity of any part of the work, even ones in which the author was not personally involved, are appropriately investigated, resolved, and the resolution documented in the literature.

Conflict of interest

None to be declared.

Acknowledgment

This research has been part of the medical doctor's thesis that was supported by Tehran University of Medical Sciences.

References

1. MacDorman MF, Declercq E, Thoma ME. Making Vital Statistics Count: Preventing U.S. Maternal Deaths Requires Better Data. *Obstet Gynecol.* 2018;131(5):759-61.
2. Sarani M, Shahraki Z, Shirazi M, Saravani S. Risk factors of maternal mortality in Sistan region: 10-year report. *Tehran University Medical Journal.* 2014;72(9).
3. Zalvand R, Tajvar M, Pourreza A, Asheghi H. Determinants and causes of maternal mortality in Iran based on ICD-MM: a systematic review. *Reprod Health.* 2019;16(1):16.
4. Khan KS, Wojdyla D, Say L, Gulmezoglu AM, Van Look PF. WHO analysis of causes of maternal death: a systematic review. *Lancet.* 2006;367(9516):1066-74.
5. Kuriya A, Piedimonte S, Spence AR, Czuzoj-Shulman N, Kezouh A, Abenhaim HA. Incidence and causes of maternal mortality in the USA. *J Obstet Gynaecol Res.* 2016;42(6):661-8.
6. Moazzeni MS. Maternal mortality in the Islamic Republic of Iran: on track and in transition. *Matern Child Health J.* 2013;17(4):577-80.
7. Shirazi M RS-BFMS. Rupture of an unscarred uterus in a nullipar pregnant women with a history of curettage: case report. *Tehran University Medical Journal.* 2015;73(2):138-42.
8. Shirazi M, Hantoush-Zadeh S, Rezaie-Keikhaie K, Pirjani R. Spontaneous Uterine Rupture and Live Fetus in 21th Week of Pregnancy with Hemorrhagic Shock Due to Placenta Percreta: A Case Report. *Case Reports in Clinical Practice.* 2016;1(1):19-21.
9. Shirazi M, Zaban MT, Gummadi S, Ghaemi M. Peptic ulcer perforation after cesarean section; case series and literature review. *BMC Surg.* 2020;20(1):110.
10. St Pierre A, Zaharatos J, Goodman D, Callaghan WM. Challenges and Opportunities in Identifying, Reviewing, and Preventing Maternal Deaths. *Obstet Gynecol.* 2018;131(1):138-42.
11. James AH, Jamison MG, Brancazio LR, Myers ER. Venous thromboembolism during pregnancy and the postpartum period: incidence, risk factors, and mortality. *American Journal of Obstetrics & Gynecology.* 2006;194(5):1311-5.
12. Bates SM, Middeldorp S, Rodger M, James AH, Greer I. Guidance for the treatment and prevention of obstetric-associated venous thromboembolism. *Journal of thrombosis and thrombolysis.* 2016;41(1):92-128.
13. Kourlaba G, Relakis J, Kontodimas S, Holm MV, Maniadas N. A systematic review and meta-analysis of the epidemiology and burden of venous thromboembolism among pregnant women. *International Journal of Gynecology & Obstetrics.* 2016;132(1):4-10.
14. Lindqvist P, Dahlbäck B, Maršál K. Thrombotic risk during pregnancy: a population study. *Obstetrics & Gynecology.* 1999;94(4):595-9.

15. Alguel G, Vormittag R, Simanek R, Kyrle PA, Quehenberger P, Mannhalter C, et al. Preeclampsia and pregnancy loss in women with a history of venous thromboembolism and prophylactic low-molecular-weight heparin (LMWH) during pregnancy. *Thrombosis and haemostasis*. 2006;96(03):285-9.
16. Bagaria SJ, Bagaria V. Strategies for diagnosis and prevention of venous thromboembolism during pregnancy. *Journal of pregnancy*. 2011;2011.
17. Reducing the Risk of Venous Thromboembolism during Pregnancy and the Puerperium [Internet]. RCOG. 2015.
18. Simpson E, Lawrenson R, Nightingale A, Farmer R. Venous thromboembolism in pregnancy and the puerperium: incidence and additional risk factors from a London perinatal database. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2001;108(1):56-60.
19. Dargaud Y, Rugeri L, Vergnes MC, Arnuti B, Miranda P, Negrier C, et al. A risk score for the management of pregnant women with increased risk of venous thromboembolism: a multicentre prospective study. *British journal of haematology*. 2009;145(6):825-35.
20. Danilenko-Dixon DR, Heit JA, Silverstein MD, Yawn BP, Petterson TM, Lohse CM, et al. Risk factors for deep vein thrombosis and pulmonary embolism during pregnancy or post partum: a population-based, case-control study. *American Journal of Obstetrics & Gynecology*. 2001;184(2):104-10.
21. Greer IA. Thrombosis in pregnancy: maternal and fetal issues. *The Lancet*. 1999;353(9160):1258-65.
22. Bates SM, Greer IA, Middeldorp S, Veenstra DL, Prabulos A-M, Vandvik PO. VTE, thrombophilia, antithrombotic therapy, and pregnancy: antithrombotic therapy and prevention of thrombosis: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines. *Chest*. 2012;141(2):e691S-e736S.
23. Jacobsen A, Skjeldestad F, Sandset P. Ante- and postnatal risk factors of venous thrombosis: a hospital-based case-control study. *Journal of Thrombosis and Haemostasis*. 2008;6(6):905-12.
24. Jacobsen AF, Drolsum A, Klow NE, Dahl GF, Qvigstad E, Sandset PM. Deep vein thrombosis after elective cesarean section. *Thrombosis research*. 2004;113(5):283-8.
25. Blondon M, Harrington L, Boehlen F, Robert-Ebadi H, Righini M, Smith N. Pre-pregnancy BMI, delivery BMI, gestational weight gain and the risk of postpartum venous thrombosis. *Thrombosis research*. 2016;145:151-6.
26. Friederich PW, Sanson B-J, Simioni P, Zanardi S, Huisman MV, Kindt I, et al. Frequency of pregnancy-related venous thromboembolism in anticoagulant factor-deficient women: implications for prophylaxis. *Annals of internal medicine*. 1996;125(12):955-60.
27. Wu O, Robertson L, Twaddle S, Lowe GD, Clark P, Greaves M, et al. Screening for thrombophilia in high-risk situations: systematic review and cost-effectiveness analysis. The Thrombosis: Risk and Economic Assessment of Thrombophilia Screening (TREATS) study. *Health Technol Assess*. 2006;10(11):1-110.
28. Sanson B-J, Lensing AW, Prins MH, Ginsberg JS, Barkagan ZS, Lavenne-Pardonge E, et al. Safety of low-molecular-weight heparin in pregnancy: a systematic review. *Thrombosis and haemostasis*.

1999;81(05):668-72.

29. Lepercq J, Conard J, Borel-Derlon A, Darmon JY, Boudignat O, Francoual C, et al. Venous thromboembolism during pregnancy: a retrospective study of enoxaparin safety in 624 pregnancies. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2001;108(11):1134-40.
30. Kearon C, Akl EA, Comerota AJ, Prandoni P, Bounameaux H, Goldhaber SZ, et al. Antithrombotic therapy for VTE disease: antithrombotic therapy and prevention of thrombosis: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2012;141(2):e419S-e96S.
31. Mirza Aghazadeh M, Pirzadeh A, Hariri AM. Correlation between Blood Groups and Venous Thromboembolism in Hospitalized Patients in Ardabil University of Medical Sciences Affiliated Centers. *Journal of Ardabil University of Medical Sciences*. 2014;14(1):71-8.
32. Keramati MR, Shakibaei H, Kheiyami MI, Ayatollahi H, Badiei Z, Samavati M, et al. Blood group antigens frequencies in the northeast of Iran. *Transfusion and apheresis science*. 2011;45(2):133-6.
33. Larsen TB, Johnsen SP, Gislum M, Moller CA, Larsen H, Sorensen HT. ABO blood groups and risk of venous thromboembolism during pregnancy and the puerperium. A population-based, nested case-control study. *J Thromb Haemost*. 2005;3(2):300-4.