Phenotype-guided subpopulation identification from single-cell sequencing data

Duanchen Sun

Oregon Health & Science University

Xiangnan Guan

Oregon Health & Science University

Amy Moran

Oregon Health & Science University

David Qian

Oregon Health & Science University

Pepper Schedin

Oregon Health & Science University

Mu-Shui Dai

Oregon Health & Science University

Alexey Danilov

City of Hope National Medical Center

Joshi Alumkal

University of Michigan

Andrew Adey

Oregon Health & Science University <https://orcid.org/0000-0001-7648-8717>

Paul Spellman

Oregon Health Science University <https://orcid.org/0000-0002-4810-0022>

Zheng Xia (✉ xiaz@ohsu.edu)

Oregon Health & Science University <https://orcid.org/0000-0003-3364-8324>

Article

Keywords: single-cell sequencing, phenotype-to-cell, cell subpopulation, data integration

Posted Date: May 5th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-493014/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Phenotype-guided subpopulation identification from single-cell sequencing data

Keywords: single-cell sequencing; phenotype-to-cell; cell subpopulation; data
integration

Duanchen Sun^{1,2}, Xiangnan Guan^{1,2}, Amy E. Moran^{3,4}, David Z. Qian⁴, Pepper Schedin^{3,4},
Mu-Shui Dai⁵, Alexey V. Danilov⁶, Joshi Alumkal⁷, Andrew Adey⁵, Paul T. Spellman^{1,4,5}, and
Zheng Xia^{1,2,4,8*}

¹ Computational Biology Program, Oregon Health & Science University, Portland, OR USA.

² Department of Biomedical Engineering, Oregon Health & Science University, Portland, OR USA.

³ Department of Cell, Developmental and Cancer Biology, Oregon Health & Science University,
Portland, OR, USA.

⁴ Knight Cancer Institute, Oregon Health & Science University, Portland, OR, USA.

⁵ Department of Molecular and Medical Genetics, Oregon Health & Science University, Portland, OR
USA.

⁶ City of Hope National Medical Center, Duarte, CA, USA.

⁷ Department of Internal Medicine, University of Michigan, Ann Arbor, MI USA

⁸ Department of Molecular Microbiology and Immunology, Oregon Health & Science University,
Portland, OR, USA.

*** Corresponding author:**

Zheng Xia, Ph.D.

Email: xiaz@ohsu.edu, Phone: 503-494-9726

Abstract

Single-cell sequencing yields novel discoveries by distinguishing cell types, states, and lineages within the context of heterogeneous tissues. However, interpreting complex single-cell data from highly heterogeneous cell populations remains challenging. Currently, most existing single-cell data analyses focus on cell type clusters defined by unsupervised clustering methods that cannot directly link cell clusters with specific biological and clinical phenotypes. Here we present Scissor, a novel approach that utilizes disease phenotypes to identify cell subpopulations from single-cell data that are most highly associated with the given phenotype. This “phenotype-to-cell” within a single-step strategy utilizes clinical information collected from bulk assays to identify the most highly phenotype-associated cell subpopulations. When applied to a lung cancer single-cell RNA-seq (scRNA-seq) dataset, Scissor identified a subset of cells associated with worse survival as well as another cell subpopulation associated with *TP53* mutation. Furthermore, in a melanoma scRNA-seq dataset, Scissor discerned a T cell subpopulation with low *PDCD1/CTLA4* and high *TCF7* expressions, which is associated with a favorable immunotherapy response. Scissor also demonstrated its broad applicability and effectiveness in interpreting Facioscapulohumeral muscular dystrophy (FSHD) and Alzheimer’s disease scRNA-seq datasets. Thus, Scissor provides a novel framework to identify biologically and clinically relevant cell subpopulations from single-cell assays by leveraging the wealth of phenotype and bulk-omics datasets.

Introduction

Single-cell sequencing technologies are revolutionizing biomedical research and clinical practice by enabling the comprehensive characterization of cells from complex tissues^{1,2}. In contrast to bulk data that measures the averaged properties of whole tissue, single-cell sequencing allows identifying cell types, states, and lineages of different cell subpopulations in a heterogeneous tissue ecosystem³⁻⁵. To recognize critical subpopulations from single-cell data, the standard approach is to perform unsupervised clustering to define cell clusters, inspect marker genes of each cluster, and assess the enrichment of the marker genes in known cell types and pathways to evaluate the importance for each cell cluster^{6,7}. However, identifying cell subpopulations that drive phenotypes, such as disease stage, tumor metastasis, treatment response, and survival outcome, is of indispensable importance since it will facilitate cell-type targeted therapies as well as prognostic biomarker discovery^{3,8}. Unfortunately, single-cell technology is not practical in large cohorts, and most single-cell experiments involve less than twenty patient samples⁹⁻¹¹, which lacks the statistical power to identify the cell subpopulations driving the phenotype of interest.

Meanwhile, valuable clinical phenotype information is widely available from big data consortia like The Cancer Genome Atlas (TCGA)¹² and the International Cancer Genome Consortium (ICGC)¹³ through a decade-long collection of clinicopathologic annotations. Clinical phenotype information is primarily collected on bulk tissue samples, especially in the form of formalin-fixed-paraffin-embedded (FFPE) samples, which are not feasible for single-cell profiling. Therefore, there is an unmet need to leverage such widely accessible and valuable phenotype information to guide cell subpopulation identification from single-cell data.

To the best of our knowledge, there is no bioinformatics tool that uses external bulk phenotypes to guide the identification of key cell subpopulations in a unified framework for single-cell data analysis. Therefore, in this study, we introduce **Single-Cell Identification of Subpopulations with bulk Sample phenOtype coRrelation (Scissor)**. By leveraging bulk data and

phenotype information, this algorithm automatically selects cell subpopulations from single-cell data that are most responsible for the differences of phenotypes. The novelty of Scissor is that it utilizes phenotype information from bulk data to identify the most highly disease-relevant cell subsets. When we applied Scissor to real datasets, the selected cells exhibited high specificity of association with the phenotype of interest. Our studies suggest that Scissor is a promising tool to explore and interpret single-cell data from a new perspective, which can shed fresh light on disease mechanisms and improve the diagnosis and treatment of diseases.

Results

Overview of phenotype-guided cell subpopulation identification method Scissor.

To utilize bulk data and phenotype information for assisting single-cell data analysis, we developed a novel algorithm, Scissor, to identify cell subpopulations from single-cell data that are most highly associated with the given phenotypes. The workflow of Scissor is shown in Fig. 1. Briefly, the three data sources for Scissor inputs are a single-cell expression matrix, a bulk expression matrix, and a phenotype of interest (Fig. 1a). The phenotype annotation of each bulk sample can be a continuous dependent variable, binary group indicator vector, or clinical survival data. The key step of Scissor is to quantify the similarity between the single-cell data and bulk data through a measurement like the Pearson correlation for each pair of cells and bulk samples. After this, Scissor optimizes a regression model on the correlation matrix with the sample phenotype (Fig. 1b). The selection of the regression model depends on the type of the input phenotype, i.e., linear regression for continuous variables, logistic regression for dichotomous variables, and Cox regression for clinical survival data. Since a small subset of cells could drive the phenotype of interest¹⁴, a sparse penalty and a graph regularization are imposed on the regression model to select similar cells that are important for the given phenotype with high confidence (Fig. 1b). Based on the signs of the estimated regression

coefficients, the cells with non-zero coefficients can be indicated as Scissor positive (Scissor+) cells and Scissor negative (Scissor-) cells, which are positively and negatively associated with the phenotype of interest, respectively (Fig. 1c). The cells with coefficients of zero are indicated as background cells. Furthermore, to control the false associations between single-cell and bulk data, we designed a reliability significance test to determine whether the chosen data is suitable for our phenotype-to-cell associations (Fig. 1d). Finally, the Scissor selected cells will be further characterized in downstream analyses, such as exploration of signature genes and functionally enriched pathways (Fig. 1e).

Scissor captured phenotype-associated cell subpopulations in simulation studies.

We first assessed the performance of Scissor on a series of simulated datasets. The simulation study was designed to test whether Scissor can recover the known phenotype-associated cell subpopulations. To achieve this, we employed Splatter¹⁵ to simulate two phenotype-specific cell subpopulations as the ground truth and one common cell type shared by the two phenotypes as well as the corresponding bulk expression data (Fig. 2a; Methods). In one simulation, the two ground-truth phenotype-specific cell subpopulations were characterized by 21 differentially expressed genes (true DEGs) between them but cannot be distinguished by standard single-cell analysis pipelines since they were located in the same cluster (Fig. 2b, c). Utilizing bulk gene expressions to guide the cell subpopulation identification, we found that Scissor recovered 90.5% and 91.8% of the ground truth cells unique to each phenotype, with low false positive rates of 2.1% and 0.9%, respectively (Fig. 2d, e). There were 22 DEGs between the two cell subpopulations identified by Scissor, including all 21 true DEGs between the two ground-truth phenotype-associated cell subpopulations (Fig. 2c). In contrast, the fold change signals of these 21 true DEGs at the bulk level were weak with values less than 1.1 (Supplementary Fig. 1a). Such improved phenotype-specific gene signal detection will facilitate the reliable downstream interrogation of different cell types and states within a heterogeneous cell population. When

performing this simulation 100 times, Scissor had an average recall of 0.88 and precision of 0.93 in identifying phenotype-associated subpopulations.

Furthermore, to check whether the performance of Scissor depends on the true DEG signals at the bulk level, we reset the expression levels of these true DEGs as the least differentially expressed genes (Methods). We found that Scissor can still identify 87.6% and 87.3% of the ground truth cells for each phenotype, with slightly higher false positive rates of 4.2% and 5.9% (Fig. 2f, g). In this setting, Scissor detected all 21 true DEGs with only two false positive DEGs (Fig. 1c). Finally, we demonstrated the robustness of Scissor on the batch-effects corrected scRNA-seq data (Supplementary Fig. 1 b-e). Overall, the results from these simulations were largely consistent with each other, indicating Scissor's capacity to identify the phenotype-associated cell subpopulations from single-cell data.

Scissor identified cell subpopulations related to the tumor and normal phenotypes from a lung cancer scRNA-seq dataset.

We then applied Scissor to a lung cancer scRNA-seq data that included tumor cells and cells from the tumor microenvironment with known cell types¹⁰ (Fig. 2h). In order to demonstrate the efficacy of our algorithm, we utilized the tumor and normal phenotypes from 577 TCGA lung adenocarcinoma (LUAD) bulk samples¹⁶ to guide Scissor analysis. We expected that, by using these data, Scissor would infer cells that were most highly associated with cancer or normal phenotype in this heterogeneous single-cell dataset. Because of the binary phenotype settings of this application, where samples with a phenotype indicator value 1 correspond to tumor samples, Scissor+ cells should be associated with the cancer cells, and Scissor- cells should be associated with the normal phenotype. Among 29,888 cells from different cell types (Fig. 2h), 361 Scissor+ cells and 534 Scissor- cells were selected by Scissor, which were associated with the tumor and normal phenotypes with high confidence (Fig. 2i). As anticipated, over 98% of Scissor+ cells were verified to be malignant cells, and the remaining were T cells and B cells

(Fig. 2j). Such a high proportion of cancer cells in Scissor+ cells cannot be selected by chance (Fig. 2k, hypergeometric test $p < 2e-16$), which confirmed that Scissor can identify the cells associated with the phenotypes of interest. As for Scissor- cells, the cell types were relatively more balanced than Scissor+ cells since it was designed to associate with more diverse non-malignant cell types (Fig. 2j). Myeloid cells and alveolar cells were the two main selected cell types, which account for 42.3% and 36.9% of total Scissor- cells, respectively. All cell types in Scissor- cells, especially the alveolar cells, are important cell types in normal lung tissue. Thus, we demonstrated that Scissor can precisely identify the most phenotype-associated cells from single-cell data with the guidance of phenotype information from bulk data.

Scissor identified a hypoxic subpopulation of lung cancer cells associated with worse survival.

Cancer cells are heterogeneous and include subpopulations such as cancer stem cells, which are known to drive tumor progression, drug resistance, and poor prognosis^{17, 18}. Therefore, we applied Scissor, guided by 471 TCGA-LUAD bulk samples with survival information¹⁶, to identify aggressive cancer cell subpopulations within 4,102 cancer cells from the same lung cancer scRNA-seq dataset¹⁰. These cells were separated into 12 clusters (Fig. 3a), which demonstrated the heterogeneous nature of the cancer cells. Out of 205 Scissor selected cells, 201 Scissor+ cells were associated with **Worse Survival** (defined as Scissor_WS cells thereafter), and only 4 Scissor- cells were associated with good survival (Fig. 3b). The Scissor_WS cells were mainly from clusters 1 and 3 (Fig. 3c). To understand the underlying transcriptional patterns of Scissor_WS cells, we compared the gene expressions of those cells with all other cells. As a result, 23 up-regulated genes and 205 down-regulated genes were differentially expressed in Scissor_WS cells over all other cells, respectively (Fig. 3d and Supplementary Table 1). Interestingly, we found that multiple important hypoxia-related genes (*GAPDH*, *ENO1*, *TPI1*, *LDHA*, *ALDOA*, *PGK1*, and *FAM162A*) were among the above 23

overexpressed genes (Fig. 3e). Functional enrichment analysis also confirmed that the hypoxia-related pathways, such as glycolysis and glucose metabolism processes, were activated in Scissor_WS cells (Fig. 3f and Supplementary Table 1). Consistently, motif analysis revealed that the *HIF1A* binding motif was the most enriched motif in the 23 up-regulated genes (Supplementary Table 1), which is a key mediator of cellular response to lowered oxygen levels and plays a critical role in adapting cancer cells' metabolism to a hypoxic microenvironment¹⁹. We also observed higher hypoxia pathway activities in an identified cell subpopulation related to higher tumor stages (Supplementary Fig. 2). Therefore, the hypoxic cell subpopulation might drive worse clinical outcomes since oxygen-starved cancer cells have a survival advantage to elevate tumor progression²⁰.

To further examine the clinical relevance of the above 23 overexpressed genes (defined as lung cancer signature; Supplementary Table 1), we chose six independent lung cancer datasets collected in PRECOG²¹ (Supplementary Table 1). We found that in 5 out of 6 datasets, the patients with higher signature scores had significantly worse survival time than the patients with lower signature scores (Fig. 3g and Supplementary Fig. 3). This observation indicated that the lung cancer signature derived from Scissor_WS cells was associated with worse survival and could provide potential drug targets for further investigation²². Among the six chosen datasets, two of them have additional clinical features. We thereby investigated whether our lung cancer signature had predictive power for patient survival beyond clinical features. To achieve this, we examined pathological stage, sex, age at diagnosis, and our lung cancer signature in these two datasets. We found that only the pathological stage and our signature were significantly associated with patient survival in univariate Cox survival analyses (Fig. 3h). Furthermore, our signature remained statistically significant in both datasets after adjusting for tumor stage in multivariable Cox survival analyses (Fig. 3i). These findings suggested that the lung cancer signature has distinct predictive power for patient survival and might serve as a novel biomarker for poor clinical prognosis.

In summary, Scissor identified an aggressive cancer cell subpopulation from the LUAD scRNA-seq data that was associated with worse survival outcomes and can be characterized by the overexpression of hypoxia-related genes. The high hypoxia activity might drive LUAD progression and thereby conferred poor outcomes to patients whose tumors contained significant numbers of such cells.

Scissor determined a cancer cell subpopulation associated with *TP53* mutation.

To further demonstrate the high flexibility of Scissor in exploring a variety of hypotheses of cell states in single-cell data, we used other phenotypical features provided by TCGA-LUAD to guide the identification of cell subpopulations within the same 4,102 lung cancer cells. Here we focused on *TP53*, one of the most commonly mutated tumor suppressor genes found in human malignancies.

We collected *TP53* mutation status (mutant or wild-type) from TCGA-LUAD as the phenotypes of 498 bulk samples. Scissor identified a total of 414 Scissor+ cells associated with the *TP53* mutant and 318 Scissor- cells associated with the wild-type (Fig. 4a). To uncover the transcriptional differences between these cells, we compared the gene expressions of Scissor+ cells with that of Scissor- cells. As a result, 337 up-regulated genes and 14 down-regulated genes were differentially expressed in Scissor+ cells over Scissor- cells (Fig. 4b and Supplementary Table 2). The 337 up-regulated genes include multiple E2F target genes and cell cycle progression-related genes, e.g., *AURKA*, *CDK1*, *CCNB2*, *EZH2*, and *TOP2A* (Fig. 4b). Functional enrichment analysis also confirmed that cell cycle-related pathways, such as hallmark E2F targets and G2M checkpoint pathways, were activated in Scissor+ cells (Fig. 4c and Supplementary Table 2). Consistently, master regulator analysis revealed that the activities of E2F transcription factors (TFs) family members E2F1 and E2F4 were both highly elevated in Scissor+ cells (Fig. 4d). In this analysis, we also found that TFs FOXM1 and MYC, which are known to be repressed by wild-type *TP53*^{23, 24}, were activated in the cell subpopulation

associated with the *TP53* mutant (Fig. 4d). MYC and FOXM1 are known to promote cell cycle progression, and FOXM1 is reported as a major predictor of unfavorable outcomes in human cancers²⁴. These observations are in line with the literature that *TP53* mutation leads to tumorigenesis by involving its impaired capability to arrest cell cycle and maintain DNA repair in response to oncogenic stimuli²⁵. Indeed, master regulator analysis confirmed that TP53 is inactivated in Scissor+ cells (Fig. 4d), which further demonstrates the capability of Scissor in identifying phenotype-associated cell subpopulations.

Furthermore, by linking those 337 up-regulated genes (defined as *TP53* mutation signature; Supplementary Table 2) with clinical outcomes, we demonstrated that the patients with higher *TP53* mutation signature scores had significantly worse survival time than the patients with lower scores (Fig. 4e). This result is consistent with the pro-proliferation roles of cell cycle-related pathways and TFs in cancer progression, which are enriched in those up-regulated genes. For the down-regulated genes, we found that major histocompatibility complex (MHC) class-related genes *HLA-A*, *B2M*, and *CD74* are down-regulated in Scissor+ cells, and only one of them can be directly identified from the bulk level gene expression (Fig. 4f, Supplementary Fig. 4). Pathway enrichment analysis also found lower allograft rejection activity (Fig. 4c), indicating an inadequate immune response in Scissor+ cells. Notably, *B2M* is a critical component of MHC class I antigen presentation. Its function in effective immunotherapy response has been validated in lung cancer mouse models²⁶, and loss-of-function mutations in *B2M* have been reported in cancer patients resistant to immunotherapy²⁷. Thus, our phenotype-guided scRNA-seq analysis implied that *TP53* mutation is likely a mechanism of resistance to checkpoint inhibitor treatment.

In summary, Scissor identified a cell subpopulation associated with *TP53* mutation. These cells were characterized by dysregulated cell proliferation and deficient MHC Class I antigen-presenting functions. Therefore, this application demonstrated the versatile abilities of Scissor in integrating a variety of phenotypes for single-cell subpopulation annotations.

Scissor identified a *PD1* low subpopulation of T cells associated with the response to melanoma immunotherapy.

Immune checkpoint therapy by blocking *CTLA4*, PD-1, and PD-L1 has achieved exciting results in a wide variety of cancers^{28, 29}. However, only a small portion of patients can benefit from current checkpoint inhibition (CPI). Thus, understanding the mechanism underlying CPI response facilitated by single-cell sequencing is critical to improving the efficacy of these therapies. Here, based on the established role of T cells in immunotherapy response, we performed Scissor on a melanoma scRNA-seq dataset to identify a T cell subpopulation related to CPI response.

We focused our analysis on 1,894 T cells from the metastatic melanoma tumor microenvironment³⁰ and collected 70 melanoma bulk patients with known immunotherapy response information from two studies^{31, 32}. In the standard scRNA-seq data analysis, these T cells were clustered into six clusters (Fig. 5a). By performing Scissor, Scissor identified 105 T cells as Scissor+ cells, which were associated with a **Favorable immunotherapy Response** (defined as Scissor_FR cells thereafter), and did not report any cells that are associated with the unfavorable immunotherapy responses (Fig. 5b). These 105 Scissor_FR cells mainly resided in clusters 2 and 3 (Fig. 5c), indicating clusters 2 and 3 were more associated with the effective responses than other clusters. To characterize the transcriptional identities of the Scissor_FR cells, we compared the gene expression of these cells with all other cells. In total, 17 up-regulated and 120 down-regulated differential expression genes were identified in Scissor_FR cells (Fig. 5d and Supplementary Table 3). The Scissor_FR cells that were associated with an effective CPI response had increased expressions of genes linked to T cell memory (*CCR7* and *SELL*) and survival (*IL7R*) as well as lower expressions of inhibitory genes (*HAVCR2*, *LAG3*, *PDCD1*, *CTLA4*) and MHC II class genes (*HLA-DRB5*, *HLA-DRB1*, *HLA-DPA1*, *HLA-DQB2*, and *HLA-DRB6*) (Fig. 5d, e and Supplementary Table 3). The Scissor_FR cells also exhibited enhanced expressions of transcript factor *TCF7* (encoding TCF1) that have

been shown to associate with favorable outcomes in CPI treatment^{33, 34} (Fig. 5e). The first most down-regulated gene in the Scissor_FR cell population was *TOX*, which is involved in CD8 T cell dysfunction³⁵ (Fig. 5e and Supplementary Table 3). Pathway enrichment analysis showed the Scissor_FR cells had higher TNFA signaling and lower activity of *CTLA4*, *PD1* signaling, and LCMV/tumor exhaustion pathways (Fig. 5f).

The above 137 differential expression genes related to effective immunotherapy response (defined as immunotherapy responsive signature; Supplementary Table 3) could be informative in predicting treatment success. We found that our signature scores were significantly higher in CPI responders than in non-responders in an independent CPI dataset (Fig. 5g, Student's *t*-test $p=5.0e-4$). Additionally, the up-regulated and down-regulated genes in our immunotherapy responsive signature were also significantly enriched in responders and non-responders (Fig. 5h, Kolmogorov-Smirnov test FDR=0.002 and 0.0085, respectively). Since there is a strong relationship between immunotherapy response and T cell states³⁶, we thereby evaluated our immunotherapy responsive signature in five types of tumor-infiltrating lymphocytes (TILs) with distinctive differentiation states. We found that LAG3-low/PD1-low effector CD8 T cells had the highest signature scores, followed by Naïve CD8, Bystander TIL, LAG3-high/PD1-high effector CD8, and exhausted CD8 T cells (Fig. 5i). Notably, our signature can significantly distinguish between PD1-low effector CD8, PD1-high effector CD8, and exhausted CD8 T cells (Fig. 5i and Supplementary Fig. 5a, b). We further performed a pseudotime analysis using our immunotherapy responsive signature on the original melanoma scRNA-seq dataset (Fig. 5a). The pseudotime trajectories revealed the relative orders of the six clusters as cluster 2->3->4->5->6->1 (Supplementary Fig. 5c). This result was consistent with the observation that the majority of Scissor_FR cells came from clusters 2 and 3 with low *PDCD1* expressions (Fig. 5c). Moreover, the PD1-low TILs also exhibit heterogeneity, containing memory-precursor-like and effector-like subsets³⁷. We found that our signature was more enriched in the memory precursor CD8 T cells than the short-lived effector T cells (Fig. 5j, Student's *t*-test $p=0.02$). This

observation indicated that Scissor_FR cells were more like PD1-low memory-precursor cells, which have higher *TCF7* expression and are associated with a good immunotherapy response.

Collectively, our Scissor analysis of a scRNA-seq melanoma dataset independently revealed a *PDCD1/CTLA4* low and *TCF7* high T cell subpopulation whose distinct transcriptome was essential to favorable response to immunotherapy. These results demonstrated that Scissor analysis of single-cell data is capable of identifying subpopulations associated with the specific phenotype even though the single-cell data itself has no such phenotype information.

Scissor identified a subpopulation of FSHD cells that was characterized by dysfunction of fibrinolysis, myogenesis, and muscle contractions.

We further applied Scissor on an Facioscapulohumeral muscular dystrophy (FSHD) single-cell dataset to explore the applicability of our method on non-cancer studies. In total, 6,899 cells derived from FSHD and healthy control samples³⁸ were analyzed by Scissor with the guidance of 35 bulk muscle biopsies (27 FSHD patients and 8 controls). These cells were initially grouped into 14 clusters in scRNA-seq data analysis (Fig. 6a). After integrating these cells and bulk data by Scissor, 579 cells were identified as Scissor+ cells, which were associated with the FSHD; and 74 cells marked as Scissor- cells were linked with the normal phenotype (Fig. 6b).

Consistent with the encoded phenotype information for Scissor in this application, 559 out of 579 Scissor+ cells (97.5%) originated from FSHD patients, and nearly 80% of Scissor- cells were from normal samples (Fig. 6c), indicating that Scissor can identify phenotype-associated subpopulations with high specificity in non-cancer cases.

Given the adequate number of the Scissor selected cells, we could directly compare the gene expression of Scissor+ cells with Scissor- cells to uncover the underlying transcriptional patterns of the identified cells with distinct phenotypes. As a result, 299 up-regulated genes and 83 down-regulated genes were differentially expressed in Scissor+ cells over Scissor- cells, respectively (Fig. 6d and Supplementary Table 4). Among these genes, we found that many

down-regulated genes in Scissor+ cells are involved in normal muscle functions (Fig. 6e), like myogenesis (*LSP1*, *MEF2C*, *MYL1*), muscle contraction (*IGFBP5*, *MYBPC1*, *MYOM1*), and actin filament or myosin-based functions (*PDLIM3*, *TTN*, *SGCD*). As for the up-regulated genes in Scissor+ cells, we found a proportion of these genes may be dysregulated due to fibrotic infiltration³⁹, which includes extracellular matrix proteins as collagen types III and IV (*COL3A1*, *COL4A1*) and fibronectin (*FN1*). FSHD region gene 1 (*FRG1*) was also over-expressed in Scissor+ cells (Fig. 6e). *FRG1* is crucial for angiogenesis and epithelial to mesenchymal transition⁴⁰ and has been reported as a candidate gene for FSHD⁴¹. The functional enrichment analysis also confirmed that Scissor+ cells were characterized by the reduction of muscle fibers as well as loss of myogenesis and muscle contraction functions (Fig. 6f and Supplementary Table 4), which is consistent with the molecular hallmarks of FSHD^{42, 43}. To further demonstrate the characteristics of the phenotype-associated cells identified by Scissor, we built an FSHD molecular signature using the 382 differentially expressed genes between the two cell subpopulations (Supplementary Table 4) and examined the signature's ability in distinguishing FSHD patients from normal controls. We found that on all three independent datasets, the enrichment scores of the FSHD molecular signature were significantly higher in FSHD patients than in normal controls (Fig. 6g, Student's *t*-test $p=5.44e-3$, $3.46e-2$, and $4.99e-2$, respectively).

Therefore, our Scissor analysis of FSHD scRNA-seq data identified disease-associated subpopulations characterized by the dysfunction of fibrinolysis, myogenesis, as well as muscle contractions, which substantiates the utility of Scissor on non-cancer studies.

Scissor identified several cell subpopulations highly associated with Alzheimer's disease.

Due to the prominent heterogeneity of cells in mammalian nervous systems, single-cell technology presents immense opportunities to disentangle cellular diversity and reveal alterations in neurological disorders. Here, we applied Scissor on three brain cell types from an

Alzheimer's disease (AD) scRNA-seq study⁴⁴ to explore the cell subpopulations that are most highly associated with the disease status and progression. Collectively, 7,432 oligodendrocytes, 1,078 oligodendrocyte progenitor cells (OPCs), and 2,171 astrocytes (Supplementary Fig. 6a) were analyzed separately by Scissor with the guidance of 14 bulk samples (7 AD patients and 7 healthy controls). For oligodendrocytes, Scissor identified 206 Scissor+ cells associated with the AD patients and 194 Scissor- cells associated with the healthy controls (Fig. 7a). When compared with the cell sources, 203 out of 206 Scissor+ cells (99%) were from AD patients, and 86% of Scissor- cells were from normal samples (Fig. 7b). With respect to OPCs, 20 Scissor+ cells and 201 Scissor- cells were identified (Fig. 7c), with 90% Scissor+ cells from AD patients and 95% Scissor- cells from normal samples (Fig. 7d). As for astrocytes, the identified cell subpopulations consisted of 179 Scissor+ cells and 14 Scissor- cells (Fig. 7e), with 83% Scissor+ cells from AD patients and 93% Scissor- cells from normal samples, respectively (Fig. 7f). Therefore, Scissor successfully associated cell subpopulations with the desired phenotype in all three brain cell types, demonstrating its ability to identify phenotype-associated cells guided by phenotypes from bulk data.

We then investigated the differentially expressed genes between Scissor+ cells and Scissor- cells within each cell type (Fig. 7g; Supplementary Fig. 6b and Supplementary Table 5). In oligodendrocytes and OPCs, multiple heat shock and chaperone genes involving proper protein folding (e.g., *HSPA1A*, *HSPB1*, *DNAJB1*, and *CRYAB*) were up-regulated in Scissor+ cells, consistent with the previous observation⁴⁵. Pathway enrichment analysis confirmed that the HSF1 pathway, a master activator of chaperone gene expression⁴⁶, is highly activated in Scissor+ cells to protect cells against protein misfolding (Fig. 7h and Supplementary Table 5). In astrocytes, most differentially expressed genes were down-regulated in Scissor+ cells, including *GRIA2*, a glutamate receptor, which increases NMDA receptor activity⁴⁷. Pathway enrichment results also suggested the impaired NMDA-dependent long-term potentiation (LTP) in Scissor+ astrocytes (Fig. 7h and Supplementary Table 5), and LTP deficit has been observed in AD⁴⁸.

350 Additionally, glial fibrillary acidic protein (*GFAP*) was coordinately up-regulated in Scissor+ cells
351 across all three cell types (Fig. 7g). It has been reported that *GFAP* expressions correlate with
352 amyloid- β (A β) plaque density in AD brain tissue⁴⁹, and elevated *GFAP* levels in the blood
353 positively correlate with cognitive impairment⁵⁰. All these results showed that Scissor indeed
354 identified cell subpopulations with characteristics consistent with the corresponding bulk
355 phenotype.

356 To further demonstrate the phenotypic associations of the cell subpopulations identified by
357 Scissor, we constructed molecular signatures based on the differentially expressed genes in
358 Scissor-identified cell subpopulations (Supplementary Table 5) and used another independent
359 AD dataset to evaluate the functions of these signatures. As a result, the enrichment scores of
360 the corresponding molecular signatures in each cell type were significantly higher in AD patients
361 than in normal controls (Fig. 7i-k, Student's *t*-test $p=2.66\text{e-}4$ for oligodendrocytes, $5.43\text{e-}5$ for
362 OPCs, and $5.45\text{e-}7$ for astrocytes). Intriguingly, we also found that our astrocytes molecular
363 signature is positively associated with the progression of AD from incipient, moderate, to severe
364 conditions (Fig. 7l, Kruskal-Wallis test $p=1.44\text{e-}2$). Thus, this Scissor+ cell subpopulation of
365 astrocytes could play a vital role in the early stages of AD, and functional alterations of this
366 subpopulation could lead to neurodegeneration.

367 Taken together, our Scissor analysis identified cell subpopulations that are most highly
368 associated with AD in three brain cell types, which could contribute to comprehending the
369 underlying pathogenesis of Alzheimer's disease and might facilitate disease diagnosis and
370 therapy.

Discussion

Identifying the phenotype-specific cell subpopulations from single-cell data can give rise to breakthroughs in understanding disease mechanisms. Multiple efforts have been made to discern disease-relevant cells from single-cell data⁵¹. For example, HoneyBadger⁵² and inferCNV⁵³ can identify cancerous cells by predicting copy number variations from scRNA-seq data. However, there are a variety of external phenotypes beyond tumor-vs-normal in cancer and non-cancer diseases, such as treatment resistance, cancer metastasis, survival outcomes, disease stages, and aging, which are widely available in bulk data. Therefore, there is a great need for further methodological progress to utilize these abundant phenotypes in single-cell data analysis. To this end, we introduce Scissor as a new computational tool to leverage the phenotype information from bulk data to identify the most highly phenotype-associated cell subpopulations.

We demonstrated the broad utilities of Scissor in a total of ten applications across different diseases, ranging from engineered simulations, cancer cells, immune cells, and FSHD cells to multiple types of brain cells in AD (Supplementary Table 6). The advantages of Scissor can be briefly summarized as follows: First, the phenotype-associated cells identified by Scissor exhibit distinct molecular properties, which involve crucial marker genes and biological processes of the given phenotypes. Second, Scissor does not require any unsupervised clustering on single-cell data, which avoids subjective decisions of cell cluster numbers or clustering resolution⁵⁴. Finally, Scissor provides a flexible framework to integrate various external phenotypes in bulk data to guide single-cell data analysis, enabling hypothesis-free identification of clinically and biologically relevant cell subpopulations.

The signs of the regression coefficients were used to infer the relation of the Scissor selected cells with the phenotypes. In real applications, we observed little proportions of negative correlations between single-cell and bulk samples (Supplementary Table 9). Based on our designed generalized criterion for proper interpretation of Scissor results (Supplementary

Table 12 and Methods), those little proportions of negative correlations did not affect our current results and interpretations. Besides, we tested Scissor's performance in various perturbed bulk samples and demonstrated that the Scissor identification results were robust to noise (Supplementary Fig. 7-9) and had no model-driven preferences to experimental noise or bias (Supplementary Fig. 11 and 12). For computational efficiency, the running time and memory usages of Scissor in real applications are acceptable, with cell numbers ranging from ~1,000 to ~30,000 (Supplementary Table 13).

The construction of the correlation matrix is a key step in Scissor to quantify the dependence or similarity between the single-cell data and bulk data. Except for the Pearson correlation applied in Scissor, we will explore other similarity measurements like entropy-based mutual information and hypothesis testing-based methods. In the future, we also plan to improve the performance and interpretability of Scissor by directly modifying its optimization model to simultaneously identify the phenotype-associated cell subpopulations and the corresponding genes that drive these associations. Furthermore, although we only conducted the integration of bulk gene expressions with scRNA-seq data, Scissor can also be applied to other single-cell measurements like chromatin accessibility⁵⁵ and DNA methylation⁵⁶. For example, Scissor may help identify the cell subpopulations based on chromatin accessibility by integrating single-cell ATAC-seq of cancer cells and immune cells with bulk ATAC-seq data from TCGA⁵⁷.

Overall, Scissor demonstrates promise for integrating single-cell data with phenotype information to dissect clinically significant subsets from heterogeneous cell populations. This strategy will boost biological discoveries and interpretation from a novel perspective in single-cell data analysis. We anticipate that Scissor will enable a broad application of widely available phenotype information on single-cell data analysis and help unravel the most disease-relevant subpopulations for cell-targeted therapies.

Methods

Phenotype-guided single-cell subpopulation identification by Scissor

The workflow of Scissor is shown in Fig. 1. Denote m and n as the cell number and the bulk sample number, respectively. The three data sources for Scissor inputs are a single-cell expression matrix with m cells, a bulk profiling data with n samples, and a sample phenotype Y of interest (Fig. 1a). Y annotates each bulk sample and can be a continuous dependent variable, binary group indicator vector, or clinical survival data. Scissor first uses quantile normalization on the single-cell and bulk expression data to remove the underlying batch effect. After this, a Pearson correlation matrix $S = (s_{ij})_{n \times m}$ is calculated for each pair of cells and bulk samples to quantify the similarity between the single-cell data and bulk data, where s_{ij} is the correlation of sample i and cell j across common genes in normalized single-cell and bulk data.

Scissor optimizes a regression model on the correlation matrix S with the sample phenotype Y (Fig. 1b). Let β denote a vector of coefficients on cells, and $l(\beta)$ denote an appropriately chosen log-likelihood function. The formula of $l(\beta)$ depends on the input phenotype Y , i.e., linear regression for continuous dependent variables, logistic regression (classification) for dichotomous variables, and Cox regression for clinical survival data (see more details in next section). Since a cell subpopulation alone could drive the phenotype of interest, we impose a sparse penalty (l_1 -norm) on the model to select the high confidence cells that are important for the given phenotype. Furthermore, the network-based penalty enforces the tightly connected nodes (cells) in the network to have more similar coefficients⁵⁸, making the phenotype-to-cell association results more consistent and interpretable. Inspired by this, we use the shared nearest neighbor graph calculated in Seurat⁵⁹ to serve as a cell-cell similarity network G and impose a corresponding graph regularization on the regression model.

Overall, Scissor is formulated as the following network regularized sparse regression model (Fig. 1b):

$$\min_{\beta} -\frac{1}{n}l(\beta) + \lambda \left\{ \alpha \|\beta\|_1 + \frac{1-\alpha}{2} \beta^T L \beta \right\},$$

where L is a symmetric normalized Laplacian matrix, which is defined as

$$L = D^{-\frac{1}{2}}(D - A)D^{-\frac{1}{2}} = I - D^{-\frac{1}{2}}AD^{-\frac{1}{2}},$$

where $A = (a_{ij})_{m \times m}$ is a binary or weighted adjacency matrix of G . a_{ij} equals one or a value ranging from 0 to 1 if cell i and j are connected in G , and $a_{ij} = 0$, otherwise. $D = (d_{ij})_{m \times m}$ is the degree matrix of G , where $d_{ii} = \sum_{j=1}^m a_{ij}$, and $d_{ij} = 0$ for $i \neq j$. The tuning parameter λ controls the overall strength of the penalty and α balances the amount of regularization for smoothness and sparsity.

The non-zero coefficients of β solved by the above optimization model are used to select the cell subpopulations associated with the phenotype of interest (Fig. 1c). According to the sign of β , we denote the selected cells by Scissor+ cells and Scissor- cells, which are positively and negatively associated with the phenotype of interest, respectively. The cells with coefficients of zero are denoted as background cells. Furthermore, to control false associations between the single-cell and bulk data, we designed a reliability significance test to determine whether the chosen data is suitable for our phenotype-to-cell associations (Fig. 1d). Finally, the Scissor selected cells will further be investigated by several downstream analyses, such as the differential expression gene analysis, functional enrichment analysis, and motif analysis, to reveal the underlying biological mechanisms of the selected cell subpopulations (Fig. 1e).

Log-likelihood functions

The formula of log-likelihood function $l(\beta)$ depends on the type of the phenotype Y . Scissor can do: (1) linear regression for continuous dependent variables; (2) classification for dichotomous variables; (3) Cox regression for clinical survival data.

In detail, denote $S_i = (s_{i1}, s_{i2}, \dots, s_{im})^T$ as the correlation coefficients for sample i across all m cells. If $Y = (y_1, y_2, \dots, y_n)^T$ is a continuous variable, the linear regression log-likelihood function is used:

$$l(\beta) = - \sum_{i=1}^n (y_i - \beta^T S_i)^2$$

If Y is a binary group indicator vector, e.g., $y_i \in \{0,1\}$, the logistic regression log-likelihood function is used:

$$l(\beta) = \sum_{i=1}^n [y_i \beta^T S_i - \log(1 + \exp(\beta^T S_i))]$$

For time-to-event outcomes subject to independent censoring, the Cox regression is considered. Let T_i be the non-negative event time, and C_i be the censoring time. Denote $\hat{T}_i = \min(T_i, C_i)$ as the observed event time or censoring time, and $\delta_i = I(T_i \leq C_i)$ as the event indicator, where $I(\cdot)$ is an indicator function. The following log-likelihood function is used:

$$l(\beta) = \sum_{i=1}^n \delta_i \left[\beta^T S_i - \log \left(\sum_{k \in R_i} \exp(\beta^T S_k) \right) \right]$$

where $R_i = \{k: \hat{T}_k \geq \hat{T}_i\}$ denotes the risk set at time $\hat{T}_i$.

Parameter tunings and implementations

The algorithm proposed by *Li et al.*⁶⁰ is used to solve the above network regularized sparse regression. In Scissor, two model parameters need to be determined. First, parameter λ controls the overall strength of the penalty. We set 100 possible λ and applied 10-fold cross-validation to select the optimal λ with the minimum averaged error. Second, parameter $\alpha \in [0, 1]$ balances the effect of the l_1 -norm and the network-based penalties. A larger α inclines to emphasize l_1 -norm to encourage sparsity, making Scissor select fewer cells than the result using a smaller α . In real applications, a fixed value of α cannot work for all kinds of datasets since different datasets could have different sensitivities to the changes of α . To select similar

cells as many as possible, we will start to search α from a small value, and the default searching list for α in Scissor is $\{0.005, 0.01, 0.05, 0.1, 0.2, 0.3, \dots, 0.9\}$. The goal of Scissor is to identify a small group of cells that are most highly correlated with the specific phenotypes with high confidence. Based on this motivation as *a priori*, we determined α using the following criteria: the number of Scissor selected cells should not exceed a certain percentage of total cells (default 20%) in the single-cell data. In each experiment, a search on the above searching list is performed from the smallest to the largest until a value of α meets the above criteria.

Reliability significance test

We designed a reliability significance test to exclude the false association between the identified cell subpopulations and bulk phenotypes (Fig. 1d). This statistical test can determine whether the inferred phenotype-to-cell associations are reliable (statistical p-value less than 0.05) or are false positives. Our motivation for the test is: if the chosen single-cell and bulk data are not suitable for the phenotype-to-cell associations, the correlations would be less informative and not well associated with the phenotype labels. Thus, the corresponding prediction performance would be poor and not be significantly distinguishable from the randomly permuted labels. Inspired by this, we used the following procedures to perform the reliability significance test:

First, we performed k-fold cross-validation (CV) on correlation matrix S and used the training sets only to estimate the coefficients of cells in Scissor. The prediction performances of the trained Scissor models were evaluated on testing sets, and an averaged evaluation measurement was obtained to serve as an actual test statistic. Second, we randomly permuted the bulk sample labels multiple times to break up the original bulk phenotype-genotype relationships. By performing the same Scissor analysis and CV evaluation using each permuted bulk data, we obtained a background distribution of the corresponding evaluation measurement quantifying the prediction performances at the random level. Finally, the actual test statistic calculated in the original data was compared to the background distribution values.

The reliability significance test p-value was the number of the permutation-based test statistics above (or below) the actual test statistic, divided by the permutation times. In this study, the evaluation measurements used in the reliability significance test are the mean squared error (MSE) for linear regression (smaller is better), the area under the ROC curve (AUC) for classification (higher is better), and the concordance index (c-index) for Cox regression (higher is better).

Then, we explored whether the reliability significance test can effectively detect the ‘false positive’ associations between the single-cell and bulk data by reporting non-significant p-values. We performed two kinds of randomizations on bulk data to generate false positive examples. After Scissor analysis of those randomized bulk data, our reliability significance test reported that the p-values for these counterexamples were all greater than 0.05 (Supplementary Table 8), indicating that these randomized bulk data were not suitable to use in Scissor to identify the corresponding phenotype-associated cells. We also performed the reliability significance test on the main applications used in this study and found that the test p-values were all less than 0.05 (Supplementary Fig. 10), indicating that the inferred phenotype-to-cell associations are reliable.

Scissor selected cells interpretations

Both Scissor+ cells and Scissor- cells are the Scissor selected cells that are most highly associated with the specified phenotypes, corresponding to the cells with the estimated coefficients greater than and less than zero, respectively. The associations between the Scissor selected cells and phenotypes depend on the model in use and should be interpreted in a context-specific manner. For both linear regression and classification models, the initial values in Y will affect the interpretations. The Scissor+ cells will be associated with the phenotypes encoded as a higher value in Y , and Scissor- cells will correspond to the phenotypes encoded as a lower value. For example, in Scissor’s application on the FSHD single-cell dataset, if the

FSHD patients are encoded as one, and the normal controls are encoded as zero in Y , Scissor+ cells will be associated with the FSHD, and Scissor- cells will be associated with the normal phenotype. If the encoding for the two phenotypes is reversed in Y , the interpretations of Scissor+ cells and Scissor- cells are reversed accordingly. For Cox regression, Scissor+ cells are always associated with worse survival, and Scissor- cells are associated with good survival.

Scissor can associate cells with phenotypes, and this kind of association is a relative concept between phenotypes. Namely, Scissor assigns which phenotype a cell is more likely associated with than the other phenotype. Considering the possible negative correlations between the single-cell and bulk samples, we can further interpret a cell by assigning it to the following two categories: if the average of a cell's correlations with all bulk samples is greater than 0, this cell is more similar to the associated phenotype. Otherwise, this cell should be interpreted as more dissimilar to the other phenotype (Supplementary Table 12). In most cases, the identified cells fall into the "more similar" category, and the negative correlations would not affect the interpretations of Scissor results.

In total, there are ten applications of Scissor in this study. The corresponding phenotype encoding and the interpretations of the Scissor selected cells were summarized in Supplementary Table 6. We showed that Scissor was capable of detecting both Scissor+ and Scissor- cells through a variety of applications. In some datasets, Scissor+ cells and Scissor- cells can be largely unbalanced. Scissor can detect more cells of one phenotype than the other phenotype depending on how strong the cells in a dataset are associated with the specific phenotypes.

Simulation setup

We used Splatter¹⁵ to simulate a single-cell dataset with 1,000 cells and 5,000 expression genes. These cells were from three simulated cell types with group probabilities of 0.8, 0.1, and 0.1, respectively, leading to one large and two small cell subpopulations. The probabilities of a

gene being differentially expressed in each of the three groups were set as 0.1, 0.01, and 0.01. Then, we assigned these cells to two different phenotypes (named phenotype I and phenotype II) to simulate the known relationships between cells and phenotypes as the ground truth. To achieve this, we set the large cell subpopulation as common cells shared by the two phenotypes and assigned the other two small cell subpopulations to each phenotype, respectively. These two small subset cells are unique to each phenotype and thus can be viewed as the ground truth phenotype-specific cell subpopulations. To simulate the expression profiles of bulk samples for each phenotype, we randomly selected 1000 cells with replacement from the cell subpopulations contained by each phenotype and then averaged the expressions. In this way, we generated bulk gene expressions of 50 samples for each phenotype.

The differential expression genes between the two cell subpopulations were called by Seurat⁵⁹ using the default Wilcoxon rank-sum test. To explore how the performance of Scissor was affected by the true DEG signals at the bulk level, we simulated another bulk data that the true DEGs are not differentially expressed. In detail, we first selected the least differentially expressed gene from bulk data and then used its expression profiles to replace the expressions of all the true DEGs. This true DEG signal-removed bulk expression data would make Scissor more challenging to capture the real phenotype-to-cell associations.

We used precision and recall measurements in supervised learning to evaluate the performance of Scissor. The precision is defined as the proportion of the Scissor-identified ground truth cells among all Scissor selected cells, and recall is the proportion of the Scissor-identified ground truth cells among all ground truth cells.

Single-cell RNA-seq data preprocessing

We applied Seurat R package (version 3.2.1)⁵⁹ functions to preprocess the scRNA-seq data used in Scissor. Similar to the standard quality control step in single-cell data analysis, we first filtered out the genes with low expressions in cells. After this, we normalized the filtered

expression matrix using the NormalizeData function with the default parameters. Furthermore, we focused on the genes that exhibited high cell-to-cell variation and identified the variable genes using the FindVariableFeatures function with the default 'vst' method. We then performed the principal components analysis on the scaled variable gene expressions. Using the first ten principal components, we constructed a shared nearest neighbor graph for cells and identified cell clusters using the FindClusters function. Finally, we applied the Uniform Manifold Approximation and Projection (UMAP)⁶¹ technique using the RunUMAP function to visualize cells in low dimensions.

Differential expression gene analysis

The differentially expressed genes in Scissor+ cells versus Scissor- cells (or all other cells in some applications) were identified using the FindMarkers function in Seurat⁵⁹ with the default Wilcoxon rank-sum test. We required the following criteria to obtain the differential expression genes: First, the gene expression between the two groups is statistically significant with a false discovery rate (FDR) less than 0.05. Second, the absolute value of expression fold change between the two groups is greater than 1.25. We used the VlnPlot function to exhibit the relative expression levels of the selected differential expression genes.

We built several Scissor-derived signatures based on the differential expression genes in each experiment. The complete lists of signature genes can be found in Supplementary Tables.

Pathway enrichment analysis

The pre-ranked version of CAMERA⁶² (function name: cameraPR) in the limma R package (version 3.42.2) was used to evaluate the enriched pathways. cameraPR needs a pre-ranked gene list according to a user-defined statistic score. This study used the differential expression genes output of the FindMarkers function to obtain the pre-ranked gene list. In detail, the statistic ranking score for each gene was calculated using the following formula:

$$\log\text{FC} * \log_{10} \frac{1}{\text{FDR} + 10^{-100}}$$

where logFC stands for the log-transformed fold change between the two groups, and FDR is the adjusted Wilcoxon rank-sum test p-value. The pathways used in CAMERA were downloaded from the Molecular Signatures Database (MSigDB, version 7.3)⁶³. The Mognol_LCMV_Tumor_Exhaustion gene set contains the up-regulated genes in exhausted T cells from both OT-I mouse model and lymphocytic choriomeningitis virus (LCMV) infected mice⁶⁴.

Survival analysis

We performed some survival analyses to examine the clinical relevance of the Scissor-derived signatures (lung cancer signature and *TP53* mutation signature). The gene set variation analysis (GSVA) algorithm⁶⁵ with the default settings, as implemented in the GSVA R package (version 1.34.0), was applied to calculate our signature score for each sample. Next, the samples were stratified into two groups based on the quantile values of the signature scores (upper quartile versus lower quartile). Survival curves of these two groups of patients were estimated by the Kaplan-Meier method with statistical significance calculated using the log-rank test.

We used the univariate Cox proportional hazard (Cox PH) model to examine the association between survival time and clinical features (Pathological Stage: Stage I=1, Stage II=2, Stage III=3; Sex: 'Male'=1, 'Female'=2; Age at diagnosis). The significant prognostic factors ($p < 0.05$) in the univariate Cox PH model were included in the subsequent multivariable Cox PH model. The Kaplan-Meier estimator, log-rank test, and Cox PH models were calculated in the survival R package (version 3.2-3).

Motif analysis

The oPOSSUM software⁶⁶ (version 3.0) was used to detect the motifs of the transcription factor binding sites for the lung cancer signature genes. We used the transcription factor binding site annotations provided by the tool (JASPAR CORE Profiles). For the parameter settings, we set 2000 base pairs in upstream and downstream sequences and kept other parameters as default values. Only motifs with a Z-score larger than ten and a Fisher score larger than seven were reported in the final motif list.

Master regulator analysis

Transcription factor activity was inferred using the master regulator inference algorithm (MARINa)⁶⁷ in the viper R package (version 1.20.0). A pre-ranked gene list with scores and a regulatory network are the two data sources required as the inputs. In this work, we used the same pre-ranked gene list calculated in the pathway enrichment analysis. The transcription factor regulome was curated from several databases as previously described⁶⁸.

Signature enrichment score calculation

A pseudo-regulon was built based on the significantly up-regulated and down-regulated genes between the cell subpopulations identified by Scissor. These genes were served as the positive and negative target genes of the constructed pseudo-regulon, respectively. The single-sample extension of MARINa⁶⁹ (function name: viper) in the viper R package (version 1.20.0) was employed to infer the activity of the constructed pseudo-regulon, which was used as the signature enrichment score for each sample.

Dataset

The descriptions and preprocessing of all datasets used in this study can be found in Supplementary Materials.

Software availability

662 The open-source Scissor R package and tutorial are available at GitHub:

663 <https://github.com/sunduanchen/Scissor>.

Acknowledgements

This work was supported by the following funding: NIH 5K01LM012877 (to Z.X.); NIH 1R21HL145426 (to Z.X.); NIH 1R01CA207377 (to D.Z.Q); NIH NIGMS MIRA R35GM124704 (to A.A.); The Medical Research Foundation of Oregon (to Z.X.). We would like to thank Weston Anderson and Ann Hill for editing the manuscript. The resources of the Exacloud high-performance computing environment developed jointly by OHSU and Intel, and the technical support of the OHSU Advanced Computing Center are gratefully acknowledged.

Author Contributions

D.S. and Z.X. conceived the idea, implemented the algorithm and performed the analyses. D.S., G.X., P.T.S. and Z.X. interpreted the results. A.M., D.Z.Q., P.S., M.D., A.D., J.A. and A.A. provided scientific insights on the applications. Z.X. supervised the study. D.S. and Z.X. wrote the manuscript with feedback from all other authors. All of the authors read and approved the final manuscript.

Competing interests

A.E. Moran discloses receipt of a sponsored research agreement from AstraZeneca. A.V. Danilov reports consultancy from Abbvie, Beigene, Celgene, Curis, Janssen, Karyopharm, Nurix, Seattle Genetics, Teva Oncology, and TG Therapeutics; research funding from Aptose Biosciences, Bristol-Myers Squibb, Gilead Sciences, and Takeda Oncology; and consultancy and research funding from AstraZeneca, Bayer Oncology, Genentech, and Verastem Oncology. J.A. Alumkal has received consulting income from Janssen Biotech, Merck Sharp & Dohme, and Dendreon; honoraria for speaker's fees from Astellas. All other authors declare no competing interests.

References

1. Zhang, Q. et al. Landscape and Dynamics of Single Immune Cells in Hepatocellular Carcinoma. *Cell* **179**, 829-845 e820 (2019).
2. Yofe, I., Dahan, R. & Amit, I. Single-cell genomic approaches for developing the next generation of immunotherapies. *Nat Med* **26**, 171-177 (2020).
3. Wagner, J. et al. A Single-Cell Atlas of the Tumor and Immune Ecosystem of Human Breast Cancer. *Cell* **177**, 1330-1345 e1318 (2019).
4. Villani, A.C. et al. Single-cell RNA-seq reveals new types of human blood dendritic cells, monocytes, and progenitors. *Science* **356** (2017).
5. Patel, A.P. et al. Single-cell RNA-seq highlights intratumoral heterogeneity in primary glioblastoma. *Science* **344**, 1396-1401 (2014).
6. Trapnell, C. et al. The dynamics and regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381-386 (2014).
7. Levine, J.H. et al. Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with Prognosis. *Cell* **162**, 184-197 (2015).
8. Miao, Y. et al. Adaptive Immune Resistance Emerges from Tumor-Initiating Stem Cells. *Cell* **177**, 1172-1186 e1114 (2019).
9. Azizi, E. et al. Single-Cell Map of Diverse Immune Phenotypes in the Breast Tumor Microenvironment. *Cell* **174**, 1293-1308 e1236 (2018).
10. Lambrechts, D. et al. Phenotype molding of stromal cells in the lung tumor microenvironment. *Nat Med* **24**, 1277-1289 (2018).
11. Guo, X. et al. Global characterization of T cells in non-small-cell lung cancer by single-cell sequencing. *Nat Med* **24**, 978-985 (2018).
12. Cancer Genome Atlas Research, N. et al. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet* **45**, 1113-1120 (2013).
13. International Cancer Genome, C. et al. International network of cancer genome projects. *Nature* **464**, 993-998 (2010).
14. Karaayvaz, M. et al. Unravelling subclonal heterogeneity and aggressive disease states in TNBC through single-cell RNA-seq. *Nat Commun* **9**, 3588 (2018).
15. Zappia, L., Phipson, B. & Oshlack, A. Splatter: simulation of single-cell RNA sequencing data. *Genome Biol* **18**, 174 (2017).
16. Cancer Genome Atlas Research, N. Comprehensive molecular profiling of lung adenocarcinoma. *Nature* **511**, 543-550 (2014).
17. Lawson, D.A. et al. Single-cell analysis reveals a stem-cell program in human metastatic breast cancer cells. *Nature* **526**, 131-135 (2015).
18. Brady, S.W. et al. Combating subclonal evolution of resistant cancer phenotypes. *Nat Commun* **8**, 1231 (2017).
19. Ryan, H.E. et al. Hypoxia-inducible factor-1alpha is a positive factor in solid tumor growth. *Cancer Res* **60**, 4010-4015 (2000).
20. Godet, I. et al. Fate-mapping post-hypoxic tumor cells reveals a ROS-resistant phenotype that promotes metastasis. *Nat Commun* **10**, 4862 (2019).
21. Gentles, A.J. et al. The prognostic landscape of genes and infiltrating immune cells across human cancers. *Nat Med* **21**, 938-945 (2015).
22. Wilson, W.R. & Hay, M.P. Targeting hypoxia in cancer therapy. *Nat Rev Cancer* **11**, 393-410 (2011).
23. Santoro, A. et al. p53 Loss in Breast Cancer Leads to Myc Activation, Increased Cell Plasticity, and Expression of a Mitotic Signature with Prognostic Value. *Cell Rep* **26**, 624-638 e628 (2019).
24. Barsotti, A.M. & Prives, C. Pro-proliferative FoxM1 is a target of p53-mediated repression. *Oncogene* **28**, 4295-4305 (2009).

25. Perri, F., Pisconti, S. & Della Vittoria Scarpati, G. P53 mutations and cancer: a tight linkage. *Ann Transl Med* **4**, 522 (2016).
26. Gettinger, S. et al. Impaired HLA Class I Antigen Processing and Presentation as a Mechanism of Acquired Resistance to Immune Checkpoint Inhibitors in Lung Cancer. *Cancer Discov* **7**, 1420-1435 (2017).
27. Sade-Feldman, M. et al. Resistance to checkpoint blockade therapy through inactivation of antigen presentation. *Nat Commun* **8**, 1136 (2017).
28. Robert, C. et al. Pembrolizumab versus Ipilimumab in Advanced Melanoma. *N Engl J Med* **372**, 2521-2532 (2015).
29. Weber, J.S. et al. Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): a randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* **16**, 375-384 (2015).
30. Tirosh, I. et al. Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq. *Science* **352**, 189-196 (2016).
31. Hugo, W. et al. Genomic and Transcriptomic Features of Response to Anti-PD-1 Therapy in Metastatic Melanoma. *Cell* **165**, 35-44 (2016).
32. Van Allen, E.M. et al. Genomic correlates of response to CTLA-4 blockade in metastatic melanoma. *Science* **350**, 207-211 (2015).
33. Chen, Z. et al. TCF-1-Centered Transcriptional Network Drives an Effector versus Exhausted CD8 T Cell-Fate Decision. *Immunity* **51**, 840-855 e845 (2019).
34. Siddiqui, I. et al. Intratumoral Tcf1(+)PD-1(+)CD8(+) T Cells with Stem-like Properties Promote Tumor Control in Response to Vaccination and Checkpoint Blockade Immunotherapy. *Immunity* **50**, 195-211 e110 (2019).
35. Scott, A.C. et al. TOX is a critical regulator of tumour-specific T cell differentiation. *Nature* **571**, 270-274 (2019).
36. Sade-Feldman, M. et al. Defining T Cell States Associated with Response to Checkpoint Immunotherapy in Melanoma. *Cell* **175**, 998-1013 e1020 (2018).
37. Kurtulus, S. et al. Checkpoint Blockade Immunotherapy Induces Dynamic Changes in PD-1(-)CD8(+) Tumor-Infiltrating T Cells. *Immunity* **50**, 181-194 e186 (2019).
38. van den Heuvel, A. et al. Single-cell RNA sequencing in facioscapulohumeral muscular dystrophy disease etiology and development. *Hum Mol Genet* **28**, 1064-1075 (2019).
39. Cooper, D. & Upadhyaya, M. Facioscapulohumeral Muscular Dystrophy (FSHD): clinical medicine and molecular cell biology. (Taylor & Francis, 2004).
40. Tiwari, A., Pattnaik, N., Mohanty Jaiswal, A. & Dixit, M. Increased FSHD region gene1 expression reduces in vitro cell migration, invasion, and angiogenesis, ex vivo supported by reduced expression in tumors. *Biosci Rep* **37** (2017).
41. Grewal, P.K., Todd, L.C., van der Maarel, S., Frants, R.R. & Hewitt, J.E. FRG1, a gene in the FSH muscular dystrophy region on human chromosome 4q35, is highly conserved in vertebrates and invertebrates. *Gene* **216**, 13-19 (1998).
42. Lassche, S. et al. Sarcomeric dysfunction contributes to muscle weakness in facioscapulohumeral muscular dystrophy. *Neurology* **80**, 733-737 (2013).
43. Banerji, C.R.S. et al. Dynamic transcriptomic analysis reveals suppression of PGC1alpha/ERRalpha drives perturbed myogenesis in facioscapulohumeral muscular dystrophy. *Hum Mol Genet* **28**, 1244-1259 (2019).
44. Grubman, A. et al. A single-cell atlas of entorhinal cortex from individuals with Alzheimer's disease reveals cell-type-specific gene expression regulation. *Nat Neurosci* **22**, 2087-2097 (2019).
45. Ashraf, G.M. et al. Protein misfolding and aggregation in Alzheimer's disease and type 2 diabetes mellitus. *CNS Neurol Disord Drug Targets* **13**, 1280-1293 (2014).

- 786 46. Neef, D.W., Jaeger, A.M. & Thiele, D.J. Heat shock transcription factor 1 as a
787 therapeutic target in neurodegenerative diseases. *Nat Rev Drug Discov* **10**, 930-944
788 (2011).
- 789 47. Yu, S.P., Sensi, S.L., Canzoniero, L.M., Buisson, A. & Choi, D.W. Membrane-delimited
790 modulation of NMDA currents by metabotropic glutamate receptor subtypes 1/5 in
791 cultured mouse cortical neurons. *J Physiol* **499** (Pt 3), 721-732 (1997).
- 792 48. Prieto, G.A. et al. Pharmacological Rescue of Long-Term Potentiation in Alzheimer
793 Diseased Synapses. *J Neurosci* **37**, 1197-1212 (2017).
- 794 49. Muramori, F., Kobayashi, K. & Nakamura, I. A quantitative study of neurofibrillary
795 tangles, senile plaques and astrocytes in the hippocampal subdivisions and entorhinal
796 cortex in Alzheimer's disease, normal controls and non-Alzheimer neuropsychiatric
797 diseases. *Psychiatry Clin Neurosci* **52**, 593-599 (1998).
- 798 50. Chatterjee, P. et al. Plasma glial fibrillary acidic protein is elevated in cognitively normal
799 older adults at risk of Alzheimer's disease. *Transl Psychiatry* **11**, 27 (2021).
- 800 51. Vieira Braga, F.A. et al. A cellular census of human lungs identifies novel cell states in
801 health and in asthma. *Nat Med* **25**, 1153-1163 (2019).
- 802 52. Fan, J. et al. Linking transcriptional and genetic tumor heterogeneity through allele
803 analysis of single-cell RNA-seq data. *Genome Res* **28**, 1217-1227 (2018).
- 804 53. inferCNV of the Trinity CTAT Project. <https://github.com/broadinstitute/inferCNV>.
- 805 54. Kiselev, V.Y., Andrews, T.S. & Hemberg, M. Challenges in unsupervised clustering of
806 single-cell RNA-seq data. *Nat Rev Genet* **20**, 273-282 (2019).
- 807 55. Satpathy, A.T. et al. Massively parallel single-cell chromatin landscapes of human
808 immune cell development and intratumoral T cell exhaustion. *Nat Biotechnol* **37**, 925-
809 936 (2019).
- 810 56. Mulqueen, R.M. et al. Highly scalable generation of DNA methylation profiles in single
811 cells. *Nat Biotechnol* **36**, 428-431 (2018).
- 812 57. Corces, M.R. et al. The chromatin accessibility landscape of primary human cancers.
813 *Science* **362** (2018).
- 814 58. Li, C. & Li, H. Network-constrained regularization and variable selection for analysis of
815 genomic data. *Bioinformatics* **24**, 1175-1182 (2008).
- 816 59. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell
817 transcriptomic data across different conditions, technologies, and species. *Nat*
818 *Biotechnol* **36**, 411-420 (2018).
- 819 60. Li, X., Xie, S., Zeng, D. & Wang, Y. Efficient l0 -norm feature selection based on
820 augmented and penalized minimization. *Stat Med* **37**, 473-486 (2018).
- 821 61. McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and
822 projection for dimension reduction. *arXiv preprint arXiv:1802.03426* (2018).
- 823 62. Wu, D. & Smyth, G.K. Camera: a competitive gene set test accounting for inter-gene
824 correlation. *Nucleic Acids Res* **40**, e133 (2012).
- 825 63. Subramanian, A. et al. Gene set enrichment analysis: a knowledge-based approach for
826 interpreting genome-wide expression profiles. *Proc Natl Acad Sci U S A* **102**, 15545-
827 15550 (2005).
- 828 64. Mognol, G.P. et al. Exhaustion-associated regulatory regions in CD8(+) tumor-infiltrating
829 T cells. *Proc Natl Acad Sci U S A* **114**, E2776-E2785 (2017).
- 830 65. Hanzelmann, S., Castelo, R. & Guinney, J. GSEA: gene set variation analysis for
831 microarray and RNA-seq data. *BMC Bioinformatics* **14**, 7 (2013).
- 832 66. Kwon, A.T., Arenillas, D.J., Worsley Hunt, R. & Wasserman, W.W. oPOSSUM-3:
833 advanced analysis of regulatory motif over-representation across genes or ChIP-Seq
834 datasets. *G3 (Bethesda)* **2**, 987-1002 (2012).
- 835 67. Lefebvre, C. et al. A human B-cell interactome identifies MYB and FOXM1 as master
836 regulators of proliferation in germinal centers. *Mol Syst Biol* **6**, 377 (2010).

- 837 68. Robertson, A.G. et al. Integrative Analysis Identifies Four Molecular and Clinical Subsets
838 in Uveal Melanoma. *Cancer Cell* **32**, 204-220 e215 (2017).
839 69. Alvarez, M.J. et al. Functional characterization of somatic mutations in cancer using
840 network-based inference of protein activity. *Nat Genet* **48**, 838-847 (2016).
841

Figure 1

Fig. 1. The workflow of phenotype-guided cell subpopulation identification method

Scissor. **a**, The inputs for Scissor are a single-cell expression matrix, a bulk expression matrix, and a corresponding phenotype of interest such as drug response or clinical information. **b**, Scissor calculates a correlation matrix and a cell-cell similarity network based on input sources, which are further integrated with the phenotype into a network regularized sparse regression model to select the most relevant cell subpopulations. **c**, The selected cells by Scissor with red and blue dots indicating the cells positively or negatively associated with the phenotype of interest. **d**, The Scissor results are evaluated by a reliability significance test to control the false associations between single-cell and bulk data. **e**, The selected cells by Scissor can be further investigated by downstream analyses.

Figure 2

Fig. 2. Validations of Scissor's effectiveness on simulation studies and a tumor-normal phenotype guided application. **a**, The UMAP visualization of the simulated cells. The orange and green dots are the ground truth cell subpopulations specific to phenotype I and II, respectively. **b**, The UMAP visualization of the clustering result on the simulated cells. **c**, The Venn diagram shows the overlaps between the true DEGs and the two sets of DEGs identified from the Scissor selected cells in two simulation scenarios. **d**, The visualization of the Scissor selected cells on the same UMAP as in **a**. The red and blue dots are the Scissor selected cells associated with phenotype I and II. **e**, The Venn diagrams show overlaps between the Scissor selected cells and the ground truth phenotype-specific cells. **f**, The UMAP visualization of the Scissor selected cells based on the true DEGs signal-removed bulk data. **g**, The Venn diagrams show overlaps between the Scissor selected cells and the ground truth cells. **h**, The UMAP visualization of 29,888 LUAD cells. The color encoding the cell types defined in the original paper. **i**, The UMAP visualization of the Scissor selected cells. The red and blue dots are cells associated with the tumor and normal phenotypes. **j**, The pie chart of the Scissor selected cells with the corresponding bar plots showing the detailed constitutions in each type defined in **h**. **k**, The Venn diagram shows a significant overlap between Scissor+ cells and cancer cells. The statistical p -value was calculated by the hypergeometric test.

Figure 3

Fig. 3. Scissor identification results on lung cancer cells guided by TCGA-LUAD survival outcomes. **a**, The UMAP visualization of 4,102 LUAD cancer cells. **b**, The UMAP visualization of the Scissor selected cells. The red and blue dots are Scissor+ (worse survival) and Scissor- (good survival) cells. **c**, The bar-plot shows the detailed constitution of Scissor+ cells in different clusters. **d**, The volcano plot of differential gene expressions in Scissor+ cells versus all other cells. Red and blue points mark the genes with significantly increased or decreased expressions in Scissor+ cells compared to other cells (FDR <0.05 and absolute fold-change > 1.25). The two vertical dashed lines represent $\pm \ln(1.25)$ fold-changes in gene expression, and the horizontal dashed line denotes FDR cutoff 0.05. **e**, The violin plots of expression levels of selected up-regulated genes in Scissor+ cells. The FDR was calculated by the Wilcoxon rank-sum test. **f**, The enrichment bar-plot of selected hypoxia-related pathways in the Reactome and Hallmark domains. **g**, The Kaplan-Meier survival curves show the clinical relevance of the lung cancer signature on four independent datasets. Tick marks indicate censoring events. A table is attached below in each panel to display the number of alive samples at given time points. The statistical *p*-values were determined by the log-rank sum test. **h**, The forest plots show the hazard ratios and 95% confidence intervals for the lung cancer signature and three additional clinical features according to the univariate Cox model. **i**, The forest plots show the hazard ratios and 95% confidence intervals for the lung cancer signature and stage information according to the multivariable Cox model.

Figure 4

Fig. 4. Scissor identification results on lung cancer cells guided by *TP53* mutation status.

a, The UMAP visualization of the Scissor selected cells. The red and blue dots are cells associated with the *TP53* mutant and wild-type phenotypes. **b**, The volcano plot of differential gene expressions in Scissor+ cells versus Scissor- cells. Red and blue points mark the genes with significantly increased or decreased expressions in Scissor+ cells (FDR <0.05 and absolute fold-change > 1.25). The two vertical dashed lines represent $\pm \ln(1.25)$ fold-changes in gene expression, and the horizontal dashed line denotes FDR cutoff 0.05. **c**, The enrichment bar-plot of selected cell cycle-related pathways in the Hallmark domain. **d**, The master regulator analysis reveals the top 10 most activated TFs in Scissor+ cells and Scissor- cells, respectively. The targets of each TF are shown as tick marks with red vertical lines representing positive targets and blue vertical lines for negative targets. Each row also illustrates the statistical *p*-value and the inferred differential activity for each transcription factor. **e**, The Kaplan-Meier survival curve shows the clinical relevance of the *TP53* mutation signature. Tick marks indicate censoring events. A table is attached below to display the number of alive patients at given time points. The statistical *p*-value was determined by the log-rank sum test. **f**, The violin plots of expression levels of selected down-regulated genes in Scissor+ cells. The FDR was calculated by the adjusted Wilcoxon rank-sum test.

Figure 5

Fig. 5. Scissor identification results on melanoma T cells. **a**, The UMAP visualization of 1,894 melanoma T cells in six clusters. **b**, The UMAP visualization of the Scissor selected cells. The red and blue dots are cells associated with immunotherapy responders and non-responders in melanoma. **c**, The bar-plot shows the distribution of Scissor+ cells across the six T cell populations. **d**, The volcano plot of differential gene expressions in Scissor+ cells versus all other cells. Red and blue points mark the genes with significantly increased or decreased expressions in Scissor+ cells compared to other cells (FDR <0.05 and absolute fold-change > 1.25). The two vertical dashed lines represent $\pm \ln(1.25)$ fold-changes in gene expression, and the horizontal dashed line denotes FDR cutoff 0.05. **e**, The violin plots show the expression levels of important immune genes in Scissor+ cells. The FDR was calculated by the Wilcoxon rank-sum test. **f**, The enrichment bar-plot shows the significantly enriched pathways in Scissor+ cells compared with all other cells (FDR <0.05). **g**, The boxplot shows the enrichment scores of the immunotherapy responsive signature in the non-responders and responders from Sade-Feldman's cohort. The statistical *p*-value was calculated by the Student's *t*-test. **h**, The GSEA enrichment plot of the up and down signature genes in the responder vs. non-responder comparison from Sade-Feldman's cohort. **i**, The boxplot shows the enrichment scores of the immunotherapy responsive signature in five types of CD8 T cells from a mouse liver tumor model. The statistical *p*-values were determined by the Student's *t*-test. **j**, The boxplot shows the enrichment scores of the immunotherapy responsive signature in the memory precursor CD8 T cells and short-lived effector CD8 T cells. The statistical *p*-value was determined by the Student's *t*-test.

Figure 6

Fig. 6. Scissor identification results on Facioscapulohumeral muscular dystrophy (FSHD) cells. **a**, The UMAP visualization of 6,899 cells derived from FSHD and control samples. **b**, The UMAP visualization of the Scissor selected cells. The red and blue dots are representing Scissor+ and Scissor- cells, associated with FSHD and control phenotypes, respectively. **c**, The bar-plot shows the detailed phenotypic constitutions of the Scissor selected cells. **d**, The volcano plot of differential gene expressions in Scissor+ cells versus Scissor- cells. Red and blue points mark the genes with significantly increased or decreased expressions in Scissor+ cells compared to Scissor- cells (FDR <0.05 and absolute fold-change > 1.25). The two vertical dashed lines represent $\pm \ln(1.25)$ fold-changes in gene expression, and the horizontal dashed line denotes FDR cutoff 0.05. **e**, The violin plots show the expression levels of selected dysregulated genes in Scissor+ cells. The FDR was calculated by the Wilcoxon rank-sum test. **f**, The enrichment bar-plot of selected muscle-related pathways in the Hallmark, GO's biological process, and cellular component domains. **g**, The boxplots show the enrichment scores of the FSHD molecular signature in the FSHD and normal controls from three independent validation datasets. The statistical *p*-values were determined by the Student's *t*-test.

Figure 7

Fig. 7. Scissor identification results on Alzheimer's disease (AD). **a**, The UMAP visualization of the Scissor selected cells on oligodendrocytes. **b**, The bar-plot shows the detailed phenotypic constitutions of the Scissor selected oligodendrocyte cells. **c**, The UMAP visualization of the Scissor selected cells on astrocytes. **d**, The bar-plot shows the detailed phenotypic constitutions of the Scissor selected astrocyte cells. **e**, The UMAP visualization of the Scissor selected cells on OPCs. **f**, The bar-plot shows the detailed phenotypic constitutions of the Scissor selected OPCs cells. **g**, The heatmap of differential gene fold-changes in Scissor+ cells versus Scissor- cells across all three brain cell types. Red and blue elements mark the genes with significantly increased or decreased expressions in Scissor+ cells (FDR <0.05 and absolute fold change > 1.25). **h**, The heatmap of enriched pathways in the Reactome domain. The red and blue elements represent the activated and repressed pathways in corresponding cell types. **i**, The boxplot shows the enrichment scores of the oligodendrocytes molecular signature in AD patients and normal controls from GSE109887. The statistical *p*-value was determined by the Student's t-test. **j**, The boxplot shows the enrichment scores of the OPCs molecular signature in AD patients and normal controls from GSE109887. The statistical *p*-value was determined by the Student's t-test. **k**, The boxplot shows the enrichment scores of the astrocytes molecular signature in AD patients and normal controls from GSE109887. The statistical *p*-value was determined by the Student's t-test. **l**, The boxplot shows the enrichment scores of the astrocytes molecular signature in control, incipient, moderate, and severe stage AD patients from GSE28146. The statistical *p*-value was determined by the Kruskal-Wallis test. The linear regression line represents the relationship between median enrichment scores and AD stages.

Figures

Figure 1

The workflow of phenotype-guided cell subpopulation identification method Scissor.

Figure 2

Validations of Scissor's effectiveness on simulation studies and a tumor-normal phenotype guided application.

Figure 3

Scissor identification results on lung cancer cells guided by TCGA-LUAD survival outcomes.

Figure 4

Scissor identification results on lung cancer cells guided by TP53 mutation status.

Figure 5

Scissor identification results on melanoma T cells.

Figure 6

Scissor identification results on Facioscapulohumeral muscular dystrophy (FSHD) cells.

Figure 7

Scissor identification results on Alzheimer's disease (AD).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [ScissorSuppMaterials.pdf](#)
- [SupplementaryTables.zip](#)