Regional trends and associated factors of childhood visual impairment: a case study in Shandong province, China

Lizhen HAN

Peking University

Jinzhu Jia (✉ jzjia@math.pku.edu.cn)

Lu Wang

Physical Examination Office of Shandong Province, China

Research

Keywords: Visual impairment, children, regional difference, spatial analysis

Posted Date: August 2nd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-49579/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Scientific Reports on August 16th, 2021.
See the published version at <https://doi.org/10.1038/s41598-021-95906-7>.

1 **Regional trends and associated factors of childhood visual impairment: a case**
2 **study in Shandong province, China**

3 Lizhen Han,¹ Jinzhu Jia,^{1,*} Lu Wang^{2,*}

4 ¹ Department of Biostatistics, School of Public Health, Peking University, Beijing, China, No. 38, Xueyuan Road,
5 Haidian District, Beijing City, 100191, China

6 ² Physical Examination Office of Shandong Province, Jinan 250014, China

7 * Correspondence

8 **Corresponding Author:**

9 Jinzhu Jia, Ph.D. (jzjia@math.pku.edu.cn), Department of Biostatistics, School of Public Health,
10 Peking University, Beijing, China, No. 38, Xueyuan Road, Haidian District, Beijing City, 100191,
11 China

12 Lu Wang, Ph.D. (luwang2716@163.com), Physical Examination Office of Shandong Province, Jinan,
13 China, No.9 Yandong New Road, Lixia District, Jinan City, Shandong Province, 250014, China

14 **Abstract**

15 **Background:** Visual impairment is a common child health problem. We hopes to analyze the
16 regional trend, related factors and spatial distribution of children's visual impairment in Shandong
17 province, to explore the spatial changes brought by time and their influencing factors, so as to
18 provide scientific basis for prevention of childhood visual impairment.

19 **Methods:** This study covers 5 complete cross-sectional surveys (Physical examination data in
20 Shandong province) from 2013 to 2017, involving about 29.24 million students. The 11th International
21 Classification of Diseases was selected as the diagnostic standard. Spatial autocorrelation and hotspot
22 analysis methods in ArcGIS software were used to analyze spatial features. The associated factors were
23 analyzed by multinomial logistic regression.

Results: The visual impairment prevalence showed a trend of decreasing first and then increasing from 2013-2017, with slight changes. In terms of regional spatial differences, Weihai and Yantai have the highest VI rates in all years, and there was a large-scale spatial aggregation phenomenon. The southern low-value clusters, however, showed a weakening year by year. Further exploration revealed that the per capita disposable income of rural households, the growth rate of gross domestic product, total retail sales of consumer goods and number of full-time teachers were verified as risk factors for regional visual impairment levels.

Conclusions: The slight rebound of the prevalence of visual impairment and the high rate in the eastern and northern regions of Shandong province need more attention. It is suggested that relevant departments should focus on the influence of regional economic and educational factors when formulating relevant strategies.

Keywords

Visual impairment, children, regional difference, spatial analysis

Introduction

Childhood is considered to be one of the most critical stages of growth and development, which has long been concerned from many aspects, such as the state, schools, families and individuals. How to create a healthier growth environment for children in the current situation (increasingly complex living environment) is an urgent problem to be solved. Among them, vision issues (such as visual impairment (VI), myopia, amblyopia, etc.) are most common and prominent. As a window for one to understand and observe the world, the importance of eyes is self-evident. In China, a populous country that has liberalized the Two-Child policy, the prevalence of myopia and other eye-related diseases among children and adolescents has been rising in recent years. For example, the VI rate with strong generality

has rapidly increased almost 2.5-fold from 23.7% to 55% (1985-2014) [1]. From a global perspective, China is also the region with the highest prevalence of low vision per million people in the surveyed area (including Africa, America, Eastern Mediterranean, Europe, South-east Asia and Western Pacific) [2]. The visual impairment has become an urgent health problem in China, which requires cooperation of various departments.

Visual impairment refers to deficits in the ability of the person to perform vision-related activities of daily living. It also reflects the burden of vision loss for the person. If the unaided distance visual acuity (VA) of worse than 6/12 ($\log\text{MAR} < 0.5$), it is judged as visual impairment. Studies have confirmed that there are many causes for the occurrence of VI, including heredity (e.g. macular dystrophies and retinitis pigmentosa), diseases (e.g. refractive error, posterior segment (retinal) diseases, corneal degenerations and optic nerve head disease) and trauma, etc. [3-5]. Compared with normal sight children, the overall quality of life (QoL) of visual impaired children is relatively lower [6]. This is not only manifested in physical health and psychological aspects (e.g. individual nutritional status [7] and depression [8]), but also in social relations and living environment [9], and its impact will be lifelong. A recent study by Jones, N and other scholars found that visual impairment will affect the corresponding life skills (e.g. shopping and cooking) when it develops into adulthood [3]. In addition, visual impaired children will have a relatively lower level of participation in sports activities (the higher the degree of VI, the lower the level of participation) [3, 10]. This phenomenon will not merely affect children's social interaction and exercise, but also give rise to a series of problems such as overweight and obesity [11, 12].

On the other hand, the relationship between visual impairment and economy is also inseparable. Research showed that for every 100% increase in gross domestic product (GDP), the risk of VI will rise

by 20%, and the risk of moderate to severe VI will even increase by 27% [1]. Meanwhile, compared with normal children, visual impaired children are more likely to come from families with relatively higher economic level [10]. In China, the direct cost per patient due to VI is US\$ 6988.6 ± 10,834.3 per year, of which 70% is direct medical expenses and only less than 30% can be reimbursed [13]. A recent German study also showed that VI and blindness have brought huge social burden (annual cost is about EU€49.6 billion) [14]. Furthermore, if patients suffer from chronic diseases such as hypertension, the risk of VI will be further increased [15]. The economic pressure brought by this multiple disease burden is obvious.

Globally, the number of visual impaired people among all ages reached about 285 million as early as 2010 [2]. Moreover, the current problem of VI in China has a clear tendency to be younger. Albeit the older age group still occupies the majority in general, the prevalence of younger group has increased even more rapidly [1]. Therefore, this article will study the evolution of childhood vision problems in Shandong province (about 100 million people), a representative province in China, based on the VI prevalence of children aged 6-12. The aim was to find out the secular trend, regional distribution characteristics and related risk factors through the analysis on the visual impaired schoolchildren from 2013 to 2017, so as to provide data support for the next policy adjustment and targeted preventive interventions by the government and relevant departments. Our ultimate goal is to create a healthier growth and living environment for children.

Material and methods

Data Source

The sample came from students' physical examination data from September to December every year in Shandong province, totaling 5 years (i.e. 2013-2017). After adjusting for age and gender, the missing

data and the error records beyond the scope of the visual chart (totaling 430,421) were eliminated. A total of 29 237 771 schoolchildren aged 6-12 years were included (person-time), covering whole primary schools in all 17 cities (137 counties) in Shandong province. The sample sizes in this data for different years ranged from 1 864 242 to 7 763 686, and the annual girl to boy ratio and urban to rural ratio approximately equaled 1:1.2 and 1:1.7, respectively.

Measurement Methods

The National Health Commission of the People's Republic of China, the Ministry of Education of the People's Republic of China (*Measures for the administration of health examination of primary and middle school students* [16]) and the Shandong Provincial Education Department (*Measures for the implementation of health examination management for primary and middle school students in Shandong province* [17]) stipulate that students in school should have a physical examination every school year. This program organized by the health administrative department at or above the county level, and rigorous training was conducted, covering all medical professionals who participated in physical examination. In addition, through the schools' teachers, they informed the students and their parents about the details of the physical examination and got their consent. After the examination, students and parents will receive feedback in the form of reports.

According to the requirements of *Health examination methods for primary and secondary schools in Shandong province*, all physical examinations procedures were carried out in accordance with regulations and using standardized professional instruments. The data involved in this study were obtained through physical measurements and did not involve laboratory examinations.

Visual acuity was measured using a standard logarithmic E chart of vision (in line with the national standard GB11533—2011, International Classification for Standards (ICS) 13.100) [18].

Students need to keep the naked eye to participate in the test. During vision inspection, students should maintain a distance of 5m from the instrument, and the sight line of the tested person should be consistent with the height of the visual chart (line "5.0"). Then, the left eye and the right eye are covered by the eye mask in turn, and the VA of the corresponding eye was detected. The reading of the smallest line that can be seen clearly by the subject was the VA of the relevant eye, and the value was recorded.

Judging Criteria for Visual Impairment

All children had been diagnosed with visual impairment according to the 11th International Classification of Diseases (ICD-11) issued by World Health Organization (WHO) in 2018 [19]. According to the actual situation in China (i.e., the visual chart used in children's physical examination does not involve the detection range of less than 6/60) and previous studies [1, 20], in this study, visual impairment was divided into mild and moderate-severe categories. Mild VI was defined as unaided distance VA of worse than 6/12 ($\log\text{MAR} < 0.5$) and that of equal to or better than 6/18 ($\log\text{MAR} \geq 0.3$) in the worse eye, moderate to severe VI was defined as unaided distance VA worse than 6/18 ($\log\text{MAR} < 0.3$) in the worse eye.

Other Data

In terms of influencing factors, seven regional associated factors (obtained from the *Shandong Statistical Yearbook* over the years) were included, covering gross domestic product (GDP), the growth rate of GDP (GRGDP), general public budget expenditure (GPBE)—state financial expenditures that are spent on public services, education, medical care and social security, etc., total retail sales of consumer goods (CGTRS)—sales amount of physical commodities for non-production and non-business purposes (including catering services), number of full-time teachers (FTT)—weighted by

the number of students in the region, representing the number of teachers per thousand students, and per capita disposable income of rural households (RPCDI). Meanwhile, in order to effectively reduce the impact of population differences in different areas, this study weighted the regional associated factors by their population (except GRGDP). In addition, based on the annual VI prevalence range and the effect of mapping, the regional VI level was divided into 5 grades from low to high (ranging from 0% to 35%, with every seven percentage points recorded as a level).

Statistical Analysis

Raw data was extracted through SQL Sever2017, and SPSS 22.0 was used to analyze the data. The measurement data were described by the mean (standard deviation), and t-tests and linear correlation were used for inter-group comparison. The counting data was expressed by the rate (%), and the comparative analysis between groups was tested by chi-square tests. Furthermore, multinomial logistic regression analysis was used to analyze population-weighted associated factors. A probability level of $p < 0.05$ represented the result with statistical significance.

Spatial Analysis

ArcGIS 10.2 software was used to analyze spatial distribution, regional variation differences, etc. In light of the distribution range of VI rates, this study artificially divided it into 5 levels. Besides, in order to find out whether there is spatial clustering and its variation trend in the Shandong province, spatial autocorrelation (Global Moran's I) and hotspot analysis (Getis-Ord G_i^*) were applied to this study.

Spatial autocorrelation (Global Moran's I) is a method for measuring spatial autocorrelation based on element locations and element values, which is used to evaluate whether the pattern expressed is clustered, dispersed, or random. It evaluates the significance of data results by calculating the Moran's I index value, z -score and p -value. If the index in the dataset tend to cluster spatially (aggregation of

similar values), the Moran's index will be positive. Conversely, the index will be negative. If positive cross-product values balance negative cross-product values, the index will be near zero [21].

The clustering of high and low values was obtained through hotspot analysis (calculation the Getis-Ord G_i^* statistic for each feature in a dataset). After a partial sum of a feature and its neighbors was compared with the sum of all features, if the local sum was significantly different from the expected local sum that it cannot be randomly generated, a statistically significant z-score will be resulted. Based on the resultant z-scores and p -values, the location where the high-value or low-value elements cluster in space can be illustrated.

The Getis-Ord local statistic is given as :

$$G_i^* = \frac{\sum_{j=1}^n w_{i,j} x_j - \bar{X} \sum_{j=1}^n w_{i,j}}{S \sqrt{\frac{[n \sum_{j=1}^n w_{i,j}^2 - (\sum_{j=1}^n w_{i,j})^2]}{n-1}}}$$

Where x_j is the attribute value for feature j , $w_{i,j}$ is the spatial weight between feature i and j , and :

$$\bar{X} = \frac{\sum_{j=1}^n x_j}{n}$$

$$S = \sqrt{\frac{\sum_{j=1}^n x_j^2}{n} - (\bar{X})^2}$$

The G_i^* statistic returned for each feature in the dataset is the z-score. For statistically significant positive z-scores, the larger the z-score is, the more intense the clustering of high values (hot spot). The opposite clustering method is called cold spot [22].

Patient and public involvement

No patients or public were involved in this study.

Results

173 The sample was from the physical examination data of primary and secondary school students in
174 Shandong province (2013-2017), and children aged 6-12 years old were selected as subjects (a total of
175 29 237 771, mean (SD) age: 9.06 (1.91) years; 53.96% were boys), which basically covers whole
176 school children in Shandong province. Among them, the sample size in these 5 years were 1 864 241, 5
177 859 099, 6 685 362, 7 065 383 and 7 763 686 respectively. Basic information of children in each city
178 can be found in Table 1.

Table 1 Information of children in different cities of Shandong province from 2013 to 2017

City	Number of children (mean age (years))				
	2013	2014	2015	2016	2017
Binzhou	- ^a	249 557(8.91)	251 354(8.93)	252 737(8.97)	272 209(8.95)
Dezhou	35 673(9.00)	357 337(8.85)	360 762(8.82)	421 942(8.84)	441 156(8.90)
Dongying	48 955(8.95)	118 448(8.91)	131 950(8.93)	140 396(9.00)	144 435(9.01)
Heze	- ^a	870 954(8.66)	919 634(8.72)	953 536(8.80)	1 025 622(8.86)
Jinan	201 006(8.77)	272 641(8.76)	403 972(8.79)	435 378(8.85)	470 622(8.87)
Jining	17 802(9.38)	532 320(8.79)	581 734(8.82)	672 818(8.88)	722 942(9.06)
Laiwu	25 985(9.20)	66 054(9.12)	60 871(9.18)	73 363(9.06)	72 568(9.06)
Liaochen	1 007(9.24)	152 313(8.56)	453 943(8.63)	523 583(8.65)	591 048(8.79)
Linyi	188 051(8.74)	618 792(8.70)	776 428(8.74)	714 024(8.73)	1 049 007(8.83)
Qingdao	482 202(8.77)	550 205(8.74)	561 041(8.76)	601 326(8.92)	615 741(9.01)
Rizhao	74 876(8.76)	192 778(8.85)	192 202(8.94)	211 653(8.94)	210 700(8.97)
Taian	157 793(9.10)	337 878(9.04)	341 652(8.99)	346 422(9.02)	359 240(8.98)

Weihai	80 748(8.80)	129 262(8.78)	132 928(8.68)	142 225(8.86)	151 096(8.98)
Weifang	112 438(8.53)	570 382(8.72)	600 495(8.86)	613 485(9.04)	636 462(9.19)
Yantai	202 825(8.99)	329 707(8.93)	322 536(8.85)	342 520(9.02)	348 904(9.10)
Zaozhuang	7 478(8.40)	270 160(8.58)	323 092(8.55)	346 292(8.68)	374 907(8.85)
Zibo	227 402(8.92)	240 331(8.94)	270 768(8.94)	273 683(9.02)	277 027(9.08)

^a The symbol ('-') represents a missing value

179 The total VI rates (T) for five years were relatively similar except for 2013 (14.87%), which
180 fluctuated around 12%. The gap between different cities, however, was quite wide, especially in Weihai
181 and Yantai, where the VI prevalence was the most prominent and has exceeding 20%. In general, VI
182 rates presented a trend of decreasing first and then rising, like a concave curve (although no
183 linear-by-linear association was detected, $p = 0.707$). In 17 different cities of Shandong province, the
184 differences of VI prevalence in each year were statistically significant, and all showed the same trend
185 (linear-by-linear association $p < 0.001$) except Weifang city ($p = 0.214$). In addition, the trends for the
186 mild (M) and moderate-severe (M-S) VI rate were similar to the total ($r_{T\&M} = 0.980$, $p = 0.003$;
187 $r_{T\&M-S} = 0.982$, $p = 0.003$), and their results were very close (about 6%). The results were shown in
188 Table 2 and Figure 1.

Table 2 Visual impairment (VI) of children in different cities of Shandong province from 2013 to 2017

City	Total VI rate (%)					χ^2 value	p -value
	2013	2014	2015	2016	2017		
Binzhou	- ^a	11.04	9.32	11.01	8.58	1325.08	< 0.001

Dezhou	7.17	11.00	10.88	11.72	14.63	4448.80	< 0.001
Dongying	15.99	18.07	18.68	19.78	18.27	380.16	< 0.001
Heze	- ^a	5.85	5.75	7.21	8.05	5657.60	< 0.001
Jinan	13.09	13.83	13.38	13.97	14.38	287.43	< 0.001
Jining	18.23	10.12	10.70	11.10	10.86	1348.03	< 0.001
Laiwu	15.97	18.33	18.02	19.79	20.57	358.02	< 0.001
Liaochen	6.95	9.85	7.68	8.50	8.38	736.83	< 0.001
Linyi	9.51	7.67	8.49	9.48	10.36	3978.22	< 0.001
Qingdao	15.14	13.89	13.95	16.08	17.09	3453.35	< 0.001
Rizhao	14.03	13.54	12.01	10.57	10.24	1729.40	< 0.001
Taian	12.57	12.29	12.97	12.51	14.95	1427.22	< 0.001
Weihai	21.92	22.53	20.42	20.71	20.47	276.64	< 0.001
Weifang	12.27	12.73	12.28	12.22	12.59	102.39	< 0.001
Yantai	22.77	21.05	20.01	22.57	22.74	1129.04	< 0.001
Zaozhuang	7.23	9.53	9.13	9.38	11.39	1350.70	< 0.001
Zibo	14.75	15.62	14.96	16.14	18.12	1440.65	< 0.001

^a The symbol ('-') represents a missing value

189 From figure 1, we found that the trends of VI rates for boys (B) and girls (G) were similar to that
190 of the whole over the past 5 years ($r_{T\&B}=0.995$, $p < 0.001$; $r_{T\&G}=0.993$, $p = 0.001$). No matter the
191 total VI prevalence, or the mild or moderate to severe VI rate, obvious gender differences were
192 detected (female rates > male rates), and the differences were statistically significant ($\chi^2_T=69727.572$,
193 $\chi^2_M=18642.317$, $\chi^2_{M-S}=49944.350$, $p < 0.001$). It is worth noting that the same results were obtained by

comparing students from urban and rural areas separately ($\chi^2_{urban}=33506.411$, $\chi^2_{rural}=39188.863$, $p < 0.001$). Besides, the single-sex curve clarified that the prevalence of mild and moderate to severe VI was similar from 2013-2017.

Compared with rural areas, urban regions had a higher VI prevalence, such as the overall rate ($\chi^2_T=313936.581$, $p < 0.001$), this gap was maintained at more than 6% every year. Meanwhile, the urban-rural gap in mild and moderate-severe morbidities were also significant ($\chi^2_M=96834.083$, $\chi^2_{M-S}=205497.682$, $p < 0.001$). However, in contrast, the upward trend of urban areas has declined in 2017. The results were shown in Figure 2.

In terms of age, there was a strong linear-by-linear association between VI prevalence and age (e.g. $r_{2015}= 0.960$, $p = 0.001$). The results (Figure 3) demonstrated that the trends in each year were relatively consistent, showing a trend that the prevalence of VI increases with age (from about 2% at the age of 6 to 30% in 12 years old), and the difference between ages was statistically significant ($\chi^2_T=2409075.899$, $p < 0.001$). Among them, the rise in 2013 was even fierce, while that in the other 4 years was relatively similar.

Spatial Analysis Results

The results of spatial analysis showed that from 2013 to 2017, the regions with high prevalence of VI among children aged 6-12 years in Shandong province were mainly concentrated in the eastern peninsula and the northern area. Likewise, the prevalence in the central region was slightly higher than that in surrounding counties. It is worth noting that two new high-value districts were added in 2017: the western (parts of Dezhou) and central (parts of Zibo) regions, and their prevalence were significantly higher than those in previous years. The results were shown in Figure 4-8.

Focusing on the regional distribution of mild and moderate-severe VI (yellow and purple dots) in

these maps, it was found that the aggregation phenomenon was prominent in the northeast of the peninsula and areas around Jiaozhou Bay. In addition, the cluster of central counties has become more and more distinct with the passage of time.

Further, this study analyzed the spatial relationships that may exist between regional differences in overall VI rates. The results (Table 3) illustrated that there was spatial aggregation of VI in Shandong province.

Table 3 Spatial autocorrelation of visual impairment (VI) among children in Shandong province from 2013-2017

Year	Moran's I index	Variance	z-score ^a	p-value
2013	0.40	0.01	4.76	< 0.001
2014	0.56	0.01	10.57	< 0.001
2015	0.53	0.01	10.32	< 0.001
2016	0.59	0.01	11.44	< 0.001
2017	0.53	0.01	10.26	< 0.001

^a $|z| > 2.58$ indicates that the probability of randomly generating this clustering pattern is less than 1%

Based on the spatial aggregation characteristics of VI in Shandong province from 2013 to 2017, the cluster map was presented after hotspot analysis. In 2013, although data were relatively scarce, a large range of high-value aggregation areas were still detected in the eastern part of Shandong peninsula (Weihai, Yantai), while the cold spot region was mainly concentrated in the central part. The distribution characteristics in the following four years were relatively consistent, and clusters with high/low values were located in the eastern/southern regions. In addition, in some years, a small

number of counties in the north/west (hot/cold spots) also showed aggregation.

Combined with the maps of various years, the range of hot spots has been fluctuating slightly, while the accumulation of cold spots has shown a trend of weakening year by year, especially in Zaozhuang and Linyi cities. The results were shown in Figure 9-13.

Influencing factors Analysis

After the collinearity diagnostics, five types of regional economic associated factors and one social influencing factor (including 137 counties in Shandong province): gross domestic product (GDP), the growth rate of GDP (GRGDP), general public budget expenditure (GPBE), total retail sales of consumer goods (CGTRS), per capita disposable income of rural households (RPCDI) and number of full-time teachers (FTT) were included in the multinomial logistic regression model. Among them, except GRGDP, the rest variables were weighted by regional population. Moreover, economic variables were recorded in units of CN ¥ 1000 per capita (approximately equal to US\$142.27), and the unit of FTT was 1 person.

Studies have confirmed that RPCDI, GRGDP and FTT were risk factors for regional VI level. RPCDI played a role in promoting the "development" of regional VI in different grades, while the latter two indexes only worked in the highest grade (compared with the level 1). Besides, CGTRS was also verified as a risk factor (only in level 3). Nevertheless, GDP and GPBE showed no statistically significant effect on the degree of VI. The results were shown in Table 4.

Table 4 Correlation between regional associated factors and visual impairment (VI) of children in Shandong province from 2013 to 2017

Regional	Factors	β	S.E.	Wald	p	$Exp(\beta)$	95% CI for $exp(\beta)$
----------	---------	---------	------	------	-----	--------------	-------------------------

VI level						Lower	Upper
						bound	bound
2	Intercept	-2.853	1.257	5.149	0.023		
	GDP	0.014	0.010	1.813	0.178	1.014	1.035
	GRGDP	0.017	0.027	0.395	0.530	1.017	1.074
	GPBE	0.045	0.107	0.174	0.676	1.046	1.289
	CGTRS	-0.001	0.023	0.004	0.950	0.999	1.044
	FTT	-0.001	0.006	0.047	0.829	0.999	1.011
	RPCDI	0.267	0.109	6.035	0.014	1.305	1.615
3	Intercept	-6.998	1.424	24.137	0.000		
	GDP	0.018	0.011	2.850	0.091	1.018	1.040
	GRGDP	0.006	0.032	0.035	0.851	1.006	1.070
	GPBE	0.014	0.111	0.017	0.897	1.014	1.260
	CGTRS	0.048	0.023	4.516	0.034	1.050	1.097
	FTT	0.001	0.008	0.008	0.929	1.001	1.016
	RPCDI	0.414	0.118	12.345	0.000	1.514	1.907
4	Intercept	-11.221	1.940	33.459	0.000		
	GDP	0.013	0.012	1.149	0.284	1.013	1.036
	GRGDP	-0.022	0.040	0.302	0.583	0.978	1.058
	GPBE	0.030	0.118	0.063	0.802	1.030	1.297
	CGTRS	0.028	0.026	1.236	0.266	1.029	1.081
	FTT	-0.001	0.012	0.005	0.943	0.999	1.022

	RPCDI	0.681	0.146	21.820	0.000	1.976	1.485	2.630
5	Intercept	-20.582	3.342	37.939	0.000			
	GDP	0.016	0.014	1.330	0.249	1.016	0.989	1.045
	GRGDP	0.116	0.055	4.345	0.037	1.123	1.007	1.252
	GPBE	-0.001	0.133	0.000	0.991	0.999	0.769	1.297
	CGTRS	0.021	0.033	0.419	0.518	1.022	0.958	1.090
	FTT	0.049	0.010	25.675	0.000	1.051	1.031	1.071
	RPCDI	0.932	0.221	17.738	0.000	2.540	1.646	3.919

Discussion

As a large country with a population of 1.4 billion, China has an equally large number of children. Since the Two-Child policy was completely liberalized in early 2016, the number of newborns has increased significantly. Meanwhile, the VI prevalence is also continuously rising [1]. In order to better deal with children's visual impairment and provide them with a healthy growth environment, this study selected children from Shandong province as the research subjects, based on comprehensive demographic and economic indicators (domestic population ranking: 2nd, economic ranking: 3rd).

The study found that the VI rate of children in Shandong province from 2013 to 2017 showed a flat U-shaped trend, with a slight decrease and increase. Considering that the data for 2013 only covered counties with better economy (the economy has a positive effect [15]), the detection rate was higher than that of other years. Excluding it, there was a slow upward trend on the whole. Compared with previous study,[1] there were obvious differences. The reason for this phenomenon may be related to the gradual steady increase in children's living environment and economic factors in recent years, so the increase in VI rate has slowed down.

In terms of gender, statistically significant differences were detected. Whether the overall VI rate or mild, moderate-severe VI rates, the results of female group were significantly higher than those of male group. This is consistent with previous research results [23-25]. Although there is no direct evidence to prove the causes of this difference, according to other studies, we guessed that it may be affected by the following factors: We think the most important factor is that Chinese girls tend to be gentle and quiet under the traditional concept. Whether in study or sports, girls will spend more time indoors than boys. Furthermore, less outdoor activities will aggravate the degree of myopia [26]. Meanwhile, the higher prevalence of dry eye syndrome in women may be one of the reasons [27]. A hypothesis about hormone action may be another reason [28]. Sex hormone levels in myopic groups were often higher than those in non-myopic groups: Women were mainly affected by follicle stimulating hormone and luteinizing hormone, while men were influenced by luteinizing hormone and testosterone, and female have the higher luteinizing hormone levels [29]. In addition, the interaction between sex and steroidogenesis enzyme genes has also been proved to be a regulator of sex hormone metabolism and high myopia risk [30].

Differences between urban and rural areas were also detected: The prevalence of the three VI was higher in cities than that in countries. The gap in economic level has been considered as the main reason for this phenomenon [15]. Albeit the economic gap between urban and rural areas in China is narrowing, "faults" still exist. Urban children living in a pleasant environment have more opportunities to contact with electronic products and even become addicted to them. Affected by this, compared with rural children, sedentary lifestyle of urban students tends to be normalized, and their visual condition also deteriorates [31, 32].

At the age level, there was a strong positive correlation between VI rate and age. With the increase

of age, the prevalence of VI among children also rose significantly. Multiple studies have confirmed this result [23, 33]. When children are young, their eyeballs are smaller and their axes are shorter. The eyes are maintained in a state of hyperopia and have a certain "hyperopia reserve". After that, with the growth of children and the influence of various factors (such as increased schoolwork burden, more screen time, etc.), hyperopia reserve is consumed prematurely, and VI problems gradually become prominent and rapidly increase, like myopia. In addition, the higher school year also plays a negative role in visual development (e.g. study and exam pressure, reduced outdoor activity time) [34].

Through the integration of the annual distribution and the spatial aggregation maps, the high-value aggregation in the eastern part of the peninsula has been confirmed. As for the specific reason: Why cluster will be formed in this region, it is still unknown. Previous studies have shown that there was a certain association between VI and overweight and obesity [35]—mainly influenced by sports activities [12, 31]. Likewise, previous research on obesity in the same population found that the high prevalence area was exactly consistent with the above results [36]. Initial speculation suggests that children in Yantai and Weihai cities may have less time for physical exercise. The relationship between VI, exercise, overweight and obesity is like a two-way closed cycle, which interacts and affects each other (i.e. visual impaired children have less physical activity, and sedentary lifestyle will lead to high prevalence of obesity; And vice versa). For a long time, measures that can promote students' physical activities have been mentioned, and policies and systems are also constantly being updated, but the results have been poor. How to effectively implement the policy and improve the current situation is the real issue to be considered in the following work. Moreover, the influence of economy and day length cannot be ignored. Research by Cui, Dongmei and other scholars pointed out that axial growth and myopia progression will decrease with the increase of day length [37]. Therefore, they may be

another cause of the high prevalence in central, eastern and northern regions.

Due to the large number of missing values in 2013, longitudinal comparison was not included. From 2014-2017, the low-value aggregation range showed a narrowing feature (although the change was slight), and it was mainly concentrated in the southern areas. In this aspect, economic changes and the above-mentioned day length are mainly considered: The economic level of the southern region is lagging behind in all years, and the side effects brought by technology (e.g. massive open online course (MOOC), multimedia class, etc.) have slightly weaker impact on the students. On the other hand, the regional economy was still on the rise, which also explains the shrinking of the aggregations to some extent.

If subdivided by gender, the distribution characteristics (including total, mild and moderate to severe VI) were highly consistent with the overall, and the results for boys and girls were similar. Therefore, this article does not further explore the gender differences in regional distribution.

This study also included a variety of representative regional influencing factors, in order to conduct a more comprehensive discussion on the reasons of the above problems. After adjusting for confounding, through multiple logistic regression analysis, a total of three risk factors and one protection factor were detected. They were: risk factors—RPCDI, GRGDP, CGTRS and FTT. When the regional VI level reached the fifth grade, they played the most significant role (except CGTRS). This means that compared with the underdeveloped areas, regions with higher RPCDI, GRGDP and FTT were more likely to develop into the regions with higher VI rate, and their risks were 2.540 (95% *CI*, 1.646-3.919), 1.123 (95% *CI*, 1.007-1.252) and 1.050 (95% *CI*, 1.004-1.097) times of those with lower VI prevalence respectively. Hence, we found that there were inextricable links between the economy and VI. When the regional economy and per capita disposable income have increased

significantly (especially in the underdeveloped rural areas), the stronger purchasing power under favorable conditions has greatly increased the affordability and consumption of electronic products, LED lamps and other items, and also accelerated the popularization of smart phones among younger people, thus making it easier to form an unfavorable eye-using environment. In addition, combined with the effects of long-term near work [38, 39] and harmful lights [40], it is not surprising that the prevalence of VI was relatively high. Moreover, the risk effect of CGTRS was also manifested in the above aspects.

As another risk factor, the number of full-time teachers should also receive more attention from the education department. This study found that the risk of a region with abundant teacher resources evolving into an area with high VI level was 1.051 times that of a region with low VI level. The schoolwork pressure caused by teachers may have contributed to this result. Although this problem has been improved in primary schools, it can't be ignored due to the difficulty of entering a higher school (limited enrollment quota). Meanwhile, as the paramount part of the teaching process, teachers' behaviors, attitudes and professional teaching methods will affect children's health [41-43]. Under the background that Chinese parents attach great importance to traditional education, children's burden is already heavy, so teachers should further optimize their behaviors and attitudes to create a health environment for the childhood growth. However, the specific reasons for this result is still unknown. We suggested that future research should focus more on the teachers, so as to further explore the specific causes of this association.

This is the first systematic study on the visual impairment of children aged 6-12 in Shandong province from 2013 to 2017, and the differences among them were expounded from various aspects. The sample involved in the study is huge, and the results are highly credible and regionally

representative. Nevertheless, this research also has some limitations: 1) The year covered by this study is only 5 years, and the data in 2013 are missing a lot. Due to the limited conditions, the data in other years cannot be supplemented, so the results of trend analysis are for reference only. 2) Family, school and other related associated factors [24, 39, 44] were not included in this research, and the impacts of schools' and families' differences among districts on VI cannot be verified. The above problems need further study.

Conclusions

In conclusion, there was a significant phenomenon of high prevalence and aggregation of visual impairment in the eastern region of Shandong province, and economic and educational factors have played a certain role in the development of regional visual impairment. The slight rebound of the VI prevalence in recent years was also worthy of vigilance. Based on the research results presented by various regions, it is suggested that relevant departments should improve children's VI status (especially in Weihai and Yantai) according to their own economic and educational conditions and relevant policies (*Healthy China 2030*, *Healthy China Action*, etc.). Moreover, it is necessary to promote the reform and implementation of quality-oriented education (e.g. 1000m race [31]) and expedite the application and popularization of assistive technologies [45]. Prevention and treatment should be combined to comprehensively improve children's living environment and reduce the VI rate.

Declarations

Acknowledgments

The authors would like to express their gratitude for the support from Maosun Fu, the professor of Department of Social Medicine and Maternal & Child Health, School of Public Health, Shandong University.

370 **Authorship contributions**

371 LH and JJ conceived and designed the study. LW provided the source data of the study. LH and JJ
372 prepared software and performed the statistical analysis. LH prepared the manuscript and interpreted
373 the data. JJ and LW assisted with the editing of the paper and provided critical comments. JJ and LW
374 revised it critically for important intellectual content. All authors read and approved the final
375 manuscript.

376 **Funding**

377 The author(s) received no financial support for the research, authorship, and/or publication of this
378 article.

379 **Competing interests**

380 The authors declare that they have no competing interests.

381 **Availability of data and materials**

382 The datasets generated and/or analysed during the current study are not publicly available due the data
383 is confidential, but are available from the corresponding author on reasonable request.

384 **Ethics approval and consent to participate**

385 Ethics approval was not available in this cross-sectional study because we did not include any data of
386 students' personal information, including name, identity information, address, telephone number, etc.

387 This study only showed the secondary aggregated data on county-level, therefore, waived off ethical
388 approval.

389 **Consent of publication**

390 Not applicable

391 **References**

- 392 1. Jan C, Xu R, Luo D, Xiong X, Song Y, Ma J, et al. Association of visual impairment with
393 economic development among Chinese schoolchildren. *JAMA Pediatr.* 2019;173(7):e190914.
394 doi: 10.1001/jamapediatrics.2019.0914.
- 395 2. Pascolini D, Mariotti SP. Global estimates of visual impairment: 2010. *Br J Ophthalmol.*
396 2012;96(5):614-8. doi: 10.1136/bjophthalmol-2011-300539.
- 397 3. Jones N, Bartlett HE, Cooke R. An analysis of the impact of visual impairment on
398 activities of daily living and vision-related quality of life in a visually impaired adult population.
399 *British Journal of Visual Impairment.* 2019;37(1):50-63. doi: 10.1177/0264619618814071.
- 400 4. Ma YY, Qu XM, Zhu XF, Xu X, Zhu JF, Sankaridurg P, et al. Age-specific prevalence of
401 visual impairment and refractive error in Children aged 3-10 years in Shanghai, China.
402 *Investigative Ophthalmology & Visual Science.* 2016;57(14):6188-96. doi:
403 10.1167/iovs.16-20243.
- 404 5. Ghaderi S, Hashemi H, Jafarzadehpur E, Yekta A, Ostadimoghaddam H, Mirzajani A, et
405 al. The prevalence and causes of visual impairment in seven-year-old children. *Clinical and*
406 *Experimental Optometry.* 2018;101(3):380-5. doi: 10.1111/cxo.12646.
- 407 6. Habib F, Irshad E. Impact of visual impairment on quality of life among adolescents. *Fwu*
408 *Journal of Social Sciences.* 2018;12(1):149-55.
- 409 7. Muurinen SM, Soini HH, Suominen MH, Saarela RKT, Savikko NM, Pitkala KH. Vision
410 impairment and nutritional status among older assisted living residents. *Archives of*
411 *Gerontology and Geriatrics.* 2014;58(3):384-7. doi: 10.1016/j.archger.2013.12.002.
- 412 8. Augestad LB. Mental Health among Children and Young Adults with Visual Impairments:
413 A Systematic Review. *Journal of Visual Impairment & Blindness.* 2017;111(5):411-25.

- 414 9. Rey-Galindo J, Rizo-Corona L, Gonzalez-Munoz EL, Aceves-Gonzalez C. Needs and
415 Use of the Information in the Environment by People with Visual Impairment. In: Bagnara S,
416 Tartaglia R, Albolino S, Alexander T, Fujita Y, editors. Proceedings of the 20th Congress of the
417 International Ergonomics Association. Advances in Intelligent Systems and Computing. 824.
418 Cham: Springer International Publishing Ag; 2019. p. 1622-33.
- 419 10. Haegele JA, Garcia JM, Healy S. The association between neighborhood factors and
420 physical activity and screen-time among youth with visual impairments. Disability and Health
421 Journal. 2019;12(3):509-13. doi: 10.1016/j.dhjo.2019.02.004.
- 422 11. Magdalena W, Urzedowicz B, Motylewski S, Zeman K, Pawlicki L. Body mass index and
423 waist-to-height ratio among schoolchildren with visual impairment A cross-sectional study.
424 Medicine. 2016;95(32):7. doi: 10.1097/md.00000000000004397.
- 425 12. Augestad LB, Jiang L. Physical activity, physical fitness, and body composition among
426 children and young adults with visual impairments: A systematic review. British Journal of
427 Visual Impairment. 2015;33(3):167-82. doi: 10.1177/0264619615599813.
- 428 13. Guan X, Fu M, Lin F, Zhu D, Vuillermin D, Shi L. Burden of visual impairment associated
429 with eye diseases: exploratory survey of 298 Chinese patients. BMJ open. 2019;9(9):e030561.
430 doi: 10.1136/bmjopen-2019-030561.
- 431 14. Chuvarayan Y, Finger RP, Koberlein-Neu J. Economic burden of blindness and visual
432 impairment in Germany from a societal perspective: a cost-of-illness study. The European
433 journal of health economics : HEPAC : health economics in prevention and care. 2019. doi:
434 10.1007/s10198-019-01115-5.
- 435 15. Yan XC, Chen L, Yan H. Socio-economic status, visual impairment and the mediating role

of lifestyles in developed rural areas of China. Plos One. 2019;14(4):18. doi:
10.1371/journal.pone.0215329.

16. National Health Commission of the People's Republic of China, China MoEotPsRo.
Measures for the administration of health examination of primary and middle school students:
National Health Commission of the People's Republic of China,
Ministry of Education of the People's Republic of China; 2008. Available from:
http://www.gov.cn/gongbao/content/2008/content_1175830.htm. Accessed 11.20, 2019.

17. Department SPE. Measures for the implementation of health examination management
for primary and middle school students in Shandong province: Shandong Provincial Education
Department
2009. Available from: http://edu.shandong.gov.cn/art/2009/6/4/art_107055_7734443.html.
Accessed 11.22, 2019.

18. China NHCotPsRo. Standard for logarithmic visual acuity charts: National Health
Commission of the People's Republic of China, Standardization Administration; 2012.
Available from:
<http://openstd.samr.gov.cn/bzgk/gb/newGbInfo?hcno=A9F9E03A346211223DE34421A85CA1C8>. Accessed 10.17, 2019.

19. Organization WH. International Classification of Diseases (ICD)-11: World Health
Organization; 2018. Available from:
<https://icd.who.int/browse11/l-m/en#/http%3a%2f%2fid.who.int%2fcd%2fentity%2f1103667651>. Accessed 10.06, 2019.

20. Colenbrander A. Visual standards: aspects and ranges of vision loss with emphasis on

- 458 population surveys. Sydney, Australia: International Congress of Ophthalmology; 2002.
- 459 21. ESRI. How spatial autocorrelation (Global Moran's I) works: ESRI; 2018. Available from:
- 460 [http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-spatial-autocor](http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-spatial-autocorrelation-moran-s-i-spatial-st.htm)
- 461 [relation-moran-s-i-spatial-st.htm](http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-spatial-autocorrelation-moran-s-i-spatial-st.htm). Accessed 10.11, 2019.
- 462 22. ESRI. How hot spot analysis (Getis-Ord Gi*) works: ESRI.; 2018. Available from:
- 463 [http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-hot-spot-analy](http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-hot-spot-analysis-getis-ord-gi-spatial-statistics.htm)
- 464 [sis-getis-ord-gi-spatial-stati.htm](http://desktop.arcgis.com/en/arcmap/10.3/tools/spatial-statistics-toolbox/h-how-hot-spot-analysis-getis-ord-gi-spatial-statistics.htm). Accessed 10.11, 2019.
- 465 23. Li T, Du LP, Du LZ. Prevalence and causes of visual impairment and blindness in Shanxi
- 466 province, China. Ophthalmic Epidemiology. 2015;22(4):239-45. doi:
- 467 10.3109/09286586.2015.1009119.
- 468 24. Yi HM, Zhang LX, Ma XC, Congdon N, Shi YJ, Pang XP, et al. Poor vision among China's
- 469 rural primary school students: Prevalence, correlates and consequences. China Economic
- 470 Review. 2015;33:247-62. doi: 10.1016/j.chieco.2015.01.004.
- 471 25. Wu JF, Bi HS, Wang SM, Hu YY, Wu H, Sun W, et al. Refractive error, visual acuity and
- 472 causes of vision loss in children in Shandong, China. The Shandong Children Eye Study.
- 473 PLoS One. 2013;8(12):e82763. doi: 10.1371/journal.pone.0082763.
- 474 26. Sherwin JC, Reacher MH, Keogh RH, Khawaja AP, Mackey DA, Foster PJ. The
- 475 association between time spent outdoors and myopia in children and adolescents: a
- 476 systematic review and meta-analysis. Ophthalmology. 2012;119(10):2141-51. doi:
- 477 10.1016/j.ophtha.2012.04.020.
- 478 27. Schaumberg DA, Nichols KK. The global sex disparity in blindness and visual impairment.
- 479 Optometry and Vision Science. 2006;83(10):700-1. doi:

480 10.1097/01.opx.0000244832.29106.10.

481 28. Miller EM. Reported myopia in opposite sex twins - a hormone hypothesis. *Optometry and*
482 *Vision Science*. 1995;72(1):34-6. doi: 10.1097/00006324-199501000-00007.

483 29. Xie H, Mao X, Yang H, Xie Z, Pan Y, Gao Y. Analysis on the relationship between
484 adolescent myopia and serum sex hormone. *Zhonghua yi xue za zhi*. 2014;94(17):1294-7.

485 30. Chen ZTY, Wang IJ, Liao YT, Shih YF, Lin LLK. Polymorphisms in steroidogenesis genes,
486 sex steroid levels, and high myopia in the Taiwanese population. *Molecular Vision*.
487 2011;17(248-50):2297-310.

488 31. Brian A, Pennell A, Haibach-Beach P, Foley J, Taunton S, Lieberman LJ. Correlates of
489 physical activity among children with visual impairments. *Disability and Health Journal*.
490 2019;12(2):328-33. doi: 10.1016/j.dhjo.2018.10.007.

491 32. He MG, Xiang F, Zeng YF, Mai JC, Chen QY, Zhang J, et al. Effect of time spent outdoors
492 at school on the development of myopia among children in China a randomized clinical trial.
493 *Jama-Journal of the American Medical Association*. 2015;314(11):1142-8. doi:
494 10.1001/jama.2015.10803.

495 33. You QS, Wu LJ, Duan JL, Luo YX, Liu LJ, Li X, et al. Factors associated with myopia in
496 school children in China: the Beijing childhood eye study. *Plos One*. 2012;7(12):10. doi:
497 10.1371/journal.pone.0052668.

498 34. Wu PC, Tsai CL, Wu HL, Yang YH, Kuo HK. Outdoor activity during class recess reduces
499 myopia onset and progression in school children. *Ophthalmology*. 2013;120(5):1080-5. doi:
500 10.1016/j.opthta.2012.11.009.

501 35. Houwen S, Hartman E, Visscher C. The relationship among motor proficiency, physical

502 fitness, and body composition in children with and without visual impairments. Research
 503 Quarterly for Exercise and Sport. 2010;81(3):290-9.

504 36. Qin W, Wang L, Xu L, Sun L, Li J, Zhang J, et al. An exploratory spatial analysis of
 505 overweight and obesity among children and adolescents in Shandong, China. BMJ open.
 506 2019;9(8):e028152. doi: 10.1136/bmjopen-2018-028152.

507 37. Cui DM, Trier K, Ribel-Madsen SM. Effect of day length on eye growth, myopia
 508 progression, and change of corneal power in myopic children. Ophthalmology.
 509 2013;120(5):1074-9. doi: 10.1016/j.ophtha.2012.10.022.

510 38. Hsu CC, Huang N, Lin PY, Fang SY, Tsai DC, Chen SY, et al. Risk factors for myopia
 511 progression in second-grade primary school children in Taipei: a population-based cohort
 512 study. British Journal of Ophthalmology. 2017;101(12):1611-7. doi:
 513 10.1136/bjophthalmol-2016-309299.

514 39. Low W, Dirani M, Gazzard G, Chan YH, Zhou HJ, Selvaraj P, et al. Family history, near
 515 work, outdoor activity, and myopia in Singapore Chinese preschool children. British Journal of
 516 Ophthalmology. 2010;94(8):1012-6. doi: 10.1136/bjo.2009.173187.

517 40. Pan CW, Wu RK, Liu H, Li J, Zhong H. Types of lamp for homework and myopia among
 518 Chinese school-aged children. Ophthalmic Epidemiology. 2018;25(3):250-6. doi:
 519 10.1080/09286586.2017.1420204.

520 41. Haegele JA, Zhu XH. Experiences of Individuals With Visual Impairments in Integrated
 521 Physical Education: A Retrospective Study. Research Quarterly for Exercise and Sport.
 522 2017;88(4):425-35. doi: 10.1080/02701367.2017.1346781.

523 42. Sit CHP, McManus A, McKenzie TL, Lian J. Physical activity levels of children in special

524 schools. Preventive Medicine. 2007;45(6):424-31. doi: 10.1016/j.ypmed.2007.02.003.

525 43. Lieberman LJ. Physical education and sport for individuals with visual impairments or
 526 deaf-blindness: Foundations of instruction. New York: American Foundation for the Blind;
 527 2015.

528 44. Ayvazoglu NR, Oh HK, Kozijb FM. Explaining physical activity in children with visual
 529 impairments: A family systems approach. Exceptional Children. 2006;72(2):235-48. doi:
 530 10.1177/001440290607200207.

531 45. Bin Tuwaym ST, Berry AB. Assistive technology for students with visual impairments: a
 532 resource for teachers, parents, and students. Rural Special Education Quarterly.
 533 2018;37(4):219-27. doi: 10.1177/8756870518773397.

534

535 Figure 1. Trends in overall and gender-specific visual impairment (VI) among children in Shandong
 536 province from 2013 to 2017

537 Figure 2. Trends in urban and rural visual impairment (VI) among children in Shandong province from
 538 2013 to 2017

539 Figure 3. Age trends of visual impairment (VI) among children in Shandong province from 2013 to
 540 2017

541 Figure 4. Spatial distribution of total visual impairment (VI) among children in Shandong province in
 542 2013

543 Figure 5. Spatial distribution of total visual impairment (VI) among children in Shandong province in
 544 2014

545 Figure 6. Spatial distribution of total visual impairment (VI) among children in Shandong province in

546 2015

547 Figure 7. Spatial distribution of total visual impairment (VI) among children in Shandong province in

548 2016

549 Figure 8. Spatial distribution of total visual impairment (VI) among children in Shandong province in

550 2017

551 Figure 9. Spatial aggregation of visual impairment (VI) among children in Shandong province in 2013

552 Figure 10. Spatial aggregation of visual impairment (VI) among children in Shandong province in 2014

553 Figure 11. Spatial aggregation of visual impairment (VI) among children in Shandong province in 2015

554 Figure 12. Spatial aggregation of visual impairment (VI) among children in Shandong province in 2016

555 Figure 13. Spatial aggregation of visual impairment (VI) among children in Shandong province in 2017

556

Figures

Figure 1

Trends in overall and gender-specific visual impairment (VI) among children in Shandong province from 2013 to 2017

Figure 2

Trends in urban and rural visual impairment (VI) among children in Shandong province from 2013 to 2017

Figure 3

Age trends of visual impairment (VI) among children in Shandong province from 2013 to 2017

Figure 4

Spatial distribution of total visual impairment (VI) among children in Shandong province in 2013. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country,

territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 5

Spatial distribution of total visual impairment (VI) among children in Shandong province in 2014. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 6

Spatial distribution of total visual impairment (VI) among children in Shandong province in 2015. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 7

Spatial distribution of total visual impairment (VI) among children in Shandong province in 2016. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 8

Spatial distribution of total visual impairment (VI) among children in Shandong province in 2017. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.m

Figure 9

Spatial aggregation of visual impairment (VI) among children in Shandong province in 2013. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 10

Spatial aggregation of visual impairment (VI) among children in Shandong province in 2014. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 11

Spatial aggregation of visual impairment (VI) among children in Shandong province in 2015. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 12

Spatial aggregation of visual impairment (VI) among children in Shandong province in 2016. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Figure 13

Spatial aggregation of visual impairment (VI) among children in Shandong province in 2017. Note: The designations employed and the presentation of the material on this map do not imply the expression of any opinion whatsoever on the part of Research Square concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. This map has been provided by the authors.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Abbreviations.docx](#)
- [Abbreviations.docx](#)