

Costs and Healthcare Utilisation of Patients With Heart Failure in Spain

Carlos Escobar (✉ escobar_cervantes_carlos@hotmail.com)

Hospital Universitario La Paz <https://orcid.org/0000-0001-5584-4735>

Luis Varela

AstraZeneca

Beatriz Palacios

Astrazeneca

Margarita Capel

AstraZeneca

Antoni Sicras

atrys health

Aram Sicras

Atrys Health

Antonio Hormigo

Centro de Salud Puerta Blanca

Roberto Alcazar

Hospital Infanta Leonor

Nicolas Manito

Hospital Universitari de Bellvitge

Manuel Botana

Hospital Universitario Lucus Augusti

Research article

Keywords: cost, DAPA-HF, healthcare, heart failure, hospitalization, medication

Posted Date: August 12th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-50044/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on October 20th, 2020. See the published version at <https://doi.org/10.1186/s12913-020-05828-9>.

Abstract

Background: To evaluate the cumulative costs and healthcare utilisation in heart failure (HF) patients in Spain. This was also analyzed among the DAPA-HF like population.

Methods: Observational, retrospective, population-based study using BIG-PAC database, which included data from specialized and primary care of people aged ≥ 18 years, from 7 autonomous communities in Spain, who received care for HF between 2015 and 2019. The healthcare and medication costs were summarized on a yearly basis starting from the index date (1st January 2015), and then cumulatively until the last year of follow-up (2019).

Results: We identified 17,163 patients with HF (year 2015: mean age 77.3 ± 11.8 years, 53.5% men, 51.7% systolic HF and 43.6% on NYHA functional class II). During the 2015-2019 period, total HF associated costs reached 15,373 Euros per person, being cardiovascular disease hospitalizations the most important determinant (75.8%), particularly HF hospitalizations (51.0%). Total medication cost accounted for 7.0% of the total cost. During this period, there was a progressive decrease of cardiovascular disease hospital costs per year (from 2,834 Euros in 2015 to 2,146 Euros in 2019, $P < 0.001$), as well as cardiovascular and diabetic medication costs. Hospitalization and medication costs were higher in the DAPA-HF like population.

Conclusions: During the 2015-2019 period, costs of patients with HF in Spain were substantial, being HF hospitalizations the most important determinant. Medication costs represented only a small proportion of total HF costs. Improving HF management, particularly through the use of those drugs that reduce HF hospitalization may be helpful to reduce HF burden.

Background

Heart Failure (HF) is a complex clinical syndrome caused by structural or functional cardiac alterations, leading to a reduced cardiac output or elevated intracardiac pressures at rest or during stress, that causes the typical symptoms such as breathlessness, or fatigue [1,2]. HF is a growing problem worldwide [3]. It has been estimated that the prevalence of HF is around 2% of the adult population in developed countries, raising to more than 10% in those individuals aged 70 years or older [4]. In Spain, it has been reported a higher prevalence of HF, reaching 5% in some studies [5,6]. Of note, it is expected that the prevalence of HF will increase in the following years, mainly due to the ageing of the population, the rise in HF risk factors (i.e. hypertension, diabetes) and the better treatment of acute cardiovascular events [3].

Despite traditional HF therapies (i.e. renin angiotensin system inhibitors, beta blockers and aldosterone antagonists), mortality and hospitalization rates remain unacceptably high [7,8]. Thus, the MAGGIC meta-analysis that included individual data on 39,372 patients with HF, from 30 cohort studies showed that 40.2% of patients died after 2.5 years of follow-up [8]. However, in the last years, the PARADIGM-HF and more recently, the DAPA-HF trials have shown that sacubitril/valsartan and dapagliflozin, respectively, have a positive impact on morbidity and mortality among patients with HF and reduced left ventricular ejection fraction [9,10].

Of note, HF represents a major and growing economic problem [3,11,12]. Studies particularly focused on HF economic burden are important as they contribute to a better understanding of the drivers and problems which may lead to the increasing HF costs [3]. Increasing the knowledge about HF costs and their determinants is important in order to ascertain how HF management can be optimized, leading to a significant decrease of HF costs [3,7,11,12]. Unfortunately, data about costs of HF in Spain are very scarce or limited to the use of specific drugs, but not focused on a comprehensive approach [13-17].

The aim of this study was to evaluate the cumulative costs and healthcare utilisation in HF patients in Spain over the last 5 years, along with the epidemiological characterization of the population at index date (1st January 2015). This was also analyzed in a population who met the most relevant inclusion criteria of the DAPA-HF trial [10] (DAPA-HF like population)

with the aim of understanding the costs associated with the management of HF with reduced ejection fraction from the DAPA-HF trial population.

Methods

Observational cohort study, comprising cross-sectional and longitudinal retrospective analyses using secondary data captured in electronic health records from 7 Spanish regions. Data sources were provided by BIG-PAC®. BIG-PAC is an electronic database that integrate information from primary and specialist care medical records. This database has been validated as an information source for studies of epidemiology, therapeutic adaptation and health/non-healthcare resource use and associated costs. It is representative of the Spanish population [18]

The HF population was defined as all patients ≥ 18 years of age and with at least one diagnosis of HF prior to the index date (1st January 2015). Type 2 diabetes (T2D) was defined as all patients ≥ 18 years of age filling a prescription of any antidiabetic medication, T2D diagnosis or HbA1c $> 7\%$ prior to index date. The Dapa-HF like population included those patients ≥ 18 years, with a HF history > 8 weeks, treatment with device therapy or standard HF treatment, NYHA class $\geq II$ within one year prior to index date and left ventricular ejection fraction $\leq 40\%$. Patients with HF hospitalization < 4 weeks prior to index date, myocardial infarction, unstable angina pectoris, stroke, transient ischemic attack, coronary revascularization or implantation of therapeutic device < 12 weeks prior to index date, were excluded from the population.

Comorbidities were searched for in all available data prior to index date, excepting for severe hypoglycemia that was considered only within 1 year before index date and cancer which was searched for up to 5-year prior to index date. A minimum of 1 year of data before index date was required. Comorbidities included cardiovascular disease (myocardial infarction, percutaneous or surgical revascularization, unstable angina, angina pectoris), HF, chronic kidney disease, atrial fibrillation, stroke (hemorrhagic, ischemic, transitory ischemic attack), peripheral artery disease, major organ specific bleeding, microvascular complications (diabetic mono-/polyneuropathy, diabetic eye complications, diabetic kidney disease), severe hypoglycemia, cancer, chronic obstructive pulmonary disease, and lower limb complications. ICD-9 and ICD-10 codes (<https://eciemaps.mscbs.gob.es>) were considered for the diagnosis of comorbidities (supplementary table 1).

The information about treatment was recorded from the registries for dispensing medicines, according to the Anatomical Therapeutic Chemical Classification System (supplementary table 1) [19]. HF treatments (angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, beta blockers, aldosterone antagonists, sacubitril/valsartan, loop diuretics, digoxin), warfarin, statins, aspirin, receptor P2Y₁₂ antagonists, calcium channel blockers, thiazides, nitrates, antidiabetic medications (SGLT-2 inhibitors, metformin, sulfonylurea, DPP-4 inhibitors, GLP-1 receptor agonists, metiglinides, glitazones, acarbose, miglitol, insulin) were recorded. The prescription of a drug in a specific patient was based only on medical criteria (clinical practice).

Prevalence, incidence and baseline characteristics (total HF population and by T2D status), including demographics, comorbidities and medications were calculated at index date (1st January 2015).

The healthcare resource use and costs and medication costs were summarized for the total HF population on a yearly basis starting from index date (1st January 2015), and then cumulatively until the end of the last year of follow up (31st December 2019). All hospital visits (total and cardiovascular events), the number of medical visits and emergency room visits and medication costs (total, cardiovascular related, HF related and diabetes related) were included for the analysis of the annual direct healthcare costs. Patients who died during follow-up had a cost of 0 allocated to the remaining duration of the study, whereas a patient leaving the database prior to data cut off was not included in the denominator for the time after leaving the database. Annual indirect non-health costs included the number of days of productivity lost due to disability.

Rates were obtained from hospital accounting, except for the medication and indirect costs which were calculated as follows, respectively: a) medical prescriptions: according to the retail price per package at the time of dispensing [20]; b)

costs for days of productivity lost: according to the mean interprofessional wage [21]. Hospital admission costs for cardiovascular events during follow-up were obtained taking into consideration daily hospital rate and number of hospital days per stay. Rates are summarized in supplementary table 2.

Statistical analysis.

Categorical variables were described by their absolute (n) and relative frequencies (%). Continuous variables were described using the mean and standard deviation. Categorical variables were compared with the Chi-square test or the Fisher exact test when appropriate. When two means were compared, the t-student test was used. Analyses of health care cost were performed for the index date with 5 year of follow-up. The cumulative mean healthcare cost was estimated and presented on a yearly basis from the index date until last year of follow-up. Health care costs were presented per patient (mean cost). A level of statistical significance of 0.05 was applied in all the statistical tests. The data were analyzed using the statistical package SPSS v22.0 (SPSS Inc., Chicago, Illinois, USA).

Results

Out of 1,7 millions of persons included in the BIG-PAC® database in 2015, 1,3 million people were attended during the 2012-2014 period, of whom 964,862 were 18 years or older. At index date, 17,598 patients had HF. As 435 patients were excluded due to inconsistent data, 17,163 patients (1.78%) comprised the HF study population (figure 1). Incidence at index date was 2.84 x 1000 patient-years.

The baseline clinical characteristics of the HF population according to the presence of T2D were presented in table 1. Overall, mean age was 77.3±11.8 years, 53.5% of patients were men, 51.7% of patients had reduced left ventricular ejection fraction HF, and the majority of patients were on NYHA functional class II (43.6%) or III (36.1%). A total of 5,815 (33.9%) patients had T2D. The presence of other comorbidities was common: 30.0% of patients had atrial fibrillation, 29.5% chronic kidney disease, 23.3% ischemic heart disease, 17.2% chronic obstructive pulmonary disease, 12.8% cancer and 9.5% previous stroke. Two thirds of patients were taking renin angiotensin system inhibitors, 68.3% beta blockers, 30.0% aldosterone antagonists, and 8.5% sacubitril/valsartan. The baseline clinical profile of patients according to the presence of T2D was also compared. The percentage of patients with left ventricular ejection fraction ≤40% was higher in patients with type 2 diabetes (53.1% vs 51.0%; P=0.015). Those patients with T2Dd had more ischemic heart disease, stroke, atrial fibrillation, peripheral artery disease, and chronic kidney disease (P<0.05 or less for all comorbidities).

Patient hospital mean cost for year was presented in table 2. In general, from 2015 to 2019, there was a progressive decrease of cardiovascular disease hospital cost per patient year (from 2,834 to 2,146 Euros, P<0.001). Overall, patient cumulative cardiovascular disease hospital mean cost reached 11,649 Euros in 2019 (supplementary table 3 and figure 2). The great burden for this cost was due to cardiorenal (HF and/or chronic kidney disease) hospitalizations (88.8% of the total cost), particularly HF (67.3% of the total cost). With regard to medication, from 2015 to 2019, diabetes medication mean cost varied from 101 to 85 Euros (P<0.001) per patient and year and HF medication mean cost from 86 to 90 Euros (P<0.001), respectively (table 2). The cumulative mean cost of diabetes medication and HF medication reached 486 and 417 Euros, respectively, in 2019 (supplementary table 3 and figure 3).

The health resources use for each year, including primary care visits, specialized visits, emergency rooms visits, hospitalization and diagnostic tests, was shown in table 3. The proportion of hospitalized patients decreased from 31.5% in 2015 to 22.9% in 2019 (P<0.001), the days for hospitalized patients due to HF from 8.3 to 7.2 days (P<0.001), and the proportion of patients that died from 9.4% to 4.9% (P<0.001), respectively. Total sanitary cost decreased from 3,700 Euros in 2015 to 2,770 Euros in 2019 (P<0.001). Including indirect costs, total cumulative patient mean costs reached 15,373 Euros in 2019, 263,852,978 Euros per total HF population (table 4).

A specific analysis was performed in the DAPA-HF like population (n=3,178). In this subpopulation, mean age was 76.9±11.7 years, 51.4% were men, and all patients had reduced left ventricular ejection fraction HF (mean left ventricular ejection

fraction $34.5 \pm 7.9\%$). The majority of patients were on NYHA functional class II (52.1%) or III (43.0%). A total of 1,314 (41.3%) patients had T2D. With regard HF medication, 77.0% of patients were taking renin angiotensin system inhibitors, 100% beta blockers, 21.0% aldosterone antagonists, and 8.8% sacubitril/valsartan. Compared with patients without diabetes, those patients with T2D were taking more renin angiotensin system inhibitors, aldosterone antagonists and sacubitril/valsartan (supplementary table 4).

With regard to patients hospital mean cost for year for this subpopulation, there was a progressive decrease of cardiovascular disease hospital cost per year (from 3,269.6 Euros in 2015 to 2,539.5 Euros in 2019, $P < 0.001$). Overall, patient cumulative cardiovascular disease hospital mean cost reached 13,775 Euros in 2019. The great burden for this cost was due to cardiorenal hospitalizations (87.7% of the total hospital cost), particularly HF (65.9% of the total hospital cost). With regard to medication, from 2015 to 2019, diabetes medication mean cost decreased from 128.5 to 74.6 Euros ($P < 0.001$) and HF medication mean cost from 112.9 to 74.7 Euros ($P < 0.001$), respectively. The cumulative mean cost of diabetes medication and HF medication reached 540 and 514 Euros, respectively, in 2019 (supplementary table 5).

Discussion

Our study showed that in Spain, during the 2015-2019 period HF associated costs were high (patient total cost of 15,373 Euros), being cardiovascular hospitalizations the most important determinant (75.8%), particularly HF hospitalizations (51.0%). Total medication cost accounted for 7% of the total HF cost. In addition, the annual cardiovascular hospitalization mean cost progressive decreased over time.

In our study, the prevalence of HF was about 1.8%. With regards to the HF population, mean age was 77 years, around half of patients had systolic HF, the majority of patients were on NYHA functional class II or III, one third had diabetes and comorbidities were common. In Spain, the studies performed in different clinical settings (hospital and outpatients) show a higher prevalence of HF [22]. However, the population-based studies report similar numbers to our study [23]. The proportion of patients with systolic HF, as well as the presence of numerous comorbidities are in line with previous studies [5,23]. As a result, our study can be considered fairly representative of the Spanish population.

With regards to HF therapies, approximately two thirds of patients were taking renin angiotensin system inhibitors and beta blockers, nearly one third aldosterone antagonists, and only 8.5% sacubitril/valsartan. The optimization of treatment of patients with HF is necessary not only to improve functional class and quality of life, but also to reduce morbidity and mortality [1,2]. These numbers are lower than those reported in HF units, but in line with those from other clinical settings [24,25]. This is very relevant, as the underuse of evidence-based HF medication is associated with a higher use of healthcare resources, particularly first and recurrent hospitalizations [26].

During the period 2015-2019, patient cumulative cardiovascular disease hospital mean cost reached 11,649 Euros. Importantly, cardiorenal hospitalizations were the most important contributor for the total cost, particularly HF hospitalizations. Overall, HF hospitalizations represent 1-2% of total admissions [7,27] and HF is the most common diagnosis in elderly hospitalized patients [28]. During the first year after diagnosis of HF, approximately half of the patients may be expected to be hospitalized at least once. In addition, readmission rates are high [7,29-31]. Of note, it has been reported that in Spain, rates of first hospitalization due to HF continue to increase, with high mortality [31]. A recent systematic review analyzed 16 cost-of-illness studies related to HF. Although there were large variations concerning cost components, the majority of them showed that hospital admission costs were the most expensive cost element. Annual costs for HF patients ranged from 868 Dollars (≈ 774 Euros) for South Korea to 25,532 Dollars ($\approx 22,760$ Euros) for Germany [3]. Other systematic review focused on economic HF burden also showed that hospitalization cost was found to be the main cost driver to the total health care cost and that the HF annual cost ranged from 908 Dollars (≈ 809 Euros) to 40,971 Dollars ($\approx 36,522$ Euros) per patient [12]. In our study, during the 2015-2019 period, HF associated costs per patient reached 15,373 Euros, in line with these studies. However, among other factors, methodological heterogeneity and specific cost items

(including treatments) accounted for in the estimations indicate that cost comparisons across publications should be made with caution [32]. Certainly, all these data confirm the high cost burden of HF.

As the most important contributor for HF cost is HF hospitalizations, the use of those drugs that have demonstrated to be beneficial in this clinical context may be very helpful in reducing total HF cost. Thus, in 2014 the PARADIGM-HF trial showed that compared with enalapril, sacubitril/valsartan significantly reduced the risk of HF hospitalization by 21% and this might have had a positive impact [9]. In our study, from 2015 to 2019, in general there was a progressive reduction of cardiovascular disease hospital cost per year, as the proportion of hospitalized patients decreased. Of note, there was only a slight increase in HF medication cost per year which is a small contributor for total HF cost. This is in line with previous studies that have shown a decline in standardized HF hospitalization rates in Europe and United States [33,34]. However, absolute numbers of HF hospital admissions are expected to increase by about 50% in the following years due to the ageing of the population [7]. As a result, new drugs are needed to improve these numbers. In 2019, the DAPA-HF trial showed that in addition to recommended therapy, dapagliflozin significantly reduced the risk of a first worsening HF event by 30% [10]. Therefore, it can be hypothesized the addition of dapagliflozin to standard HF therapy may contribute to reduce HF costs.

Other contributors to total HF cost included primary care visits, specialized visits, and diagnostic tests. It has been reported that a better integrated hospital primary care HF program is associated with a significant reduction of readmission for HF and mortality [35]. In addition, moving to case management at home rather than outpatient cardiology clinic follow-up may also reduce healthcare costs [36]. Therefore, transition to an integrated management of HF patients is necessary to reduce HF burden.

A recent meta-analysis estimated the 1, 2, 5 and 10-year survival to be 87%, 73%, 57% and 35%, respectively, among HF patients [37]. HF hospitalization is an independent predictor for increased HF mortality [7,29,30]. In our study the proportion of patients who died decreased from 9.4% in 2015 to 4.9% in 2019, in line with the decrease in hospitalization rates. Although there is much room for progress, it is likely that the improvement in HF management during these years may have had a positive impact.

A specific analysis was performed in the DAPA-HF like population. In the DAPA-HF trial, the addition of dapagliflozin resulted in a significant reduction of HF hospitalizations, death from cardiovascular causes, and death from any cause, regardless the presence of diabetes [10]. In our study, in the DAPA-HF like population, all patients had reduced left ventricular ejection fraction HF and the majority of patients were on NYHA functional class II or III. Compared with the overall HF population, patients were taking more renin angiotensin system inhibitors and beta blockers. Although in these patients there was a decrease of patients hospital mean cost for year, these were higher than in the overall HF population. Thus, cumulative cardiovascular disease hospital cost reached 13,775 Euros (vs 11,649 Euros in the overall HF population). The great burden for this cost was due to cardiorenal hospitalizations (87.7% of the total hospital cost), particularly HF (65.9% of the total hospital cost). Therefore, to reduce HF cost burden in the DAPA-HF like population is of utmost importance to improve the HF management. As the great majority of these patients were taking renin angiotensin system inhibitors and beta blockers, the prescription of newer HF drugs, such as dapagliflozin, could be of particular benefit in the reduction of HF costs [10].

This study has some limitations that should be commented. This was an observational cohort study that used secondary data from electronic health records. As a result, only indirect causality can be provided. However, the high number of patients included, as well as the robustness of the data may allow to determine the value of the study. On the other hand, although data came from 7 Spanish regions, previous studies have shown that these data are representative of the Spanish population [18].

Conclusion

During the 2015-2019 period, costs of patients with HF in Spain were high, being cardiovascular hospitalizations the most important determinant, particularly HF hospitalizations. Medication costs were responsible for only a small proportion of

total HF costs. Costs and healthcare resources use were even higher in the DAPA-HF like population. Improving HF management, particularly through the use of those drugs that reduce HF hospitalization may be helpful to reduce HF burden.

Abbreviations

HF: heart failure.

T2D: type 2 diabetes

Declarations

- **Ethics approval and consent to participate:** This study was approved by the Investigation Ethics Committee of Consorci Sanitari from Terrassa on 16th December 2019. This was a secondary data study and data were fully anonymized and dissociated from patients. Therefore, there was no need for providing informed consent in Spain.
- **Consent for publication:** All authors have provided comments and consent for publication.
- **Availability of data and material:** This was a secondary data study using BIG PAC® database.
- **Competing interests:** None.
- **Funding:** This study was fully funded by AstraZeneca.
- **Authors' contributions:** All authors have contributed to the study design, result review and manuscript preparation.
- **Acknowledgements:** None.

References

1. Yancy CW, Jessup M, Bozkurt B, et al. 2017 ACC/AHA/HFSA Focused Update of the 2013 ACCF/AHA Guideline for the Management of Heart Failure: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Failure Society of America. *Circulation*. 2017;136:e137-e161.
2. Ponikowski P, Voors AA, Anker SD, et al. 2016 ESC Guidelines for the Diagnosis and Treatment of Acute and Chronic Heart Failure. *Eur Heart J*. 2016;37(27):2129-2200.
3. Lesyuk W, Kriza C, Kolominsky-Rabas P. Cost-of-illness studies in heart failure: a systematic review 2004-2016. *BMC Cardiovasc Disord*. 2018;18(1):74.
4. van Riet EE, Hoes AW, Wagenaar KP, Limburg A, Landman MA, Rutten FH. Epidemiology of heart failure: the prevalence of heart failure and ventricular dysfunction in older adults over time. A systematic review. *Eur J Heart Fail*. 2016;18:242-252.
5. Sayago-Silva I, García-López F, Segovia-Cubero J. Epidemiology of heart failure in Spain over the last 20 years. *Rev Esp Cardiol (Engl Ed)*. 2013;66:649-656.
6. Conde-Martel A, Hernández-Meneses M. Prevalence and prognostic meaning of comorbidity in heart failure. *Rev Clin Esp*. 2016;216:222-228.
7. Groenewegen A, Rutten FH, Mosterd A, Hoes AW. Epidemiology of heart failure [published online ahead of print, 2020 Jun 1]. *Eur J Heart Fail*. 2020;10.1002/ehhf.1858.
8. Pocock SJ, Ariti CA, McMurray JJ, et al. Predicting survival in heart failure: a risk score based on 39,372 patients from 30 studies. *Eur Heart J*. 2013;34(19):1404–1413.
9. McMurray JJ, Packer M, Desai AS, et al. Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N Engl J Med*. 2014;371:993-1004.

10. McMurray JJV, Solomon SD, Inzucchi SE, et al. Dapagliflozin in Patients with Heart Failure and Reduced Ejection Fraction. *N Engl J Med*. 2019;381(21):1995-2008.
11. Lee H, Oh SH, Cho H, Cho HJ, Kang HY. Prevalence and socio-economic burden of heart failure in an aging society of South Korea. *BMC Cardiovasc Disord*. 2016;16(1):215.
12. Shafie AA, Tan YP, Ng CH. Systematic review of economic burden of heart failure. *Heart Fail Rev*. 2018;23(1):131-145.
13. Romero-González G, Ravassa S, González O, et al. Burden and challenges of heart failure in patients with chronic kidney disease. A call to action. *Nefrologia*. 2020;40(3):223-236.
14. Merino M, Jiménez M, Manito N, et al. The social return on investment of a new approach to heart failure in the Spanish National Health System. *ESC Heart Fail*. 2020;7(1):130-137.
15. Hernández-Madrid A, Lu X, Tsintzos SI, et al. Heart failure hospitalization reduction and cost savings achieved by improved delivery of effective biventricular pacing: economic implications of the OLE study under the US setting. *Clinicoecon Outcomes Res*. 2019;11:385-393.
16. Cainzos-Achirica M, Capdevila C, Vela E, et al. Individual income, mortality and healthcare resource use in patients with chronic heart failure living in a universal healthcare system: A population-based study in Catalonia, Spain. *Int J Cardiol*. 2019;277:250-257.
17. Manito Lorite N, Rubio-Rodríguez D, González Costello J, et al. Economic analysis of intermittent intravenous outpatient treatment with levosimendan in advanced heart failure in Spain. *Rev Esp Cardiol (Engl Ed)*. 2020;73(5):361-367.
18. Sicras-Mainar A, Enríquez JL, Hernández I, Sicras-Navarro A, Aymerich T, León M. Validation and representativeness of the Spanish BIG-PAC database: integrated computerized medical records for research into epidemiology, medicines and health resource use (Real World Evidence). *Value Health*. 2019;(Supplement 3):S734.
19. The Anatomical Therapeutic Chemical Classification System with Defined Daily Doses (ATC/DDD): World Health Organization. Available at: <https://www.who.int/classifications-/atcddd/en/>. Last accessed: 10 June 2020.
20. General Council of Official Pharmacists' Colleges of Spain. Available at: <https://botplusweb.portalfarma.com>.
21. Statistics National Institute. 2017. Average labor profit according to age and gender. Accessed: 10 May 2020 Available at: <https://www.ine.es/dynt3/inebase/index.htm?padre=4563&capsel=4563>.
22. Anguita Sánchez M, Crespo Leiro MG, de Teresa Galván E, Jiménez Navarro M, Alonso-Pulpón L, Muñiz García J; PRICE Study Investigators. Prevalence of heart failure in the Spanish general population aged over 45 years. The PRICE Study. *Rev Esp Cardiol*. 2008;61:1041-1049.
23. Farré N, Vela E, Clèries M, et al. Real world heart failure epidemiology and outcome: A population-based analysis of 88,195 patients. *PLoS One*. 2017;12:e0172745.
24. Crespo-Leiro MG, Segovia-Cubero J, González-Costello J, et al.; project research team.. Adherence to the ESC Heart Failure Treatment Guidelines in Spain: ESC Heart Failure Long-term Registry. *Rev Esp Cardiol*. 2015;68:785-793.
25. Greene SJ, Butler J, Albert NM, DeVore AD, Sharma PP, Duffy CI, et al. Medical Therapy for Heart Failure With Reduced Ejection Fraction: The CHAMP-HF Registry. *J Am Coll Cardiol*. 2018;72:351-366.
26. Sicras Mainar A, Navarro Artieda R, Ibáñez Nolla J. Economic impact of heart failure according to the effects of kidney failure. *Rev Esp Cardiol*. 2015;68:39-46.
27. Alla F, Zannad F, Filippatos G. Epidemiology of acute heart failure syndromes. *Heart Fail Rev*. 2007;12:91–95.
28. Braunwald E. The war against heart failure: the Lancet lecture. *Lancet*. 2015;385:812-824.
29. Nichols GA, Reynolds K, Kimes TM, Rosales AG, Chan WW. Comparison of risk of re-hospitalization, all-cause mortality, and medical care resource utilization in patients with heart failure and preserved versus reduced ejection fraction. *Am J Cardiol* 2015;116:1088–1092.
30. Shah KS, Xu H, Matsouaka RA, et al. Heart failure with preserved, borderline, and reduced ejection fraction: 5-year outcomes. *J Am Coll Cardiol*. 2017;70:2476–2486.

31. Fernández-Gassó L, Hernando-Arizaleta L, Palomar-Rodríguez JA, Abellán-Pérez MV, Hernández-Vicente Á, Pascual-Figal DA. Population-based Study of First Hospitalizations for Heart Failure and the Interaction Between Readmissions and Survival. *Rev Esp Cardiol (Engl Ed)*. 2019;72:740-748.
32. Oliva J, Jorgensen N, Rodríguez JM. Carga socioeconómica de la insuficiencia cardíaca: revisión de los estudios de coste de la enfermedad. *Pharmacoecon Span Res Artic* 2010;7:68–79.
33. Schmidt M, Ulrichsen SP, Pedersen L, Bøtker HE, Sørensen HT. Thirty-year trends in heart failure hospitalization and mortality rates and the prognostic impact of co-morbidity: a Danish nationwide cohort study. *Eur J Heart Fail* 2016;18:490–499.
34. Ziaeeian B, Kominski GF, Ong MK, Mays VM, Brook RH, Fonarow GC. National differences in trends for heart failure hospitalizations by sex and race/ethnicity. *Circ Cardiovasc Qual Outcomes* 2017;10:e003552.
35. Comín-Colet J, Verdú-Rotellar JM, Vela E, et al. Efficacy of an integrated hospital-primary care program for heart failure: a population-based analysis of 56,742 patients. *Rev Esp Cardiol (Engl Ed)*. 2014;67(4):283-293.
36. van Voorst H, Arnold AER. Cost and health effects of case management compared with outpatient clinic follow-up in a Dutch heart failure cohort. *ESC Heart Fail*. 2020;7(3):1136-1144.
37. Jones NN, Roalfe AK, Adoki I, Hobbs FD, Taylor CJ. Survival of patients with chronic heart failure in the community: a systematic review and meta-analysis. *Eur J Heart Fail* 2019;21:1306–1325.

Tables

Table 1. Baseline clinical characteristics of the heart failure population at index date 1st January 2015 according to the presence of type 2 diabetes.

	No T2D 66.1%)	(n=11,348;	T2D (n=5,815; 33.9%)	Total HF (n=17,163; 100%)	P*
Biodemographic data					
Age, years	77.4±12.6		77.2±10.1	77.3±11.8	0.236
≥85 years, n (%)	6,308 (55.6)		2,648 (45.5)	8956 (52.2)	<0.001
Gender, male, n (%)	6,052 (53.3)		3,135 (53.9)	9,187 (53.5)	0.470
Systolic blood pressure (mmHg)	128.7±21.9		131.8±22.6	129.8±22.2	<0,001
Heart failure data					
NYHA functional class, n (%)					
I	1,321 (11.6)		654 (11.3)	1,975 (11.5)	
II	5,008 (44.1)		2,476 (42.6)	7,484 (43.6)	0.001
III	4,068 (35.9)		2,135 (36.7)	6,203 (36.1)	
IV	395 (3.5)		268 (4.6)	663 (3.9)	
Left ventricular ejection fraction, %	44.2±10.6		41.5±12.4	43.3±11.3	<0.001
≤40%, n (%)	5,217 (51.0)		2,798 (53.1)	8,015 (51.7)	0.012
>40 - <50%, n (%)	801 (7.8)		428 (8.1)	1,229 (7.9)	0.012
≥50%, n (%)	4,217 (41.2)		2,039 (38.7)	6,256 (40.4)	0.012
Laboratory data					
eGFR, ml/min/1.73 m ²	76.1±20.4		72.3±20.4	74.8±20.5	<0.001
HbA1c, %	5.2±0.9		7.3±0.8	6.1±1.4	<0.001
Comorbidities					
Cardiovascular disease, n (%)					
Ischemic heart disease	2174 (19.1)		1834 (31.5)	4006 (23.3)	<0.001
Myocardial infarction	1,350 (11.9)		1,126 (19.4)	2,466 (14.4)	<0.001
CABG	122 (1.1)		132 (2.3)	254 (1.5)	0.001
PCI with stent	285 (2.5)		243 (4.2)	528 (3.1)	<0.001
Unstable angina	414 (3.7)		384 (6.6)	798 (4.7)	<0.001
Angina pectoris	686 (6.1)		493 (8.5)	1,179 (6.9)	<0.001
Stroke	1,015 (8.9)		622 (10.7)	1,637 (9.5)	<0.001
Ischemic stroke	779 (6.9)		467 (8.0)	1,246 (7.3)	0.005
Hemorrhagic stroke	72 (0.6)		28 (0.5)	100 (0.6)	0.213
Transitory ischemic attack	256 (2.3)		173 (3.0)	429 (2.5)	0.004
Atrial Fibrillation	3,339 (29.4)		1,805 (31.0)	5,144 (30.0)	0.029
Peripheral artery disease	500 (4.4)		326 (5.6)	826 (4.8)	0.001

Chronic kidney disease	2,819 (24.8)	2,236 (38.5)	5,055 (29.5)	<0.001
Microvascular complications	0	1,810 (31.1)	1,810 (10.5)	<0.001
Diabetic mono-/polyneuropathy	0	380 (6.5)	380 (2.2)	<0.001
Diabetic eye complications	0	1,598 (27.5)	1,849 (10.8)	<0.001
Diabetic foot/peripheral angiopathy	0	251 (4.3)	264 (1.5)	<0.001
Diabetic kidney disease	0	468 (8.1)	521 (3.0)	<0.001
Diabetes with complications	0	3,207 (55.2)	3,452 (20.1)	<0.001
Severe hypoglycemia	0	461 (7.9)	461 (2.7)	<0.001
Other comorbidities, n (%)				
Cancer	1,465 (12.9)	733 (12.6)	2,198 (12.8)	0.572
Chronic obstructive pulmonary disease	1,928 (17.0)	1,027 (17.7)	2,955 (17.2)	0.270
Lower limb amputations	37 (0.3)	108 (1.9)	145 (0.8)	<0.001
Major organ specific bleeding	111 (1.0)	177 (3.0)	288 (1.7)	<0.001
Therapies				
Heart failure medications, n (%)				
Renin angiotensin system inhibitors	6,973 (61.5)	4,452 (76.6)	11,425 (66.6)	<0.001
Angiotensin-converting enzyme inhibitors	3,354 (29.6)	1,986 (34.2)	5,340 (31.1)	<0.001
Angiotensin receptor blockers	3,619 (31.9)	2,466 (42.4)	6,085 (35.5)	<0.001
Beta blockers	7,509 (66.2)	4,218 (72.5)	11,727 (68.3)	<0.001
Loop-diuretics	7,416 (65.4)	4,645 (79.9)	12,061 (70.3)	<0.001
Aldosterone antagonists	3,424 (30.2)	1,728 (29.7)	5,152 (30.0)	0.537
Sacubitril/valsartan	987 (8.7)	477 (8.2)	1,464 (8.5)	0.272
Digoxin	799 (7.0)	398 (6.8)	1,197 (7.0)	0.632
Other cardiovascular medications, n (%)				
Low dose aspirin	3,017 (26.6)	2,037 (35.0)	5,054 (29.5)	<0.001
Receptor P2Y12 antagonists	1,177 (10.4)	634 (10.9)	1,811 (10.6)	0.284
Statins	5,928 (52.2)	3,238 (55.7)	9,166 (53.4)	<0.001
Antihypertensives	2,415 (21.3)	1,478 (25.4)	3,893 (22.7)	<0.001
Dihydropyridines CCB	1,564 (13.8)	1,027 (17.7)	2,591 (15.1)	<0.001
Thiazides	520 (4.6)	268 (4.6)	788 (4.6)	0.938
Non-hydropyridines CCB	504 (4.4)	290 (5.0)	794 (4.6)	0.107
Nitrates	1,169 (10.3)	788 (13.6)	1,957 (11.4)	<0.001
Warfarin	2,438 (21.5)	1,464 (25.2)	3,902 (22.7)	<0.001
Diabetes medications, n (%)	0	5,383 (92.6)	5,383 (31.4)	<0.001

Metformin	0	3,862 (66.4)	3,862 (22.5)	<0.001
Sulfonylurea	0	1,952 (33.6)	1,952 (11.4)	<0.001
DPP4 inhibitors	0	1,416 (24.4)	1,416 (8.3)	<0.001
SGLT-2 inhibitors	1 (0%)	2 (0)	3 (0)	<0.001
GLP-1 receptor agonists	0	57 (1.0)	57 (0.3)	<0.001
Metiglinides	0	319 (5.5)	319 (1.9)	<0.001
Glitazones	0	57 (1.0)	57 (0.3)	<0.001
Acarbose	0	88 (1.5)	88 (0.5)	<0.001
Insulin	0	1,367 (23.5)	1,367 (8.0)	<0.001

T2D: type 2 diabetes; HF: heart failure; eGFR: estimated glomerular filtration rate; CABG: coronary artery bypass graft; PCI: percutaneous coronary intervention; CCB: calcium channel blockers; DPP4: dipeptidyl peptidase 4; SGLT-2: sodium-glucose Cotransporter-2; GLP-1: glucagon-like peptide-1.

*p values comparing no T2D vs T2D

Table 2. Patients hospital mean cost for year*.

	2015		2016		2017		2018		2019		Cumulative cost in 2019
	mean	SD									
Total hospital cost											
CVD	2,834	5,212	2,416	5,065	2,268	4,839	1,985	4,937	2,146	4,947	11,649
Cardiorenal	2,536	4,859	2,155	4,565	2,001	4,433	1,759	4,329	1,896	4,460	10,346
HF	1,967	4,094	1,657	3,814	1,476	3,609	1,297	3,505	1,447	3,604	7,842
CKD	569	2,173	498	2,024	525	2,057	463	1,943	450	1,968	2,504
MI	98	812	87	800	79	760	70	776	83	761	416
Stroke	138	970	126	942	132	977	108	940	113	887	616
PAD	63	740	49	619	57	682	48	660	54	602	271
Medication cost											
Total medication	230	425	216	426	227	426	204	432	206	421	1,083
Diabetes medication	101	336	97	336	118	324	85	308	85	317	486
HF medication	86	130	79	131	76	203	86	268	90	313	417
CVD medication	44	96	40	91	33	80	33	86	31	339	181

*In Euros.

CVD: cardiovascular disease; HF: heart failure; CKD: chronic kidney disease; cardiorenal: HF and/or CKD; MI: myocardial infarction; PAD: peripheral artery disease.

Table 3. Health resources use for each year per patient.

	2015		2016		2017		2018		2019	
	mean	SD								
Primary care visits, mean (SD)	12.7	15.0	10.9	14.2	9.6	13.9	8.8	14.3	7.7	13.0
Laboratory requests, mean (SD)	0.6	1.2	0.6	1.2	0.6	1.2	0.7	1.6	1.0	2.1
Radiology and other tests, mean (SD)	0.6	1.2	0.6	1.2	0.6	1.2	0.7	1.6	1.0	2.2
Specialized visits, mean (SD)	2.1	4.2	1.3	4.9	1.1	4.8	1.2	4.9	1.1	4.8
Emergency rooms visits, mean (SD)	0.8	2.5	0.7	3.3	0.7	3.1	0.5	2.3	0.5	3.3
Hospitalization										
Days, mean (SD)	5.9	10.8	5.0	10.5	4.7	10.0	4.1	10.3	4.5	10.3
Hospitalized patients, n (%)	5,399 (31.5)		4,557 (26.6)		4,227 (24.6)		3,628 (21.1)		3,923 (22.9)	
Days for hospitalized patients due to heart failure only, mean (SD)	8.3	10.7	7.7	10.6	7.5	10.1	6.9	10.1	7.2	10.2
Frequency of hospitalization, n (%)										
0	11,764 (68.5)		12,606 (73.5)		12,936 (75.4)		13,535 (78.9)		13,240 (77.1)	
1	4,243 (24.7)		3,565 (20.8)		3,225 (18.8)		2,780 (16.2)		2,914 (17.0)	
2	1,001 (5.8)		855 (5.0)		867 (5.1)		724 (4.2)		850 (5.0)	
3+	156 (0.9)		137 (0.8)		137 (0.8)		124 (0.7)		160 (0.9)	
Disability										
Days of disability, mean (SD)	0.4	5.9	0.4	6.0	0.6	10.9	0.4	5.7	0.4	5.1
Patients with disability, n (%)	158 (0.9)		132 (0.8)		134 (0.8)		140 (0.8)		190 (1.1)	
Average days of sick leave (disability only), mean (SD)	43.8	43.5	47.4	49.5	77.2	96.9	50.5	38.5	36.7	32.6
Mortality, n (%)	1,608 (9.4)		1,259 (7.3)		927 (5.4)		893 (5.2)		839 (4.9)	

All years calculated with 17.163 patients

Table 4. Patients total mean cost for year and cumulative cost in 2019*.

	2015		2016		2017		2018		2019		Cumulative cost in 2019
	mean	SD									
Cost of primary care visits	308	362	264	344	231	336	214	345	186	315	1,202
Cost of laboratory requests	20	39	19	38	18	37	24	50	32	69	113
Cost of radiology and other tests	24	46	22	45	21	45	28	59	37	80	131
Cost of specialized visits	195	392	123	467	108	451	108	462	101	452	635
Cost of emergency rooms visits	89	291	81	394	79	363	57	276	61	382	368
Cost of Hospitalization	2,834	5,212	2,416	5,065	2,268	4,839	1,985	4,937	2,146	4,947	11,649
Cost of medication	230	425	216	426	227	426	204	432	206	421	1,083
Sanitary Cost	3,700	5,623	3,141	5,516	2,920	5,347	2,620	5,153	2,770	5,181	15,151
Indirect Cost/Sick Leave	41	597	37	606	61	1.104	42	578	41	520	222
Total Cost	3,741	5,665	3,178	5,553	2,981	5,482	2,662	5,202	2,811	1,276	15,373

*In Euros.

Figures




Figure 1

Flowchart cost population (2015).




Figure 2

Patient cumulative hospital mean cost*. *In Euros. CVD: cardiovascular disease; HF: heart failure; CKD: chronic kidney disease; cardiorenal: HF and/or CKD; MI: myocardial infarction; PAD: peripheral artery disease




Figure 3

Patient cumulative medication mean cost*. *In Euros. CVD: cardiovascular disease; HF: heart failure.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarytable3.docx](#)
- [Supplementarytable1.doc](#)
- [Supplementarytable2.docx](#)
- [Supplementarytable4.docx](#)
- [Supplementarytable5.docx](#)