

Adolescent BMI and Early-Onset Type 2 Diabetes Among Ethiopian Immigrants and Their Descendants: A Nationwide Study.

Maya Simchoni

Israel Defense Forces

Uri Hamiel

Tel Aviv Sourasky Medical Center

Orit Pinhas-Hamiel

Edmond and Lily Safra Children's Hospital

Inbar Zucker

Tel Aviv University Sackler Faculty of Medicine

Tali Cukierman-Yaffe

Sheba Medical Center at Tel Hashomer

Miri Lutski

Tel Aviv University Sackler Faculty of Medicine

Estela Derazne

Tel Aviv University Sackler Faculty of Medicine

Zivan Beer

Israel Defense Forces

Doron Behar

igentify

Lital Keinan-Boker

Tel Aviv University Sackler Faculty of Medicine

Ofri Mosenzon

Hadassah Medical Center

Dorit Tzur

Israel Defense Forces

Amon Afek

Sheba Medical Center at Tel Hashomer

Amir Tirosh

Sheba Medical Center at Tel Hashomer

Itamar Raz

Hadassah Medical Center

Gilad Twig (✉ gilad.twig@gmail.com)

Original investigation

Keywords: Adolescents, BMI, Early-onset, Epidemiology, Immigration, Type 2 diabetes

Posted Date: September 23rd, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-50057/v2>

License:   This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published on October 6th, 2020. See the published version at <https://doi.org/10.1186/s12933-020-01143-z>.

Abstract

Background: We assessed in a nationwide cohort the association between adolescent BMI and early-onset (<40 years) type 2 diabetes among Israelis of Ethiopian origin.

Methods: Normoglycemic adolescents (range 16-20 years old), including 93,806 native Israelis (≥ 3 rd generation in Israel) and 27,684 Israelis of Ethiopian origin, were medically assessed for military service between 1996 and 2011. Weight and height were measured. Data were linked to the Israeli National Diabetes Registry. Incident type 2 diabetes by December 31, 2016 was the outcome. Cox regression models stratified by sex and BMI categories were applied.

Results: 226 (0.29%) men and 79 (0.18%) women developed diabetes during 992,980 and 530,814 person-years follow-up, respectively, at a mean age of 30.4 and 27.4 years, respectively. Among native Israeli men with normal and high (overweight and obese) BMI, diabetes incidence was 9.5 and 62.0 (per 10 5 person-years), respectively. The respective incidences were 46.9 and 112.3 among men of Ethiopian origin. After adjustment for sociodemographic confounders, the hazard ratios for type 2 diabetes among Ethiopian men with normal and high BMI were 3.4 (2.3-5.1) and 15.8 (8.3-30.3) respectively, compared to third-generation Israelis with normal BMI. When this analysis was limited to Israeli-born Ethiopian men, the hazard ratios were 4.4 (1.7-11.4) and 29.1 (12.9-70.6), respectively. Results persisted when immigrants of other white Caucasian origin were the reference; and among women with normal, but not high, BMI.

Conclusions: Ethiopian origin is a risk factor for early-onset type 2 diabetes among young men at any BMI, and may require selective interventions.

Background

Diabetes mellitus is a major cause of morbidity and mortality. The global prevalence of type 2 diabetes among adults over age 18 years has exceeded 8.5%. Over 468 million are estimated to be affected(1); the mean age of diagnosis has decreased over the last couple decades.(2) Early-onset (< age 40 years) type 2 diabetes is associated with a worse clinical course and premature development of complications.(3) Several risk factors are associated with early-onset diabetes, among them adolescent obesity, certain ethnic backgrounds, immigration, and lower socioeconomic position. Additionally, there are some discrepancies between adults with type 2 diabetes from different ethnicities and risk of cardiovascular morbidity.(4,5)

Severe obesity in childhood is a risk factor for cardiovascular morbidity(6), and among children in the US, BMI at baseline preceded higher insulin levels at follow-up during childhood, with blacks and whites showing similar patterns of this one-directional relationship.(7) Immigrants from low- and middle-income countries including those of African and other ancestries have been shown to have a disproportionately high risk for type 2 diabetes, which may be evident at a lower body-mass index (BMI). (8) Among African immigrants to the US (8,9) and to Europe (10) significantly higher risk for pre-diabetes

and diabetes have been shown, compared to the native populations. Nevertheless, evidence among adolescent immigrants from Africa is limited, as this population is underrepresented in epidemiological studies and is frequently considered a homogenous group despite large ethnic variability.(11,12) As a destination for immigration from several regions worldwide, including Ethiopia, Israel provides an opportunity to address the risk of diabetes among persons of East African origin. Moreover, systematic documentation is available of adolescent health status for over five decades. Here we assessed the risk for early-onset type 2 diabetes among Israeli adolescents of Ethiopian origin compared to native Israelis and compared to other immigrant populations to Israel. Our goal was to assess whether the incidence of early-onset type 2 diabetes is higher in Israelis of Ethiopian origin than among native Israelis, and to specifically assess whether this relation differs according to adolescent BMI, sex, and immigration status.

Materials And Methods

Study population

This was a retrospective cohort study. At age 17 years, Israeli adolescents are approached to undergo a medical evaluation prior to mandatory military service. The study sample comprised all Israeli male and female adolescents who were medically evaluated for military service between ages 16-20 years from January 1, 1996 for men, and January 1, 1997 for women, until December 31, 2011. We excluded examinees with missing BMI data (n= 3,291, 2.6%) and those with a history of diabetes or dysglycemia at study entry (n=325, 0.3%) that was solely based on reports by primary care physicians. We also excluded those who died before the establishment of the Israel National Diabetes Registry (INDR) in 2012 (n=378; 0.3%) as these individuals could not be reported to the registry (See supplementary Figure S1). The study sample included 121,490 examinees for whom we had continuous follow-up from adolescence until diabetes onset, death, or December 31st, 2016, whichever came first. Data of an additional 121,997 male adolescents who immigrated to Israel from former countries of the Union of Soviet Socialist Republics (USSR) during the same years as the immigration from Ethiopia were analyzed as part of sensitivity analyses. The Israel Defense Forces Medical Corps Institutional Review Board approved this study.

The Israel National Diabetes Registry (INDR) and diagnosis of diabetes

The primary outcome of the study was incident type 2 diabetes that was diagnosed either during military service, or later in life as recorded by the INDR. All health medical organizations in Israel have been requested by law to annually report prevalent cases of diabetes to the INDR since 2012. These organizations provide medical care to nearly 100% of permanent residents in Israel. The INDR retrospectively recorded the date of diabetes diagnosis for all individuals diagnosed in Israel between January 1, 2000 and December 31, 2011, and prospectively recorded all new incidences of diabetes from January 1, 2012 onwards. Data of the INDR were linked to the IDF database using the participants' national ID number, to enable the linkage of body weight and height measured at adolescence with diabetes incidence recorded later in life. Diabetes was reported to the INDR when one or more of the following criteria were met in the previous year of the report to the registry: (i) glycated hemoglobin

(HbA1c) $\geq 6.5\%$; (ii) serum glucose concentrations of ≥ 200 mg/dL in two tests performed at an interval of at least one month; (iii) ≥ 3 purchases of glucose lowering medications in different months. Data regarding prescribed diabetes medications were used to classify diabetes as type 1, type 2, or uncertain type, as reported previously (13) (for details see the supplementary appendix). The sensitivity of the INDR for the detection of type 2 diabetes is 95%. (13) The INDR also includes weight and height measurements that were documented at routine clinic visits during the year of diabetes diagnosis. Measurements regarding weight and height at adulthood among those who did not develop diabetes were unavailable. Individuals who were reported to the INDR as having type 2 diabetes, but without a formal date of diagnosis were considered missing in the time-to-event analysis if they were first reported in the first year of the registry, 2012; and were included only in logistic regression models. For individuals without a formal date, who were first reported in 2013 or later, we assigned the date of diagnosis as July 1st of the year before the report year to the INDR, as described previously. (14)

Data collection and study variables

Age at examination and year of birth were treated as continuous variables. The health examination was performed by military physicians who reviewed the participants' medical records and provided diagnostic codes when applicable. The physical examination included measurements of weight and height (barefoot and in underwear) by trained medics using a beam balance and stadiometer. Measurements were recorded and rounded to the nearest 0.5 kg for weight and one cm for height. BMI was calculated (weight in kilograms divided by height squared in meters) and categorized into 3 subgroups: < 18.5 kg/m² (underweight), 18.5-24.9 kg/m² (normal BMI), and ≥ 25 kg/m² (high BMI). Because at age 17 adolescents have completed $>98\%$ of their growth potential, we used the world health organization (WHO) BMI categories intended for adults. (15) Individuals whose medical assessment revealed no history of major operation or comorbidity and who had no illness that requires a chronic medical treatment were defined as having unimpaired health. Data regarding education and residential socioeconomic status (SES) were obtained from governmental ministries, and were used as previously reported (16); details are provided in the supplementary appendix. Cognitive performance at adolescence was assessed by a general intelligence test that was delivered to all individuals at study entry, and was categorized (low/medium/high) as reported previously. (17) Notably, this variable was identified as an independent risk factor for early-onset type 2 diabetes in a subpopulation of this cohort (18). Examinees were classified as of Ethiopian origin if they or their fathers were born in Ethiopia, whereas Israeli-born individuals whose fathers and grandfathers were born in Israel were classified as native Israelis. (19) All individuals of Ethiopian origin were either immigrants from Ethiopia or were born in Israel to immigrants from Ethiopia (second-generation). In a similar manner, we defined in a sensitivity analysis the region of origin of immigrants or offspring of immigrants from former countries of the USSR. (20)

Statistical analysis

Prior to undertaking analysis, due to clinical plausibility, we decided to stratify the cohort by sex and adolescent BMI (underweight, normal, and high BMI). Analysis of the entire cohort demonstrated a

significant interaction term between sex and origin ($p=0.038$), and a borderline significant interaction term between BMI and origin ($p=0.086$).

The incident rate of type 2 diabetes was calculated per person-year of follow-up for native Israelis and Israelis of Ethiopian origin per a given BMI group. Kaplan-Meier survival curves were computed for the BMI percentile categories, with 95% confidence intervals. Cox proportional hazard models were used to estimate the hazard ratios (HRs) and 95%-confidence intervals (CIs) for incident diabetes, considering native Israelis with normal BMI as the reference group. Covariates were added in a stepwise manner to a model adjusted for age and birth year. Variables that were significant ($p<0.05$) in the minimally adjusted model were included in the final multivariable analysis. The assumption of proportionality of hazards was visually confirmed for all variables.

We conducted several sensitivity analyses that were limited to men, given the smaller sample of women in this cohort. (i) The analysis was restricted to those with unimpaired health status at enrollment in order to minimize residual confounding by co-existing morbidities. (ii) The analysis was restricted to second-generation, Israeli-born men of Ethiopian origin, using the same reference group as the main analysis. (iii) We assessed the risk for type 2 diabetes that was related to age of immigration among immigrants from Ethiopia with normal adolescent BMI. (iv) Diabetes risk of men from former countries of the USSR were compared to the main study groups, since immigration to Israel from this region occurred in parallel to immigration from Ethiopia to Israel; 1990-2000 vs. 1985-2000. To better assess the contribution of immigration status to diabetes risk, analysis was stratified to immigrants (USSR immigrants were the reference) and second-generation Israeli-born (native Israelis were the reference). (v) Logistic regression models were applied to account for cases with missing date of type 2 diabetes diagnosis. (vi) We assessed potential misclassification of type 2 diabetes by accounting for cases of diabetes of uncertain type, or by using a stricter definition of type 2 diabetes with a positive predicted value of over 98%,⁽¹³⁾ in which a case of type 2 diabetes is defined only if reported to the INDR in two separate years.

Results

Baseline characteristics of the men and women in this study are presented separately in Table 1. The age at study entry was similar across BMI groups in both sexes. Adolescents of Ethiopian origin of both sexes had lower residential SES than did native Israelis. Among Ethiopian Israelis of both sexes, most of those in the high BMI groups were born in Israel.

Incident Type 2 Diabetes among Israeli Men of Ethiopian Origin

During 992,980 person-years there were 226 incident cases of type 2 diabetes in men: 89 (0.52%) in those of Ethiopian origin and 137 (0.22%) in native Israelis. The mean ages at diabetes diagnosis were 30.3 and 30.4 years, respectively ($p=0.9$). The crude incidence rates for type 2 diabetes (per 10^5 person-years) in Ethiopian vs. native Israeli men were 17.3 vs. 5.1 in the underweight group, 46.9 vs. 9.5 in the

normal BMI group, and 112 vs. 62 in the high BMI group (Table 2). The mean BMI values that were measured at the time when diabetes diagnoses were first reported to the INDR were lower in Ethiopian than native Israeli men for the normal and high adolescent BMI categories (Supplementary Figure S2A). Most native (69.4%), but not Ethiopian (49.0%), Israelis were with overweight or obesity at diabetes diagnosis ($p=5.2 \times 10^{-7}$).

Kaplan-Meier survival analysis curves for men are shown in Figure 1 (A, C, E). The unadjusted HRs for type 2 diabetes were consistently higher in men of Ethiopian origin than in native Israeli men (Table 2). Further adjustment for sociodemographic data (education level, residential SES status, and cognitive performance) yielded the following HRs in Ethiopian vs. native Israelis (reference group, native Israelis with normal BMI): 1.2 (0.6-2.3) vs. 0.5 (0.2-1.3) in underweight, 3.4 (2.3-5.1) vs. 1 (reference) in normal BMI, and 15.8 (8.3-30.3) vs. 6.9 (4.9-9.9) in the high BMI group. The results persisted when the study sample was limited to persons with unimpaired health at baseline to minimize confounding by pre- or coexisting morbidities (Table 2).

Point estimates for incident diabetes were directly related to the time interval from immigration. Point estimates were higher among those who immigrated to Israel before age 6 years (HR=4.6, 95%CI= 2.6-8.2) than for those who immigrated after age 12 years (HR=3.1, 95%CI= 1.6-5.9) (supplementary Table S3). Also, the HR for diabetes among Israelis who were born in Ethiopia was lower than among second-generation Israelis of Ethiopian origin – 2.8 (2.0-3.9) vs. 4.0 (2.2-7.3) (native Israelis were the reference group; the model was adjusted for birth year, age at study entry, education level, cognitive score, and BMI (continuous); supplementary Table S3). When the analysis was limited to men of Ethiopian origin who were born in Israel, the HRs were 4.4 (1.7-11.4) in the normal BMI group and 29.1 (12.0-70.6) in the high BMI group compared to native Israelis with normal BMI (Table 2). In contrast, risk for diabetes was comparable among Israeli-born men of USSR origin to that of native Israelis at any BMI stratum (Figure 2A). For immigrants from Ethiopia, the adjusted HR to develop type 2 diabetes was 3.9 (2.8-5.5) compared to immigrants from the USSR (Figure 2B). This risk was more pronounced among immigrants from Ethiopia with normal BMI at study entry [HR=5.1 (3.4-7.8)] when immigrants from the USSR with normal BMI served as the reference (Figure 2B).

The results were accentuated when logistic regression models were applied, and with the inclusion of incident cases of type 2 diabetes for whom the date of diagnosis was missing (n=64; supplementary Table S1). Odds ratios (ORs) were calculated for the development of type 2 diabetes, using native Israelis with normal BMI as the reference group. The adjusted odds ratio (OR) for Israelis of Ethiopian origin with normal BMI was 4.1 (2.9-5.9). Among individuals with high BMI, the OR for those of Ethiopian origin was 18.8 (10.5-33.6), and for native Israelis 7.4 (5.4-10.2); this difference between the culture groups was statistically significant ($p=0.028$). Accounting for misclassification of diabetes diagnosis, either by including cases of uncertain diabetes type or by applying a stricter definition of type 2 diabetes with over 98% positive predicted value, yielded similar results (supplementary Table S2).

Incident Type 2 Diabetes among Israeli Women of Ethiopian Origin

During 530,814 person-years there were 79 cases of incident type 2 diabetes among women: 29 (0.27%) in those of Ethiopian origin and 50 (0.15%) in native Israelis. The mean ages at diabetes diagnosis were 28.3 and 27.3 years, respectively. Notably, high BMI among female adolescents of Ethiopian origin was rare, only 6 persons, in the first decade of the study. The crude incident rate for type 2 diabetes (per 10⁵ person-years) in Ethiopian vs. native Israeli women were 10.5 vs. 0 in the underweight group, 23.7 vs. 5.7 in the normal BMI group, and 54.0 vs. 55.8 in the high BMI group (Table 2). These differences were well apparent in Kaplan-Meier survival analysis (Figure 1B, D, F). The BMI recorded at the time of entrance to the INDR had no consistent pattern. Women of Ethiopian origin who developed type 2 diabetes in the normal BMI group gained more weight than did their native counterparts, whereas the opposite was observed for the high BMI group (Supplementary Figure S2B). Among women with normal BMI in adolescence, the HR for type 2 diabetes was 4.2 (2.2-8.1) for those of Ethiopian origin compared to their native peers; and 2.4 (1.1-5.1) in the adjusted model. The latter was attenuated to borderline significance when adjusted logistic regression models were applied to include incident cases without the date of diabetes diagnosis (n=47; OR=1.8, 95%CI 0.98-3.2, p=0.058; supplementary Table S1). HRs of Ethiopian and native Israeli women with high adolescent BMI were comparable.

Discussion

The main findings of this nationwide longitudinal study include the following: (1) For Israeli adolescent males of Ethiopian origin with normal or high BMI, the risk for incident early-onset diabetes is 3-4-fold higher than for native Israeli males. This difference is unlikely to be accounted by differential weight gain in adulthood and was not observed among immigrants to Israel from the former USSR countries, who arrived to Israel at the same period as immigration from Ethiopia. (2) Diabetes risk is accentuated in second-generation Israeli-born men of Ethiopian origin. Point estimates were stable following adjustment to sociodemographic background and extensive sensitivity analyses.

A solid body of evidence indicates an association of black ancestry with increased diabetes risk compared to white Caucasians. Examples are the disproportionate increased risk for early-onset diabetes among persons of African origin in the UK,(21–23) Canada,(24) and the US.(25) HRs for developing type 2 diabetes for middle-aged African Americans (2,26–30) and Ethiopian immigrants to Israel,(31,32) compared to native populations, were reported in the range of 1.3 to 2.9. Among young adults and adolescents, point estimates ranged between 2.1 (30) and 6.2.(33,34) Methodological differences relating to the nature of population selection, the level of adjustment to confounders, and the wide age range at study entry may account for the differences observed. Additionally, in several US studies, African Americans were aggregated into a homogenous group despite diversity in ethnic and cultural background.(11) In this respect, our study is population-based, with a narrow age range at study entry, and a homogenous ethnic origin. In addition, a series of sociodemographic variables were included that were systematically obtained without a selection bias, given the mandatory nature of the medical screening.

Repeated evidence supports an association between genetic polymorphism and diabetes risk among persons of African ancestry. The thrifty gene hypothesis (35) suggests that certain populations have alleles that are adaptive to the feast and famine cycles of paleolithic human existence. These may promote rapid weight gain that may be advantageous during times of limited food availability but may be metabolically deleterious at times of continuing affluence. In agreement, several examples of novel and African ancestry-specific disease loci have been discovered in association with insulin resistance and inflammation. For example, polymorphisms in tumor necrosis factor-alpha (TNF- α),(36) or the anti-inflammatory adipokine, adiponectin,(37) were more prevalent among persons of African ancestry and were linked to an increased risk for type 2 diabetes. Notably, young African Americans, who largely represent West African ancestry, have been shown to have an increased risk to develop diabetes regardless of BMI,(30) similar to Israelis of Ethiopian origin. This suggests that the increased genetic predisposition for type 2 diabetes risk is largely driven by Pan-African ancestry. This contrasts with the genetic predisposition for end-stage renal disease that affects people of West African, but not Ethiopian origin.(38)

We report higher risk for type 2 diabetes among second-generation Israeli-born Ethiopian men than among those who immigrated; this association correlated with the time interval from immigration. Our findings corroborate previous evidence that diabetes risk correlates with the time since immigration among immigrants from various ethnic backgrounds to Western countries,(39,40) and is higher among second- than first-generation immigrants by 2- to 6-fold.(39,41) Similarly, African immigrants to the US were shown to be more likely to develop pre-diabetes or diabetes than African Americans, despite their having a higher level of education and lower BMI.(8) Notably, no association was found between diabetes incidence and immigration status regarding immigration from the former USSR, which occurred in parallel to the Ethiopian immigration. Additionally, previous studies underscored lifestyle changes among Ethiopian immigrants to Israel, who changed their diet to a more Western one,(42) and became less physically active and with more obesity (19) in a manner that correlated with the time since immigration. (43)

Our findings have implications to other high-income countries that are recipients of immigrants from Ethiopia. In the US, evidence supports the need to disaggregate the African American category into its ethnic subcommunities based on differences in genetic admixture, culture, and metabolic risk.(8,9,44) Of note, as of 2018, the US population included over 730,000 immigrants from East Africa. Nearly half of them were from Ethiopia, and they are considered to be the fastest-growing sub-African community in the US. This trend is more evident in some European countries such as Germany, where African immigrants constitute nearly 1% of the population.(45) The observation that the prevalence of high BMI among Israeli-born Ethiopian male adolescents has tripled over the last decade and reached nearly 20%(19) echoes with US data,(46) and predicts an increased burden of type 2 diabetes in young adults from this population. The lower prevalence of type 2 diabetes in Ethiopia, by over 3-fold, compared to Israel, the US, and most other Western countries, further emphasizes the contribution of immigration to diabetes risk at young ages in high-income countries.(1) Importantly, undiagnosed diabetes among young adults may be over 10-fold more prevalent among blacks than whites, as exemplified in the US population.(34)

Collectively, these data emphasize the importance of clinical awareness, education for healthy lifestyle, and reduction of other preventable risk factors, especially in regard to adolescent obesity, for Ethiopian and possibly other East African immigrants.

This study has several limitations. First, only a single BMI measurement, at study entry, was available; and early longitudinal measurements throughout childhood and later during adulthood were unavailable. Nevertheless, BMI measurements among those who developed type 2 diabetes support that differential excess weight gain among Israelis of Ethiopian origin could not explain the results; this concurs with previous reports.⁽³¹⁾ Additionally, the time interval from medical assessment at late adolescence to the mean age of diabetes onset was approximately 10 years. This period has the potential to confound the results due to lifestyle manners of the participants such as diet or exercise, but also constitutes a reasonable time frame for initiating intervention strategies. Second, we lacked data regarding pre-diabetes and undiagnosed diabetes, which may be especially prominent in a young population. Third, information on lifestyle measures including dietary intake, physical activity, and smoking were not available to us. Fourth, we lacked baseline information regarding plasma glucose, cholesterol, and other measures of adiposity such as waist circumference. Of note, BMI is considered the preferable method for screening adolescents according to the U.S. Preventive Task Force,⁽⁴⁷⁾ and our dataset was based on a systematic medical screening process that enabled mitigating confounding by coexisting morbidities. Finally, we are underpowered to discuss the association between BMI and type 2 diabetes among young women with high BMI due to the small number of incident cases. However, findings in women with normal BMI resembled those in men. The strengths of this study include its systematic data collection at a narrow age range, the absence of population selection bias, and linkage between two nationwide databases and outcome cases of type 2 diabetes that were predominantly diagnosed before age 30 years. The ability to compare the findings among second generation Ethiopian descent Israelis with those of immigrants from other regions to Israel in this large sample size is advantageous.

Conclusions

Adolescent males of Ethiopian origin are at increased risk for early-onset type 2 diabetes even if their BMI is within the normal range; and those with overweight and obesity are at considerably greater risk. These findings emphasize the need for tight medical follow-up, aimed at reducing concomitant risk factors including obesity prevention in particular, education for healthy lifestyle, and early intervention medical therapy as needed.

Abbreviations

BMI – Body Mass Index; CI – Confidence Interval; HR – Hazard Ratio; SD – Standard Deviation; SES – Socio-economic Status; USSR – Union of Soviet Socialist Republics.

Declarations

Ethics approval and consent to participate: The study was approved by the Israel Defense Forces Medical Corps Institutional Review Board. Committee reference number: 2014-1476.

The IDF review board waived the need for informed consent and researches strictly maintained the anonymity and privacy of the participants.

Consent for publication: Not applicable.

Availability of the data and materials: Data sharing is not applicable for this study.

Competing interests: The authors declare that they have no competing interests.

Funding: This study was supported by a research grant from the medical corps Israel Defense Forces (Israel) award to GT (Award No. 4440985218).

Author's contributions: GT conceived and designed the study, analyzed and interpreted the data, wrote the first draft, and revised the paper incorporating contributions from coauthors. He had access to all the data, and decided on submission. MS and UH analyzed and interpreted the data, wrote the first draft and decided on submission. IZ, ML, and DT conducted database management quality assurance, interpreted the data, contributed to the discussion, and critically revised the manuscript. ED conducted the statistical analysis, interpreted the data, and critically revised the manuscript. OPH interpreted the data, contributed to the discussion, and critically revised the manuscript. AA, AT, ZB, DH, TCY, OM, IR, and TKB contributed to the discussion and critically revised the manuscript. G.T. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Acknowledgment: The authors thank Prof Hertzal Gerstein for helpful discussions that were made on earlier versions of this manuscript, and Ms. Cindy Cohen for her excellent editorial assistance.

References

1. International Diabetes Federation. IDF Diabetes Atlas [Internet]. 9th ed. Brussels; 2019 [cited 2020 Apr 28]. Available from: <https://www.diabetesatlas.org>
2. Menke A, Casagrande S, Geiss L, Cowie CC. Prevalence of and Trends in Diabetes among Adults in the United States, 1988-2012. Vol. 314, JAMA - Journal of the American Medical Association. American Medical Association; 2015. p. 1021–9.
3. Lascar N, Brown J, Pattison H, Barnett AH, Bailey CJ, Bellary S. Type 2 diabetes in adolescents and young adults. Jan 1, 2018 p. 69–80.
4. Owusu Adjah ES, Bellary S, Hanif W, Patel K, Khunti K, Paul SK. Prevalence and incidence of complications at diagnosis of T2DM and during follow-up by BMI and ethnicity: A matched case-control analysis. Cardiovasc Diabetol [Internet]. BioMed Central Ltd.; 2018 May 15 [cited 2020 Sep 11];17(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/29764436/>

5. Owusu Adjah ES, Ray KK, Paul SK. Ethnicity-specific association of BMI levels at diagnosis of type 2 diabetes with cardiovascular disease and all-cause mortality risk. *Acta Diabetol* [Internet]. Springer-Verlag Italia s.r.l.; 2019 Jan 28 [cited 2020 Sep 11];56(1):87–96. Available from: <https://pubmed.ncbi.nlm.nih.gov/30167871/>
6. Bendor CD, Bardugo A, Pinhas-Hamiel O, Afek A, Twig G. Cardiovascular morbidity, diabetes and cancer risk among children and adolescents with severe obesity [Internet]. Vol. 19, *Cardiovascular Diabetology*. BioMed Central; 2020 [cited 2020 Sep 14]. Available from: </pmc/articles/PMC7293793/?report=abstract>
7. Zhang T, Zhang H, Li Y, Li S, Fernandez C, Bazzano L, et al. Long-term Impact of Temporal Sequence from Childhood Obesity to Hyperinsulinemia on Adult Metabolic Syndrome and Diabetes: The Bogalusa Heart Study. *Sci Rep* [Internet]. Nature Publishing Group; 2017 Feb 23 [cited 2020 Sep 11];7. Available from: </pmc/articles/PMC5322533/?report=abstract>
8. O’connor MY, Thoreson CK, Ricks M, Courville AB, Thomas F, Yao J, et al. Worse cardiometabolic health in African immigrant men than African American Men: Reconsideration of the healthy immigrant effect. *Metab Syndr Relat Disord* [Internet]. Mary Ann Liebert Inc.; 2014 Aug 1 [cited 2020 Apr 28];12(6):347–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24814168>
9. Ukegbu UJ, Castillo DC, Knight MG, Ricks M, Miller B V, Onumah BM, et al. Metabolic syndrome does not detect metabolic risk in African men living in the U.S. *Diabetes Care*. 2011 Oct;34(10):2297–9.
10. Chilunga FP, Henneman P, Meeks KA, Beune E, Requena-Méndez A, Smeeth L, et al. Prevalence and determinants of type 2 diabetes among lean African migrants and non-migrants: the RODAM study. *J Glob Health* [Internet]. University of Edinburgh; 2019 Dec 1 [cited 2020 Apr 28];9(2):020426–020426. Available from: <http://jogh.org/documents/issue201902/jogh-09-020426.pdf>
11. Commodore-Mensah Y, Himmelfarb CD, Agyemang C, Sumner AE. Cardiometabolic health in African immigrants to the United States: A call to re-examine research on African-descent populations [Internet]. Vol. 25, *Ethnicity and Disease*. ISHIB; 2015 [cited 2020 Apr 17]. p. 373–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26675140>
12. Boise L, Tuepker A, Gipson T, Vigmenon Y, Soule I, Onadeko S. African refugee and immigrant health needs: Report from a Community-Based House Meeting Project. *Prog Community Heal Partnerships Res Educ Action*. 2013;7(4):369–78.
13. Bendor CD, Bardugo A, Zucker I, Cukierman-Yaffe T, Lutski M, Derazne E, et al. Childhood Pancreatitis and Risk for Incident Diabetes in Adulthood. *Diabetes Care* [Internet]. 2020 Jan [cited 2020 Jan 10];43(1):145–51. Available from: <http://care.diabetesjournals.org/lookup/doi/10.2337/dc19-1562>
14. Twig G, Zucker I, Afek A, Cukierman-Yaffe T, Bendor CD, Derazne E, et al. Adolescent Obesity and Early-Onset Type 2 Diabetes. *Diabetes Care* [Internet]. American Diabetes Association; 2020 Apr 22 [cited 2020 Apr 23];dc191988. Available from: <http://care.diabetesjournals.org/lookup/doi/10.2337/dc19-1988>
15. Greulich WW. *RADIOGRAPHIC ATLAS OF SKELETAL DEVELOPMENT OF THE HAND AND WRIST- Second Edition* [Internet]. Vol. 91, *California Medicine*. BMJ Publishing Group; 1959 [cited 2020 Sep

- 14]. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1577858/>
16. Furer A, Afek A, Orr O, Gershovitz L, Landau Rabbi M, Derazne E, et al. Sex-specific associations between adolescent categories of BMI with cardiovascular and non-cardiovascular mortality in midlife. *Cardiovasc Diabetol* [Internet]. BioMed Central Ltd.; 2018 Jun 5 [cited 2020 Sep 14];17(1):80. Available from: <https://cardiab.biomedcentral.com/articles/10.1186/s12933-018-0727-7>
17. Twig G, Tirosh A, Derazne E, Haklai Z, Goldberger N, Afek A, et al. Cognitive function in adolescence and the risk for premature diabetes and cardiovascular mortality in adulthood. *Cardiovasc Diabetol* [Internet]. BioMed Central Ltd.; 2018 Dec 5 [cited 2020 Mar 23];17(1):154. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30518353>
18. Twig G, Gluzman I, Tirosh A, Gerstein HC, Yaniv G, Afek A, et al. Cognitive function and the risk for diabetes among young men. *Diabetes Care* [Internet]. American Diabetes Association Inc.; 2014 Nov 1 [cited 2020 Mar 23];37(11):2982–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25092683>
19. Hamiel U, Pinhas-Hamiel O, Vivante A, Bendor C, Bardugo A, Afek A, et al. Impact of Immigration on Body Mass Index and Blood Pressure Among Adolescent Males and Females: A Nationwide Study. *Hypertens (Dallas, Tex 1979)* [Internet]. Lippincott Williams and Wilkins; 2019 Dec 1 [cited 2020 Jan 10];74(6):1316–23. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/31630574>
20. Twig G, Yaniv G, Levine H, Leiba A, Goldberger N, Derazne E, et al. Body-Mass Index in 2.3 Million Adolescents and Cardiovascular Death in Adulthood. *N Engl J Med* [Internet]. 2016 Jun 23 [cited 2018 Dec 12];374(25):2430–40. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27074389>
21. Haines L, Wan KC, Lynn R, Barrett TG, Shield JPH. Rising incidence of type 2 diabetes in children in the U.K. *Diabetes Care*. 2007 May;30(5):1097–101.
22. Khanolkar AR, Amin R, Taylor-Robinson D, Viner R, Warner J, Stephenson T. Ethnic Minorities Are at Greater Risk for Childhood-Onset Type 2 Diabetes and Poorer Glycemic Control in England and Wales. *J Adolesc Heal* [Internet]. Elsevier USA; 2016 Sep 1 [cited 2020 Apr 17];59(3):354–61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27426206>
23. Candler TP, Mahmoud O, Lynn RM, Majbar AA, Barrett TG, Shield JPH. Continuing rise of Type 2 diabetes incidence in children and young people in the UK. *Diabet Med* [Internet]. Blackwell Publishing Ltd; 2018 Jun 1 [cited 2020 Sep 11];35(6):737–44. Available from: <https://pubmed.ncbi.nlm.nih.gov/29460341/>
24. Tenkorang EY. Early onset of type 2 diabetes among visible minority and immigrant populations in Canada. *Ethn Heal* [Internet]. Routledge; 2017 May 4 [cited 2020 Apr 17];22(3):266–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27771960>
25. Narayan KMV, Boyle JP, Thompson TJ, Sorensen SW, Williamson DF. Lifetime Risk for Diabetes Mellitus in the United States. *J Am Med Assoc*. 2003 Oct 8;290(14):1884–90.
26. Brancati FL, Whelton PK, Kuller LH, Klag MJ. Diabetes mellitus, race, and socioeconomic status: A population-based study. *Ann Epidemiol*. Elsevier Inc.; 1996 Jan 1;6(1):67–73.

27. Brancati FL, Kao WHL, Folsom AR, Watson RL, Szklo M. Incident Type 2 Diabetes Mellitus in African American and White Adults. *JAMA* [Internet]. 2000 May 3 [cited 2019 Aug 10];283(17):2253. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10807384>
28. Bancks MP, Kershaw K, Carson AP, Gordon-Larsen P, Schreiner PJ, Carnethon MR. Association of modifiable risk factors in young adulthood with racial disparity in incident type 2 diabetes during middle adulthood. *JAMA - J Am Med Assoc* [Internet]. American Medical Association; 2017 Dec 26 [cited 2020 Mar 21];318(24):2457–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29279935>
29. Wei GS, Coady SA, Goff DC, Brancati FL, Levy D, Selvin E, et al. Blood pressure and the risk of developing diabetes in African Americans and Whites: ARIC, CARDIA, and the Framingham Heart Study. *Diabetes Care* [Internet]. 2011 Apr [cited 2020 Apr 4];34(4):873–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21346180>
30. Wei GS, Coady SA, Reis JP, Carnethon MR, Coresh J, D'Agostino RB, et al. Duration and Degree of Weight Gain and Incident Diabetes in Younger Versus Middle-Aged Black and White Adults: ARIC, CARDIA, and the Framingham Heart Study. *Diabetes Care* [Internet]. 2015 Nov [cited 2019 Aug 7];38(11):2042–9. Available from: <http://care.diabetesjournals.org/lookup/doi/10.2337/dc14-2770>
31. Jaffe A, Giveon S, Wulffhart L, Oberman B, Freedman L, Ziv A, et al. Diabetes among Ethiopian Immigrants to Israel: Exploring the Effects of Migration and Ethnicity on Diabetes Risk. Tordjman KM, editor. *PLoS One* [Internet]. Public Library of Science; 2016 Jun 14 [cited 2018 Dec 1];11(6):e0157354. Available from: <https://dx.plos.org/10.1371/journal.pone.0157354>
32. Reuven Y, Dreihier J, Shvartzman P. The prevalence of diabetes, hypertension and obesity among immigrants from East Africa and the former Soviet Union: A retrospective comparative 30-year cohort study. *Cardiovasc Diabetol*. BioMed Central Ltd.; 2016 May 5;15(1).
33. Dabelea D, Mayer-Davis EJ, Saydah S, Imperatore G, Linder B, Divers J, et al. Prevalence of Type 1 and Type 2 Diabetes Among Children and Adolescents From 2001 to 2009. *JAMA* [Internet]. 2014 May 7 [cited 2019 Aug 10];311(17):1778. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24794371>
34. The NS, Richardson AS, Gordon-Larsen P. Timing and Duration of Obesity in Relation to Diabetes: Findings from an ethnically diverse, nationally representative sample. *Diabetes Care* [Internet]. 2013 Apr 1 [cited 2019 Aug 10];36(4):865–72. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23223352>
35. NEEL J V. Diabetes mellitus: a “thrifty” genotype rendered detrimental by “progress”? *Am J Hum Genet*. Elsevier; 1962 Dec;14(4):353–62.
36. Ayelign B, Genetu M, Wondmagegn T, Adane G, Negash M, Berhane N. TNF- α (-308) Gene Polymorphism and Type 2 Diabetes Mellitus in Ethiopian Diabetes Patients. *Diabetes, Metab Syndr Obes Targets Ther* [Internet]. Dove Medical Press Ltd.; 2019 Nov 28 [cited 2020 Apr 19];Volume 12:2453–9. Available from: <https://www.dovepress.com/tnf-alpha-minus308-gene-polymorphism-and-type-2-diabetes-mellitus-in-e-peer-reviewed-article-DMSO>

37. Riestra P, Gebreab SY, Xu R, Khan RJ, Bidulescu A, Correa A, et al. Gender-specific associations between ADIPOQ gene polymorphisms and adiponectin levels and obesity in the Jackson Heart Study cohort. *BMC Med Genet* [Internet]. BioMed Central Ltd.; 2015 Aug 20 [cited 2020 Mar 21];16(1):65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26290432>
38. Behar DM, Shlush LI, Maor C, Lorber M, Skorecki K. Absence of HIV-associated nephropathy in Ethiopians. *Am J Kidney Dis* [Internet]. 2006 Jan [cited 2019 May 23];47(1):88–94. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0272638605014940>
39. Peled A, Gordon B, Twig G, Mendlovic J, Derazne E, Lisnyansky M, et al. Immigration to Israel during childhood is associated with diabetes at adolescence: a study of 2.7 million adolescents. *Diabetologia* [Internet]. Springer Verlag; 2017 Nov 1 [cited 2020 Mar 22];60(11):2226–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28821907>
40. Pasupuleti SSR, Jatrana S, Richardson K. Effect of nativity and duration of residence on chronic health conditions among Asian immigrants in Australia: A longitudinal investigation. *J Biosoc Sci* [Internet]. Cambridge University Press; 2016 May 1 [cited 2020 Mar 22];48(3):322–41. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26139212>
41. Afable-Munsuz A, Mayeda ER, Pérez-Stable EJ, Haan MN. Immigrant Generation and Diabetes Risk Among Mexican Americans: The Sacramento Area Latino Study on Aging. *Am J Public Health*. American Public Health Association; 2014 Apr 10;104(S2):S243–50.
42. Regev-Tobias H, Reifen R, Endevelt R, Havkin O, Cohen E, Stern G, et al. Dietary acculturation and increasing rates of obesity in Ethiopian women living in Israel. *Nutrition* [Internet]. 2012 Jan [cited 2019 Aug 4];28(1):30–4. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S089990071100089X>
43. Meydan C, Twig G, Derazne E, Tzur D, Gordon B, Shamiss A, et al. The immigration effect on obesity and overweight in Israeli Jewish male adolescents born 1970-1993. *Ann Epidemiol* [Internet]. 2014 Jun [cited 2019 Aug 4];24(6):424–31. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S104727971400074X>
44. Sewali B, Harcourt N, Everson-Rose SA, Leduc RE, Osman S, Allen ML, et al. Prevalence of cardiovascular risk factors across six African Immigrant Groups in Minnesota. *BMC Public Health*. BioMed Central Ltd.; 2015 Apr 22;15(1).
45. Statistisches B. Foreign population by place of birth and selected citizenships [Internet]. 2019 [cited 2020 Apr 29]. Available from: <https://www.destatis.de/EN/Themes/Society-Environment/Population/Migration-Integration/Tables/foreigner-place-of-birth.html>
46. Popkin BM, Udry JR. Adolescent Obesity Increases Significantly in Second and Third Generation U.S. Immigrants: The National Longitudinal Study of Adolescent Health. *J Nutr* [Internet]. Oxford University Press (OUP); 1998 Oct 1 [cited 2020 Feb 8];128(4):701–6. Available from: <https://academic.oup.com/jn/article/128/4/701/4728868>
47. Grossman DC, Bibbins-Domingo K, Curry SJ, Barry MJ, Davidson KW, Doubeni CA, et al. Screening for Obesity in Children and Adolescents. *JAMA* [Internet]. 2017 Jun 20 [cited 2019 May

Tables

Table 1: Baseline characteristics of the study population according to origin and adolescent BMI groups.

	Native Israelis			Ethiopian Israelis		
	Underweight BMI < 18.5	Normal BMI 18.5 ≤ BMI < 25	High BMI BMI ≥ 25	Underweight BMI < 18.5	Normal BMI 18.5 ≤ BMI < 25	High BMI BMI ≥ 25
Men						
Total number	7,435	42,600	10,736	5,307	10,708	1,035
Israeli-born (%)	100	100	100	32.5	24.6	59.7
Mean age (years ±SD)	17.3 ±0.5	17.3 ±0.5	17.4 ±0.6	17.7 ±0.7	17.8 ±0.8	17.5 ±0.6
Mean BMI (kg/m ² ±SD)	17.5 ±0.8	21.3 ±1.7	28.5 ±3.3	17.3 ±0.9	20.6 ±1.6	28.4 ±3.3
Mean weight (kg ±SD)	53.2 ±4.8	64.7 ±7.2	86.7 ±12.3	51.0 ±4.7	59.8 ±6.6	84.8 ±11.9
Mean height (cm ±SD)	173.9 ±6.8	174.0 ±6.7	174.1 ±6.8	171.5 ±6.8	170.3 ±6.6	172.9 ±7.1
Full education (%)	92.0	92.5	92.1	90.8	91.4	92.3
Low SES (%)	43.2	43.6	46.9	35.1	36.1	35.8
Medium SES (%)	35.2	35.4	36.2	62.0	60.4	61.1
High SES (%)	21.6	21.0	16.9	2.9	3.5	3.1
Low cognitive score (%)	23.2	20.6	25.1	65.6	63.3	64.5
Medium cognitive score (%)	62.2	64.1	63.5	34.2	36.3	34.8
High cognitive score (%)	14.2	15.4	11.4	0.2	0.3	0.7
Women						
Total number	4,685	23,170	5,180	3,114	6,507	1,013
Israeli-born (%)	100	100	100	36.5	34.5	51.6
Mean age (years ±SD)	17.1 ±0.3	17.1 ±0.3	17.2 ±0.4	17.6 ±0.7	17.6 ±0.7	17.5 ±0.7
Mean BMI (kg/m ² ±SD)	17.5 ±0.8	21.3 ±1.7	28.5 ±3.4	17.2 ±0.9	20.9 ±1.7	28.1 ±3.0
Mean weight (kg ±SD)	46.6 ±4.1	56.0 ±6.1	75.1 ±10.9	44.3 ±4.0	53.1 ±5.8	72.3 ±9.9

Mean height (cm ±SD)	163.0 ±6.3	162.1 ±6.1	162.2 ±6.4	160.4 ±6.0	159.1 ±6.1	160.1 ±6.7
Full education (%)	98.3	98.5	97.3	93.1	92.8	94.8
Low SES (%)	16.5	17.7	20.0	33.9	34.1	30.7
Medium SES (%)	49.3	49.8	52.8	64.2	62.8	65.5
High SES (%)	34.2	32.5	27.2	1.9	3.1	3.8
Low cognitive score (%)	14.6	12.7	18.3	62.1	62.5	60.7
Medium cognitive score (%)	73.6	74.3	72.5	37.6	37.2	38.5
High cognitive score (%)	11.8	13.0	9.2	0.3	0.3	0.8

SES: socioeconomic status (according to residence)

Table 2: Hazard ratios for type 2 diabetes incidence among Israelis of Ethiopian origin and native Israelis.

	Native Israelis			Ethiopian Israelis		
	Underweight	Normal BMI	High BMI	Underweight	Normal BMI	High BMI
Men						
Number of diabetes incidences	5	52	80	12	65	12
Mean follow-up (years; ± SD)	13.3 ±4.7	12.8 ±4.6	12.1 ±4.5	13.1 ±4.6	13.0 ±4.6	10.4 ±3.8
Cumulative follow-up (person-years)	98,635.6	546,719.6	129,126.1	69,269.4	138,542.1	10,687.5
Incident rate (per 10 ⁵ person-years)	5.1	9.5	62.0	17.3	46.9	112.3
Mean age at diagnosis (years; ± SD)	28.7 ±5.2	30.9 ±4.6	30.2 ±4.2	30.6 ±2.9	30.8 ±4.2	27.5 ±3.5
HR (unadjusted)	0.51	Reference	7.24	1.78	4.88	17.93
95% CI	0.20-1.26		5.11-10.27	0.95-3.33	3.39-7.03	9.55-33.66
p-value	0.15		1.2×10 ⁻²⁸	0.07	1.6×10 ⁻¹⁷	2.7×10 ⁻¹⁹
HR (adjusted*)	0.52	Reference	6.93	1.18	3.41	15.83
95% CI	0.21-1.31		4.86-9.88	0.60-2.29	2.29-5.06	8.29-30.25
p-value	0.17		1.3×10 ⁻²⁶	0.63	1.4×10 ⁻⁹	6.3×10 ⁻¹⁷
Men with unimpaired health						
Number of new diabetes cases	3	36	53	9	44	11
HR (adjusted*)	0.46	Reference	6.91	1.26	3.04	20.15
95% CI	0.14-1.50		4.49-10.65	0.59-2.66	1.89-4.87	10.01-40.56
p-value	0.20		1.7×10 ⁻¹⁸	0.55	4.1×10 ⁻⁶	3.9×10 ⁻¹⁷
Second-generation (Israeli-born) Ethiopians						
Number of new diabetes incidences	5	52	80	2	5	6

HR (adjusted*)	0.52	Reference	6.96	2.25	4.42	29.10
95% CI	0.21-1.31		4.88-9.93	0.54-9.43	1.71-11.42	11.99-70.61
p-value	0.17		1.1×10^{-26}	0.27	2.1×10^{-3}	9.2×10^{-14}
Women						
Number of diabetes incidences	0	16	34	4	19	6
Mean follow-up (years; \pm SD)	12.4 \pm 4.2	12.2 \pm 4.3	11.8 \pm 4.2	12.2 \pm 4.1	12.3 \pm 4.2	11.0 \pm 3.9
Cumulative follow-up (person-years)	58,210.7	282,466.6	60,968.6	38,017.4	80,045.6	11,104.7
Incident rate (per 10^5 person-years)	-	5.7	55.8	10.5	23.7	54.0
Mean age at diagnosis (years; \pm SD)	-	26.1 \pm 4.4	27.8 \pm 3.7	24.6 \pm 4.4	29.8 \pm 4.0	26.1 \pm 4.7
HR (unadjusted)	-	Reference	10.15	1.87	4.17	10.79
95% CI	-		5.60-18.39	0.62-5.59	2.15-8.12	4.22-27.60
p-value	-		2.1×10^{-14}	0.26	2.6×10^{-5}	6.9×10^{-7}
HR (adjusted*)	-	Reference	9.66	1.13	2.40	7.04
95% CI	-		5.32-17.54	0.36-3.56	1.13-5.09	2.61-18.94
p-value	-		8.9×10^{-14}	0.08	0.02	1.1×10^{-4}

HR: hazard ratio

Figures





Figure 1

Associations between origin and cumulative type 2 diabetes incidence by adolescent BMI categories. Kaplan-Meier curves were plotted for each adolescent BMI category, for males and females separately. The dashed line represents the native Israelis with normal BMI group for each sex.

(A) Israeli born men



(B) Immigrant men




Figure 2

The association between male adolescent BMI and early onset type 2 diabetes among immigrants from another origin. This association was tested among immigrants from the Union of Soviet Socialist Republics (USSR), whose immigration to Israel paralleled the immigration from Ethiopia (1990-2000 vs. 1985-2000). Each analysis included three categories: all BMI groups, normal BMI (18.5 to 25 kg/m²), and high BMI (≥25 kg/m²). When the association was tested among Israeli-born men (A), the reference group for each analysis was native Israelis; and when it was tested for immigrant men (B), the reference group was USSR origin. Models were adjusted for BMI as a continuous variable in addition to the adjustments of the main analysis (see Table 2).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [SupplementsV32June24.docx](#)