

# A comparative study of the characteristics and clinical significance of extracellular traps of neutrophils in gingival samples of periodontitis and gingivitis patients

Fei Zhang

Linyi Central Hospital

Xi-Mei Yang

Linyi Central Hospital

Shu-Yu Jia (✉ [syjr29@163.com](mailto:syjr29@163.com))

linyi central hospital <https://orcid.org/0000-0002-9054-8584>

---

## Research article

**Keywords:** periodontitis; gingivitis; neutrophils; extracellular traps of neutrophils

**Posted Date:** September 12th, 2019

**DOI:** <https://doi.org/10.21203/rs.2.14324/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

# Abstract

**Background:** We sought to compare the characteristics and clinical significance of extracellular traps of neutrophils in gingival samples of periodontitis and gingivitis. **Methods:** The clinical indexes of gingival samples from periodontitis and gingivitis were measured; the expression of TNF- $\alpha$  and IL-8 were measured by real-time fluorescent quantitative PCR; the expression of TLR-8 and MMP-9 were measured by Western blotting assays. Chemotaxis, phagocytosis and phagocytic activity of neutrophils were measured. **Results:** The periodontal clinical indexes of periodontitis and gingivitis samples were significantly higher than those of healthy people ( $P < 0.05$ ). There were significant differences in BOP, PD and CAL between periodontitis and gingivitis samples ( $P < 0.05$ ), but no differences in gingival index (GI) ( $P > 0.05$ ). Compared with the healthy group, the expressions of TNF- $\alpha$  and IL-8 in the periodontitis group and the gingivitis group increased significantly ( $P < 0.05$ ), and TNF- $\alpha$  in the gingivitis group was significantly lower than that in the healthy group ( $P < 0.05$ ). The expression of IL-8 in the periodontitis group was significantly higher than that in the gingivitis group ( $P < 0.05$ ); the expression of TLR-8 and MMP-9 in the periodontitis group was different from that in the gingivitis group and the healthy people group, and the expression of TLR-8 and MMP-9 in the gingivitis group was significantly different from that in the healthy people group ( $P < 0.05$ ); the neutrophil mobility index in healthy people was  $(3.02 \pm 0.53)$ , and that in the periodontitis group was  $(2.21 \pm 0.13)$ , and  $(2.31 \pm 0.12)$  in the gingiva. There were significant differences between the periodontitis group and the gingivitis group ( $P < 0.05$ ,  $P < 0.05$ ). The phagocytic ability and activity of neutrophils in healthy people were  $(392.03 \pm 20.04)$ ,  $(89.34 \pm 10.12)$ , the periodontitis group was  $(360.32 \pm 18.23)$ ,  $(80.02 \pm 10.21)$ , and the gingivitis group was  $(359.32 \pm 20.11)$ ,  $(78.93 \pm 17.32)$ , respectively. There were significant differences between the periodontitis group and the gingivitis group and healthy people group ( $P < 0.05$ ,  $P < 0.05$ ). **Conclusion:** The chemotaxis of neutrophils in gingival samples of periodontitis and gingivitis patients was decreased, the phagocytosis ability and activity of neutrophils were reduced, and the release of extracellular trap-releasing inducible factors TNF- $\alpha$  and IL-8 also declined.

## Background

Periodontitis is an infectious disease caused by plaque microorganisms and their metabolites(1). The main clinical manifestations are loss of periodontal attachments and gradual destruction and absorption of the alveolar bone(2). In this process, the immune response to the pathogenic microorganism plays an important role. The gum is one of the periodontal tissues (the gum, periodontal ligament, alveolar bone, and cementum) and is directly exposed to the mouth and consists of keratinized epithelium and connective tissue covering the alveolar bone and roots. Gum disease is a disease that is confined to the gum tissue, the most common being chronic marginal gingivitis. Plaque microorganisms are the initiating factor of gingivitis and periodontitis. Among them, non-adherent gingival plaque is closely related to the occurrence and development of periodontitis and is considered to be the "development frontier" of periodontitis, which is related to the rapid destruction of the alveolar bone. A large number of activated immune cells cannot only trigger the host's immune response to periodontal pathogen infection, but also

release a cascade of inflammatory factors, including various acute phase reaction proteins, cytokines, etc., leading to the destruction of periodontal soft tissue and hard tissue(3). Neutrophils have a rod-shaped nucleus or lobulated nucleus and fine particles. The cytoplasm is colorless or reddish. It has chemotaxis, phagocytosis and bactericidal action. When the body is attacked by pathogens, neutrophils are under the action of chemokines. They quickly gather near the site of infection, enter the infected tissue through the vessel wall, and kill pathogenic microorganisms(4) by phagocytosis and release of bactericidal particles. Neutrophil extracellular reticular traps (NETs) are a new defense mechanism(5), and unlike apoptosis and necrosis, release a variety of bactericidal substances to remove pathogenic microorganisms, including neutrophil primary granule release such as protease or secondary particle bacteriostatic peptide LL-37, etc.; tertiary particles such as protease-containing matrix metalloproteinase 9 (MMP-9), etc(6, 7). Many factors can stimulate neutrophils to produce neutrophil extracellular traps. For example, plasma TNF- $\alpha$  and interleukin-8 can also stimulate neutrophil release of NETs(8, 9).

## Methods

### 1.1 General Information

Between July 2015 and September 2016, we randomly selected 27 patients (14 males and 13 females) with periodontitis (CP), 17 patients (7 males and 10 females) with gingivitis (cg) and 20 patients (8 males and 12 females) with no periodontal disease (PH). Subjects were 24-60 years old with an average age of  $(33.24 \pm 6.93)$  years. Periodontitis and gingivitis were diagnosed according to the diagnostic criteria of the Periodontal Disease Association

The inclusion criteria were: at least 20 teeth in the mouth, 4-6 teeth with severe periodontitis in all periodontal patients; no periodontal treatment in the preceding 6 months; no smoking history in the preceding 6 months; no history of head and neck radiotherapy; no antibiotics, phenytoin sodium, cyclosporine, calcium channel blockers, oral contraceptives, atropine and other drugs within the month; no serious systemic diseases and serious infections in other areas. Pregnant or lactating women were excluded. The study protocol was approved by the Ethics Committee of Linyi Central Hospital. All study participants provided written informed consent.

### 1.2 Experimental methods

#### 1.2.1 Determination of clinical indicators

The researchers who received professional training in periodontal examination and passed the standard conformance test ( $Kappa=0.73$ ) used the Williams probe to detect periodontal clinical indicators in all subjects, including the gingival index (GI), detection of bleeding on probing (BOP), depth of detection (PD), and clinical attachment level (CAL). Among them, PD and CAL were detected at 6 sites on the buccal (lip) side and at the distal, middle and proximal sites of the lingual side.

#### 1.2.2 Collection of peripheral blood samples

Two mL of peripheral venous blood of all subjects was extracted and placed in an anticoagulation tube containing K2EDTA. Peripheral blood samples were collected by centrifugation (4000 r/min, 6 min) at 4°C. The supernatant was placed in a sterile Eppendorf tube and stored at -80 ° C until use.

### **1.2.3 Extraction of total RNA from peripheral blood samples and RT-PCR determination**

One ml of Trizol was added in the serum and repeatedly pipetted, and the cell lysate was transferred to a 1.5 ml Eppendorf tube, left for 5 min, and fully lysed. Two hundred µL of chloroform was added per 1 ml of Trizol, mixed for 15 s, and left at room temperature for 3 min. After centrifugation at 4° C 12000 g for 15 min, the upper aqueous phase was aspirated and transferred to another new Eppendorf tube. Then, 0.5 ml of isopropanol was added per 1 ml of Trizol, mix, and left at room temperature for 10 min. After centrifugation at 42000g of 4°C for 10 min, the supernatant was discarded, and RNA was precipitated on the bottom of the tube. One ml of 75% ethanol was added per 1 ml of Trizol and mix vigorously. After centrifugation at 4° C at 7500 g for 5 min, the supernatant was discarded. RNA was precipitated at room temperature for 5-10 min, dissolved in 30 µL of EPC treated water, and stored at -80°C for use. RNA purity and concentration were measured by spectrophotometry at A260 and A260/280. Total RNA was extracted using a total RNA rapid extraction kit and reverse transcribed with Super M-MLV reverse transcriptase using miRNA-specific reverse transcription primers (Table 1). RT-PCR was performed in a real-time quantitative system. The amplification conditions are shown in Table 2, the primer design is shown in Table 3, and the reaction system is shown in Table 4.

### **1.2.4 Western blotting assays**

The expression of TLR-8 and MMP-9 in cells was detected by Western blotting assays. The protein was resolved by a 4%-10% polyacrylamide gel electrophoresis and the gel was transferred to a PVDF membrane using SDS polyacrylamide gel electrophoresis. After blocking with 5% non-fat dry milk for 1 h at room temperature, the membrane was first incubated with 1:500 diluted primary antibody overnight, and after wash with TBS-T buffer, the membrane was incubated with secondary antibody for 1 h. Following rinse with TBS-T solution, enhanced chemiluminescence was carried out. The PVDF membrane was placed in a developing solution in a dark room and exposed to an X-ray film for exposure and development. Protein densitometry was performed using Image 1.6 software.

### **1.2.5 Preparation of leukocyte suspension**

After extracting 3 mL of venous blood from all subjects, we used heparin as an anticoagulant to precipitate red blood cells, and white blood cells were taken. After wash,  $1 \times 10^6$  / mL white blood cell suspension was prepared with RPMI1640 medium containing 10% bovine serum.

### **1.2.6 Determination of neutrophil chemotaxis**

Using agarose glass plate method: agarose solution was prepared with 1640 solution, and after addition of 10% inactivated AB serum, the glass plate was poured . Each group consisted of three circular holes

arranged in a straight line with a diameter and a pitch of 2.5 mm. Five  $\mu\text{L}$  of a white cell suspension ( $1 \times 10^6/\text{mL}$ ) was added in triplicates to the mesopores and *Escherichia coli* was added to the side wells for 24 hours, and 5  $\mu\text{L}$  of the filtrate was used as the chemokine, and the other side was treated with 5  $\mu\text{L}$  of the 1640 solution. The agarose glass plate was placed in a 37 ° C incubator, incubated with saturated humidity and 5%  $\text{CO}_2$  for 4 h, removed, fixed with methanol and formaldehyde. After agarose gel electrophoresis, the gel was stained with Wright, and measured under an optical microscope. The distance a cell moved from the edge of the hole to the hole on either side was measured.

### 1.2.7 Determination of phagocytic function of neutrophils

*Staphylococcus aureus* was grown on agar slants for 24 h, colonies were washed with sterile isotonic saline, washed twice with PBS, and the cells were suspended at  $5 \times 10^7/\text{ml}$  by specific concentration method. Three drops of heparinized blood were taken on the concave slide, and three drops of *Staphylococcus aureus* suspension were added and after thorough mixing, they were placed in a sealed wet box. After incubation at 37 ° C for 30 min, the box was shaken once every 10 min; 1 drop was taken with a capillary suction tube, pushed onto a slide on a glass slide, fixed in methanol, stained with Giemsa, and counted by light microscopy.

### 1.3 Data processing

Data were expressed as mean  $\pm$  standard deviation. One-way analysis of variance was used for comparison between groups.  $P < 0.05$  was considered to have a significant statistical difference.

## Results

### 2.1 Clinical indicators

The clinical indicators in the groups are shown in Table 5. The periodontal clinical indexes of periodontitis and gingivitis samples were significantly higher than those of healthy people ( $P < 0.05$ ). Comparison of periodontitis with gingivitis samples showed that there were significant differences in the blood index, depth of detection and CAL ( $P < 0.05$ ), and there was no difference in the gingival index ( $P > 0.05$ ).

### 2.2 RNA expression measurement results

The purity of extracted total RNA was measured by agarose gel electrophoresis, and the results are shown in Fig. 1. As can be seen from the figure, RNA was of high purity and can be used for PCR amplification.

The expression levels of RNA in each group were measured by RT-PCR. The results are shown in Table 6. The results showed that the expression of TNF- $\alpha$  and IL-8 in the periodontitis group and the gingivitis group increased significantly compared with the healthy control group ( $P < 0.05$ ), and the expression of

TNF-alpha in the gingivitis group was significantly lower than that in the periodontitis group ( $P < 0.05$ ), and the expression of IL-8 was significantly higher than that in the periodontitis group ( $P < 0.05$ ).

### 2.3 Western blotting results

The plasma expression levels of TLR-8 and MMP-9 were determined by Western blotting assays. The results are shown in Fig. 2 and Table 7. The expression of TLR-8 and MMP-9 in the periodontitis group was different from that in the gingivitis group and the healthy control group, and the difference between the gingivitis group and the healthy control group was significant ( $P < 0.05$ ). The expression of TLR-8 and MMP-9 in the periodontitis group was higher than that in the gingivitis group, but there was no significant difference ( $P > 0.05$ ). It indicated that the neutrophil chemotactic function in periodontitis and gingivitis patients was significantly lower than that in healthy subjects.

### 2.4 Neutrophil chemotaxis

The neutrophil shift index was ( $3.02 \pm 0.53$ ) in healthy subjects, ( $2.21 \pm 0.13$ ) in the periodontitis group, and ( $2.31 \pm 0.12$ ) in the periodontitis group. There were significant differences between the periodontitis group and the gingivitis group and the healthy control group ( $P < 0.05$ ,  $P < 0.05$ ). This indicated that the chemotactic function of neutrophils in patients with periodontitis and gingivitis is significantly lower than that in healthy subjects.

### 2.5 Determination of neutrophil phagocytosis

The phagocytic ability and activity of neutrophils were ( $392.03 \pm 20.04$ ), and ( $89.34 \pm 10.12$ ) in healthy subjects, and ( $360.32 \pm 18.23$ ) and ( $80.02 \pm 10.21$ ) in the periodontitis group, and ( $359.32 \pm 20.11$ ) and ( $78.93 \pm 17.32$ ) in the gingivitis group. There were significant differences between the periodontitis group and the gingivitis group and the healthy control group ( $P < 0.05$ ,  $P < 0.05$ ). The results showed that the phagocytic ability and activity of neutrophils in periodontitis and gingivitis patients were significantly lower than those in healthy subjects.

## Discussion

Periodontal disease is a chronic inflammatory disease of the tooth-supporting tissue caused by plaque infection, including gingivitis and periodontitis. The main manifestation is periodontal support tissue destruction(10).

In 2004, Brinkmann et al. first proposed another method for killing pathogenic microorganisms, neutrophil extracellular reticular traps (NETs), which differ from apoptosis and necrosis(11). NETs are closely related to human diseases and play a unique role in the development of disease(12, 13). In most cases, the production of NETs is conducive to the body's innate immunity(14), but recent studies have found that the network is also involved in the pathological process of human disease. When a mesh is produced, a large amount of autoantigens(15) is produced. If the removal is not timely, a large amount of antigens will accumulate, which may induce the body to produce a large amount of autoantibodies, leading to

autoimmune diseases. Neural network research focuses on infectious diseases and autoimmune diseases, but it is also closely related to thrombotic diseases (acute myocardial infarction)(16).

NETs is a newly discovered neutrophil bactericidal mechanism that activates neutrophils to release chromatin and then combines with the gingival sulcus to form a new defense network against gingival plaque bacteria(17). It has been reported(18, 19) that under the induction of IL-8, LPS and interferon, the neutrophil nucleus is deformed, the chromatin is homogeneous, the nuclear membrane is broken, the nuclear material is directly in contact with the particles, forming a mixture, and finally the cell membrane is broken, NETs are released. This study found that compared with the healthy control group, the expression of TNF- $\alpha$  and IL-8 in the periodontitis group and the gingivitis group was significantly increased ( $P < 0.05$ ), and TNF- $\alpha$  in the gingivitis group was significantly lower than that in the periodontitis group. The expression of IL-8 was significantly higher in the xx group ( $P < 0.05$ ) than in the periodontitis group ( $P < 0.05$ ).

MMP-8 secreted by neutrophils, also known as neutrophil collagenase, is one of the most important MMPs in inflammatory periodontal tissues. Type I collagen fibers are the main components of the periodontal ligament and alveolar bone. Studies have shown that the main role of MMP-8 is to destroy type I collagen fibers. MMP-8 is also present in normal periodontal tissues. In the absence of inflammation, the ratio of MMP-8 and its endogenous inhibitor (TIMP-1) is close to 1:1, which is involved in normal metabolism of the extracellular matrix. However, when pathogens invade, the expression of MMP-8 is abnormally regulated by cytokines, which disrupts the balance between MMP-8 and TIMP-1, leading to damage of type I collagen in periodontal tissues. This study showed that there was a difference in the expression of TLR-8 and MMP-9 between the periodontitis group and the gingivitis group and the healthy control group, and the difference between the gingivitis group and the healthy control group was significant ( $P < 0.05$ ). The expression of TLR-8 and MMP-9 in the periodontitis group was higher than that in the gingivitis group, but there was no significant difference ( $P > 0.05$ ), indicating that the chemotactic function of neutrophils in patients with periodontitis and gingivitis is significantly lower than that in healthy controls.

Neutrophils are important cells to prevent periodontal infection. The study on the role of neutrophils in juvenile periodontitis shows that chemotaxis of neutrophils decreases and the phagocytosis function of neutrophils is normal. This report proves that the weakening of neutrophil chemotaxis makes juvenile periodontitis patients susceptible to specific local bacteria and causes rapid destruction of the periodontal tissue(20). This study indicated that there were significant differences between the periodontitis group and the gingivitis group and healthy control group ( $P < 0.05$ ). The results showed that the neutrophil chemotactic function of periodontitis and gingivitis patients was significantly lower than that of healthy controls. There were significant differences between the periodontitis group and the gingivitis group and the healthy control group ( $P < 0.05$ ,  $P < 0.05$ ). The results showed that the phagocytic ability and activity of neutrophils in periodontitis and gingivitis patients were significantly lower than those in healthy controls.

In summary, the chemotaxis of neutrophils in gingival samples of periodontitis and gingivitis decreased, the ability of phagocytosis and activity of neutrophils were reduced, and the release of extracellular trap-releasing inducible factors TNF-alpha and IL-8 declined.

## List Of Abbreviations

neutrophil extracellular reticular traps: NETs

matrix metalloproteinase 9: MMP-9

periodontal disease: PH

gingival index: GI

bleeding on probing: BOP

depth of detection: PD

clinical attachment level: CAL

## Declarations

**Ethics approval and consent to participate** The study protocol was approved by the Ethics Committee of Linyi Central Hospital. All study participants provided written informed consent.

**Consent for publication:** All study participants provided written informed consent.

**Availability of data and material** Data could be available to readers at reasonable request from the corresponding author .

**Competing interests:** The authors declare that they have no competing interests.

**Funding:** No funding was received for this study.

**Authors' contributions** FZ contributed to the conception and design of the study; XY contributed to the acquisition of data; SJ performed the experiments; SJ contributed to the analysis of data; FZ wrote the manuscript; All authors reviewed and approved the final version of the manuscript.

**Acknowledgements:** None.

## References

1. Hajishengallis G. Periodontitis: from microbial immune subversion to systemic inflammation. Nature reviews Immunology. 2015;15(1):30-44.

2. Eke PI, Dye BA, Wei L, Slade GD, Thornton-Evans GO, Borgnakke WS, et al. Update on Prevalence of Periodontitis in Adults in the United States: NHANES 2009 to 2012. *Journal of periodontology*. 2015;86(5):611-22.
3. Zambon JJ, Haraszthy VI, Hariharan G, Lally ET, Demuth DR. The Microbiology of Early-Onset Periodontitis: Association of Highly Toxic *Actinobacillus actinomycetemcomitans* Strains With Localized Juvenile Periodontitis. *Journal of periodontology*. 1996;67 Suppl 3S:282-90.
4. Jaillon S, Galdiero MR, Del Prete D, Cassatella MA, Garlanda C, Mantovani A. Neutrophils in innate and adaptive immunity. *Seminars in immunopathology*. 2013;35(4):377-94.
5. Zawrotniak M, Rapala-Kozik M. Neutrophil extracellular traps (NETs) - formation and implications. *Acta biochimica Polonica*. 2013;60(3):277-84.
6. Hermosilla C, Caro TM, Silva LM, Ruiz A, Taubert A. The intriguing host innate immune response: novel anti-parasitic defence by neutrophil extracellular traps. *Parasitology*. 2014;141(11):1489-98.
7. Manda A, Pruchniak MP, Arazna M, Demkow UA. Neutrophil extracellular traps in physiology and pathology. *Central-European journal of immunology*. 2014;39(1):116-21.
8. Demers M, Wagner DD. Neutrophil extracellular traps: A new link to cancer-associated thrombosis and potential implications for tumor progression. *Oncoimmunology*. 2013;2(2):e22946.
9. White PC, Chicca IJ, Cooper PR, Milward MR, Chapple IL. Neutrophil Extracellular Traps in Periodontitis: A Web of Intrigue. *Journal of dental research*. 2016;95(1):26-34.
10. Buduneli N, Kinane DF. Host-derived diagnostic markers related to soft tissue destruction and bone degradation in periodontitis. *Journal of clinical periodontology*. 2011;38 Suppl 11:85-105.
11. Warnatsch A, Ioannou M, Wang Q, Papayannopoulos V. Inflammation. Neutrophil extracellular traps license macrophages for cytokine production in atherosclerosis. *Science*. 2015;349(6245):316-20.
12. Arazna M, Pruchniak MP, Zycinska K, Demkow U. Neutrophil extracellular trap in human diseases. *Advances in experimental medicine and biology*. 2013;756:1-8.
13. Vorobjeva NV, Pinegin BV. Neutrophil extracellular traps: mechanisms of formation and role in health and disease. *Biochemistry Biokhimiia*. 2014;79(12):1286-96.
14. Park SY, Shrestha S, Youn YJ, Kim JK, Kim SY, Kim HJ, et al. Autophagy Primes Neutrophils for Neutrophil Extracellular Trap Formation during Sepsis. *American journal of respiratory and critical care medicine*. 2017;196(5):577-89.
15. Sorensen OE, Borregaard N. Neutrophil extracellular traps - the dark side of neutrophils. *The Journal of clinical investigation*. 2016;126(5):1612-20.
16. Ginley B, Emmons T, Sasankan P, Urban C, Segal BH, Sarder P. Identification and characterization of neutrophil extracellular trap shapes in flow cytometry: SPIE; 2017.
17. Lightfoot YL, Kaplan MJ. Disentangling the role of neutrophil extracellular traps in rheumatic diseases. *Current opinion in rheumatology*. 2017;29(1):65-70.
18. Stakos DA, Kambas K, Konstantinidis T, Mitroulis I, Apostolidou E, Arelaki S, et al. Expression of functional tissue factor by neutrophil extracellular traps in culprit artery of acute myocardial

infarction. European heart journal. 2015;36(22):1405-14.

19. Meher AK, Spinosa M, Davis JP, Pope N, Laubach VE, Su G, et al. Novel Role of IL (Interleukin)-1beta in Neutrophil Extracellular Trap Formation and Abdominal Aortic Aneurysms. Arteriosclerosis, thrombosis, and vascular biology. 2018;38(4):843-53.
20. Sochalska M, Potempa J. Manipulation of Neutrophils by Porphyromonas gingivalis in the Development of Periodontitis. Frontiers in cellular and infection microbiology. 2017;7:197.

## Tables

**Table 1 Reverse transcription system**

5×PrimeScript Buffer	4μL
1×PrimeScript RT Enzyme Mix I	1μL
Oligo dT Primer (50μM)	1μL (25pmol)
Random 6 mers (100μM)	1μL (50pmol)
Total RNA	1μL
RNase Free dH2O	12μL

**Table 2 PCR conditions**

step	temperature	time
Pre-denaturation	95°C	10 min
transsexual	94°C	30 s 40 cycles
Annealing/extension	60°C	1 min
Melting curve analysis		
	95°C	15 s
	60°C	1 min
	94°C	15 s
	60°C	15 s

**Table 3 Sequences of primers used in the study**

RNA	Upstream	Downstream
TNF- $\alpha$	5'-TGTTCTTACACCCCTCCCCTTTT-3'	5'-TATAAGTGTGACCCGGCTGAGAA-3'
IL-8	5'-GGCCGAUUGUGAACAUGGATT-3'	5'-UCCAUGUUCACAAUCGGCCGC-3'
$\beta$ -actin	5'-TGGCACCCAGCACAATGAA-3'	5'-CTAAGTCATAGTCGCGCTAGAAGCA-3'

**Table 4 PCR system**

5xPrimeEx Taq TM II (2x)	12.5 $\mu$ L
PCR Forward Primer (10 $\mu$ M)	1 $\mu$ L 0.4 $\mu$ M
PCR Reverse Primer (10 $\mu$ M)	1 $\mu$ L 0.4 $\mu$ M
DNA template	2 $\mu$ L
dH <sub>2</sub> O (sterilized distilled water)	8.5 $\mu$ L
Total	25 $\mu$ L

**Table 5 Clinical indicators of the study population**

Group	Periodontal index			
	GI	Bop positive (%)	PD	CAL
PH	0	0	1.66 $\pm$ 0.21	1.66 $\pm$ 0.21
CP	2.25 $\pm$ 0.23	37.43 $\pm$ 0.21	4.34 $\pm$ 0.30	4.91 $\pm$ 0.20
CG	2.26 $\pm$ 0.15	28.34 $\pm$ 0.22	2.25 $\pm$ 0.12	2.21 $\pm$ 0.13
F	49.932	50.043	29.943	48.039
P	<0.05	<0.05	<0.05	<0.05

**Table 6 RNA expression measurement results**

Group	TNF- $\alpha$ ng/mL	IL-8 ng/mL
PH	0	0
CP	40.21 $\pm$ 10.03	102.32 $\pm$ 15.34
CG	26.31 $\pm$ 5.02	114.92 $\pm$ 20.11
F	49.323	59.182
P	<0.05	<0.05

**Table 7 TLR-8 and MMP-9 protein expression levels**

Group	TLR-8	MMP-9
PH	0.91±0.04	0.89±0.12
CP	0.89±0.03	0.82±0.04
CG	0.45±0.11	0.52±0.02
F	89.043	93.045
P	<0.05	<0.05


**Table 8 Neutrophil movement index results**

Group	Mobile index
PH	3.02±0.53
CP	2.21±0.13
CG	2.31±0.12
F	90.485
P	<0.05a

**Table 9 Neutrophil phagocytosis results**


Group	Phagocytic ability	Phagocytic activity
PH	392.03±20.04	89.34±10.12
CP	360.32±18.23	80.02±10.21
CG	359.32±20.11	78.93±17.32
F	89.043	93.045
P	<0.05	<0.05

## Figures



**Figure 1**

RNA agarose gel electrophoresis



**Figure 2**

Western blotting results