Multi-scale determinants of small mammal and lizard dimensions of diversity across Amazonian insular forest fragments

Ana Filipa Palmeirim (✉ anafilipapalmeirim@gmail.com)

University of East Anglia <https://orcid.org/0000-0003-3931-0578>

Fabio Farneda

Universidade Federal de Santa Maria

Marcus Vinicius Vieira

Universidade Federal do Rio de Janeiro

Carlos A. Peres

University of East Anglia

Research Article

Keywords: beta-diversity, community-weighted mean trait values, environmental filter, functional diversity, functional redundancy, habitat loss and fragmentation, island biogeography, phylogenetic diversity

Posted Date: May 12th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-503411/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Landscape Ecology on August 13th, 2021. See the published version at <https://doi.org/10.1007/s10980-021-01311-w>.

Abstract

Context

Hydropower development is one of the primary drivers of habitat loss and insular fragmentation. Yet, studies quantifying such effects on biodiversity are mostly limited to taxonomic metrics, often overlooking species identity and their functional and phylogenetic roles.

Objectives

We examined taxonomic, functional and phylogenetic responses to habitat loss and insular fragmentation of small mammals and lizards within (α) and between sites (β).

Methods

Both taxa were surveyed across 25 islands within the Balbina Hydroelectric Reservoir, and adjacent continuous forest, in Central Amazonia. Each dimension of diversity was related to spatial and habitat variables. We also examined functional composition using community-weighted mean trait values, and community redundancy using functional uniqueness. β -diversity was partitioned into their richness (β_{rich}) and replacement (β_{repl}) components.

Results

Functional and phylogenetic α -diversities of both taxa mirrored the taxonomic dimension, all of which increased with forest area. Individual small mammal (body mass and matrix tolerance), and lizard traits (heliothermic mode and habitat type) were also predicted by forest area. For both taxa, species functional uniqueness decreased with forest area, and all dimensions of β -diversity were predominantly partitioned in β_{rich} .

Conclusions

The environmental filter created by forest area resulted in the low conservation value associated with small forest islands, only occupied by a small set of species comprised by generalist lizards and matrix-tolerant small mammals. On the other side, large forest sites ensured ecosystem resilience to disturbance. To maintain ecosystem integrity, creating a myriad small island over large expanses of floodwaters should be avoided in future hydropower development.

Introduction

Habitat loss, fragmentation and subsequent degradation are primary drivers of biodiversity loss worldwide (Sala et al. 2000; Haddad et al. 2015). In the tropics, hydropower development is a major cause of habitat loss and fragmentation, often creating vast archipelagic landscapes, in which forest islands are isolated within a uniformly hostile open-water matrix (Jones et al. 2016). In these archipelagic landscapes, island taxa typically experience a novel hyper-disturbance regime, resulting in drastic shifts in species diversity and community composition through species extinction and turnover (Cosson et al. 1999; Gibson et al. 2013; Benchimol and Peres 2015a). Despite the growing number of recent studies showing the pervasive ecological impacts of hydroelectric dams, only a few have employed functional and phylogenetic metrics (Jones et al. 2016). Nevertheless, taxonomic, functional and phylogenetic dimensions of diversity can be simultaneously affected by habitat disturbance (e.g., Aguirre et al. 2016, Salgado-Luarte et al. 2018), while their study provides complementary information (Graham and Fine 2008).

Functional traits reflect environmental preferences and associated behaviours that directly influence individual performance (Violle et al. 2007). Functional diversity therefore incorporates species trait variation that potentially affects species performance, fitness and ecological functions (Weiss and Ray 2019). Phylogenetic diversity integrates the shared ancestry of species in terms of both the amount of evolutionary history and degree at which species are phylogenetically related (Webb et al. 2002; Veron et al. 2017). Since functional traits that allow species to persist in the environment tend to be evolutionarily conserved (i.e. positive phylogenetic signal: Safi et al. 2011), phylogenetic and functional diversity tend to be correlated (Tucker et al. 2018).

Across fragmented landscapes, larger habitat fragments tend to harbour more complete species assemblages, eventually accumulating more species traits (Dainese et al. 2015; Farneda et al. 2018). This results in the functional redundancy of those assemblages which are also more resilient to disturbance (Petchey et al. 2007; Ricotta et al. 2016). At the other extreme of the habitat fragmentation gradient, consisting of small habitat fragments harbouring depauperate species assemblages, the combination of the remaining species traits is therefore likely to be more functionally unique. In particular, local extinctions of functionally unique species at those smaller fragments may disproportionately affect ecosystem functioning (Elmqvist et al. 2003).

Improving our understanding on how species diversity is organized and maintained following habitat loss and fragmentation further requires going beyond α -diversity (within-site) metrics, thereby considering species diversity between sites (β). While taxonomic β -diversity reflect differences in species composition between sites (Whittaker 1972), functional β -diversity discriminates change between communities of taxa with either similar or different functions in the system (e.g., Ricotta and Burrascano 2008), and phylogenetic β -diversity distinguishes between phylogenetically basal and terminal changes between communities, branch points occurring early and late in a phylogeny (Graham and Fine 2008). Disentangling the contribution of the two components of either of the β -diversity dimensions – richness differences and replacement – further enables inference about the mechanisms of community assembly

of habitat loss and fragmentation, forest area

is reduced thereby creating an environmental filter under which certain species, traits and genetic lineages are expected to be successively extirpated from smaller sites (Cardoso et al. 2014; Carvalho et al. 2012).

Here, we examine multi-scale effects of habitat loss and insular fragmentation on the diversity of assemblages of two major vertebrate taxa – small mammals (marsupials and rodents) and lizards – in the Central Amazonian archipelagic landscape of the Balbina Hydroelectric Reservoir, and nearby continuous forest. These vertebrate groups are ecologically diversified in neotropical forests, spanning across the vertical forest strata and occupying multiple trophic niches (Ávila-Pires 1995; Paglia et al. 2012). Both groups are characterized by relatively low dispersal abilities due to their reduced size and non-volant habits. The varied roles played by both groups in natural ecosystems – predators, prey, grazers, seed dispersers and commensal species – are critical for ecosystem functioning, including forest regeneration (Terborgh et al. 2001). Small mammal and lizard assemblages were surveyed across a gradient of habitat loss and fragmentation comprised of 25 islands of different sizes, degrees of isolation and configurations, in addition to control continuous forest sites. We examined patterns and predictors of taxonomic, functional and phylogenetic diversity, trait-level responses and functional redundancy of small mammals and lizard assemblages. Additionally, to elucidate processes underlying community (dis)assembly resulting from habitat loss and fragmentation, we examined the partition of β -diversity in its two components. Across the insular landscape, we tested the following hypotheses:

(1) Taxonomic, functional and phylogenetic α -diversity of both taxa increase with island size and habitat quality (i.e. low-intensity edge effects and higher tree species richness) and decrease in more isolated forest islands (MacArthur and Wilson 1967; Meynard et al. 2011).

(2) Traits allowing small mammal and lizard species to persist in small, isolated and habitat-degraded islands will be favoured over the gradient of fragmentation (Devictor et al. 2008). In particular, we expect those traits to be related to higher dispersal ability (e.g. larger body size and overall matrix tolerance) and generalist habits in terms of both habitat (e.g. ability to use of open-habitat areas) and diet (Farneda et al. 2015; Wang et al. 2010).

(3) Impoverished assemblages of small islands have higher mean species uniqueness compared to larger islands because the few remaining species have more unique combinations of traits (Ricotta et al. 2016).

(4) Following a reduction in forest area, environmental filtering promotes selective extinction of species, increasing differences in β -diversity more due the richness-component (Carvalho et al. 2012). The importance of the richness-component is further expected to increase for larger differences in forest area (Cardoso et al. 2014).

Material And Methods

Study area

Loading [MathJax]/jax/output/CommonHTML/fonts/TeX/fontdata.js

Surveys were carried out in the insular landscape of the Balbina Hydroelectric Reservoir and its immediate surroundings in Central Brazilian Amazonia (1°48'S, 59°29'W; Fig. 1). This landscape was created following the construction of Balbina Dam in 1986 on the Uatumã River, a left-bank tributary of the Amazon River. This dam flooded an area of 312,900 ha of pristine primary forest, within the 443,772-ha hydroelectric reservoir (FUNCATE/INPE/ANEEL 2000). Given the undulating topography, its former hilltops were converted into 3,546 land-bridge islands of varying sizes and configuration. Many dead relics of emergent trees are still standing across the reservoir lake, as the submerged primary forest was never clear-cut. Vegetation mostly consists of dense closed-canopy *terra firme* forest, but many small islands were strongly affected by edge-related windfalls and ephemeral wildfires, which occurred during a late-1997 to early-1998 El Niño drought (Benchimol and Peres 2015b). The area within and around the former left bank of the Uatumã river has been legally protected since 1990 by the 942,786-ha Uatumã Biological Reserve, the largest reserve in its category in Brazil, further contributing to the current low human disturbance in this area. Mean annual temperature and mean annual rainfall in this region are 28°C and 2,376 mm (IBAMA 1997).

Small mammal and lizard assemblages were sampled on 25 islands (Fig. 1), 24 of which included surveys of both taxa. Due to logistic limitations, surveys of either small mammals or lizards were conducted on two additional islands. Island selection aimed to achieve a wide range of spatial configurations with island size ranging from 0.55 to 14,660 ha (mean $\pm$ SD: 199 ± 344 ha; Supporting Information Table S1) and isolation distances from each focal island to the nearest continuous forest in the mainland, ranging from 44 to 11,872 m ($4,351 \pm 3,386$ m). As a baseline, we also surveyed four control continuous forest (CF) regions. In one of the four CF regions (CF₂), trapping transects used to survey small mammals and lizards were placed in both riparian ($N = 5$ transects) and non-riparian areas ($N = 5$). Due to logistic limitations, riparian transects could not be placed in any other CF region. Given the high microhabitat specialization of lizards (Vitt and Caldwell 2014), for this group we considered riparian and non-riparian transects of CF₂ as two different CF sites: CF₂ and CF₂ – riparian. We therefore surveyed small mammals and lizards in four and five CF sites, respectively. Given the wide distribution of CF sites in the mainland, this number of CF sites was considerate adequate to obtain a representative sample of the assemblages of both taxa occupying the reservoir area before flooding (Fig. 1).

Small mammal and lizard surveys

Surveys were carried out in two seasons, the first from April to November 2014 and the second from April to November 2015. Each season consisted of a trapping session of 16 consecutive nights/days. To minimise the impact any eventual seasonal variation, sites were surveyed in the same order in both field seasons. Small mammals were sampled using traplines containing nine stations of live traps (hereafter, LTs) followed by an array of three pitfall units, and lizards were sampled using shorter transects consisting of the same array of three pitfalls. Each LT station was placed 20-m apart from others and included two Sherman traps (23 × 9 × 8 cm, H. B. Sherman Traps, Inc., Tallahassee, Florida) and one wire mesh trap (30 × 17.5 × 15 cm, Metal Miranda, Curitiba, Paraná). At each LT station, one trap was set on the ground, one in the understorey (< 1.5 m high), and one in the (sub)canopy (> 10 m high). Traps of

different types were placed alternatively on the ground and in the understory across consecutive stations, but only Sherman traps were placed in the canopy due to logistic limitations. The forest canopy stratum was sampled by placing a trap in a mobile wood platform that was left in contact with the canopy tree branch (Lambert et al. 2006). LTs were baited with a mix of bananas, peanut powder, sardines and oatmeal. Pitfall traps (plastic buckets of 100L) were also spaced by 20-m intervals and connected by a plastic drift fence 50-cm high and 10-cm underground, with 10 m of fence extending beyond the two terminal pitfalls.

The number of transects placed at different sites varied according to their area. This allowed us to increase sampling size (i.e., number of individuals recorded) in larger forest sites, where trap density and consequently the probability of an individual passing near a trap are lower. Due to spatial restrictions on islands smaller than 2 ha and those between 2 and 10 ha, alternative smaller transects were placed containing three LT stations followed by an array of one pitfall, and by six LT stations followed by an array of two pitfalls, respectively. Larger islands were sampled by as many as four transects, according to their size classes: 10 to 50 ha (one transect), 50 to 200 ha (two), 200 to 500 ha (three), and > 500 ha (four transects). CF sites were sampled by either five (lizards: CF₂, CF_{2-riparian} and CF₄), six (both taxa: CF₁ and CF₃), 10 (small mammals: CF₂) and 12 transects (small mammals: CF₄). Due to logistic restrictions, CF₂ and CF₄ were only sampled during either the first (2014) or second (2015) field season, respectively, which reflects the larger number of transects placed therein (for further details on sampling effort per site, see Table S1).

In total, we surveyed small mammals across 79 trapping transects and lizards across 71 transects, that were accordingly nested within 29 and 30 sampling sites. All traps were inspected daily. All small mammal individuals recorded were weighted, measured, collected a tissue sample (i.e., ear tip for small mammals and tail tip for lizards) and tagged (Fish and Small Animal Tag, size 1; National Band and Tag Co., Newport, Kentucky), so that any subsequent recaptures could be distinguished. During the second field season, we also collected tissue samples of lizards for genetic analyses by carefully removing the tail tip of every individual. This allowed us to distinguish individuals that had been previously captured. As recaptures represented a small fraction of the total number of records (< 2%), all lizard individuals recorded during either the first or second field season were assumed to have been captured only once. Although tissue samples were collected from both small mammals and lizards, we were not always able to identify at the species-level records at all sites the following small mammals: *Proechimys* spp. (*P. cuvieri* and *P. guyanensis*) and *Oecomys* spp. 1 (*O. roberti* and *O. bicolor*); and lizards: *Leposoma* spp. (*L. percarinatum* and *L. guianense*), *Norops* spp. (*N. planiceps* and *N. chrysolepis*), and *Kentropyx* spp. (*K. calcarata* and *K. altamazonica*). For both taxa, congeners within these genera are ecologically very similar (Ávila-Pires 1995; Jones et al. 2009), so we further refer to those taxa as *ecospecies*. To streamline, we hereafter use *species* to refer to both species and *ecospecies*. This study followed the guidelines provided by the American Society of Mammalogists (Sikes 2016), the American Society of Ichthyologists and Herpetologists (Beaupre et al. 2004) and the institutional animal care and use committee of the Federal University of Rio de Janeiro (CEUA-CCS UFRJ 2010).

Spatial and habitat variables

For each surveyed site, we considered (1) island size (area), as the total island area of each focal island ($\log_{10} x$); (2) island shape (shape), indicating the intensity of edge effects, measured as the perimeter length to area ratio of each focal island ($\log_{10} x$); (3) a proximity (prox) metric to other islands and CF sites, following McGarigal et al. (2012), consisting of the sum of all island areas divided by the squared sum of edge-to-edge distances from each focal island to all islands within a buffer of 500 m (for small mammals) and 250 m (lizards) ($\log_{10} x + 1$; Palmeirim et al. 2017, 2018); (4) the nearest Euclidian distance to the mainland (dist); and (5) the number of tree species (s_{tree}), calculated by Benchimol and Peres (2015b) from floristic surveys in 0.25-ha forest plots within each focal island or mainland site. Spatial variables were calculated using ArcMap 10.1 (ESRI 2018) based on high-resolution multi-spectral RapidEye imagery (5-m resolution with 5-band colour imagery) of the entire study landscape. The habitat variable S_{tree} was obtained from Benchimol and Peres (2015b). Whenever included in the analysis, CF sites were assigned with area values one order of magnitude larger than the largest island (146,600 ha). To streamline, we use 'forest area' to refer to both island area and CF site area.

Species traits and phylogeny

Functional traits were selected on the basis of their ecological meaning and interpretation within the context of foraging behaviour and resilience of ecosystems (Dreiss et al. 2015). Small mammal traits -- body mass, vertical forest strata, trophic level and degree of matrix tolerance -- were obtained from field measurement, available databases (Paglia et al. 2012; Wilman et al. 2014) and grey literature (Malcolm 1991). Lizard traits -- body length, thermoregulation mode, type of habitat used and range of prey size -- were obtained from Martins (1991) reporting data captured in the same landscape (both within islands and in the surrounding mainland forest), whenever available. We supplemented information on lizard traits by adding data from a comprehensive review of the Brazilian Amazonian lizard fauna (Ávila-Pires 1995). In particular, we did not consider vertical forest strata used by lizards because detection of arboreal species may be underestimated due to the sole use of pitfalls as sampling method. Small mammal vertical forest strata and lizard thermoregulation mode and type of habitat used were the only categorical variables that were further transformed into continuous variables in all subsequent analysis, except in those regarding hypothesis 2 examining different modalities within of each categorical variable. Information on individual traits is detailed in Table S2 (see Tables S3 and S4 for trait values of small mammal and lizard species, respectively).

Phylogenetic trees were obtained using small mammal (Upham et al. 2019) and lizard data (Tonini et al. 2016) available in the VertLife platform (2020). This platform provides a tool that we used to produce distributions of phylogenetic trees containing our recorded species. This tool first trimmed the data to a subset of trees ($N = 100$), and then sampled trees from a chosen pseudo-posterior distribution. For species that were not identified at the species-level, we chose the species with the core of its geographic distribution most centred in our study area, following the distribution maps of IUCN Red List (2020)

Loading [MathJax]/jax/output/CommonHTML/fonts/TeX/fontdata.js *L. guianense*, *N. planiceps*, *K. calcarata*).

Phylogenetic data was unavailable for one lizard species (*Cercosaura oshaughnessyi*, updated to *Prionodactylus oshaughnessyi*), for which we considered a co-generic species (*Cercosaura eigenmanni*). Based on the subset of phylogenetic trees obtained for each taxon, we generated an averaged tree using the 'averageTree' function (see averaged phylogenetic trees in Fig. S1), from 'phytools' R package (Revell 2012). Phylogenetic species distances were obtained from the average tree by computing the cophenetic distances for the hierarchical clustering using the 'cophenetic' function from R (R Development Core Team 2017).

Data analysis

We summed the number of small mammal and lizard records obtained for each sampling site during the two field seasons, and within sampling sites where more than one trapping transect was deployed. Sampling effort sufficiency was previously assessed for each taxon and considered satisfactory, except in the case of lizards at two small islands where only one species was recorded. Despite the lack of sampling representativeness at those two small islands, we still retained those islands in the analyses because trap density therein was much higher (~ 0.75 traps/ha) than in large islands (> 100 ha, $0.01 - 0.06$ traps/ha) and CF sites (~ 0.001 traps/ha; Table S1) (for detailed explanation on the methods used to evaluate sampling sufficiency and the corresponding values, see Palmeirim et al. 2017, 2018). We excluded two singleton records of two small mammal species (*Echimys chrysurus* and *Makalata didelphoides*), which are known to be poorly detected by the traps used in this study. Due to differences in sampling effort between sites, species abundance was previously standardized according to sampling effort applied to each site. All data analyses were performed in R (R Development Core Team 2017).

α -Diversity analysis

Following Ricotta et al. (2016), taxonomic (TD), functional (FD) and phylogenetic (PD) α -diversities were correspondingly given by the Simpson and Rao's quadratic entropy functional and phylogenetic indices, considering relative species abundances. Higher values of Simpson index, ranging between 0 and 1, indicate higher species richness and evenness. Rao's quadratic entropy index takes into account the differences (trait- and phylogenetic-based variance) between species pairs (Botta-Dukát 2005). Both Simpson and Rao's indices are based on the same species pairwise distances and relative species abundances so that Rao's is considered a generalization of Simpson (Botta-Dukát 2005). TD, FD and PD were obtained using 'SYNCSA' R package (Debastiani and Pillar 2012).

To test our first hypothesis, we first checked pairwise correlations between spatial and habitat metrics. When considering all sampled sites (i.e., including CF sites), forest area and the proximity index were highly correlated ($r > 0.80$), further preventing us from including both variables in the same models. We therefore decided to first analyse the effects of only forest area applying Generalized Linear Models (GLMs). This GLM approach is useful for community data because it can be applied to non-normal response variables typically used in community-level modelling (Warton et al. 2015). Species-area relationships in habitat islands typically follow a power law (Matthews et al. 2016). We accounted for this potential non-linear relationship by fitting GLMs both including and excluding the quadratic term of

‘forest area’, and then comparing their Akaike Information Criteria (AIC: Burnham and Anderson 2002). The quadratic term of ‘forest area’ was retained in the models regarding all the dimensions of lizard diversity (Table S5) and included in subsequent analyses considering these response variables. Non-linear relationships could also be accounted for by using Generalised Additive Models, which are a non-parametric extension of GLMs, and do not require a linear relationship between the response and the explanatory variables. We previously additionally performed GAMs using a Gaussian error structure, using the ‘mgcv’ R package (Wood and Wood 2015). The results obtained with GAM, however, were the same as those obtained using GLMs. We therefore decided to retain the simplest statistical approach, GLMs, which further allowed us to account for potential non-linear relationships (Matthews et al. 2016). Given that two of the 20 small mammal and three of the 17 lizard species could only be identified at the genus-level, possibly inflating FD or PD over TD, we ran these models with area and its quadratic term considering response variables without these species (Table S6). As no differences in the results were observed, we ran models including all species in subsequent analysis.

We then analysed the combined effects of spatial and habitat variables – area, prox, shape, dist and s_{tree} – on the α -dimensions of small mammal and lizard diversity using GLMs. This second modelling was restricted to the 25 islands surveyed, thereby excluding CF sites. To control for additional high levels of variable inter-dependence, we performed a Pearson correlation matrix. When excluding CF sites, shape was highly correlated with area ($r > 0.75$), but consistently explained less variance than area (Table S7) and was removed from subsequent analyses. A candidate model set was further constructed, using all combinations of the four explanatory variables retained (including the null model with only intercept and residual variance, without explanatory variables), and models were ranked based on their AIC values corrected for small sample sizes (AICc: Burnham and Anderson 2002), using the ‘MuMIn’ R package (Bartoń 2016). A model-averaging approach was then performed to account for model uncertainty in multi-model inference, using all alternative models (Burnham and Anderson 2002).

Functional trait composition

Patterns of functional composition were analysed individually considering each small mammal and lizard trait and calculating its community-weighted mean value (CWM: Lavorel et al. 2008). CWMs were obtained considering relative species abundance using the ‘functcomp’ function of the R package FD (Laliberté and Legendre 2010). By incorporating species relative abundance, CWM ensures assessments of shifts in mean trait values across the range of spatial and habitat-related variables considered (Lavorel et al. 2008). As before, we first analysed forest area effects on each species trait, for each taxon considering all sampled sites, by applying GLMs using a Gaussian error structure. Again, to account for any potential non-linear relationship, all GLMs were fitted both including and excluding the quadratic term of ‘forest area’, and its AIC was compared (Burnham and Anderson 2002). The quadratic ‘forest area’ term was retained in models regarding lizards ‘body length’ (Table S8) and included in all subsequent analyses considering this response variable. Then, we analysed the combined effects of all four uncorrelated spatial and habitat variables (area, shape, dist and s_{tree}) restricting the analysis to the 25 surveyed

islands. A model-averaging approach was then performed as above, to account for model uncertainty in multi-model inference, using all alternative models (Burnham and Anderson 2002).

Functional species uniqueness

To understand importance of each small mammal and lizard species in supporting a certain function at each sampled site and whether this varied over the range of forest areas, we calculated species-level functional uniqueness (K_i). K_i is given by the mean (functional) distance of a given species from all other species in the assemblage in terms of their functional traits weighted by species abundance (Ricotta et al. 2016). Higher K_i values indicate a greater distinction of a given species from all other species in the assemblage. The K_i calculations were performed with the 'uniqueness' function provided by Ricotta et al. (2016). Then, to test our hypothesis that species uniqueness increases in smaller forest islands, we considered the average of K_i across all species in a given site and related it to forest area by performing a GLM with a Gaussian distribution and retaining the 'forest area' quadratic term whenever this improved model fit (i.e., $\Delta AIC > 2$) (Table S9). To further improve model fitting, we removed from the analyses examining variation in averaged species uniqueness two clear outliers of small mammals (islands 4 and 9) and one lizard (island 8). When related to forest area, these three islands presented much lower K_i . For example, average small mammal K_i ($\pm$ SD) was 0.32 ± 0.11 , while these values for islands 4 and 9 were 0.08 and 0.02, respectively. After removing these outliers, the R^2 increased from 0.02 to 0.40. Likewise, averaged lizards K_i was 0.39 ± 0.06 , island 8 had 0.26 and its removal increased the R^2 -value from 0.02 to 0.18. Thus, these outliers likely corresponded to highly discrepant observations. To investigate whether averaged K_i per site differed significantly between small mammals and lizards, we considered all sites in which both taxa were co-sampled (24 islands and 4 CF sites) and applied a paired t -test. Secondly, to investigate whether individual species uniqueness followed the pattern observed for averaged species uniqueness across the range of forest areas, we analysed individual species K_i across the range of forest areas. To do so, we considered six small mammal and five lizard species occurring in at least 15 sites, to reach sampling sufficiency, and applied a GLM with a Gaussian distribution, and retained the 'forest area' quadratic term whenever this improved model fits (Table S10).

β -Diversity analysis

Our fourth hypothesis, regarding processes underlying community (dis)assembly across the range of forest areas, was examined using β -diversity partitioning into its richness differences (β_{rich}) and replacement (β_{rep}) components (Carvalho et al. 2012, 2013). Calculations of PD and FD were based on Faith (1992) and Petchey and Gaston (2002), respectively, which measure PD and FD of a community as the total branch length of a tree linking all species represented in such community. For each dimension, β -diversity was then partitioned by computing the species incidence data using a sites $\times$ species matrix with the Jaccard dissimilarity index. Functional and phylogenetic pairwise dissimilarity matrices in all analyses were calculated using Gower's distance. We did not use the functional space (instead of functional dendrograms) as this requires the number of species at each site to be higher than the number of traits (Baselga and Orme 2012), and that would preclude us from including smaller islands. We

Loading [MathJax]/jax/output/CommonHTML/fonts/TeX/fontdata.js | beta-diversity whenever another diversity

metric is used (Maire et al. 2015). β -diversity analyses were conducted using the 'beta' function of the R package BAT (Cardoso et al. 2015). Finally, the correlation between differences in forest area and β -diversity and its components were analysed using a Pearson correlation test using the 'cor.test' function from R (R Development Core Team 2017).

Results

Based on 65,520 trap-nights and 5,447 trap-days, we recorded a total of 884 small mammals (20 species: 10 marsupials and 10 rodents) and 1,123 lizards (17 species), respectively. Islands on average ($\pm$ SD) harboured 5.8 ± 3.8 small mammal species (1–16 species) and 5.2 ± 3.2 lizard species (1–11), while more species-rich assemblages were observed in mainland continuous forest sites (small mammals: 5.0 ± 2.5 species, 9–16; lizards: 9.3 ± 1.4 species, 8–11).

α -Diversity

TD, FD and PD of both small mammals and lizards increased significantly ($P < 0.005$) with forest area (Fig. 2; Table S11). Considering the additional spatial (prox and dist) and habitat-related variables (S_{tree}) and restricting the analysis to only 25 surveyed islands, all three dimensions of small mammal diversity were also consistently predicted by changes in forest area (Table S12, S13).

Functional trait composition

Considering all sampling sites, changes in functional trait composition of small mammals and lizards were partly associated with forest area. In particular, larger-bodied small mammal species ($P = 0.010$) exhibiting low matrix tolerance ($P = 0.034$) primarily occupied larger forest areas (Fig. 3a-b; Table S14). Lizard assemblages in large forest areas had larger body length, (area: $P = 0.015$, area²: $P = 0.006$) (Fig. 3c), were mostly heliophobic in terms of thermoregulation ($P = 0.071$, Fig. 3d), and more likely to use riparian habitats (stream banks and waterlogged areas, $P = 0.020$) rather than open-habitats (forest edges and clearings, $P = 0.071$) (Fig. 3e; Table S14). When accounting for the combined effects of spatial and habitat variables across islands only, matrix tolerance of mammals increased on more isolated islands (lower prox, $P = 0.065$) (Tables S15 and S16). Larger-bodied lizard species (longer body length, $P = 0.050$) consuming larger prey ($P = 0.054$) were associated with higher tree diversity, while the positive effect of island area was further reinforced on lizards occupying riparian habitats ($P = 0.036$) (Tables S15 and S16).

Functional uniqueness

Functional uniqueness (K_i) revealed important functional roles of some small mammal and lizard species over the full range of forest areas. For example, *M. murina* was the most functionally unique small mammal, which was consistent across most of the sites where they were detected (Fig. 4a). For lizards, several species presented high functional uniqueness across the spectrum of forest areas (e.g., *Xantusia*, *A. reticulata* and *T. exilis*; Fig. 4b) and their K_i was on average ($\pm$ SD) higher ($0.392 \pm$

0.064) than that of small mammals (0.317 ± 0.111 ; $t = -3.505$, $P = 0.002$). Nevertheless, mean K_i increased towards smaller areas for both small mammals ($P = 0.009$ and, for the quadratic term, $P = 0.058$) and lizards ($P = 0.036$) (Fig. 5; Table S17). When considering the six small mammal and five lizard species occurring in at least 15 survey sites, this trend was confirmed for the marsupial *Philander opossum* ($P = 0.001$) and the lizards *Leposoma* sp. ($P = 0.013$), *Ameiva ameiva* ($P = 0.034$ and, for the quadratic term, $P = 0.017$) and *Plica umbra* ($P = 0.015$) (Table S18).

β-Diversity

Overall, β_{rich} had a greater contribution to beta-diversity than the β_{repl} -component, with the relative contribution of each component varying further according to differences in forest area (Fig. 6). For small mammals, β_{rich} was positively correlated with differences in forest area for all dimensions ($r = 0.58$, $P < 0.001$), while β_{repl} was negatively correlated but only for TD ($r = -0.41$, $P < 0.001$; Fig. 6; Table S19). In fact, for similar-sized areas (i.e., low Δ forest area), taxonomic and functional β_{repl} was, correspondingly, higher and similar than β_{rich} (Fig. 6a, b). For lizards, both taxonomic β_{rich} ($r = 0.10$, $P = 0.041$) and β_{repl} ($r = 0.11$, $P = 0.020$) were negatively correlated with differences in forest area, while there was a positive correlation between both functional β_{rich} ($r = 0.12$, $P = 0.010$) and phylogenetic β_{repl} ($r = 0.19$, $P = 0.0001$) with differences in forest area (Fig. 6; Table S19). Only for contrasting forest areas, β_{rich} and β_{repl} of the phylogenetic dimension were similar (Fig. 6e).

Discussion

Insular habitat fragmentation following river damming rapidly drives local species extinctions (Gibson et al. 2013, Jones et al. 2016). Here we show that such taxonomic loss is followed by severe decay in both functional and phylogenetic diversity for both small mammals and lizards. Species persistence in smaller forest islands depended on traits related to dispersal ability, larger body size and higher matrix tolerance in small mammals; and longer-bodied lizards, particularly those species that were habitat generalists characterized by a heliophile thermoregulation mode and using open-areas (edges and clearings) as preferred habitats. Also, despite the overall higher species uniqueness associated with lizards, species in both taxa became more functionally unique on smaller forest islands. Across the gradient of fragmentation, both assemblages were predominantly shaped by environmental filtering, as noted by the overall prevalence of the richness-component of β -diversity. For small mammals, however, species extinction and colonization across either small or large sites could still occur randomly or due to random processes or environmental heterogeneity that was not considered in this study (higher taxonomic β_{repl} where Δ forest area was particularly low; Fig. 6).

Forest area effects on α -diversity and individual species traits

All three metrics of α -diversity (taxonomic, functional and phylogenetic) in small mammals and lizards Loading [MathJax]/jax/output/CommonHTML/fonts/TeX/fontdata.js isistent with our expectations (Petchey et al.

2007, Meynard et al. 2011). Our results further support that evolutionary time allowed for greater accumulation of trait changes in both small mammals and lizards (Tucker et al., 2018), so that considering either taxonomic, functional or phylogenetic diversity may be sufficient to predict others (Safi et al. 2011, Dreiss et al. 2015). A consistently positive relationship between TD, FD and PD with forest area was similarly observed in sub-tropical reservoir islands for bird (Si et al. 2017) and ant assemblages (Zhao et al. 2020), and for small mammals across the Brazilian Atlantic Forest biome (Bovendorp et al. 2019).

Individual trait responses further suggest that small mammal species performance across the insular fragmented landscape is mainly related to their ability to persist in habitat-degraded small islands, rather than isolated sites. For instance, larger forest areas were more likely to support larger-bodied small mammal species. Community body mass averaged 140 g on the four small islands (< 2 ha), but 401 g at the four continuous forest sites. This is at odds with our initial hypothesis that larger body size confers higher physical endurance in traversing the open-water matrix, which finds support in diverse vertebrate taxa, including small mammals (Cosson et al. 1999). An alternative hypothesis is that larger-bodied rodents and marsupials were selectively extirpated at smaller sites, which were insufficient to sustain their larger home range areas and wider habitat requirements (Lindsted et al. 1986). Also, in a study carried out on nearly the same set of islands, two of the three larger-bodied rodent species included in mid-large mammal surveys (*Myoprocta acouchy* and *Cuniculus paca*) were absent on the smallest islands (Benchimol and Peres 2015c). This reinforces our findings that the high-end of the 'small-mammal' size spectrum is selectively extirpated on small islands. This finding is particularly interesting given the relatively limited body mass range of these species (10–1,300 g), which corresponds to two orders of magnitude. The same body mass range is observed between the smallest (acouchi, *Myoprocta* spp.) and the largest species (tapir, *Tapirus* spp.) that are typically surveyed in neotropical mid-large mammal assemblages (Paglia et al. 2012). Therefore, our results are in agreement with the size-selective defaunation gradient observed at larger scales elsewhere (e.g. Canale et al. 2012), which has direct impacts to key ecological interactions (e.g., seed dispersal and insectivory) and ecosystem functioning (Dirzo et al. 2014). A similar but less marked trend was observed for lizards. Perhaps, as the home ranges of these heterotherm species are much smaller than those of mammals for any given body mass (Tamburello et al. 2015), small islands could still support a wide range of lizard body sizes (see Fig. 3d).

Matrix tolerance of small mammals further increased towards smaller forest sites indicating that species that can use non-forest matrix areas are able to thrive on smaller islands, which are further characterized by lower tree species richness, and vice-versa for forest-dependent species. Small mammal local extinctions in small forest patches are offset by the incursion of open-habitat species in southern Amazonia (Palmeirim et al. 2020) and the Atlantic Forest (Estavillo et al. 2013). Therefore, matrix tolerance has the potential to predict changes in small mammal communities following habitat loss and fragmentation.

Responses of individual lizard traits reinforced our functional α -diversity findings and corroborate a at-degraded small islands were comprised of

habitat and trophic generalists (see Palmeirim et al. 2017). As expected, species traits associated with persistence were related to greater tolerance to a wider spectrum of environmental conditions, which reflects their heliothermic mode and preference for more desiccated forest edges and clearings, in addition to their ability to disperse across the archipelagic landscape, as expected by their larger body length (Schoener and Schoener 1984). Moreover, edge effects are stronger and canopy gaps are proportionally more frequent on smaller islands (Benchimol and Peres 2015b). In these extreme habitat conditions, only large-bodied heliothermic lizards, which frequently venture into open-area habitats, are able to persist (Lima et al. 2001, Silva et al. 2014). Elsewhere in the tropics, lizard functional diversity declines under habitat disturbance (Berriozabal-Islas et al. 2017), but different species diverge in their responses to altered microclimatic conditions (Lehtinen et al. 2003, Gardner et al. 2007, Suazo-Ortuño et al. 2018).

Functional uniqueness

While certain traits alone may not confer functional uniqueness, their combination can be functionally unique particularly at smaller forest sites. For example, the scansorial marsupial *P. opossum* exhibits high degree of matrix tolerance and is particularly functional unique at small forest sites where fewer species persist. Likewise, the lizard *P. umbra* behaves as a heliophobe but has a long body length. As forest area decreases, these species play a key role by shaping the functional diversity of these communities (Ricotta et al. 2016). These results are consistent with increases in community-level functional redundancy in increasingly larger forest areas (e.g., Bovendorp et al. 2019, Farneda et al. 2020). Also, this was particularly interesting for small mammal species that were, on average, less functionally unique than lizards, and thus more functionally redundant at larger forest sites. As forest area is reduced, many small mammal species are gradually extirpated, but their ecological functions may be similarly performed by some of the remaining species (Safi et al. 2011). Functional redundancy has been widely documented in tropical mammals (Safi et al. 2011, Oliveira et al. 2016, Bovendorp et al. 2019), bats (Farneda et al. 2020), and birds (de Coster et al. 2015). In an assemblage in which species share similar traits, if one species goes extinct, it is likely that the remaining species are still able to perform the same functions (Rosenfeld 2002), which is apparently the case of small mammals, particularly at larger forest sites in Balbina.

Species (dis)assembly processes

Taxonomic, functional and phylogenetic β -diversity of both taxa were mostly partitioned into their β_{rich} -component, suggesting that environmental filtering has shaped small mammal and lizard communities in the insular Balbina landscape. Similar findings were observed for taxonomic and functional diversity of birds and lizards in land-bridge islands in China (Si et al. 2015), and bats in primary forest patches embedded in second-growth in the Brazilian Amazon (Farneda et al. 2018). Other studies considering phylogenetic diversity reported the stronger role of deterministic processes in structuring tree (Yang et al. 2015) and bat assemblages across different ecoregions (Aguirre et al. 2016). In our study, deterministic processes shaping small mammal and lizard assemblages, mediated by forest area, are additionally supported by findings on individual trait responses and functional uniqueness of small mammal and

Given the comparison between similar-sized sites (i.e., small-small and large-large), lizards mostly responded to the environmental filter. This implies that even small differences in forest area are important in structuring lizard assemblages. Small mammals, however, did not consistently respond to modest variation in forest area, for which taxonomic β_{repl} values, which were lower than those of β_{rich} , either indicate stochasticity in colonization-extinction rates or the effects of environment heterogeneity not fully accounted for in this study. For example, habitat quality on small islands is associated with the intensity of edge effects which is further related to their location within the reservoir and previous fire history (Benchimol and Peres 2015b). The higher functional redundancy of small mammals compared to lizards also supports the notion that small mammals are generally less affected by environmental gradients. Moreover, although stochasticity can promote limiting similarity (MacArthur and Levins 1967; Tilman 2004), the potential stochasticity associated with taxonomic diversity of small mammals across similar-sized sites did not necessarily result in co-existence of functionally dissimilar species. Instead, both small mammals and lizards played similar functional roles and shared similar phylogeny at similar-sized sites, but while the subset of lizard species persisting was the same, that of small mammals varied from site to site.

Conservation implications

Given the surrogacy in the high predictability of TD, FD and PD by forest area, particularly at the local (α) scale, any of these three dimensions of diversity can be used to assess the co-effects of habitat loss and fragmentation (Safi et al. 2011, Dreiss et al. 2015). Yet different aspects of functional diversity provide additional insights. In particular, the use of individual functional trait allows more general predictive ability of which species are likely to go extinct in the aftermath of insular fragmentation, and their consequences for ecosystem functioning. Also, as highlighted by patterns of functional uniqueness, the cost associated with individual species extinctions on ecosystem functioning escalated with greater severity of insular fragmentation. Nevertheless, given the inherent taxon-specific differences in responses, we urge caution when considering a single taxon as a surrogate (Wang et al. 2010). For instance, given the higher small mammal functional redundancy at larger forest sites, their persistence therein is particularly important to ensure ecosystem resilience to perturbation (Díaz et al. 2013). In addition, small mammal species composition fluctuates from site to site, even if their sizes are similar, so the conservation importance of small to mid-sized islands is high for small mammals, but much less so for lizards.

Over and above the taxon-specific differences above, our study demonstrates that small, habitat-degraded islands, harbour reduced taxonomic, functional and phylogenetic diversity of both small mammals and lizards compared to large islands and continuous forests sites. To preserve the multiple dimensions of vertebrate diversity, and thus maintain the health of the ecosystem, setting-aside large forest tracks should be imperative. This is especially relevant to mega-dams in lowland tropical forests, which typically involve inundation over extensive areas (Fearnside 2013). In Balbina in particular, 90% of all 3,546 islands are relatively small (< 100 ha). Avoiding the creation of myriad small islands should be part of the main guidelines towards sustainable hydropower development. Properly incorporating such

guidelines within country-level legislation would be a major policy challenge to preclude regional scale biodiversity collapse and its associated ecosystem services.

Declarations

Acknowledgements

We thank all 15 volunteers/field assistants who assisted in the field, M. Benchimol and D. Storck-Tonon for their assistance with landscape metrics, the Reserva Biológica do Uatumã and its staff for logistical support, and R. de Fraga and M.N.F. da Silva for their help with species identification. This study was funded by the Amazon Region Protected Areas Program (ARPA), Amazonas Distribuidora de Energia S.A., and Associação Comunidade Waimiri Atroari; Rufford Foundation (grant number 13675-1); Idea Wild; and a NERC grant (NE/J01401X/1) awarded to CAP. AFP was funded by European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No. 792678. FZF was funded by Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) scholarship, and MVV was by Fundação de Amparo à Pesquisa do Estado do Rio de Janeiro (FAPERJ, grant E-203.045/2017) and Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq, grants 426.925/2018-1, 312465/2019-0).

Conflicts of interest/Competing interests

The authors declare no conflict of interest nor competing interest.

Ethics approval

Whenever live captures could not be identified in the field, a maximum of five voucher specimens per species per site were collected during the first season and deposited at the Mammal and Herpetofauna Collections of the Instituto Nacional de Pesquisas da Amazônia (INPA), in Manaus, Brazil. This study was approved by the appropriate Brazilian government agency (SISBIO license No. 39187-4).

Consent to participate

All authors agree in participating in this manuscript.

Consent for publication

All authors agree with the publication of this manuscript.

Availability of data and material

Data available from previous publications: <https://doi.org/10.1007/s00442-018-4114-6> (Palmeirim et al. 2018), for small mammal data, and <https://doi.org/10.1016/j.biocon.2017.08.002> (Palmeirim et al. 2017), for lizard data.

Code availability

The code used to run the analysis of this study is available upon request.

Authors' contributions

AFP and FZF conceived the ideas; AFP, CAP and MVV defined the sampling design; AFP collected the data; AFP and FZF analysed the data; AFP led the writing of the manuscript. All authors contributed critically to the drafts and gave final approval for publication.

References

- Aguirre LF, Montaña-Centellas FA, Gavilanez MM, Stevens RD (2016) Taxonomic and phylogenetic determinants of functional composition of Bolivian bat assemblages. *PloS One* 11:e0158170. <https://doi.org/10.1371/journal.pone.0158170>
- Ávila-Pires TC (1995) Lizards of brazilian amazonia (Reptilia: Squamata). *Zoologische verhandelungen* 299:1-706
- Bartoń K (2016) MuMIn: Multi-Model inference. R package version 1.15.6.
- Baselga A, Orme CDL (2012) betapart: an R package for the study of beta diversity. *Methods Ecol Evol* 3:808-812. <https://doi.org/10.1111/j.2041-210X.2012.00224.x>
- Beaupre SJ, Jacobson ER, Lillywhite HB, Zamudio K (2004) Guidelines for use of live amphibians and reptiles in field and laboratory research. American Society of Ichthyologists and Herpetologists, USA
- Benchimol M, Peres CA (2015a) Widespread forest vertebrate extinctions induced by a mega hydroelectric dam in lowland Amazonia. *PloS One* 10:e0129818. <https://doi.org/10.1371/journal.pone.0129818>
- Benchimol M, Peres CA (2015b) Edge-mediated compositional and functional decay of tree assemblages in Amazonian forest islands after 26 years of isolation. *J Ecol* 103:408-420. <https://doi.org/10.1111/1365-2745.12371>
- Benchimol M, Peres CA (2015c) Predicting local extinctions of Amazonian vertebrates in forest islands created by a mega dam. *Biol Conserv* 187:61-72. <https://doi.org/10.1016/j.biocon.2015.04.005>
- Berriozabal-Islas C, Badillo-Saldaña LM, Ramírez-Bautista A, Moreno CE (2017) Effects of habitat disturbance on lizard functional diversity in a tropical dry forest of the Pacific Coast of Mexico. *Trop Conserv Sci* 10:1940082917704972. <https://doi.org/10.1177/1940082917704972>
- Botta-Dukát Z (2005) Rao's quadratic entropy as a measure of functional diversity based on multiple traits. *J Veg Sci* 16:533-540. <https://doi.org/10.1111/j.1654-1103.2005.tb02393.x>

Bovendorp RS, Brum FT, McCleery RA, Baiser B, Loyola R, Cianciaruso MV, Galetti M (2019) Defaunation and fragmentation erode small mammal diversity dimensions in tropical forests. *Ecography* 42:23-35. <https://doi.org/10.1111/ecog.03504>

Burnham KP, Anderson DR (2002) A practical information-theoretic approach. Model selection and multimodel inference, 2nd ed. Springer, New York.

Canale GR, Peres CA, Guidorizzi CE, Gatto CAF, Kierulff MCM (2012) Pervasive defaunation of forest remnants in a tropical biodiversity hotspot. *PLoS One* 7:e41671. <https://doi.org/10.1371/journal.pone.0041671>

Cardoso P, Rigal F, Carvalho JC (2015) BAT–Biodiversity Assessment Tools, an R package for the measurement and estimation of alpha and beta taxon, phylogenetic and functional diversity. *Methods Ecol Evol* 6:232-236. <https://doi.org/10.1111/2041-210X.12310>

Cardoso P, Rigal F, Carvalho JC, Fortelius M, Borges PAV, Podani J, Schmera D (2014) Partitioning taxon, phylogenetic and functional beta diversity into replacement and richness difference components. *J Biogeogr* 41:749-761. <https://doi.org/10.1111/jbi.12239>

Carvalho JC, Cardoso P, Gomes P (2012) Determining the relative roles of species replacement and species richness differences in generating beta-diversity patterns. *Glob Ecol Biogeogr* 21:760-771. <https://doi.org/10.1111/j.1466-8238.2011.00694.x>

CEUA-CCS UFRJ (2010) Comissão de Ética no Uso de Animais em Pesquisas da Universidade Federal do Rio de Janeiro. <https://www.ccs.ufrj.br/conteudos/ceua> Accessed 1 November 2019

Cosson JF, Ringuet S, Claessens O et al (1999) Ecological changes in recent land-bridge islands in French Guiana, with emphasis on vertebrate communities. *Biol Conserv* 91:213-222. [https://doi.org/10.1016/S0006-3207\(99\)00091-9](https://doi.org/10.1016/S0006-3207(99)00091-9)

Dainese M, Lepš J, de Bello F (2015) Different effects of elevation, habitat fragmentation and grazing management on the functional, phylogenetic and taxonomic structure of mountain grasslands. *Perspect. Plant Ecol Evol Syst* 17:44-53. <https://doi.org/10.1016/j.ppees.2014.09.002>

De Coster G, Banks-Leite C, Metzger JP (2015) Atlantic forest bird communities provide different but not fewer functions after habitat loss. *Proc. Royal Soc. B* 282:20142844. <https://doi.org/10.1098/rspb.2014.2844>

Debastiani VJ, Pillar VD (2012) SYNC-SA—R tool for analysis of metacommunities based on functional traits and phylogeny of the community components. *Bioinformatics* 28:2067-2068. <https://doi.org/10.1093/bioinformatics/bts325>

Devictor V, Julliard R, Jiguet F (2008) Distribution of specialist and generalist species along spatial Loading [MathJax]/jax/output/CommonHTML/fonts/TeX/fontdata.js s 117:507-514. <https://doi.org/10.1111/j.0030->

Díaz S, Purvis A, Cornelissen JH, et al (2013) Functional traits, the phylogeny of function, and ecosystem service vulnerability. *Ecol Evol* 3:2958-2975. <https://doi.org/10.1002/ece3.601>

Dirzo R, Young HS, Galetti M, Ceballos G, Isaac NJ, Collen B (2014) Defaunation in the Anthropocene. *Science* 345:401-406. <https://doi.org/10.1126/science.1251817>

Elmqvist T, Folke C, Nyström M, Peterson G, Bengtsson J, Walker B, Norberg J (2003) Response diversity, ecosystem change, and resilience. *Front Ecol Environ* 1:488-494. [https://doi.org/10.1890/1540-9295\(2003\)001\[0488:RDECAR\]2.0.CO;2](https://doi.org/10.1890/1540-9295(2003)001[0488:RDECAR]2.0.CO;2)

ESRI (2018) ArcMap 10.1. Environmental Systems Research Institute Inc., Redlands, CA, USA.

Estavillo C, Pardini R, da Rocha PLB (2013) Forest loss and the biodiversity threshold: an evaluation considering species habitat requirements and the use of matrix habitats. *PLoS One* 8:e82369. <https://doi.org/10.1371/journal.pone.0082369>

Farneda FZ, Rocha R, López-Baucells A, Groenenberg M, Silva I, Palmeirim JM, Bobrowiec PED, Meyer CFJ (2015) Trait-related responses to habitat fragmentation in Amazonian bats. *J Appl Ecol* 52:1381-1391. <https://doi.org/10.1111/1365-2664.12490>

Farneda FZ, Rocha R, López-Baucells A, et al (2018) Functional recovery of Amazonian bat assemblages following secondary forest succession. *Biol Conserv* 218:192-199. <https://doi.org/10.1016/j.biocon.2017.12.036>

Farneda FZ, Grelle CE, Rocha R, Ferreira DF, López-Baucells A, Meyer CF (2020) Predicting biodiversity loss in island and countryside ecosystems through the lens of taxonomic and functional biogeography. *Ecography* 43:97-106. <https://doi.org/10.1111/ecog.04507>

Fearnside PM (2013) Hidrelétricas na Amazônia: Impactos e tomada de decisão. pp. 43-60; 87-91. Val AL, Santos GM (eds.). Grupo de Estudos Estratégicos Amazônicos (GEEA), Tomo VI. Instituto Nacional de pesquisas da Amazônia (INPA), Manaus, Amazonas

FUNCATE/INPE/ANEEL (2000) Mapeamento por satélite das áreas inundadas por reservatórios de hidrelétricas brasileiras. Unpublished Report. Convênio FUNCATE/INPE/ANEEL, São Paulo

Gardner TA, Ribeiro-Junior MA, Barlow JOS, Ávila-Pires TCS, Hoogmoed MS, Peres CA (2007) The value of primary, secondary, and plantation forests for a neotropical herpetofauna. *Conserv Biol* 21:775-787. <https://doi.org/10.1111/j.1523-1739.2007.00659.x>

Gibson L, Lynam AJ, Bradshaw CJ et al (2013) Near-complete extinction of native small mammal fauna 25 years after forest fragmentation. *Science* 341:1508-1510. <https://doi.org/10.1126/science.1240495>

Graham CH, Fine PV (2008) Phylogenetic beta diversity: linking ecological and evolutionary processes across space in time. *Ecol Lett* 11:1265-1277. <https://doi.org/10.1111/j.1461-0248.2008.01256.x>

Guisan A, Edwards Jr TC, Hastie T (2002) Generalized linear and generalized additive models in studies of species distributions: setting the scene. *Ecol Model* 157:89-100. [https://doi.org/10.1016/S0304-3800\(02\)00204-1](https://doi.org/10.1016/S0304-3800(02)00204-1)

Haddad NM, Brudvig LA, Clobert J et al (2015) Habitat fragmentation and its lasting impact on Earth's ecosystems. *Sci Adv* 1:e1500052. <https://doi.org/10.1126/sciadv.1500052>

IBAMA 1997. Plano de Manejo Fase I: Reserva Biológica do Uatumã. Eletronorte/IBAMA. Brasília/DF/Brazil. <http://www.icmbio.gov.br>. Accessed 1 January 2019

IUCN 2020. The IUCN Red List of Threatened Species. Version 2020-2. <https://www.iucnredlist.org>. Accessed 1 June 2020

Jones IL, Bunnefeld N, Jump AS, Peres CA, Dent DH (2016) Extinction debt on reservoir land-bridge islands. *Biol Conserv* 199:75-83. <https://doi.org/10.1016/j.biocon.2016.04.036>

Jones KE, Bielby J, Cardillo M, et al (2009) PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals: Ecological Archives E090-184. *Ecology* 90:2648-2648. <https://doi.org/10.1890/08-1494.1>

Laliberté E, Legendre P (2010) A distance-based framework for measuring functional diversity from multiple traits. *Ecology* 91:299-305. <https://doi.org/10.1890/08-2244.1>

Lambert TD, Malcolm JR, Zimmerman BL (2006) Amazonian small mammal abundances in relation to habitat structure and resource abundance. *J Mammal* 87:766-776. <https://doi.org/10.1644/05-MAMM-A-261R1.1>

Lavorel S, Grigulis K, McIntyre S, et al (2008) Assessing functional diversity in the field—methodology matters! *Funct Ecol* 22:134-147. <https://doi.org/10.1111/j.1365-2435.2007.01339.x>

Lehtinen RM, Ramanamanjato JB, Raveloarison JG (2003) Edge effects and extinction proneness in a herpetofauna from Madagascar. *Biodivers Conserv* 12:1357-1370. <https://doi.org/10.1023/A:1023673301850>

Lima A, Suárez F, Higuchi I (2001) The effects of selective logging on the lizards *Kentropyx calcarata*, *Ameiva ameiva* and *Mabuya nigropunctata*. *Amphibia-Reptilia* 22:209-216. <https://doi.org/10.1163/15685380152030436>

Lindstedt SL, Miller BJ, Buskirk SW (1986) Home range, time, and body size in mammals. *Ecology* 67:413-418. <https://doi.org/10.2307/1938584>

MacArthur RH, Wilson EO (1967) The theory of island biogeography. Press Princeton, USA

MacArthur RH, Levins R (1967) The limiting similarity, convergence, and divergence of coexisting species. *Am Nat* 101:377-385

Maire E, Grenouillet G, Brosse S, Villéger S (2015) How many dimensions are needed to accurately assess functional diversity? A pragmatic approach for assessing the quality of functional spaces. *Glob Ecol Biogeogr* 24:728-740. <https://doi.org/10.1111/geb.12299>

Malcolm JR (1991) The small mammals of Amazonian forest fragments: patterns and process. PhD Thesis. University of Florida

Martins M (1991) The lizards of Balbina, Central Amazonia, Brazil: a qualitative analysis of resource utilization. *Stud Neotrop Fauna Environ* 26:179-190. <https://doi.org/10.1080/01650529109360851>

Matthews TJ, Guilhaumon F, Triantis KA, Borregaard MK, Whittaker RJ (2016) On the form of species–area relationships in habitat islands and true islands. *Glob Ecol Biogeogr* 25:847-858. <https://doi.org/10.1111/geb.12269>

McGarigal K, Cushman SA, Ene E (2012) FRAGSTATS v4: Spatial Pattern Analysis Program for Categorical and Continuous Maps. <https://www.umass.edu/landeco/research/fragstats/fragstats.html>. Accessed 01 Jan 2015

Meynard CN, Devictor V, Mouillot D, Thuiller W, Jiguet F, Mouquet N (2011) Beyond taxonomic diversity patterns: how do α , β and γ components of bird functional and phylogenetic diversity respond to environmental gradients across France? *Glob Ecol Biogeogr* 20:893-903. <https://doi.org/10.1111/j.1466-8238.2010.00647.x>

Oliveira BF, Machac A, Costa GC, Brooks TM, Davidson AD, Rondinini C, Graham CH (2016) Species and functional diversity accumulate differently in mammals. *Glob Ecol Biogeogr* 25:1119-1130. <https://doi.org/10.1111/geb.12471>

Paglia AP, Fonseca GAB, Rylands AB et al (2012) Lista Anotada dos Mamíferos do Brasil. *Occasional Papers in Conservation Biology* 6. Conservation International, Arlington, 2nd ed 76p.

Palmeirim AF, Santos-Filho M, Peres AC (2020) Marked decline in forest-dependent small mammals following habitat loss and fragmentation in an Amazonian deforestation frontier. *PLoS One* 15:e0230209. <https://doi.org/10.1371/journal.pone.0230209>

Palmeirim AF, Benchimol M, Vieira MV, Peres CA (2018) Small mammal responses to Amazonian forest islands are modulated by their forest dependence. *Oecologia* 187:191-204. <https://doi.org/10.1007/s00442-018-4114-6>

- Palmeirim AF, Vieira MV, Peres CA (2017) Non-random lizard extinctions in land-bridge Amazonian forest islands after 28 years of isolation. *Biol Conserv* 214:55-65. <https://doi.org/10.1016/j.biocon.2017.08.002>
- Petchey OL, Evans KL, Fishburn IS, Gaston KJ (2007) Low functional diversity and no redundancy in British avian assemblages. *J Anim Ecol* 76:97-108. <https://doi.org/10.1111/j.1365-2656.2007.01271.x>
- R Development Core Team (2017) R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org/>.
- Revell LJ (2012) phytools: an R package for phylogenetic comparative biology (and other things). *Methods Ecol Evol* 3:217-223. <https://doi.org/10.1111/j.2041-210X.2011.00169.x>
- Ricotta C, Burrascano S (2009) Testing for differences in beta diversity with asymmetric dissimilarities. *Ecol Indic* 9:719-724. <https://doi.org/10.1016/j.ecolind.2008.09.003>
- Ricotta C, de Bello F, Moretti M, Caccianiga M, Cerabolini BE, Pavoine S (2016) Measuring the functional redundancy of biological communities: a quantitative guide. *Methods Ecol Evol* 7:1386-1395. <https://doi.org/10.1111/2041-210X.12604>
- Rosenfeld JS (2002) Functional redundancy in ecology and conservation. *Oikos* 98:156-162. <https://doi.org/10.1034/j.1600-0706.2002.980116.x>
- Safi K, Cianciaruso MV, Loyola RD, Brito D, Armour-Marshall K, Diniz-Filho JAF (2011) Understanding global patterns of mammalian functional and phylogenetic diversity. *Phil Trans R Soc B* 366:2536-2544. <https://doi.org/10.1098/rstb.2011.0024>
- Sala OE, Chapin FS, Armesto JJ et al (2000) Global biodiversity scenarios for the year 2100. *Science* 287:1770-1774. <https://doi.org/10.1126/science.287.5459.1770>
- Salgado-Luarte C, Escobedo VM, Stotz GC, Rios RS, Arancio G, Gianoli E (2019) Goat grazing reduces diversity and leads to functional, taxonomic, and phylogenetic homogenization in an arid shrubland. *Land Degrad Dev* 30:178-189. <https://doi.org/10.1002/ldr.3208>
- Schoener A, Schoener TW (1984) Experiments on dispersal: short-term floatation of insular anoles, with a review of similar abilities in other terrestrial animals. *Oecologia* 63:289-294
- Si X, Baselga A, Ding P (2015) Revealing beta-diversity patterns of breeding bird and lizard communities on inundated land-bridge islands by separating the turnover and nestedness components. *PLoS One* 10:e0127692. <https://doi.org/10.1371/journal.pone.0127692>
- Si X, Cadotte MW, Zeng D et al (2017) Functional and phylogenetic structure of island bird communities. *J Anim Ecol* 86:532-542. <https://doi.org/10.1111/1365-2656.12650>

- Sikes RS (2016) 2016 Guidelines of the American Society of Mammalogists for the use of wild mammals in research and education. *J Mammal* 97:663-688. <https://doi.org/10.1093/jmammal/gyw078>
- Silva DJ, Santos-Filho M, Canale GR (2014) The importance of remnant native vegetation of Amazonian submontane forest for the conservation of lizards. *Braz. J Biol* 74:523-528. <https://doi.org/10.1590/bjb.2014.0095>
- Suazo-Ortuño I, Benítez-Malvido J, Marroquín-Páramo J, Soto Y, Siliceo H, Alvarado-Díaz J (2018) Resilience and vulnerability of herpetofaunal functional groups to natural and human disturbances in a tropical dry forest. *Forest Ecol Manag* 426:145-157. <https://doi.org/10.1016/j.foreco.2017.09.041>
- Tamburello N, Côté I, Dulvy NK (2015). Energy and the scaling of animal space use. *The American Naturalist* 186:196-211. <https://doi.org/10.1086/682070>
- Terborgh J, Lopez L, Nuñez P et al (2001) Ecological meltdown in predator-free forest fragments. *Science* 294:1923-1926. <https://doi.org/10.1126/science.1064397>
- Tilman D (2004) Niche tradeoffs, neutrality, and community structure: a stochastic theory of resource competition, invasion, and community assembly. *Proc Natl Acad Sci* 101:10854-10861. <https://doi.org/10.1073/pnas.0403458101>
- Tonini JFR, Beard KH, Ferreira RB, Jetz W, Pyron RA (2016) Fully-sampled phylogenies of squamates reveal evolutionary patterns in threat status. *Biol Conserv* 204:23-31. <https://doi.org/10.1016/j.biocon.2016.03.039>
- Tucker CM, Davies TJ, Cadotte MW, Pearse WD (2018) On the relationship between phylogenetic diversity and trait diversity. *Ecology* 99:1473-1479. <https://doi.org/10.1002/ecy.2349>
- Upham NS, Esselstyn JA, Jetz W (2019) Inferring the mammal tree: Species-level sets of phylogenies for questions in ecology, evolution, and conservation. *PLoS Biol* 17:e3000494. <https://doi.org/10.1371/journal.pbio.3000494>
- Veron S, Davies TJ, Cadotte MW, Clergeau P, Pavoine S (2017) Predicting loss of evolutionary history: Where are we? *Biol Rev* 92:271-291. <https://doi.org/10.1111/brv.12228>
- VertLife (2020) VertLife: Phylogeny subsets. <https://vertlife.org/phylosubsets/>. Accessed 1 June 2020
- Villéger S, Grenouillet G, Brosse S (2013) Decomposing functional β -diversity reveals that low functional β -diversity is driven by low functional turnover in European fish assemblages. *Glob Ecol Biogeogr* 22:671-681. <https://doi.org/10.1111/geb.12021>
- Violle C, Navas ML, Vile D, Kazakou E, Fortunel C, Hummel I, Garnier E (2007) Let the concept of trait be functional! *Oikos* 116:882-892. <https://doi.org/10.1111/j.0030-1299.2007.15559.x>

- Vitt LJ, Caldwell JP (2014) *Herpetology: an introductory biology of amphibians and reptiles*. Elsevier, Norman
- Warton DI, Blanchet FG, O'Hara RB, Ovaskainen O, Taskinen S, Walker SC, Hui FKC (2015) So many variables: joint modeling in community ecology. *Trends Ecol Evol* 30:766–779
- Wang Y, Bao Y, Yu M, Xu G, Ding P (2010) Nestedness for different reasons: the distributions of birds, lizards and small mammals on islands of an inundated lake. *Divers Distrib* 16:862-873
- Webb CO, Ackerly DD, McPeck MA, Donoghue MJ (2002) Phylogenies and community ecology. *Annu Rev Ecol Evol Syst* 33:475-505
- Weiss KC, Ray CA (2019) Unifying functional trait approaches to understand the assemblage of ecological communities: synthesizing taxonomic divides. *Ecography* 42:2012-2020
- Whittaker RH (1972) Evolution and measurement of species diversity. *Taxon* 21:213-251
- Wilman H, Belmaker J, Simpson J, de la Rosa C, Rivadeneira MM, Jetz W (2014) EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals: *Ecological Archives* E095-178. *Ecology* 95:2027-2027
- Wood S, Wood MS (2015) Package 'mgcv'. R package version, 1, 29
- Yang J, Swenson NG, Zhang G et al (2015) Local-scale partitioning of functional and phylogenetic beta diversity in a tropical tree assemblage. *Sci Rep* 5:12731
- Zhao Y, Dunn RR, Zhou H, Si X, Ding P (2020) Island area, not isolation, drives taxonomic, phylogenetic and functional diversity of ants on land-bridge islands. *J Biogeogr* 47:1627-1637

Figures

Figure 1

(a) Sites where small mammals and lizards were surveyed within the Balbina Hydroelectric Reservoir of Central Brazilian Amazonia: 25 islands (in red and highlighted by a buffer if relatively small) and four continuous forest regions in the mainland (CF1, CF2, CF3 and CF4; indicated by red rectangles); asterisks indicate islands where only small mammals (*) or lizards (**) were sampled. (b) Aerial view of the Balbina archipelagic landscape (photo credit: E. Venticinque) including the aquatic matrix; (c) Sampling design of each transect: an array of three pitfall-traps followed by nine live-trap stations (i.e., Sherman and wire-mesh traps), each of which including three traps deployed on the forest floor, understory and (sub)canopy; distances between traps are indicated in the figure.

Figure 2

Relationships between taxonomic (a, d), functional (b, e) and phylogenetic diversity (c, f) with forest area – islands ($\log_{10} x$) and continuous forest sites (CF) – for small mammals (upper panels) and lizards (lower panels) at the Balbina reservoir landscape. Taxonomic diversity is given by the Simpson index and functional and phylogenetic diversities are given by the Rao Quadratic entropy indices. Lines are the model adjusted for the strongest relationships ($P \leq 0.05$), and shaded areas represent the 95% confidence region. Boxplots for CF sites indicate the median, 1st and 3rd quartiles, and minimum and maximum values of species richness and abundance.

Figure 3

Relationship between community-weighted mean (CWM) trait values and forest area – islands ($\log_{10} x$) and continuous forest sites (CF) – for small mammals and lizards in the Balbina reservoir landscape. Results are shown for statistically significant functional traits. Small mammal traits include (a) body mass and (b) matrix tolerance; lizard traits include (c) body length, (d) heliothermic mode (categories: heliophobe and heliophile) and (e) habitat type (categories: creeks/swamps, edges/clearings, and terra firme forest). Lines represent the model adjusted for the strongest relationships (approx. $P \leq 0.05$), and shaded areas represent the 95% confidence region. Results for all traits are provided in Table S14.

Figure 4

Species-level functional uniqueness for small mammals (a) and lizards (b) at 29 and 30 sampled sites, respectively, in the Balbina Reservoir landscape. Values of functional uniqueness are indicated by the colour of each dot representing the presence of a certain species at a certain site. Islands are ordered in ascending size; CF indicates continuous forest sites. Sites where only one species was recorded were excluded from the analysis (lizards: $N = 4$).

Figure 5

Relationship between mean species-level functional uniqueness and forest area ($\log_{10} x$) for small mammals (a) and lizards (b) in the Balbina landscape. Sites where only one species was recorded were

excluded from the analysis (lizards: N = 4). Lines represent the model adjusted for the stronger relationships (approx. $P \leq 0.05$), and shaded areas represent the 95% confidence region.

Figure 6

Relationship between each of the two components of taxonomic (a, d), functional (b, e) and phylogenetic β -diversity (c, f) – richness differences (β_{rich}) and replacement (β_{repl}) – and differences in forest area ($\log_{10} x$) for all pairwise combinations of sites. We considered 29 and 30 sites where small mammals (upper panels) and lizards (lower panels) were surveyed, respectively, across the Balbina Reservoir landscape. Circles are sized according to the area of the smaller site within each site pair. Trend lines are indicated for all relationships, and shaded areas represent the 95% confidence region.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [supp.mat..docx](#)