Association of ABO Polymorphisms and Cancer/Cardiocerebrovascular Disease: a Meta-analysis

Yanxia Li

Tianjin Medical University Cancer Institute and Hospital

Luyang Liu

Tianjin Medical University Cancer Institute and Hospital

Yubei Huang

Tianjin Medical University Cancer Institute and Hospital

Hong Zheng

Tianjin Medical University Cancer Institute and Hospital

Lian Li (✉ lilian@tmu.edu.cn)

Tianjin Medical University Cancer Institute and Hospital <https://orcid.org/0000-0001-6963-558X>

Research article

Keywords: ABO gene polymorphism, rs505922, rs657152, Cardiocerebrovascular diseases, Cancer, Meta-analysis

Posted Date: September 12th, 2019

DOI: <https://doi.org/10.21203/rs.2.14412/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: ABO gene polymorphisms have been reported to be associated with the risk of multiple cancers and cardiocerebrovascular diseases. However, the results remained controversial. In this study, we conducted a systematic review and meta-analysis to clarify the association between two SNPs (rs505922 and rs657152) in ABO gene and cancers/cardiocerebrovascular diseases.

Method: All eligible case-control studies come from PubMed, Embase and Web of Science up to Jan. 1, 2019. Pooled odds ratios (ORs) with 95% confidence intervals (CIs) were used to assess the corresponding associations. Sensitivity analysis, publication bias assessment, and heterogeneity test were performed using STATA 12.0.

Results: A total of eighteen articles involving twenty-nine case-control populations were included according to inclusion and exclusion criteria. Eleven populations (16,929 cases and 23,941 controls) were used to evaluate the relationship between rs505922 and overall cancers and nine populations (22,275 cases and 71,549 controls) were included to assess the association between rs505922 and cardiocerebrovascular diseases. The results showed a significant association between the rs505922 polymorphism and cancers (CvsT: OR=1.13, 95% CI=1.04-1.22, P=0.003), and cardiocerebrovascular diseases (OR=1.36, 95%CI=1.19-1.57, P<0.001). Four populations (5,158 cases and 7,021 controls) were included to evaluate association between rs657152 and cancers and five populations (8,105 cases and 6,712 controls) were included to estimate the relationship between rs657152 and cardiocerebrovascular diseases. The result of meta-analysis reveals that rs657152 was significantly associated with cancers (OR=1.16, 95%CI=1.09-1.24, P<0.001) and cardiocerebrovascular diseases (OR=1.54, 95%CI=1.24-1.92, P<0.001).

Conclusion: Our study suggested that ABO polymorphisms may serve as a risk factor of cancers and cardiocerebrovascular diseases.

Background

Epidemiological studies have demonstrated that ABO blood groups were associated with several chronic inflammation related diseases, including cancers and cardiocerebrovasclar diseases [1–3]. The ABO blood group system is composed of complex carbohydrate structures that are biosynthesized by A- and B- transferases encoded by ABO gene. ABO gene is located at the 9q34 region of the chromosome and encodes enzyme glycosyltransferase with two main allele (A and B), a specific glycosyltransferase catalyzes the covalent attachment of N-acetylgalactosamine or D-galactose to a common precursor side chain (H determinant), ultimately to an A or B antigen [4]. However, the relationship between histoblood group antigens and inflammation related diseases remains unknown and the regulatory mechanisms underlying ABO expression was still unclear. Several SNPs in the ABO gene have been suggested to be associated with increased risks to the development of cancer and cardiocerebrovasclar disease by genome-wide association studies (GWAS) and candidate gene studies [5–8]. Particularly, the most widely investigated SNPs of the ABO gene were rs505922 and rs657152 [9–12]. Meta-analysis that evaluated the relationship between rs505922 SNP and overall cancer have already been reported by Duan et al [13] four years ago. However, several new findings of ABO gene SNPs have been reported in recent years and the relationship between SNPs in ABO gene and cancer/cardiocerebrovascular diseases risk was still unclear. Here, we have performed a meta-analysis including the newly published studies and evaluated the associations between polymorphisms of ABO gene and cancers/cardiocerebrovascular diseases.

Methods

Search strategy

Two investigators (Yanxia Li and Luyang Liu) performed a systematic literature search in three databases: PubMed, Web of Science and Embase to identify relevant articles published from the initial to Jan 1, 2019. The following search terms were used either separately or in combination: “SNP” “Polymorphism” and “Polymorphism, Single Nucleotide” “rs505922, rs657152” and “neoplasms” “carcinoma” “tumor” “cancer” and “vascular diasease” “cardiovascular disease” “cardiac-cerebral disease” and “ABO Blood-Group System” “Lewis Blood-Group System” “ABO”. Once suitable studies were singled out from the search results, other potentially relevant articles were identified by cross-references within eligible studies. The references of each identified articles were also searched manually to identify eligible studies.

Selection criteria

The inclusion criteria for these studies were as follows: 1) Studies evaluating the association between rs505922/rs657152 variants and cancers or cardiocerebrovascular diseases; 2) The study design was a case-control study in humans; 3) Researches containing universal allele and genotype data; 4) Studies written in English. The exclusion criteria were as follows: 1) Studies did not describe the association of ABO gene polymorphism with cancers or cardiocerebrovascular diseases; 2) Systematic reviews or articles focusing on animals; 3) Studies that did not provide usable data for meta-analysis.

Data extraction and quality assessment

Two investigators (Yanxia Li and Luyang Liu) independently extract data and verify the accuracy of the data. The following information was extracted from each article: first authors, publication years, ethnicity, cancer types, and study design (hospital- or population-based), sample size of subject, number of cases and controls for each genotype, OR and 95% CI in allele model. To assess the quality of the study, we used the Newcastle-Ottawa Scale (NOS) with a nine-star system [14]; this scale assesses the quality of cohort and case-control studies. The highest score of NOS is nine stars: four stars for the selection process, three stars for exposure/outcome, and two stars for comparability. A score of seven or above was considered to be high-quality study.

Statistical analysis

We used odds ratios (ORs) and 95% confidence intervals (CIs) to assess the relationship between each SNP and the risk of cancers or cardiocerebrovascular diseases. The association was examined using allele model. The significance of the pooled OR was determined by the Z-test. Subgroup analyses were conducted according to cancer types, ethnic groups and sources of controls. Heterogeneity between articles was identified with Q-test and I^2 index, I^2 values of >50% indicated heterogeneity among studies. If there was heterogeneity (I^2 >50%) between studies, we used a random effect model (DerSimonian-Laird method), otherwise we used a fixed effect model (Mantel-Haenszel method). Sensitivity analysis was performed to assess the effects of individual studies on pooled results and the stability of the results. We used Egger's and Begg's test to evaluate the publication bias, with a P >0.05 considered to be evidence for no potential publication bias. Trim and fill method was also applied in detecting publication bias. All the tests were two-sided and P <0.05 was considered to be statistically significant. Stata version 12.0 SE was applied to carry out the statistical analysis.

Results

Study characteristics

After retrieving the database, a total of 545 records [PubMed (n = 146), Web of Science (n = 149), EMBASE (n = 250)] were obtained. According to the inclusion/exclusion criteria, eighteen articles were included and the detailed flowchart of study selection process was presented in Fig. 1. Eleven studies reported the association between rs505922 and cancer risk [1, 2, 8–10, 15–20] and four studies reported the association between rs657152 and cancer risk [9, 10, 19, 21]. Six studies included nine populations reported the association of cardiocerebrovascular diseases with rs505922 [11, 12, 22–25], and three studies with rs657152 [11, 12, 24]. Detailed characteristics and genotype distribution of included articles for two SNPs were shown in Table 1 and Table 2. According to the source of control groups, nine studies were population-based (PB), six studies were hospital-based (HB), and two studies were population and hospital-based (PB/HB) control. For SNP rs505922, ten studies were from Caucasian population [1, 2, 8, 10, 12, 15, 16, 20, 22, 24], four studies were from Asian [9, 17, 23, 25] and three studies were from Mixed and African [11, 18, 19]. For the rs657152 polymorphism, there were four studies originating from Caucasian [10, 12, 21, 24], and three studies were from Asian and African or Mixed population [9, 11, 19]. Each study was scored based on the Newcastle-Ottawa Scale (NOS) and detailed study qualities were presented in Suppl Table 1 and Suppl Table 2.

Meta-analysis of rs505922 polymorphism

Meta-analysis was conducted to estimate the associations between rs505922 and cancer risk (Fig. 2a; Suppl Table 3) in 16,929 cancer cases and 23,941 controls. The rs505922 polymorphism was significantly associated with an increased cancers risk in the allele model (OR = 1.13, 95%CI = 1.04–1.22, P = 0.003). Subgroup analysis was conducted based on ethnicity, type of cancer, and source of control. The association between rs505922 and cancer risk was identified in Asian population subgroup (OR = 1.27, 95%CI = 1.10–1.48, P = 0.002), Mixed population subgroup (OR = 1.23, 95%CI = 1.16–1.31, P <0.001), Pancreatic cancers subgroup (OR = 1.23, 95%CI = 1.16–1.31, P <0.001), and Hospital based control groups (OR = 1.30, 95%CI = 1.12–1.51, P = 0.003) (Suppl Table 4; Suppl Figure 1). However, no significant association was observed in Caucasian population (OR = 1.05, 95%CI = 0.97–1.13, P = 0.232). Sensitivity analyses were conducted by omitting each individual article to measure its specific effect on the pooled ORs (Suppl Figure 2a). Sensitivity analysis plot indicated that no single study significantly affected the combined OR of SNP loci. Because of the heterogeneity of the research, we use the random effect model (allele model: I^2 = 79.1%) (Suppl Table 3). No significant publication bias was observed in any studies of SNPs (Fig. 4a; Suppl Table 3). After the applying the trim and fill method, there is no change in the OR value after the combination, also indicating that the original result is stable (Suppl Figure 3).

We also performed a meta-analysis to evaluate the association between rs505922 SNP and cardiocerebrovascular diseases (Fig. 2b; Suppl Table 3). The rs505922 SNP was significantly associated with an increased cardiocerebrovascular diseases risk in the allele model (C/T: OR = 1.36, 95%CI = 1.19–1.57, P <0.001). Subgroup analysis indicated that rs505922 was associated with a significantly higher risk of cardiocerebrovascular diseases in Caucasian population subgroup (OR = 1.39, 95%CI = 1.19–1.64, P <0.001, allele model), African population subgroup (OR = 1.52, 95%CI

= 1.18–1.96, $P = 0.001$), Hospital based control (OR = 1.14, 95%CI = 1.05–1.23, $P = 0.003$) and Population based control (OR = 1.54, 95%CI = 1.18–2.02, $P = 0.002$) (Suppl Table 4; Suppl Figure 4). However, no significant association was observed in Asian subgroup (OR = 1.21, 95%CI = 0.67–2.19, $P = 0.524$) (Suppl Table 4; Suppl Figure 4). Sensitivity analysis showed the results of this study were stable (Suppl Figure 2b). Begg's and Egger's test did not found significant publication bias among the enrolled studies (Fig. 4b; Suppl Table 3). After applying the trim and fill method, no new literature has been added, indicating that the result is stable (Suppl Figure 5).

Meta-analysis of rs657152 polymorphisms

Four studies reported the association between rs657152 and cancers risk. Our results showed that rs657152 was significantly associated with increased cancers risk in allele model (OR = 1.16, 95%CI = 1.09–1.24, $P < 0.001$) (Fig. 3a; Suppl Table 3). Sensitivity analysis revealed no significant influence on the pooled OR by any individual study (Suppl Figure 6a). We conducted Begg's and Egger's tests to assess the publication bias for these studies (Fig. 4c; Suppl Table 3). No evidence of publication bias was detected among the enrolled studies as shown in Table 3. According to the results of trim and fill method, the result was stable (Suppl Figure 7).

We also evaluated the relationship between rs657152 and cardiocerebrovascular diseases. A significant association between rs657152 and cardiocerebrovascular diseases under allele model were observed (OR = 1.54, 95%CI = 1.24–1.92, $P < 0.001$) (Fig. 3b; Suppl Table 3). Sensitivity analysis suggested that there was no significant change in the overall outcomes after removing any of the studies (Suppl Figure 6b). The Begg's and Egger's tests also showed that no meaningful publication bias were found (Fig. 4d; Suppl Table 3). The result of the trim and fill method proves that the result was stable (Suppl Figure 8).

Discussion

In this study, we conducted a meta-analysis to clarify the relationship between ABO SNPs (rs505922 and rs657152) and cancer/cardiocerebrovascular diseases risk. Our results showed that these two SNPs were associated with cancer risk and also increased the risk of cardiocerebrovascular diseases.

Regarding to rs505922 SNP, our results showed that the variant type of rs505922 could increase the risk of overall cancer, suggesting a potential predictive ability of this SNP for cancer risk. When we conducted a subgroup analysis of rs505922 based on cancer sites and ethnicity, we found that there was no significant association between rs505922 and Non-pancreatic cancer subgroup or Caucasian subgroup. This result may be due to heterogeneity of cancer types or insufficient statistical power. Therefore, further studies with large samples size are warrant to evaluate the association between the rs505922 polymorphisms in Non-pancreatic cancers. On the other side, our data also showed that rs505922 was related to cardiocerebrovascular diseases. In subgroup analysis, we found that there was no association between rs505922 and cardiocerebrovascular diseases in Asian subgroup. Only two studies with small population were included in this analysis, further studies are needed in Asian population. The SNP is located in the first intron region of ABO gene, the protective T allele of rs505922 is in complete linkage disequilibrium ($r^2 = 1.0$) with the O allele, is marks of O allele. However, the regulatory mechanism underlying the expression of histoblood group antigens was unclear.

For rs657152, our study demonstrated that this SNP was associated with cancer/ cardiocerebrovascular diseases risk. However, only seven studies [9–12, 19, 21, 24] were reported and most of studies were conducted in Caucasian population. Therefore, more studies with different ethnic background and larger sample size are needed in the future. rs657152 is located in the intron area of ABO, thus it may affect gene transcription and shearing [23]. In addition, rs657152 has been found to be associated with several biological molecule, including LDL cholesterol [26], liver derived alkaline phosphatas [27], and IL-6 [28]. This implied that rs657152 may affect the occurrence and development of disease by influencing these biological molecule. Further subsequent functional studies are warrant.

The underlying mechanism for the relationship between ABO blood group and cancer risk is still poorly understood [29]. It is reported that blood type may affect the progression and expansion of malignant tumors by altering the systemic inflammatory response [30]. Recent studies reported an association between polymorphisms at the ABO gene locus and circulating levels of tumour necrosis factor-alpha [31], soluble intercellular adhesion molecule (ICAM)-1 [32, 33], E-selectin [34, 35], and P-selectin [33]. These adhesion molecules were important mediators of chronic inflammation and immune cell recruitment [36]. They may provide a biological basis for the postulated influence of ABO on cancer survival, by linking ABO blood group and tumour initiation and spread [29]. In addition, some research have shown that the structure of certain tumor antigens was similar to the structure of antigens of ABO blood group system. Smith and Prieto [37] suggested the Forssmann antigen which predominant in stomach and colon tumors, was almost structurally identical to the A antigen determinant. Blood group A carrier may have diminished tumor immune response due to reduced ability to recognize and attack tumor cells [38].

There was some evidence linking ABO blood group and cardiocerebrovascular diseases. Jenkins, P. V. et al. reported an association between ABO blood types and von Willebrand factor (vWF) and factor VIII (FVIII), both of which play crucial roles in the coagulation pathway [39]. Higher levels of vWF and FVIII has been observed in non-O blood type than O blood type [40]. Therefore, type O blood may be a risk factor for bleeding [41]. In addition, the non-O blood group has been shown to be correlated with higher total cholesterol and LDL-C levels [42], and the latest study proposed

that approximately 10% of the effect of ABO blood group on coronary artery disease (CAD) susceptibility was mediated by plasma cholesterol levels[43].

Limitations in this study should be mentioned. First, the studies included in our meta-analysis were limited to published reports and English language studies. Unpublished reports or those published in non-English language studies were not included in the analysis. It would limit our sample size and publication bias might exist. Second, both of the hospital based and population based case-control studies were included in our study. Therefore, selection bias would exist compared to the meta-analysis only included population based case-control studies. Third, the limited number of published studies may influence the reliability of our results. Finally, the lack of original data limited further evaluations of the potential gene-gene and gene-environment interactions.

Conclusion

In summary, the results of our meta-analysis revealed that rs505922 and rs657152 were correlated with an increased cancers risk. Due to most of the studies were conducted in pancreatic cancer type and Caucasian populations, further studies in multiple cancer types and multiple ethnic populations are needed. In addition, our meta-analysis also revealed that rs505922 and rs657152 associated with cardiocerebrovascular diseases. However, owing to limited number of studies, further studies with larger samples size are warrant. This study can provide clues for further exploration of novel biomarkers with cancer/cardiocerebrovascular early-warning function.

Abbreviations

PB: population based, HB: hospital based, NOS: Newcastle-Ottawa Scale.

Declarations

Ethics approval and consent to participate:

Not applicable.

Consent for publication:

Not applicable.

Availability of data and material:

Not applicable.

Competing interests:

The authors declare no conflicts of interest.

Funding:

This work was supported by the National Natural Science Foundation of China (81320108022, 81470153, and 81502877), Tianjin Science and Technology Committee Foundation (18YFZCSY00520 and 16JCYBJC26600), the National Key Research and Development Program of China (2016YFC1302703).

Author' Contributions:

Study conception and design: Lian Li and Hong Zheng. Acquisition, analysis, or interpretation of data: Yanxia Li and Luyang Liu. Statistical analysis: Yanxia Li and Yubei Huang. Drafting of the manuscript: Yanxia Li and Luyang Liu. Administrative, technical, or material support: Hong Zheng. Study supervision: Lian Li and Hong Zheng. All authors were involved in writing the paper and had final approval of the submitted and published versions.

Acknowledgements:

Not applicable.

References

- [1] Krawczyk M, Mihalache F, Hoblinger A, et al., Pancreatic cancer risk variant ABO rs505922 in patients with cholangiocarcinoma. *World J Gastroenterol.* 2011; 17(41):4640–2.
- [2] Gates M A, Xu M, Chen W Y, et al., ABO blood group and breast cancer incidence and survival. *Int J Cancer.* 2012; 130(9):2129–37.
- [3] Dentali F, Sironi A P, Ageno W, et al., Non-O blood type is the commonest genetic risk factor for VTE: results from a meta-analysis of the literature. *Semin Thromb Hemost.* 2012; 38(5):535–48.
- [4] Yamamoto F, Clausen H, White T, et al., Molecular genetic basis of the histo-blood group ABO system. *Nature.* 1990; 345(6272):229–33.
- [5] Ling X, Zheng Y, Tao J, et al., Association study of polymorphisms in the ABO gene with ischemic stroke in the Chinese population. *BMC Neurology.* 2016; 16:146.
- [6] Heit J A, Armasu S M, Asmann Y W, et al., A genome-wide association study of venous thromboembolism identifies risk variants in chromosomes 1q24.2 and 9q. *J Thromb Haemost.* 2012; 10(8):1521–31.
- [7] Gao M, Tang H, Zheng X, et al., Association analysis of GWAS and candidate gene loci in a Chinese population with coronary heart disease. *Int J Clin Exp Med.* 2015; 8(5):7497–7506.
- [8] Markt S C, Shui I M, Unger R H, et al., ABO blood group alleles and prostate cancer risk: Results from the breast and prostate cancer cohort consortium (BPC3). *Prostate.* 2015; 75(15): 1677–81.
- [9] Xu H-L, Cheng J-R, Zhang W, et al., Re-evaluation of ABO gene polymorphisms detected in a genome-wide association study and risk of pancreatic ductal adenocarcinoma in a Chinese population. *Chinese Journal of Cancer.* 2014; 33(2):68–73.
- [10] Duell E J, Bonet C, Munoz X, et al., Variation at ABO histo-blood group and FUT loci and diffuse and intestinal gastric cancer risk in a European population. *Int J Cancer.* 2015; 136(4):880–93.
- [11] Hernandez W, Gamazon E R, Smithberger E, et al., Novel genetic predictors of venous thromboembolism risk in African Americans. *Blood.* 2016; 127(15):1923–9.
- [12] Reilly M P, Li M, He J, et al., Identification of ADAMTS7 as a novel locus for coronary atherosclerosis and association of ABO with myocardial infarction in the presence of coronary atherosclerosis: two genome-wide association studies. *Lancet.* 2011; 377(9763):383–92.
- [13] Duan Y F, Zhu F, Li X D, et al., Association between ABO gene polymorphism (rs505922) and cancer risk: a meta-analysis. *Tumour Biol.* 2015; 36(7):5081–7.
- [14] Stang A, Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol.* 2010; 25(9):603–5.
- [15] Rizzato C, Campa D, Pezzilli R, et al., ABO blood groups and pancreatic cancer risk and survival: results from the PANcreatic Disease ReseArch (PANDoRA) consortium. *Oncol Rep.* 2013; 29(4):1637–44.
- [16] Poole E M, Gates M A, High B A, et al., ABO blood group and risk of epithelial ovarian cancer within the Ovarian Cancer Association Consortium. *Cancer Causes Control.* 2012; 23(11):1805–10.
- [17] Nakao M, Matsuo K, Hosono S, et al., ABO blood group alleles and the risk of pancreatic cancer in a Japanese population. *Cancer Science.* 2011; 102(5):1076–80.
- [18] Wolpin B M, Kraft P, Gross M, et al., Pancreatic cancer risk and ABO blood group alleles: results from the pancreatic cancer cohort consortium. *Cancer Res.* 2010; 70(3):1015–23.
- [19] Li D, Duell E j, Yu K, et al., Pathway analysis of genome-wide association study data highlights pancreatic development genes as susceptibility factors for pancreatic cancer. *Carcinogenesis.* 2012; 33(7):1384–90.
- [20] Willis J A, Olson S H, Orlow I, et al., A replication study and genome-wide scan of single-nucleotide polymorphisms associated with pancreatic cancer risk and overall survival. *Clin Cancer Res.* 2012; 18(14):3942–51.

- [21] Rizzato C, Campa D, Giese N, et al., Pancreatic cancer susceptibility loci and their role in survival. *PLoS One*. 2011; 6(11):e27921.
- [22] Williams F M, Carter A M, Hysi P G, et al., Ischemic stroke is associated with the ABO locus: the EuroCLOT study. *Ann Neurol*. 2013; 73(1):16–31.
- [23] Zhang H, Zhang Z, Zhang J, et al., Fine-mapping of ABO gene identifies two novel SNPs associated with large artery atherosclerotic stroke in a Chinese han population. *Mol Neurobiol*. 2017; 54(3):2107–2113.
- [24] Tregouet D A, Heath S, Saut N, et al., Common susceptibility alleles are unlikely to contribute as strongly as the FV and ABO loci to VTE risk: results from a GWAS approach. *Blood*. 2009; 113:5298–5303.
- [25] Li H, Cai Y, and Xu A D, Association study of polymorphisms in the ABO gene and their gene-gene interactions with ischemic stroke in Chinese population. *J Clin Lab Anal*. 2017; 32(4):e22329.
- [26] Teupser D, Baber R, Ceglarek U, et al., Genetic regulation of serum phytosterol levels and risk of coronary artery disease. *Circ Cardiovasc Genet*. 2010; 3(4):331–9.
- [27] Yuan X, Waterworth D, Perry J R B, et al., Population-based genome-wide association studies reveal six loci influencing plasma levels of liver enzymes. *American Journal of Human Genetics*. 2008; 83(4):520–528.
- [28] Naitza S, Porcu E, Steri M, et al., A genome-wide association scan on the levels of markers of inflammation in Sardinians reveals associations that underpin its complex regulation. *PLoS Genetics*. 2012; 8(1):e1002480.
- [29] Franchini M, Favaloro E J, Targher G, et al., ABO blood group, hypercoagulability, and cardiovascular and cancer risk. *Crit Rev Clin Lab Sci*. 2012; 49(4):137–49.
- [30] Grivnenikov S I, Greten F R, and Karin M, Immunity, inflammation, and cancer. *Cell*. 2010; 140:883–99.
- [31] Melzer D, Perry J R, Hernandez D, et al., A genome-wide association study identifies protein quantitative trait loci (pQTLs). *PLoS Genet*. 2008; 4(5):e1000072.
- [32] Pare G, Chasman D I, Kellogg M, et al., Novel association of ABO histo-blood group antigen with soluble ICAM–1: results of a genome-wide association study of 6,578 women. *PLoS Genet*. 2008; 4(7):e1000118.
- [33] Barbalic M, Dupuis J, Dehghan A, et al., Large-scale genomic studies reveal central role of ABO in sP-selectin and sICAM–1 levels. *Hum Mol Genet*. 2010; 19(9):1863–72.
- [34] Paterson A D, Lopes-Virella M F, Waggott D, et al., Genome-wide association identifies the ABO blood group as a major locus associated with serum levels of soluble E-selectin. *Arterioscler Thromb Vasc Biol*. 2009; 29(11):1958–67.
- [35] Qi L, Cornelis M C, Kraft P, et al., Genetic variants in ABO blood group region, plasma soluble E-selectin levels and risk of type 2 diabetes. *Hum Mol Genet*. 2010; 19(9):1856–62.
- [36] Franchini M, Liumbruno G M, and Lippi G, The prognostic value of ABO blood group in cancer patients. *Blood Transfus*. 2016; 14(5):434–40.
- [37] Smith D F and Prieto P A, Forssmann antigen. In: Roitt IM, Delves PH (eds). *Encyclopedia of Immunology*. Academic Press. 1992:591–2.
- [38] S RYuzhalin A E and Kutikhin A G, ABO and Rh blood groups in relation to ovarian, endometrial and cervical cancer risk among the population of South-East Siberia. *Asian Pac J Cancer Prev*. 2012; 13(10):5091–6.
- [39] Jenkins P V and O'Donnell J S, ABO blood group determines plasma von Willebrand factor levels: a biologic function after all? *Transfusion*. 2006; 46(10):1836–44.
- [40] Gill J C, Endres-Brooks J, Bauer P J, et al., The effect of ABO blood group on the diagnosis of von Willebrand disease. *Blood*. 1987; 69(6):1691–5.
- [41] Dentali F, Sironi A P, Ageno W, et al., Relationship between ABO blood group and hemorrhage: a systematic literature review and meta-analysis. *Semin Thromb Hemost*. 2013; 39(1):72–82.
- [42] Silbernagel G, Chapman M J, Genser B, et al., High intestinal cholesterol absorption is associated with cardiovascular disease and risk alleles in ABCG8 and ABO: evidence from the LURIC and YFS cohorts and from a meta-analysis. *J Am Coll Cardiol*. 2013; 62(4):291–9.

[43] Chen Y, Chen C, Ke X, et al., Analysis of circulating cholesterol levels as a mediator of an association between ABO blood group and coronary heart disease. Circ Cardiovasc Genet. 2014; 7(1):43–8.

Tables

SNP loci	Year	First author	Ethnicity	Cancer/disease	Control*	Sample size (case/control)	Case			Control			OR(95%CI)
							Hom [†]	Het [†]	Hom	Hom	Het	Hom	
							wild	variant		wild	variant		
rs505922	2015	SC. Markt	Caucasian	Prostate cancer	-	2774/4443	1172	1226	376	1772	2070	601	0.95(0.89-1.02)
	2015	E. Duell	Caucasian	Gastric cancer	PB	365/1284	137	172	49	533	541	188	1.06(0.89-1.26)
	2014	H. Xu	Asian	Pancreatic cancer	PB	256/548	69	124	63	186	246	115	1.24(1.00-1.53)
	2013	C.Rizzato	Caucasian	Pancreatic cancer	HB/PB	1028/2257	342	518	163	860	1055	336	1.13(1.01-1.26)
	2012	E.Poole	Caucasian	Ovarian cancer	HB/PB	5233/6838	2222	2407	603	2987	3044	806	1.02(0.97-1.08)
	2012	D. Li	Mixed	Pancreatic cancer	HB	3851/3934	-	-	-	-	-	-	1.21(1.13-1.30)
	2012	J. Willis	Caucasian	Pancreatic cancer	HB	385/149	-	-	-	-	-	-	1.65(1.20-2.26)
	2012	M.Gates	Caucasian	Breast cancer	PB	1138/1090	489	505	144	471	487	132	1.02(0.90-1.15)
	2011	M.Krawczyk	Caucasian	Cholangiocarcinoma	PB	180/350	84	68	28	154	146	50	0.97(0.74-1.27)
	2011	M. Nakao	Asian	Pancreatic cancer	HB	185/1465	38	101	46	428	745	292	1.31(1.06-1.63)
	2010	B.Wolpin	Mixed	Pancreatic cancer	PB	1534/1583	511	752	271	657	705	221	1.28(1.16-1.42)
rs657152	2015	E. Duell	Caucasian	Gastric cancer	PB	365/1284	-	-	-	-	-	-	1.05(0.89-1.23)
	2014	H. Xu	Asian	Pancreatic cancer	PB	256/548	71	124	61	199	240	108	1.29(1.05-1.60)
	2012	D. Li	Caucasian	Pancreatic cancer	HB	3851/3934	-	-	-	-	-	-	1.19(1.12-1.27)
	2011	C. Rizzato	Caucasian	Pancreatic cancer	PB	686/1255	199	357	130	437	591	227	1.11(0.96-1.27)

Table1. Characteristics of the studies included in the meta-analysis of the association between SNPs and cancers

*PB: population based control, HB: hospital based control; †Hom: homozygous, Het: heterozygote

Table2. Characteristics of the studies included in the meta-analysis of the association between SNPs and cardiocerebrovascular diseases

SNP loci	Year	First author	Ethnicity	Cancer/disease	Control*	Sample size (case-control)	Case			Control			OR(95%CI)
							Hom [†]	Het [†]		Hom	Het		
							Hom			Hom			
							wild variant			wild variant			
rs505922	2017	H. Li	Asian	Ischemic stroke	PB	991/1002	511	406	74	657	306	39	1.64(1.41-1.90)
	2017	H. Zhang	Asian	Large artery atherosclerotic stroke	PB	644/642	146	325	173	129	322	191	0.90(0.77-1.05)
	2016	W. Hernandez	African	Venous Thrombosis	HB	146 /432	-	-	-	-	-	-	1.52(1.20-2.00)
	2013	FM.Williams-a [‡]	Caucasian	Ischemic stroke	HB	4092/8383	-	-	-	-	-	-	1.06(1.01-1.14)
	2013	FM.Williams-b [‡]	Caucasian	Ischemic stroke	HB	8443 /54810	-	-	-	-	-	-	1.07(1.03-1.11)
	2011	MP. Reilly	Caucasian	Myocardial infarction	HB	5783/3644	-	-	-	-	-	-	1.20(1.13-1.28)
	2009	DA.Trégouet-a [§]	Caucasian	Venous thromboembolism	PB	419/1228	97	209	113	519	559	150	2.01(1.71-2.35)
	2009	DA.Trégouet-b [§]	Caucasian	Venous thromboembolism	PB	1150/801	299	575	276	339	364	98	1.79(1.57-2.04)
	2009	DA.Trégouet-c [§]	Caucasian	Venous thromboembolism	PB	607/607	177	302	128	265	272	70	1.66(1.41-1.95)
rs657152	2016	W. Hernandez	African	Venous Thrombosis	HB	146 /432	-	-	-	-	-	-	1.39(1.10-1.80)
	2011	MP. Reilly	Caucasian	Myocardial infarction	HB	5783/3644	-	-	-	-	-	-	1.19(1.12-1.27)
	2009	DA.Trégouet-a [§]	Caucasian	Venous thromboembolism	PB	419/1228	89	208	122	472	579	177	1.91(1.63-2.24)
	2009	DA.Trégouet-b [§]	Caucasian	Venous thromboembolism	PB	1150/801	276	575	299	318	373	110	1.77(1.55-2.02)
	2009	DA.Trégouet-c [§]	Caucasian	Venous thromboembolism	PB	607/607	171	302	134	249	280	78	1.58(1.34-1.86)

*PB: population based control, HB: hospital based control; †Hom: homozygous, Het: heterozygote; a[‡] represent the MOnica Risk, Genetics, Archiving and Monograph(MORGAM) and the Wellcome Trust Case Control Consortium 2(WTCCC2) population, b[‡] represent MetaStroke population; a[§] represent GWAS population from 4 different French centers, b[§] represent MARseille THrombosis Association study (MARTHA) population, c[§] represent FARIVE that is a multicenter case-control study

Figures

Figure 1

Flow chart of the study selection process

A **B** **Figure 2**

Forest plot of the relationship between rs505922 polymorphisms and cancer (a) and cardiocerebrovascular disease (b) risk (allele model and random-effect model).

A **B** **Figure 3**

Forest plot of the relationship between rs657152 polymorphisms and cancer (a) and cardiocerebrovascular disease(b) risk (allele model and random-effect model).

Figure 4

Begg's funnel plot of publication bias test for rs505922 and rs657152(allele model and random-effect model); Each point represents a separate study for the indicated association between rs505922 and cancer (a) and cardiocerebrovascular disease(b), rs657152 and cancer (c) and cardiocerebrovascular disease(d)risk, respectively.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [PRISMA2009checklist.doc](#)
- [ABOmetasupplemental.doc](#)