

Flexor Hallucis Longus Hypertrophy Secondary to Achilles Tendon Tendinopathy: An MRI-based Case-Control Study

Stephan Wirth

Uniklinik Balgrist

Octavian Andronic (✉ octavian.andronic@balgrist.ch)

Uniklinik Balgrist <https://orcid.org/0000-0002-3743-7033>

Fabian Aregger

Uniklinik Balgrist

Anna Jungwirth-Weinberger

Uniklinik Balgrist

Thorsten Jentsch

Uniklinik Balgrist

Andreas Hecker

Uniklinik Balgrist

Research article

Keywords: Achilles tendon tendinopathy, flexor hallucis longus hypertrophy, chronic achilles tendon rupture, flexor hallucis longus transfer, Achilles tendon MRI

Posted Date: October 19th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-51795/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at European Journal of Orthopaedic Surgery & Traumatology on February 8th, 2021. See the published version at <https://doi.org/10.1007/s00590-021-02891-8>.

Abstract

Background: The purpose of this study was to outline an indirect sign of advanced Achilles tendinopathy on magnetic resonance imaging (MRI), based on the hypothesis that these patients would present with secondary hypertrophy of the flexor hallucis longus muscle (FHL).

Methods: MRI scans of Achilles tendon were analyzed retrospectively in two cohorts. The study group consisted of consecutive patients presenting with clinical signs of Achilles tendinopathy and no previous surgeries, whilst the control group were patients that had an MRI due to other reasons and no signs of tendinopathy. Two parameters from two muscle bellies were measured and compared on axial MRI scans 4 to 5 cm above the ankle joint line at the level of greatest thickness: area and diameter of the triceps surae (TS) and of the FHL muscle. Ratios (FHL/TS) were calculated for area (Ar) and diameter (Dm) measurements. Interobserver agreement was analyzed. A receiver operating characteristic (ROC) curve was created for both ratios to assess potential cut-off points to differentiate between the groups.

Results: A total of 60 patients for each study group were included. Both ratios, Ar(FHL/TS) and Dm(FHL/TS) showed significant higher values in the tendinopathy group ($p < 0.001$). There were strong to very strong intraclass correlation coefficients (ICC=0.75-0.93). A diameter ratio Dm (FHL/TS) of 2.0 or higher had a sensitivity of 49% and specificity of 90% for concomitant Achilles tendinopathy.

Conclusion: In our patient cohort, FHL hypertrophy was observed in patients with Achilles tendinopathy as a possible compensatory mechanism. Measuring a diameter ratio Dm(FHL/TS) of 2.0 or higher on an axial MRI, may be indicative as an indirect sign of functional deterioration of the Achilles tendon.

Background

Chronic Achilles tendinopathy represents a common disease that can affect professional athletes(1) and also people engaging occasionally in sports or even having a sedentary lifestyle.(2) In 2011, its incidence in a Dutch population was described as 2 per 1000 patients registered by a general practitioner (GP).(3) In an histological study, 34% of the Achilles tendon specimens of spontaneously ruptured tendons were showing signs of tendinopathy, suggesting that the acute lesions also have a predisposing degeneration. (4) This further supports the importance of timely diagnosing and delivering the appropriate management.

First line of treatment is usually conservative and may take up months to demonstrate effectiveness. This includes reduction of load, stretching as well as eccentric muscle strengthening and may lead to a favorable outcome in 71% of the cases.(5) However, as many as 25% of patients fail conservative therapy and might benefit from surgery.(6) Surgical procedures, mostly consisting of debridement with direct repair, are considered when conservative treatment is not successful after an average period of 3 to 6 months.(7, 8) A variety of surgical options have been described without a definitive consensus on the best technique.(9)

If there is advanced damage of the tendon(6), a tendon transfer is a recommended option. (10-13) One of the most widely used is the flexor hallucis longus (FHL) transfer.(14) Evidence demonstrated a hypertrophy of the FHL after performing a tendon transfer.(15, 16)Furthermore, recent biomechanical studies provide important evidence supporting FHL as transfer option, as FHL transfer significantly increases load to failure of Achilles tendons.(17)

In our practice, there was a clinical observation that the FHL hypertrophy may even occur preoperatively in cases with severe Achilles tendinopathy as a compensatory mechanism. The purpose of this study was to outline this finding on magnetic resonance imaging (MRI), based on the hypothesis that patients with Achilles tendinopathy would present with secondary hypertrophy of the flexor hallucis longus muscle (FHL).

Methods

Consecutive MRI scans between 01/2016 and 01/2017, that assessed the calf region and included the Achilles tendon, were extracted from the local institutional database. All patients that were included signed a written consent form. The study was approved by our institutional ethical review board and by the local Ethics Commission (BASEC Nr. 2018-00098). The study was carried out in accordance with the World Medical Association Declaration of Helsinki.(18)

Patient selection

Inclusion criteria for the tendinopathy group were patients that exhibited radiographic signs of Achilles tendon tendinopathy as defined by previous studies,(19, 20) which included combinations of the following: intratendinous signal alterations, interstitial or insertional tears, peritendinous edema, loss of the physiological anterior concave margin of the tendon, irregular mucoid deposition, intratendinous multifocal speckled appearance with changes in volume. Only imaging of sufficient quality that included sagittal and axial views (T1 as well as T2 weighted/ STIR (short tau inversion recovery)/ with and without fat suppression) were included

The control group included subjects without any documented complaints regarding the Achilles function and presented a normal radiographic morphology of the Achilles tendon on MRI. Exclusion criteria in both groups were relevant morphologic changes of the upper ankle joint, the hindfoot or the calf muscles as well as history of relevant previous surgery or injury of the lower limb.

Radiographic Measurement Strategy

Axial MRI scans (slice thickness 6mm) were analyzed on a level 4 to 5 cm above the ankle joint line at the level of greatest thickness for each muscle belly. This height was chosen as it was consistently representative of a good portion of muscle belly for both FHL and Triceps surae/Soleus. A sagittal view was then used to verify the height (**Fig. 1**). Next, for measurement of the diameter of the TS, a perpendicular line was drawn to the frontal plane through the midline of the muscle belly. The FHL

diameter was measured as the distance from the medial corner of the fibula to the posteromedial corner of the FHL muscle (**Fig. 2**), identified as the location where the neurovascular bundle runs along the FHL. Measurements were made in both groups by two independent investigators (**Fig 3**). Next, ratios of the area - Ar(FHL/TS) and of the diameter - Dm(FHL/TS) were calculated.

Statistical Analysis

Data was mainly non-normally distributed. Medians and interquartile ranges (IQR) are given. Spearman correlation analysis tested for interobserver agreement. The Wilcoxon rank sum test was used to compare measurements between groups. Intraclass correlation coefficients (ICC) for individual agreements with F tests were calculated for interrater agreement. We also provide minimal detectable change and standard error of measurement values. A logistic regression model was fitted to account for the potential confounder age, sex, and side. A receiver operating characteristic (ROC) curve was created for both quotients to assess potential cut-off points. A post hoc power analysis using a test comparing two independent means yielded a sufficient power (power=0.99 for the quotient of the diameter of FHL divided by TS using the mean and standard deviations (SD) for both groups (2.16 (SD 0.62) and 1.71 (SD 0.32) (n=120)). Stata/IC (version 13.1; StataCorp LP, College Station, TX, USA) was used.

Results

Demographics of the study participants are represented in **Table 1**. The mean height of measurement was 4.5cm above the ankle joint in both groups.

The diameter ratio Dm (FHL/TS) showed significantly higher values in the tendinopathy group than the control group, $p < 0.001$ (median=2.0 [IQR=0.8] vs 1.7 [0.3]) (**Table 2**). Similar results were obtained by calculating the ratio regarding the area Ar (FHL/TS), which also showed significantly higher values in the tendinopathy group than in the control group (1.8 [1.3] vs 1.3 [0.7]), $p < 0.001$.

A strong to very strong ICC regarding all measurements was determined (0.75-0.93) (**Table 3**).

In the tendinopathy group there were more males and overall older patients than in the control group ($p < 0.001$). This issue was then addressed in a logistic regression model that fitted results depending on age, gender and affected side.

As such, there was still a significant odd's ratio=9.56 [95% confidence interval 2.46-37.22], $p = 0.001$) for an increased ratio of diameter FHL/TS in the tendinopathy group (**Table 2**).

Diameter values together with its calculated quotient Dm (FHL/TS) showed less variation as demonstrated by reduced dispersion on the boxplots (**Fig 4**). Therefore, it was this value that was chosen for the calculation of a potential radiographic tool. For the calculation of a suitable cut-off point for the diameter ratio Dm (FHL/TS), a receiver operating characteristic (ROC) analysis was undertaken (**Fig. 5**). A value of 1.6 or above was most sensitive to detect a positive correlation with concomitant tendinopathy

with a sensitivity of 83% and specificity of 45%. On the other hand, a value of 2.0 or above, provided a higher specificity of 90%. The latter value was chosen as cut-off margin.

Discussion

This is the first study reporting evidence for FHL hypertrophy that occurs in cases of severe chronic Achilles tendinopathy even before a transfer is employed. Also, the study provides a new objective radiographic tool that might help in the surgical decision-making and should be the base for future studies that will correlate it to clinical outcome.

Previous studies evaluated the reliability of ultrasound examination(21) and compared clinically symptomatic and asymptomatic Achilles tendons which found tendinopathy in 32% of the asymptomatic volunteers.(22) This demonstrates the further need for additional parameters in assessing the functionality of the Achilles tendon. Oksanen et al. found FHL hypertrophy in over 50% of patients after performing a FHL transfer.(15) Similar findings were observed and reported on postoperative MRI imaging.(15, 23, 24) We believe the hypertrophy may also occur in case of functional deterioration of the triceps surae due to chronic Achilles tendinopathy as a compensatory mechanism.

Another area of need for an additional objective tool is the indication for surgical treatment of a chronic Achilles tendinopathy which is a matter of current debate. Rahm et al. suggested to perform FHL transfer after failure of local debridement, failure of free tendon grafting in case of a tendon defect or advanced fatty infiltration of the TS.(6) A minimal defect of 50% of the tendon was determined as an indication for the need of a tendon transfer by several authors (7, 10, 11, 25) Lin et al.(26) presented an algorithm for the treatment of chronic Achilles tendinopathy based on the presence or absence of stumps on preoperative MRI and the defect gap measured intra-operatively. FHL transfer was considered when the tendon stumps did not have enough integrity.(4) Clinical results were reported to be good to excellent after FHL transfer for multiple pathologies of the Achilles tendon, including extended degenerative aberrations, partial defects and muscle dysfunction of the TS.(12, 27). Very good ankle plantar flexion strength and an overall clinical success rate of over 70% have been reported. (28-30)

As such, the management for Achilles tendinopathy experiences a variety of treatment strategies and lacks standardized criteria as it depends on subjective considerations of the caretaker.(31) The observed FHL hypertrophy in patients with signs of chronic Achilles tendinopathy provides evidence that further supports its functional synergism and as such, another argument for choosing an FHL transfer. Using the new ratio of diameter D_m (TS/FHL), an additional radiographic sign can be potentially employed in the surgical decision-making. Future clinical prospective studies should correlate this parameter to clinical and functional outcome.

Another subject of discussion may be to question the MRI as diagnostic tool for Achilles tendinopathy. MRI can be time consuming and expensive.(32) Although it was shown that it is not necessary for diagnosing acute injuries, in atraumatic cases with inconclusive clinical findings (normal or equivocal

Thompson test, nonpalpable gap between tendon edges, known chronic rupture and other), it has decisive importance for diagnosis.(33)

Obviously, there are several limitations that should be addressed. The design has a weakness regarding study groups that were not matched on age, BMI or comorbidities, which are possible confounders. This matter was addressed by performing a multiple regression analysis. Another issue is the cross-sectional measurement of the muscle belly at a specific location, that may be not representative of the total muscular volume. Future studies may consider potential confounders height, weight and length of the free Achilles tendon (distance from calcaneus to tip of soleus muscle) in their analyses to account for different anatomical configurations in each individual (e.g. the left patient in Figure 3 may have a longer Achilles tendon than the right patient, potentially leading to a smaller soleus volume 4-5 cm above the ankle joint line).

Conclusion

In our patient cohort, flexor hallucis longus hypertrophy was observed in the Achilles tendinopathy group, probably as a compensatory mechanism. Measuring a diameter ratio D_m (FHL/TS) of 2.0 or higher on an axial MRI, may be indicative as an indirect sign of functional deterioration of the Achilles tendon.

Future studies should correlate this finding to clinical outcome, which may potentially aid in the surgical decision-making and the employment of a tendon transfer.

Declarations

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

The study was approved by our institutional ethical review board and by the Cantonal Ethics Commission of Zurich (BASEC Nr. 2018-00098). All patients have signed a written consent form.

CONSENT TO PUBLISH

Consent to publish was obtained from all relevant study participants.

AVAILABILITY OF DATA AND MATERIALS

Data was deposited safely in a local repository according to the ethical commission regulations. REDCAP software was used for storage and limited access to the principal investigators to protect patient personal information.

COMPETING INTERESTS

All authors declare that they have no conflict of interest.

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

AUTHORS' CONTRIBUTIONS

All authors must have made substantial contributions to all four categories established by the International Committee of Medical Journal Editors (<http://www.icmje.org>) to include: (1) conception and design, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be published, and (4) agree to be accountable for all aspects of the work if questions.

S.W.: Conceptualization, Project Administration, Resources, Software, Supervision, Writing – Original Draft Preparation, Writing – Review & Editing

O.A.: Investigation, Formal Analysis, Methodology, Validation, Writing – Review & Editing, Statistical work.

F.A.: Data acquisition, Formal Analysis, Validation, Writing – Review & Editing

A.J.: Investigation, Formal Analysis, Methodology, Validation, Writing – Review & Editing **T.J.:** Investigation, Methodology, Statistical Expertise, Validation, Writing – Review & Editing

A.H.: Conceptualization, , Project Administration, Resources, Software, Supervision, Writing – Review & Editing

ACKNOWLEDGEMENTS

None to declare.

References

1. Chan JJ, Chen KK, Sarker S, Hasija R, Huang HH, Guzman JZ, et al. Epidemiology of Achilles tendon injuries in collegiate level athletes in the United States. *International orthopaedics*. 2020;44(3):585-94.
2. Aström M, Gentz CF, Nilsson P, Rausing A, Sjöberg S, Westlin N. Imaging in chronic achilles tendinopathy: a comparison of ultrasonography, magnetic resonance imaging and surgical findings in 27 histologically verified cases. *Skeletal radiology*. 1996;25(7):615-20.
3. de Jonge S, van den Berg C, de Vos RJ, van der Heide HJ, Weir A, Verhaar JA, et al. Incidence of midportion Achilles tendinopathy in the general population. *Br J Sports Med*. 2011;45(13):1026-8.
4. Kannus P, Jozsa L. Histopathological changes preceding spontaneous rupture of a tendon. A controlled study of 891 patients. *J Bone Joint Surg Am*. 1991;73(10):1507-25.
5. Roche AJ, Calder JD. Achilles tendinopathy: A review of the current concepts of treatment. *Bone Joint J*. 2013;95-B(10):1299-307.

6. Rahm S, Spross C, Gerber F, Farshad M, Buck FM, Espinosa N. Operative treatment of chronic irreparable Achilles tendon ruptures with large flexor hallucis longus tendon transfers. *Foot & ankle international*. 2013;34(8):1100-10.
7. Shakked RJ, Raikin SM. Insertional Tendinopathy of the Achilles: Debridement, Primary Repair, and When to Augment. *Foot Ankle Clin*. 2017;22(4):761-80.
8. Lohrer H, David S, Nauck T. Surgical treatment for achilles tendinopathy – a systematic review. *BMC musculoskeletal disorders*. 2016;17(1):207.
9. Abubeih H, Khaled M, Saleh WR, Said GZ. Flexor hallucis longus transfer clinical outcome through a single incision for chronic Achilles tendon rupture. *International orthopaedics*. 2018;42(11):2699-704.
10. Baumbach SF, Braunstein M, Mack MG, Massen F, Bocker W, Polzer S, et al. [Insertional Achilles tendinopathy : Differentiated diagnostics and therapy]. *Unfallchirurg*. 2017;120(12):1044-53.
11. Will RE, Galey SM. Outcome of single incision flexor hallucis longus transfer for chronic achilles tendinopathy. *Foot Ankle Int*. 2009;30(4):315-7.
12. Amlang MH, Rosenow M, Rammelt S, Heineck J, Zwipp H. [Transfer of the flexor hallucis longus to replace the Achilles tendon: indications, technique and results]. *Unfallchirurg*. 2008;111(7):499-506.
13. Martin RL, Manning CM, Carcia CR, Conti SF. An outcome study of chronic Achilles tendinosis after excision of the Achilles tendon and flexor hallucis longus tendon transfer. *Foot Ankle Int*. 2005;26(9):691-7.
14. Staggers JR, Smith K, de CNC, Naranje S, Prasad K, Shah A. Reconstruction for chronic Achilles tendinopathy: comparison of flexor hallucis longus (FHL) transfer versus V-Y advancement. *International orthopaedics*. 2018;42(4):829-34.
15. Oksanen MM, Haapasalo HH, Elo PP, Laine HJ. Hypertrophy of the flexor hallucis longus muscle after tendon transfer in patients with chronic Achilles tendon rupture. *Foot and ankle surgery : official journal of the European Society of Foot and Ankle Surgeons*. 2014;20(4):253-7.
16. Heikkinen J, Lantto I, Piilonen J, Flinkkila T, Ohtonen P, Siira P, et al. Tendon Length, Calf Muscle Atrophy, and Strength Deficit After Acute Achilles Tendon Rupture: Long-Term Follow-up of Patients in a Previous Study. *J Bone Joint Surg Am*. 2017;99(18):1509-15.
17. Schmidtberg B, Johnson JD, Kia C, Baldino JB, Obopilwe E, Cote MP, et al. Flexor Hallucis Longus Transfer Improves Achilles Tendon Load to Failure in Surgery for Non-Insertional Tendinopathy: A Biomechanical Study. *JBJS*. 2019;101(16).
18. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *Jama*. 2013;310(20):2191-4.
19. Shalabi A. Magnetic resonance imaging in chronic Achilles tendinopathy. *Acta radiologica Supplement*. 2004(432):1-45.
20. Schweitzer ME, Karasick D. MR Imaging of Disorders of the Achilles Tendon. *American Journal of Roentgenology*. 2000;175(3):613-25.

21. Schneebeli A, Del Grande F, Vincenzo G, Cescon C, Barbero M. Test-retest reliability of echo intensity parameters in healthy Achilles tendons using a semi-automatic tracing procedure. *Skeletal radiology*. 2017;46(11):1553-8.
22. Khan KM, Forster BB, Robinson J, Cheong Y, Louis L, Maclean L, et al. Are ultrasound and magnetic resonance imaging of value in assessment of Achilles tendon disorders? A two year prospective study. *Br J Sports Med*. 2003;37(2):149-53.
23. Hahn F, Meyer P, Maiwald C, Zanetti M, Vienne P. Treatment of chronic achilles tendinopathy and ruptures with flexor hallucis tendon transfer: clinical outcome and MRI findings. *Foot Ankle Int*. 2008;29(8):794-802.
24. DeCarbo WT, Bullock MJ. Midsubstance Tendinopathy, Surgical Management. *Clin Podiatr Med Surg*. 2017;34(2):175-93.
25. Singh A, Calafi A, Diefenbach C, Kreulen C, Giza E. Noninsertional Tendinopathy of the Achilles. *Foot Ankle Clin*. 2017;22(4):745-60.
26. Lin Y, Yang L, Yin L, Duan X. Surgical Strategy for the Chronic Achilles Tendon Rupture. *BioMed Research International*. 2016;2016:1416971.
27. Walther M, Dorfer B, Ishak B, Dreyer F, Mayer B, Roser A. [Reconstruction of extended defects of the Achilles tendon using a flexor hallucis longus tendon transfer.]. *Oper Orthop Traumatol*. 2011.
28. Longo UG, Ronga M, Maffulli N. Achilles Tendinopathy. *Sports Med Arthrosc Rev*. 2018;26(1):16-30.
29. Maffulli N, Oliva F, Maffulli GD, Buono AD, Gougoulas N. Surgical management of chronic Achilles tendon ruptures using less invasive techniques. *Foot Ankle Surg*. 2018;24(2):164-70.
30. Hunt KJ, Cohen BE, Davis WH, Anderson RB, Jones CP. Surgical Treatment of Insertional Achilles Tendinopathy With or Without Flexor Hallucis Longus Tendon Transfer: A Prospective, Randomized Study. *Foot & ankle international*. 2015;36(9):998-1005.
31. Kader D, Saxena A, Movin T, Maffulli N. Achilles tendinopathy: some aspects of basic science and clinical management. *Br J Sports Med*. 2002;36(4):239-49.
32. Tocci SL, Madom IA, Bradley MP, Langer PR, DiGiovanni CW. The diagnostic value of MRI in foot and ankle surgery. *Foot & ankle international*. 2007;28(2):166-8.
33. Garras DN, Raikin SM, Bhat SB, Taweel N, Karanjia H. MRI is unnecessary for diagnosing acute Achilles tendon ruptures: clinical diagnostic criteria. *Clinical orthopaedics and related research*. 2012;470(8):2268-73.

Tables

Table 1. Demographics of continuous and categorical data (n=120)

Tendinopathy (median [IQR])			
Variable	Yes (n=60)	No (n=60)	P-value
Age (y)	60 (16)	38 (31)	<0.001*
Gender			0.006†
Females	24 (40%)	39 (65%)	
Males	36 (60%)	21 (35%)	
Side			0.144†
Right	34 (57%)	26 (43%)	
Left	26 (43%)	34 (57%)	

*Wilcoxon rank sum test

†Chi-squared test

Abbreviations: IQR (interquartile range), y (years)

Table 2. Measurements comparing Achilles tendinopathy group to controls

Variable	Tendinopathy (median [IQR])		P-value*	OR (95% CI)	P-value†
	Yes (n=60)	No (n=60)			
Diameter (mm)					
FHL	29.5 (6.8)	26.5 (5.0)	<0.001	1.24 (1.06-1.44)	0.006
TS	14.8 (4.5)	16.0 (3.5)	0.006	0.82 (0.70-0.96)	0.015
FHL/TS	2.0 (0.8)	1.7 (0.3)	<0.001	9.56 (2.46-37.22)	0.001
Area (mm ²)					
FHL	545.0 (122.3)	453.3 (131.5)	0.080	1.01 (1.00-1.01)	0.002
TS	274.5 (182.0)	331.0 (140.5)	<0.001	1.00 (0.99-1.12)	0.236
FHL/TS	1.8 (1.3)	1.3 (0.7)	<0.001	2.52 (1.33-4.78)	0.005

*Wilcoxon rank sum test

†Wald test in a logistic regression model adjusting for age, sex, and side.

Abbreviations: IQR (interquartile range), OR (odds ratio), % (percent), CI (confidence interval), FHL (flexor hallucis longus), TS (triceps surae)

Table 3. Intraclass correlation coefficient.

Variable		ICC (95% CI)	MDC	SEM	P-value*
Diameter (mm)					
	FHL	0.75 (0.66-0.82)	2.55	0.92	<0.001
	TS	0.93 (0.90-0.95)	2.25	0.81	<0.001
	FHL/TS	0.87 (0.82-0.91)	0.53	0.19	<0.001
Area (mm ²)					
	FHL	0.80 (0.72-0.85)	149.35	53.88	<0.001
	TS	0.83 (0.76-0.88)	144.64	52.18	<0.001
	FHL/TS	0.83 (0.77-0.88)	1.14	0.41	<0.001

*F test for ICC.

Abbreviations: ICC (intraclass correlation coefficient), % (percent), CI (confidence interval), MDC (minimal detectable difference), SEM (standard error measurement), FHL (flexor hallucis longus), TS (triceps surae)

Note: Strength of correlation: 0.00-0.19=very weak; 0.20-0.39=weak; 0.40-0.59=moderate; 0.60-0.79=strong; 0.80-1.00=very strong

Figures




Figure 1

The height was determined on the sagittal plane by calculating the distance from the upper ankle joint line. The red arrow marks the neurovascular bundle.




Figure 2

Measurement of diameter and area of FHL and TS muscle bellies.




Figure 3

An example of two participants (left from Achilles tendinopathy group and on the right-side control group). Obvious quantitative differences can be observed.




Figure 4

Ratios of diameter and area compared between study groups. The boxplots represent medians of the ratios of FHL/TS (blue for diameter and gray for area).




Figure 5

Receiver operating characteristic (ROC) curve assessing diameter ratio FHL/TS. The area under the curve (AUC) is 0.70 (95 percent [%] confidence interval [CI] 0.65-0.81). Since the lower bound of the 95% CI is >0.50, it may be assumed that the model is suitable (i.e. superior to a random one). At a chosen cut-off point of a ratio of 2.0 and above, there was a sensitivity of 47% and specificity of 90% for concomitant signs of Achilles tendon tendinopathy.