Suicidal ideation, attempt, and determinants among medical students

Getachew Tesfaw Desalegn (✉ getachewtesfaw@gmail.com)

University of Gondar College of Medicine and Health Sciences

Mesele Wondie

University of Gondar College of Medicine and Health Sciences

Saron Dereje

Gondar University Hospital

Adanech Addisu

Gondar University Hospita

Research article

Keywords: Suicidal Ideation, Attempt, Medical Students, Prevalence

Posted Date: September 17th, 2019

DOI: <https://doi.org/10.21203/rs.2.14532/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background Suicidal ideation and attempt are highly prevalent among medical students compared to the general population and negatively impacts the quality of life, physical, and **mental** well being of students. However, research into suicidal ideation and attempt among medical students in low and middle-income countries are limited. Therefore, this study aimed to explore the **suicide** ideation and attempt and associated factors among medical students in Ethiopia. **Methods** An institution based cross-sectional study was conducted from March to June 2019 at the University of Gondar among medical students. A simple random sampling technique used to get study units by their identification card. **Suicide** ideation and attempt were assessed by using World Health Organization Composite International Diagnostic Interview (CIDI) to assess **suicide** ideation and attempt. Data were collected using self-administered questionnaires. Binary logistic regression analysis was used to identify associated factors between suicidal ideation and attempt with the explanatory variables at $p\text{-value} < 0.05$ was considered statistically significant. **Results** The prevalence of suicidal ideation and attempt among study participants was found to be 14% and 7.4% with 95% CI (10.9, 18.1) and (5.1, 10.2), respectively. In multivariate logistic regression; female sex (AOR: 5.21, 95% CI: 2.42, 11.20), depression (AOR: 10.12, 95% CI: 4.80, 21.52), current khat chewing (AOR: 4.46, 95% CI: 1.32, 15.02), and poor social support (AOR: 4.46, 95% CI: 1.43, 13.87) were factors significantly associated with suicidal ideation. Being female (AOR: 8.08, 95% CI: 3.04, 21.39), depression (AOR: 10.66, 95% CI: 4.01, 28.01) and history of **mental** illness (AOR: 5.53, 95% CI: 1.20, 25.50) were factors significantly associated with suicidal attempt. **Conclusion** In the current study, the prevalence of suicidal ideation among medical students was low compared to other studies, but the suicidal attempt was high. The University of Gondar better to instate a screening program for every medical student for early diagnosis and intervention of **suicide** ideation and attempt.

Background

Suicide is the act of deliberately causing one's own death (1). Suicidal ideation is thought of serving as the agent of one's own death while the suicidal attempt is self-injurious behavior with a non-fatal outcome accompanied by implicit evidence that person's intended to die (2, 3).

Globally, the rate of suicide has increased by 60% in the past 45 years and about one million people die from suicide every year. It is the second leading cause of death among young people and it is greater in males than in females. And Suicidal act results from social, cultural, psychological, biological, and environmental factors can interact to lead suicidal behavior (4, 5).

Suicide ideation, attempt, and psychological distress are common in university students

(6-10). The global prevalence of suicidal ideation and attempt from 36 college students were 23.3% and 3.2%, respectively (7). Mental distress among medical students and professionals are widespread (11). Suicidal behavior is a mental health problem affecting a remarkable proportion of medical students (12) and it is highly affected in medical students compared with the general population (13, 14). Most medical

students experienced greater risk for suicidal ideation and thoughts of dropping out of medical school (15). Physicians are one of the highest risk groups for suicide and which seems that this problem starting from medical school (2, 16-19)

The magnitude of suicidal ideation among medical students by including 13 Western and non-Western countries ranged from 1.8% to 53.6% (20). The prevalence of suicidal ideation and attempt among medical students in South Africa were 32.3% and 3.2%, respectively (21). In Ethiopia, the prevalence of suicidal ideation among the University students was 19.9%, but no study regarding suicidal attempts (13).

Suicidal ideation and depression are strong predictors of attempted suicide in medical students (4, 14). The rate of suicidal ideation and attempt among female medical students were higher than male medical students (5, 11). Different factors could play a role in the development of suicidal behavior. These include depression, burnout, sleep disorders, family history of mental illnesses, previous psychiatric disorders, years of study, gender, substance use, poor social support, lived alone, feeling of neglect by parents, something valued lost, breaking of a steady love relationship, and poor physical health (12, 13, 18, 20-22). The impact of suicidal behavior among medical students includes substance use, unhealthy peer relationship, fixation with death or violence, poor academic performance, dropout, and suicide attempt (23-25).

Even though suicidal behavior has a high prevalence among medical students and most of available research evidence comes from developed countries, little attention is given in lower-middle-income and lower-income countries (26). To the best of our knowledge, there has been no published study on suicidal ideation, attempt, and associated factors among medical students in Ethiopia. This study, therefore, aimed to investigate the magnitude and determinants of suicidal ideation and attempt among medical students at the University of Gondar with a view of informing the development of interventions.

Methods

Study design and period: An institution based cross-sectional study was conducted at the University of Gondar College of Medicine and Health Sciences, March 2019, Northwest Ethiopia.

Study Setting: The University of Gondar is one of the oldest Universities in Ethiopia. It was established in 1954 as the Public Health College and Training Center. It has 12 departments, including schools of medicine. The study was conducted on 1397 medical students. The University of Gondar, CHMS is located in Gondar town, which is 728 km far from the capital city Addis Ababa.

Study population: Undergraduate medical students at the University of Gondar College of Medicine and Health Sciences included in the sample and excluded critically ill students.

Sample size calculation and sampling procedures: The sample size was calculated by using the single population proportion formula with a 95% CI, a 5% margin of error, and suicidal ideation, and attempt of

50% because of no published work in our country. Assuming a 10% non-response rate, 423 students were recruited randomly by using the simple random sampling technique. The number of medical students in the college with their identification number taken from the UoG CMHS registrar office; the required sample was selected through lottery methods. The lists of dormitory students took from the UoG CMHS student's union dormitory affairs.

Data collection tools

Data were collected using self-administered questionnaire which contained suicidal ideation and attempt as the dependent variable and several other explanatory variables that included socio-demographic factors, social support, clinical factors, and substance use factors. Data on social support was assessed by Oslo social support scale (27). It has a three items social support scale (OSS-3) which provided a brief measure of social support and functioning and it is considered to be one of the best predictors of mental health. It covered different fields of social support by measuring the number of people the respondent feels close to the interest and concern shown by others, and the ease of obtaining practical help from others. Social support was collected by the Oslo 3-item social support scale which had a 3-item questionnaire commonly used to assess social support and used in several studies. The sum score scale ranges from 3 to 14, and had three broad categories: "poor support" 3-8, "moderate support" 9-11, and "strong support" 12-14(27).

Depression was assessed by using Patient Health Questionnaire (PHQ-9) and its score was greater or equal to 10; (5–9 mild depression, 10–14 moderate depression, 15–19 moderately severe depression, and 20–27 severe depression). PHQ-9 screening tool has nine items and was validated in Ethiopia with sensitivity of 86% and specificity of 67% (28).

Suicide ideation and attempt were measured according to the WHO questionnaire. If the respondent provided a "Yes" answers to the question, "Have you ever seriously thought about committing or attempted suicide, respectively?" they were considered to have had suicidal ideation or attempt, respectively (29).

Data processing and analysis

Data were entered into Epi-info version 7 software after checking for completeness and exported to SPSS version 20 for analysis. Univariate analysis was done to see the association of each independent variable with the dependent variable. Those variables a P-value less than 0.2 were entered into the multivariate logistic regression model to identify the effect of each independent variable with the outcome variables. The strength of the association presented by the adjusted odds ratio with a 95% Confidence interval, and a P-value less than 0.05 was considered statistically significant.

Socio-demographic characteristics

A total of 393 participants was included in the study with a response rate of 92.9%. The mean age of the respondents was 22.16 (± 1.86) years. Out of the participants, 241 (61.3%) were male; 254 (64.6%) were

the age of greater than 25 years, and 304 (77.4%) were coming from the urban residence. Nearly three-fourth (72%) of the participants was Orthodox Christian; and more than half (55.5%) their cumulative grade was between 2.76 to 3.5. The median monthly income of the participants was 1000 Birr and ranges from 50 to 6036 Ethiopian Birr (table 1).

Clinical and social characteristics

Of the participants, 50 (12.7%) had a family history of mental illness, 13 (3.3%) had a history of mental illness, 33 (8.4%) had a family history of suicidal attempt, and 122 (31%) had depression. Regarding social factors, 104 (26.5%), and 202 (51.4%) of the respondents had poor and moderate social supports (table 2).

Substance use characteristics

At the movement, 168 (42.7%) were drinking alcohol, 51 (13%) were chewing Chat, and 37 (9.4%) were taking tobacco. Of the participants, 205 (52.5%) were drinking alcohol, 66 (16.8%) were chewing Chat, and 45 (11.5%) had been taking tobacco ever use respectively (table3).

The prevalence of suicidal ideation and attempt

This study showed that the prevalence of suicidal ideation and attempt of the participants were 14% and 7.4%, with a 95% CI (10.9, 18.1) and (5.1, 10.2), respectively. Out of the participants, 29 (7.4%) had a suicidal plan (table 4).

Factors associated with suicidal ideation and attempt

Among all covariates, Female sex, depression, current and ever Chat chewing, current drinking alcohol, and smoking cigarettes, history of mental illness, family history of mental illness, and social support had less than 0.2 P-values in the bi-variable logistic regression and considered as the multivariable logistic regression model. The multivariate analysis suggested that the female sex was more than five times (95%; CI: 2.42, 11.20) more likely to have suicidal ideation compared with the male sex. Depression was ten times (95%; CI: 4.80, 21.52) more risky to develop suicidal ideation compared to counterparts. At the moment, chewing khat was 4.46 times (95%; CI: 1.32, 15.02) more risky for suicidal ideation compared to students who did not chew khat and having poor social support more than four times (95%; CI: 1.43, 13.87) to develop suicidal ideation compared to good social support (table 5). On the other hand, female sex was more than eight times (95%; CI: 3.04, 21.39) to develop suicidal attempt compared with counterparts and depression also has 10.66 times (95%; CI: 4.01, 28.01) higher risk to develop suicidal attempt compared to students who did not have depression. Finally, having a history of mental illness was about 5.53 times (95%; CI: 1.20, 25.50) more likely to increase suicidal attempt compared to counterparts (table 5).

Discussion

In the current study, the magnitude of suicidal ideation, attempt, and their possible association with various factors were assessed among medical students in Ethiopia for the first time. The prevalence of lifetime suicidal ideation and attempt were found to be 14% and 7.4%, respectively. Regarding prevalence of suicide ideation, our result is consistent with those of reported magnitude other studies carried out among medical students in Taiwan, China, Australia, and Turkey, in the prevalence estimated at 11.5%, 17.7%, 11.3%, 12%, respectively (22, 30, 31).

On the other hand, this finding is higher than those of the studies done in China, Pakistan, and Germany the magnitude estimated at 7.5%, (5, 14, 32). However, the prevalence of suicidal ideation in the current study is lower than the study done in South Africa among medical students 32.3% (21), in India 53.6% (33), Turkey 27.3%, and Austria 37.8%, (31) and the USA 29.9% (34). The variation may be due to distinctions study designs, sample size, and the socio-cultural variations between Ethiopia, and the other countries. In addition to the above, the difference was suicide ideation and reactions to suicide among medical students were cultural factors of the countries (31).

Regarding factors associated with suicidal ideation, female sex had five times more risk of suicidal ideation compared with male medical students. This was supported by those of other studies carried out a meta-analysis among the university students in China female students had higher suicidal thoughts than male students(10), in Pakistan female medical students a greater risk of suicidal ideation than male students(5), and another meta-analysis was done among medical students and medical professionals, psychological distress was higher for female physicians and medical students compared with males (11). Female physicians have a greater risk of suicide than other women (2).

In this study, medical students with depression were strongly associated with suicidal ideation. The odds of suicidal ideation were ten times higher who had depression than those without depression. Our results are consistent with findings reported in other studies that have shown medical students' depression was the most common predictor of suicidal ideation (11, 14).

In this study, current khat chewing was 4.46 times more risky for the development of suicidal ideation compared to students who had no suicidal ideation. Which was supported by other studies done among the university students of Ethiopia, Current khat chewing was nearly two times increased suicide (13), having drug misuse among medical students were a strong predictor of suicidal thoughts (20). Suicide risks among those who abuse substances were high in different studies (2).

Poor social support increased more than four times among medical students compared to their counterparts in this study. This is in line with a study done in Ethiopia (13). Lack of social support was two times risk of suicidal ideation among the University students in Ethiopia compared to students had moderate or good social support. Suicidal ideation was highly prevalence among people who had less social support from their family, friends, and other relatives(35).

On the other hand, our result of suicidal attempt was in line with those of other studies conducted among medical students. For example, a study in South Africa medical students reported magnitude was 6.2%

(21) and a cross cultural investigation among similar study participants in Turkey the prevalence of attempt estimated at 6.4% (33). Similarly, the prevalence of suicidal attempt among study participants was higher than those of other studies conducted in United Arab Emirates (30), Germany (32), Delhi (33), china (9) the prevalence was estimated at 1.8%, 1.4, 2.6%, 2.7%, respectively.

Our study revealed that the female sex was eight folds of attempts suicide among medical students compared to their counterparts. This is supported by a study done in Nepal, girls were nearly two folds of more likely to attempt suicide compared to boys (36). There were different studies documented in Uganda, South Africa, and Iran, females were more risky to have suicide attempt compared to males (37-39). Women attempt suicide or suicidal thoughts three folds as often as males (2).

In this study, depression and other history of mental illness were strong predictors for suicidal attempt. Depression is more than ten times risky for suicidal attempt compared with those who had no depression and history of mental disorders was more than fifth folds of suicidal attempt compared to students who had no history of mental illness. These were agreed from the previous studies done in South Africa (21), and (12). Previous diagnosis of depression or psychiatric disorders had strong correlations with suicidal attempt (21) and suicidal behavior or attempt was higher in medical students who have depression (12). Depression disorders accounts for 80% of suicidal attempt while previous psychiatric disorders risk for suicide is three to twelve times that of non-psychiatric patients, but the degree of risk varies depending on sex, age, diagnosis, and treatment (2). Since the questionnaire is self administered the participants may not give genuine response. A cross-sectional study design cannot permit conclusions for some variables, for example, to decide whether social anxiety symptoms are risks for or consequence. This study also was not included sixth year medical students so this is not representative all medical students. This finding is likely only to hint at the complex interactions between suicidal ideation and attempt with explanatory variables (risk factors).

Conclusions

The prevalence of suicidal ideation among medical students was low compared to other studies, but the suicidal attempt was high. Female sex, depression, current khat chewing, and poor social support were factors significantly associated with suicidal ideation while female sex, depression, and history of mental illness were factors significantly associated with suicidal attempt. The University of Gondar and Ministry of Health better to instate a screening program for every medical student for early diagnosis and intervention of suicide ideation and attempt. Further research on risk factors of suicidal ideation and attempt should be conducted to strengthen and broaden these findings.

Abbreviations

CI	Confidence Interval
CMHS	College of Medicine and Health Science
Epi-info	Epidemiological Information
QOL	Quality of Life
OR	Odds Ratio
SPSS	Statistical Package for Social Sciences
UAE	United Arab Emirates
UOG	University of Gondar
USA	United States of America

Declarations

Ethical approval and consent to participate

Ethical approval was obtained from the Institutional Review Board (IRB) of the University of Gondar Department of Psychiatry. The objectives and demands of the study were explained carefully. Participants who agreed to participate gave written consent. Confidentiality and anonymity were assured to the participants. The respondents did not write their name at the time of responding the questions. Selected participants for the study were informed of the general purpose, possible risks, and benefits. Participants were participated voluntarily and had the right not to answer any of the questions. To ensure confidentiality, participants' data were linked to a code number and registered.

Funding

The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors. The funder has no role in collection, analysis and interpretation of data and in writing the manuscript

Availability of data and materials

The dataset during and/or analyzed during the current study available from the corresponding author on reasonable requests

Authors' contributions

GT conceived the study and was involved in the study design, reviewed the article, analysis, report writing and drafted the manuscript. MW, SD, and AA were involved in the study design, analysis and drafted the manuscript. Both authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Consent for publication

Not applicable

Acknowledgments

First of all, we would like to thank the University of Gondar, College of Medicine and Health Sciences, Department of Psychiatry for organizing this research program. Secondly, our heartfelt thanks also extend to the University of Gondar, College of Medicine and Health Sciences, registrar office for their assistance to get the data of students to determine sample size and their help during a data collection time. Finally, we would like to thank our study participants for their cooperation in providing the information.

References

1. Kanchan T, Menezes RG. Suicidal hanging in Manipal, South India–victim profile and gender differences. *Journal of forensic and legal medicine*. 2008;15(8):493-6. DOI:[10.1016/j.jflm.2008.05.004](https://doi.org/10.1016/j.jflm.2008.05.004).
2. Sadock B, Ruiz P. Kaplan & Sadock's synopsis of psychiatry: behavioral sciences: Walters Kluwer; 2015.
3. Menezes RG, Subba S, Sathian B, Kharoshah MA, Senthilkumaran S, Pant S, et al. Suicidal ideation among students of a medical college in Western Nepal: a cross-sectional study. *Legal Medicine*. 2012;14(4):183-7. DOI:[10.1016/j.legalmed.2012.02.004](https://doi.org/10.1016/j.legalmed.2012.02.004)
4. Organization WH. Preventing suicide: A global imperative: World Health Organization; 2014.
5. Osama M, Islam MY, Hussain SA, Masroor SMZ, Burney MU, Masood MA, et al. Suicidal ideation among medical students of Pakistan: a cross-sectional study. *Journal of forensic and legal medicine*. 2014;27:65-8. DOI:[10.1016/j.jflm.2014.08.006](https://doi.org/10.1016/j.jflm.2014.08.006)
6. Eskin M, Sun J-M, Abuidhail J, Yoshimasu K, Kujan O, Janghorbani M, et al. Suicidal behavior and psychological distress in university students: a 12-nation study. *Archives of suicide research*. 2016;20(3):369-88. DOI:[10.1080/13811118.2015.1054055](https://doi.org/10.1080/13811118.2015.1054055)
7. Mortier P, Cuijpers P, Kiekens G, Auerbach R, Demyttenaere K, Green J, et al. The prevalence of suicidal thoughts and behaviours among college students: a meta-analysis. *Psychological medicine*. 2018;48(4):554-65. DOI:[10.1017/S0033291717002215](https://doi.org/10.1017/S0033291717002215)
8. Wang L, He CZ, Yu YM, Qiu XH, Yang XX, Qiao ZX, et al. Associations between impulsivity, aggression, and suicide in Chinese college students. *BMC public health*. 2014;14(1):551. DOI:[10.1186/1471-2458-14-551](https://doi.org/10.1186/1471-2458-14-551)
9. Yang L-S, Zhang Z-H, Sun L, Sun Y-H, Ye D-Q. Prevalence of suicide attempts among college students in China: a meta-analysis. *PLoS one*. 2015;10(2):e0116303.

<https://doi.org/10.1371/journal.pone.0116303>

10. Li Z-Z, Li Y-M, Lei X-Y, Zhang D, Liu L, Tang S-Y, et al. Prevalence of suicidal ideation in Chinese college students: a meta-analysis. *PLoS One*. 2014;9(10):e104368.
<https://doi.org/10.1371/journal.pone.0104368>
11. Onyishi M, Talukdar D, Sanchez R, Olaleye A, Medavarapu S. Prevalence of Clinical Depression among Medical Students and Medical Professionals: A Systematic Review Study. *Arch Med*. 2016;8:6. Doi: 10.21767/1989-5216.1000178
12. Coentre R, Faravelli C, Figueira ML. Assessment of depression and suicidal behaviour among medical students in Portugal. *International journal of medical education*. 2016;7:354-363.
doi:10.5116/ijme.57f8.c468
13. Dachew BA, Bifftu BB, Tiruneh BT, Anlay DZ, Wassie MA. Suicidal thoughts among university students in Ethiopia. *Annals of General Psychiatry*. 2016;17(1):1.<http://dx.doi.org/10.1037/t03506-000>
14. Sobowale K, Zhou AN, Fan J, Liu N, Sherer R. Depression and suicidal ideation in medical students in China: a call for wellness curricula. *International journal of medical education*. 2014;5:31.
DOI:[10.5116/ijme.52e3.a465](https://doi.org/10.5116/ijme.52e3.a465)
15. Dyrbye LN, Harper W, Durning SJ, Moutier C, Thomas MR, Massie Jr FS, et al. Patterns of distress in US medical students. *Medical teacher*. 2011;33(10):834-9. doi:10.3109/0142159X.2010.531158.
16. Shanafelt TD, Balch CM, Dyrbye L, Bechamps G, Russell T, Satele D, et al. Special report: suicidal ideation among American surgeons. *Archives of surgery*. 2011;146(1):54-62.
doi:10.1001/archsurg.2010.292.
17. Schernhammer ES, Colditz GA. Suicide rates among physicians: a quantitative and gender assessment (meta-analysis). *American Journal of Psychiatry*. 2004;161(12):2295-302.
DOI:[10.1176/appi.ajp.161.12.2295](https://doi.org/10.1176/appi.ajp.161.12.2295)
18. Dyrbye LN, Thomas MR, Massie FS, Power DV, Eacker A, Harper W, et al. Burnout and suicidal ideation among US medical students. *Annals of internal medicine*. 2008;149(5):334-41.
DOI:[10.7326/0003-4819-149-5-200809020-00008](https://doi.org/10.7326/0003-4819-149-5-200809020-00008)
19. Moutinho Coentre R, Luisa Figueira M. Depression and suicidal behavior in medical students: a systematic review. *Current Psychiatry Reviews*. 2015;11(2):86-101.
DOI: [10.2174/1573400510666140807005141](https://doi.org/10.2174/1573400510666140807005141)
20. Coentre R, Góis C. Suicidal ideation in medical students: recent insights. *Advances in medical education and practice*. 2018;9:873-880. Doi:10.2147/AMEPS162626
21. Van Niekerk L, Scribante L, Raubenheimer P. Suicidal ideation and attempt among South African medical students. *SAMJ: South African Medical Journal*. 2012;102(6):372-3.
doi:10.7196/SAMJ.5503.
22. Fan AP, Kosik RO, Mandell GA, Tran DTP, Cheng HM, Chen CH, et al. Suicidal ideation in medical students: who is at risk? *Annals of the Academy of Medicine-Singapore*. 2012;41(9):377.

23. Tyssen R, Vaglum P, Grønvold NT, Ekeberg Ø. Suicidal ideation among medical students and young physicians: a nationwide and prospective study of prevalence and predictors. *Journal of affective disorders*. 2001;64(1):69-79. [https://doi.org/10.1016/S0165-0327\(00\)00205-6](https://doi.org/10.1016/S0165-0327(00)00205-6)
24. Midtgaard M, Ekeberg Ø, Vaglum P, Tyssen R. Mental health treatment needs for medical students: a national longitudinal study. *European Psychiatry*. 2008;23(7):505-11. DOI: 10.1016/j.eurpsy.2008.04.006.
25. Mandal A, Ghosh A, Sengupta G, Bera T, Das N, Mukherjee S. Factors affecting the performance of undergraduate medical students: A perspective. *Indian journal of community medicine: official publication of Indian Association of Preventive & Social Medicine*. 2012;37(2):126-129. DOI:10.4103/0970-0218.96104
26. Breet E, Goldstone D, Bantjes J. Substance use and suicidal ideation and behaviour in low-and middle-income countries: a systematic review. *BMC public health*. 2018;18(1):549. DOI:10.1186/s12889-018-5425-6.
27. Bøen H. Characteristics of senior centre users–and the impact of a group programme on social support and late-life depression. *Norsk epidemiologi*. 2012;22(2).
28. Gelaye B, Williams MA, Lemma S, Deyessa N, Bahretibeb Y, Shibre T, et al. Validity of the patient health questionnaire-9 for depression screening and diagnosis in East Africa. *Psychiatry research*. 2013;210(2):653-61. DOI: 10.1016/j.psychres.2013.07.015.
29. Kessler RC, Üstün TB. The world mental health (WMH) survey initiative version of the world health organization (WHO) composite international diagnostic interview (CIDI). *International journal of methods in psychiatric research*. 2004;13(2):93-121.
30. Amiri L, Voracek M, Yousef S, Galadari A, Yammahi S, Sadeghi M-R, et al. Suicidal behavior and attitudes among medical students in the United Arab Emirates. *Crisis*. 2013. Doi: 10.1027/0227-5910/a000170.
31. Eskin M, Voracek M, Stieger S, Altinyazar V. A cross-cultural investigation of suicidal behavior and attitudes in Austrian and Turkish medical students. *Social psychiatry and psychiatric epidemiology*. 2011;46(9):813-23. DOI:10.1007/s00127-010-0254-7
32. Wege N, Muth T, Li J, Angerer P. Mental health among currently enrolled medical students in Germany. *Public Health*. 2016;132:92-100. Doi: 10.1016/j.puhe.2015.12.014.
33. Goyal A, Kishore J, Anand T, Rathi A. Suicidal ideation among medical students of Delhi. *J Ment Health Human Behavior*. 2012;17(1):60-9.
34. MacLean L, Booza J, Balon R. The impact of medical school on student mental health. *Academic Psychiatry*. 2016;40(1):89-91. Doi:10.1007/s40596-015-0301-5.
35. Endo G, Tachikawa H, Fukuoka Y, Aiba M, Nemoto K, Shiratori Y, et al. How perceived social support relates to suicidal ideation: A Japanese social resident survey. *International Journal of Social Psychiatry*. 2014;60(3):290-8. <https://doi.org/10.1177/0020764013486777>

36. Pandey AR, Bista B, Dhungana RR, Aryal KK, Chalise B, Dhimal M. Factors associated with suicidal ideation and suicidal attempts among adolescent students in Nepal: Findings from Global School-based Students Health Survey. PloS one. 2019;14(4):e0210383. doi: 10.1371/journal.pone.0210383.
37. Rudatsikira E, Muula AS, Siziya S, Twa-Twa J. Suicidal ideation and associated factors among school-going adolescents in rural Uganda. BMC psychiatry. 2007;7(1):67.DOI: 10.1186/1471-244X-7-67.
38. Meehan S-A, Broom Y. Analysis of a national toll free suicide crisis line in South Africa. Suicide and Life-Threatening Behavior. 2007;37(1):66-78.
39. Lari AR, Joghataei MT, Adli YR, Zadeh YA, Alaghebandan R. Epidemiology of suicide by burns in the province of Isfahan, Iran. Journal of burn care & research. 2007;28(2):307-11.

Tables

Table 1 Socio demographic characteristics of respondents among medical students at University of Gondar, northwest Ethiopia, 2019(n=393).

Variables	Categories	Frequency	Percent
Sex	Male	241	61.3
	Female	152	38.7
Age	19-24	139	35.4
	≥25	254	64.6
Religion	Orthodox	283	72.0
	Muslim	41	10.4
	Protestant	54	13.7
	Other*	15	3.8
Marital status	Married	25	6.4
	Single	368	93.6
Ethnicity	Amhara	205	52.1
	Oromo	91	23.3
	Tigrie	56	14.2
	Other**	41	10.4
Residency	Urban	304	77.4
	Rural	89	22.6
Year of study	1 st	28	7.1
	2 nd	73	18.6
	3 rd	107	27.2
	4 th	106	27.0
	5 th	79	20.1
CGPA	2.00-2.75	59	15.0
	2.76-3.5	218	55.5
	>3.5	116	29.5
Monthly income	50-1800	323	82.2
	>1800	70	17.8

s

Note: others*: Catholic, waqefeta, atheist

Others**: Gurage, Somali, Afar

Table 2 Clinical and social characteristics of respondents among medical students at University of Gondar, northwest Ethiopia, 2019 (n=393).

Variables	Categories	Frequency	Percent
Family history of mental illness	Yes	50	12.7
	No	343	87.3
History of mental illness	Yes	13	3.3
	No	380	96.7
History of chronic illness	Yes	25	6.4
	No	368	93.6
Family history of Suicide attempt	Yes	33	8.4
	No	360	91.6
Depression	No	271	69
	Yes	122	31
Social support	Poor	104	26.5
	Medium	202	51.4
	Good	87	22.1

Table3. Substance use characteristics of respondents among medical students at University of Gondar, Northwest Ethiopia, 2019(n=393).

Variables	Categories	Ever use		Current use	
		Frequency	Percent	Frequency	Percent
Alcohol	No	188	47.8	225	57.3
	Yes	205	52.5	168	42.7
Chat	No	327	83.2	342	87
	Yes	66	16.8	51	13
Cigarette	No	348	88.5	356	90.6
	Yes	45	11.5	37	9.4s
Others*	No	387	98.5	386	98.2
	Yes	6	1.5	7	1.8

NB: Other*: Cannabis and Shisha

Table 4 Suicidal ideation, plan, and attempt characteristics of respondents among medical students at University of Gondar, Northwest Ethiopia, 2019(n=393).

Variables	Categories	Frequency	Percent
Lifetime suicidal ideation	Yes	55	14.0
	No	338	86.0
Suicidal plan	Yes	29	7.4
	No	364	92.6
Lifetime suicidal attempt	Yes	29	7.4
	No	364	92.6

Table 5 Bi-variable and multivariate analysis of suicidal ideation among medical students at the University of Gondar, Northwest Ethiopia, 2019 (N=393).

Variables	Categories	Suicidal ideation		COR (95%, CI)	AOR (95%, CI)	P-value
		Yes	No			
Sex	Male	26	215	1	1	0.001
	Female	29	123	1.95(1.10,3.46)	5.21(2.42,11.20)	
Family history of mental illness	Yes	14	36	2.86(1.42,5.76)	1.88(0.80,4.41)	0.149
	No	41	302	1	1	
History of mental illness	Yes	4	9	2.87(0.85,9.66)	2.40(0.57,10.15)	0.23
	No	51	329	1	1	
Depression	Yes	40	82	8.32(4.38,15.84)	10.12(4.80, 21.52)	0.001
	No	15	256	1	1	
Current alcohol	Yes	32	136	2.07(1.16,3.68)	1.57(0.76,3.24)	0.39
	No	23	202	1	1	
Current khat	Yes	16	35	3.55(1.80,7.00)	4.46(1.32, 15.02)	0.01
	No	39	303	1	1	
Current cigarette	Yes	10	27	2.56(1.16,5.64)	0.53(0.13, 2.19)	0.39
	No	45	311	1	1	
Social support	Poor	25	79	5.20(1.89,14.23)	4.46(1.43, 13.87)	0.01
	Moderate	25	177	2.31(0.86,6.27)	1.95(0.67, 5.81)	0.22
	Good	5	82	1	1	

Table 6 Bi-variable and multivariate analysis of suicidal attempt among medical students at University of Gondar, Northwest Ethiopia, 2019(n=393).

Variables	Categories	Suicidal attempt		COR(95%, CI)	AOR(95%, CI)	P-value
		Yes	No			
Sex	Male	10	231	1	1	0.001
	Female	19	133	3.30(1.49,7.30)	8.08(3.04,21.39)	
Family history of mental illness	Yes	7	43	2.38(0.96,5.89)	1.58(0.54,4.61)	0.40
	No	22	321	1	1	
History of mental illness	Yes	4	9	6.31(1.81,21.93)	5.53(1.20,25.50)	0.02
	No	25	355	1	1	
Depression	Yes	21	101	6.83(2.93,15.92)	10.66(4.01, 28.01)	0.001
	No	8	263	1	1	
Ever alcohol use	Yes	20	185	2.15(0.95,4.84)	0.89(0.16, 4.96)	0.89
	No	9	179	1	1	
Ever Chat use	Yes	9	57	2.42(1.05,5.60)	0.68(0.03,12.91)	0.80
	No	20	307	1	1	
Ever cigarette use	Yes	7	38	2.73(1.09,6.81)	1.37(0.07,25.86)	0.83
	No	22	326	1	1	
Current alcohol use	Yes	18	150	2.33(1.07,5.09)	2.39(0.42, 13.35)	0.32
	No	11	214	1	1	
Current Chat use	Yes	8	43	2.84(1.19,6.81)	2.20(0.09,51.70)	0.84
	No	21	321	1	1	
Current cigarette use	Yes	6	31	2.80(1.06,7.40)	1.45(.06,36.00)	0.82
	No	23	333	1	1	