Fetal Cardiac Parameters for Predicting Postnatal Operation Type of Fetuses With Tetralogy of Fallot

Suyeon Park

Hallym University Sacred Heart Hospital <https://orcid.org/0000-0002-4199-5011>

Hye-Sung Won

Asan Medical Center

Rina Kim

Jeju National University Hospital

Mijin Kim

Asan Medical Center

Jeong Jin Yu

Asan Medical Center

Chun Soo Park

Asan Medical Center

Tae-Jin Yun

Asan Medical Center

Yewon Jung

Chungnam National University Hospital

Usamah Al Harbi

Asan Medical Center

Mi-Young Lee (✉ poptwinkle@hanmail.net)

University of Ulsan College of Medicine, Asan Medical Center

Research

Keywords: fetal echocardiography, pulmonary artery, tetralogy of Fallot

Posted Date: May 18th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-524340/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Cardiovascular Ultrasound on February 21st, 2022. See the published version at <https://doi.org/10.1186/s12947-022-00274-5>.

Abstract

Background: To assess fetal cardiac parameters predictive of postnatal operation type in fetuses with tetralogy of Fallot (TOF).

Methods: Echocardiographic data obtained in the second and third trimesters were retrospectively reviewed for fetuses diagnosed with TOF between 2014 and 2018 at Asan Medical Center. The following fetal cardiac parameters were analyzed: 1) pulmonary valve annulus (PVA) z-score, 2) right pulmonary artery (RPA) z-score, 3) aortic valve annulus (AVA) z-score, 4) pulmonary valve peak systolic velocity (PV-PSV), 5) PVA/AVA ratio, and 6) RPA/descending aorta (DAo) ratio. These cardiac parameters were compared between a primary corrective surgery group and a palliative shunt operation followed by complete repair group.

Results: A total of 100 fetuses with TOF were included. Only one neonatal death occurred. Ninety patients underwent primary corrective surgery and 10 neonates underwent a multistage surgery. The PVA z-score, RPA z-score, and RPA/DAo ratio measured in the second trimester and the PVA z-score, RPA z-score, and PVA/AVA ratio measured in the third trimester were significantly lower in the multistage surgery group, while the PV-PSV as measured in both trimesters were significantly higher in the multistage surgery group.

Conclusion: Fetal cardiac parameters are useful for predicting the operation type necessary for neonates with TOF.

Background

Tetralogy of Fallot (TOF) is the most common form of cyanotic heart disease, with an incidence of one per 3600 live births and affecting 5–7% of infants with congenital heart diseases [1] [2, 3]. The diagnosis of TOF is commonly made during the fetal period with a high degree of accuracy [1, 3]. TOF is described by four classic anatomical characteristics: outlet ventricular septal defect (VSD) with anterior malalignment, overriding aorta, pulmonary stenosis (PS), and right ventricular hypertrophy. According to the degree of right ventricular outflow tract (RVOT) obstruction, TOF can manifest as a variety of clinical presentations ranging from asymptomatic detection to early postnatal cyanosis requiring urgent treatment. Ultimately, TOF with the exception of pulmonary atresia with VSD can be divided into two types depending on the size of the pulmonary artery (PA) and the type of surgery. Type 1 patients, who have normal-sized PA and mild PS, usually undergo primary corrective surgery in the first year of life [4, 5] [6]. However, type 2 patients, who have small-sized PA and severe cyanosis, require multistage surgery consisting of, first, a palliative shunt operation that is subsequently followed by complete repair in later months [4–6].

In the fetal period, information on predictive factors to determine the operation type necessary in neonates with TOF is limited. In this study, we evaluated the cardiac parameters of fetuses prenatally diagnosed and postnatally confirmed to have TOF to support the prediction of postnatal operation type.

Materials And Methods

This was a retrospective cohort study of fetuses prenatally diagnosed and postnatally confirmed to have TOF at Asan Medical Center, Seoul, Korea between January 2014 and December 2018. The study protocol was approved by the institutional review board of Asan Medical Center (approval no. 2020 – 0463). Formal consent was not required because it was a retrospective study. All prenatal ultrasonographic examinations were performed using a WS80A (Samsung Medison Co., Ltd., Seoul, Korea) or a Voluson E8 or E10 Expert (GE Healthcare Austria GmbH & Co. OG, Zipf, Austria) with a 2- to 6-MHz transabdominal probe. The inclusion criteria were patients with simple TOF who were followed up longitudinally with at least twice by echocardiographic evaluations in the second and third trimesters, respectively, and who were born at our center. Patients with combined minor cardiac anomalies, such as right aortic arch, aberrant right or left subclavian artery, or persistent left superior vena cava, were also included. Meanwhile, patients with pulmonary atresia with VSD, an absence of pulmonary valve syndrome, or other associated complex cardiac anomalies were excluded.

Relevant prenatal and postnatal data collected for this study included the maternal age; gestational age (GA) at diagnosis, follow-up, and delivery; rate of preterm birth; birth weight; Apgar score at one minute and five minutes; associated chromosomal and extracardiac anomalies; and occurrence of neonatal death.

Fetal echocardiography was conducted in all cases. The following fetal cardiac parameters were analyzed: 1) pulmonary valve annulus (PVA) z-score, 2) right PA (RPA) z-score, 3) aortic valve annulus (AVA) z-score, 4) pulmonary valve peak systolic velocity (PV-PSV), 5) PVA/AVA ratio, and 6) RPA/descending aorta (DAo) ratio (modified McGoon ratio). Valve annuli were measured from hinge point to hinge point at the location of maximal expansion in the long-axis view of the heart during peak systole (Figs. 1A and 1C) [7, 8]. The RPA was measured in either the three-vessel view or oblique short-axis view, yielding a longitudinal view of the PA and its bifurcation (Fig. 1B). Pulsed Doppler was adopted to assess the PV-PSV in the ductal arch view or right long-axis view (Fig. 1A and 1D). Doppler velocity was recorded with the angle of insonation not exceeding 15° and with at least five uniform waveforms required for the measurements. The DAo was measured in the aortic arch view at the level of the diaphragm (Fig. 1E). Z-scores were calculated using normative data adjusted to GA [7]. During the study period, several experts in fetal echocardiography performed examinations using the same methodology. A single investigator (S. P.) retrospectively re-measured the following fetal cardiac parameters twice using the stored images and the mean values of such were used for the analysis.

All patients underwent either single-stage (primary corrective surgery) or multistage surgery (palliative shunt operation followed by complete repair). The choice of operation was made by pediatric cardiologists and pediatric cardiac surgeons, depending on the severity of the RVOT obstruction, including clinical symptoms (e.g., cyanosis and spell events) and the McGoon ratio and the Nakata index as measured by postnatal transthoracic echocardiography or cardiac computed tomography (CT). The McGoon ratio was calculated as the sum of diameter of the left PA (LPA) and RPA at the prebranching

point divided by the diameter of the DAo at the level of the diaphragm, while the Nakata index is the sum of the area of the LPA and RPA divided by the body surface[9, 10].

We compared fetal cardiac parameters according to the type of surgery—namely, single- or multistage surgery. Furthermore, we conducted a subgroup analysis in the single-stage surgery group according to the need for a secondary procedure or reoperation due to subsequent RVOT obstruction. We also compared the cardiac parameters between the two groups.

Statistical analysis

Descriptive statistics were calculated, with continuous data presented as means $\pm$ standard deviations or medians (interquartile ranges) and categorical variables presented as numbers (percentages). Parametric testing was used to compare data with normal distributions and comparisons were performed using the Student's *t*-test. A receiver-operating characteristic (ROC) curve was used to obtain the cutoff value of each fetal cardiac parameter. The indicator of accurate prediction was the area under the curve (AUC). The cutoff value of each parameter was determined according to sensitivity and specificity. Data were assessed with the IBM SPSS Statistics version 26.0 software program (IBM Corp., Armonk, NY). *P*-values < 0.05 were considered to be statistically significant.

Results

A total of 100 patients met the inclusion criteria for this study and the baseline characteristics are demonstrated in Table 1. Twelve neonates (12.0%) showed clinical symptoms related to RVOT obstruction immediately after birth, thus requiring medical support. All included patients underwent either single- (N = 90) or multistage surgery (N = 10). Among the 40 fetuses who underwent karyotyping, all showed normal results. Among the 31 fetuses undergoing multiplex ligation-dependent probe amplification for the detection of microdeletion syndromes, three were confirmed to have 22q11.2 deletion. 24 fetuses had right aortic arch and only one among CATCH 22 fetus had right aortic arch. Extracardiac anomalies were found in eight patients (8.0%)—these included renal diseases in three, gastrointestinal diseases in two, brain anomaly in one, spinal anomaly in one, and VACTERL association in one patient(s), respectively.

Table 1
Prenatal and postnatal characteristics of the fetuses with tetralogy of Fallot

Variable	Study population (N = 100)
Maternal age (years)	32 (23–43)
GA at diagnosis (weeks)	23.4 (19.6–27.6)
GA at follow-up (weeks)	32.5 (28.5–38.2)
GA at delivery (weeks)	39 (35.5–40.6)
Preterm birth	6 (6.0)
Birth weight (g)	3075 (1960–4182)
Apgar score at one minute (≥ 7)	90 (90)
Apgar score at five minutes (≥ 7)	100 (100)
Abnormal karyotype	0/40 (0)
22q11.2 deletion	3/31 (9.7)
Extracardiac anomalies	8 (8.0)
Neonatal death	1 (1.0)
Data are presented as medians (ranges) or numbers (percentages). GA, gestational age.	

Ninety patients underwent single-stage surgery at a median age of five months, while 10 underwent multistage surgery; in this latter group, the palliative shunt operation was conducted at a median age of three months and complete repair was performed at a median age of 10 months. Only one neonatal death occurred in an intrauterine growth-restricted newborn delivered at 38.3 weeks of gestation, who died as a result of sepsis after surgical correction.

Comparison of parameters according to operation type

Fetal cardiac parameters measured in the second and third trimesters are described in Tables 2 and 3, respectively, where they are classified into the single- and multistage surgery groups. The PVA z-score, RPA z-score, and RPA/DAo ratio measured in the second trimester and the PVA z-score, RPA z-score, and PVA/AVA ratio measured in the third trimester were significantly lower in the multistage surgery group, while the PV-PSV as measured in both trimesters were significantly higher in the multistage surgery group.

Table 2
Comparison of parameters according to operation type at the second trimester

Parameter	Study population (N = 100)	Single-stage surgery (N = 90)	Multi-stage surgery (N = 10)	P-value
PVA z-score	-3.1 ± 1.6	-3.0 ± 1.6	-4.3 ± 1.4	0.014
RPA z-score	-0.5 ± 1.1	-0.5 ± 1.1	-1.2 ± 0.7	0.029
AVA z-score	0.6 ± 1.2	0.6 ± 1.2	0.4 ± 1.4	0.625
PV-PSV (m/s)	1.1 ± 0.2	1.1 ± 0.2	1.3 ± 0.2	0.001
PVA/AVA	0.77 ± 0.15	0.78 ± 0.14	0.70 ± 0.20	0.152
RPA/DAo	0.73 ± 0.14	0.75 ± 0.13	0.62 ± 0.11	0.004
Data are expressed as means ± standard deviations. PVA, pulmonary valve annulus; RPA, right pulmonary artery; AVA, aortic valve annulus; PV-PSV, pulmonary valve peak systolic velocity; DAo, descending aorta.				

Table 3
Comparison of parameters according to operation type at the third trimester

Parameter	Study population (N = 100)	Single-stage surgery (N = 90)	Multi-stage surgery (N = 10)	P-value
PVA z-score	-3.2 ± 1.3	-3.0 ± 1.1	-4.9 ± 1.3	< 0.001
RPA z-score	-0.5 ± 0.8	-0.5 ± 0.8	-1.1 ± 0.8	0.022
AVA z-score	1.0 ± 1.0	1.0 ± 1.0	1.1 ± 1.0	0.732
PV-PSV (m/s)	1.3 ± 0.3	1.2 ± 0.3	1.5 ± 0.2	0.002
PVA/AVA	0.75 ± 0.13	0.77 ± 0.12	0.60 ± 0.10	< 0.001
RPA/DAo	0.71 ± 0.10	0.72 ± 0.10	0.63 ± 0.10	0.058
Data are expressed as means ± standard deviations. PVA, pulmonary valve annulus; RPA, right pulmonary artery; AVA, aortic valve annulus; PV-PSV, pulmonary valve peak systolic velocity; DAo, descending aorta.				

The ROC curves for the abovementioned meaningful parameters are presented in Fig. 2. The AUC values of all these parameters were greater than 0.700 and the PVA z-score measured in the third trimester achieved the highest AUC value of all (0.886). The optimal cutoff value of the PVA z-score measured in the third trimester was - 3.8 (sensitivity = 90.0% and specificity = 81.1%), while the optimal cutoff value of

the RPA/DAo ratio measured in the second trimester was 0.63 (sensitivity = 70.0% and specificity = 77.8%).

Subgroup analysis

The single-stage surgery group was further analyzed in accordance with the need for a secondary procedure or reoperation after the primary corrective surgery (Table 4). Of all patients in the group, 82 (91.1%) did not require a second procedure or operation because no complications occurred, while the remaining eight patients (8.9%) underwent balloon valvuloplasty or reoperation because of subsequent PA narrowing at the median age of 27 months. The PVA z-score and RPA z-score measured in both trimesters, PVA/AVA ratio in the third trimester, and RPA/DAo ratio in the second trimester were significantly different between the two groups. However, there was no significant difference in the PV-PSVs between the two groups when measured in either the second or third trimesters.

Table 4
Comparison of parameters according to the need for second procedure or reoperation

Parameter		Observation (N = 82)	Procedure or reoperation (N = 8)	P-value
PVA z-score	Second trimester	-2.9 ± 1.6	-4.1 ± 1.0	0.049
	Third trimester	-2.8 ± 1.0	-4.4 ± 0.9	< 0.001
RPA z-score	Second trimester	-0.4 ± 1.1	-1.2 ± 0.0	0.031
	Third trimester	-0.4 ± 0.7	-1.2 ± 1.0	0.004
PV-PSV (m/s)	Second trimester	1.1 ± 0.2	1.2 ± 0.2	0.342
	Third trimester	1.2 ± 0.3	1.2 ± 0.2	0.927
PVA/AVA	Third trimester	0.78 ± 0.11	0.65 ± 0.13	0.003
RPA/DAo	Second trimester	0.76 ± 0.13	0.64 ± 0.11	0.021
Data are expressed as means ± standard deviations.				
PVA, pulmonary valve annulus; RPA, right pulmonary artery; PV-PSV, pulmonary valve peak systolic velocity; AVA, aortic valve annulus; DAo, descending aorta.				

Discussion

Our study demonstrates that several cardiac parameters could have predictive values for determining the postnatal operation type for TOF fetuses with the largest number of patients. The PVA z-score, RPA z-score, and PV-PSV in the second and third trimesters; PVA/AVA ratio in the third trimester; and RPA/DAo ratio in the second trimester are useful markers for the prediction of the postnatal operation type. Particularly, a PVA z-score of - 3.2 or less in the second trimester and - 3.8 or less in the third trimester; an RPA z-score of - 0.5 or less in the second trimester and - 0.6 or less in the third trimester; and a PV-PSV of

1.2 m/s or greater in the second trimester and 1.4 m/s or greater in the third trimester were associated with a high probability of requiring a palliative shunt operation followed by complete repair, whereas when these values were conversely above or below the mentioned cutoff, only single-stage surgery was required. Furthermore, we also suggest that a PVA/AVA ratio of 0.63 or less in the third trimester, and an RPA/DAo ratio—the new cardiac parameter suggested for the first time in the fetal period—of 0.63 or less in the second trimester may predict the need for a multistage surgery.

Our findings are consistent with those of previous studies that evaluated prenatal echocardiographic markers that may predict the outcomes of fetuses with TOF[11–13] [14]. Several studies reported that fetuses who required neonatal intervention had lower PVA z-scores and PVA/AVA ratios[11, 13] [14], and other research suggested that a PV-PSV of 144.5 cm/s or greater measured at 34 to 38 weeks of gestation supported the accurate prediction of early intervention and placement of transannular patches in 23 TOF fetuses [12]. However, in contrast with previous reports that did not report any significance of fetal RPA z-score in the prediction of postnatal outcomes of TOF fetuses[12, 13], our study found that the RPA z-score as measured in the second and third trimesters could also be a marker for predicting the postnatal operation type.

The RPA/DAo ratio is derived from the neonatal McGoon ratio, which is calculated using neonatal CT angiography and is used for quantifying the degree of PA hypoplasia[9] [10]. In the neonatal period, pediatric cardiologists commonly rely on this ratio to determine the best operation type for neonates with TOF. Hence, in this study, we created a new cardiac parameter by modifying the neonatal McGoon ratio, in which we only included the RPA rather than the sum of the LPA and RPA. RPA can be easily measured in the three-vessel view because it originates at the right angle and runs behind the ascending aorta, whereas the LPA runs the same course as the main PA, making it challenging to differentiate between the LPA and ductus arteriosus. Furthermore, as this study was retrospective in nature, many cases did not have LPA images available in their records.

During gestation, the affected structures in fetuses with TOF may change progressively[15]. For this reason, echocardiographic evaluations performed at later stages of pregnancy could contribute more so to elucidating key differences than those measured at earlier stages. However, our results did not fully correlate with this concept. This may be because accurate measurements are sometimes difficult to collect at advanced GA in that ultrasonographic scans may fail to provide accurate information because of poor image quality due to fetal positioning or ossification of the fetal chest. Furthermore, retrospective nature of the study potentially affects this discordancy.

All infants underwent either single- or multistage surgery, with all but one of the patients included in this study surviving. As such, we confirmed an excellent prognosis associated with TOF. Our study also found that several infants who underwent single-stage surgery developed secondary pulmonary obstruction, requiring balloon valvuloplasty or surgical correction. A previous study reported that the PVA z-score could predict the need for reintervention after primary surgical correction[14]. In the present study, we found that not only the PVA z-score but also the RPA z-score, PVA/AVA ratio, and RPA/DAo ratio are

valuable parameters in predicting the need for a second procedure or reoperation. Meanwhile, the PV-PSV as measured in both trimesters did not show statistical significance; however, because there were only eight patients who required further intervention, it might have been difficult to obtain significant findings. Future studies should be conducted involving larger numbers of patients.

This study has several strengths. First, it included the largest sample size to date compared to other similar studies at single center [12] [16] [17]. Second, the data were collected longitudinally from the second trimester to the third trimester and demonstrated that changes in cardiac parameters occurred with advancing GA. Through these longitudinally collected data, we compared various fetal cardiac parameters at each trimester and suggested the cutoff values of various parameters relevant at each trimester for predicting the type of surgical operation required. Third, we only included postnatally confirmed TOF, excluding pulmonary atresia with VSD and other diseases that cause outflow obstruction, to avoid heterogeneity of the study population. In addition, we suggested a new parameter (the modified McGoon ratio) for predicting the operation type for fetuses with TOF.

However, this study also has several limitations. First, it was a retrospective study conducted at a single center. In addition, the analysis was limited to only include patients with available images. Furthermore, only one investigator selected the most appropriate image and measured the cardiac parameters. As such, selection bias was possible.

Conclusion

In conclusion, PVA z-score, RPA z-score, PV-PSV, PVA/AVA ratio, and RPA/DAo ratio demonstrate the potential to be useful parameters for predicting the necessary operation type for neonates with TOF. This information is clinically important because it can facilitate prenatal counselling and provide accurate data on the prognosis of neonates with TOF.

Abbreviations

TOF: Tetralogy of Fallot; PVA: Pulmonary valve annulus; RPA: Right pulmonary artery; AVA: Aortic valve annulus; PV-PSV: Pulmonary valve-peak systolic velocity; DAo: Descending Aorta; VSD: Ventricular septal defect; PS: Pulmonary stenosis; RVOT: Right ventricular outflow tract; PA: Pulmonary artery; GA: Gestational age; ROC: Receiver-operating characteristic; AUC: Area under the curve.

Declarations

Ethical approval and consent to participate

All data were anonymously analysed without individual patient consent owing to the retrospective nature of the study. This study protocol was approved by the institutional review board of Asan Medical Center (approval no. 2020-0463).

Consent for publication

Not applicable

Availability of data and materials

The data and materials in this study are available from the corresponding author on request.

Competing interests

No potential conflict of interest was reported by the authors

Funding

None.

Authors' contributions

Suyeon Park; designed and performed experiments, analyzed data, and wrote the paper. Mi-Young Lee; supervised the research and final approval of the version to be published. Hye-Sung Won; supervised the research. Mijin Kim, Jeong Jin Yu, Chun Soo Park, and Tae-Jin Yun; conceptualized the research and reviewed the manuscript. Rina Kim, Yewon Jung, and Usamah Al Harbi; evaluated prenatal data and performed statistics.

Acknowledgments

None

References

1. Mitchell SC, Sellmann AH, Westphal MC, Park J: **Etiologic correlates in a study of congenital heart disease in 56,109 births.** *Am J Cardiol* 1971, **28**(6):653-657.
2. Poon LC, Huggon IC, Zidere V, Allan LD: **Tetralogy of Fallot in the fetus in the current era.** *Ultrasound Obstet Gynecol* 2007, **29**(6):625-627.
3. Galindo A, Mendoza A, Arbues J, Graneras A, Escribano D, Nieto O: **Conotruncal anomalies in fetal life: accuracy of diagnosis, associated defects and outcome.** *Eur J Obstet Gynecol Reprod Biol* 2009, **146**(1):55-60.

4. Ooi A, Moorjani N, Baliulis G, Keeton BR, Salmon AP, Monro JL, Haw MP: **Medium term outcome for infant repair in tetralogy of Fallot: Indicators for timing of surgery.** *Eur J Cardiothorac Surg* 2006, **30**(6):917-922.
5. Steiner MB, Tang X, Gossett JM, Malik S, Prodhan P: **Timing of complete repair of non-ductal-dependent tetralogy of Fallot and short-term postoperative outcomes, a multicenter analysis.** *J Thorac Cardiovasc Surg* 2014, **147**(4):1299-1305.
6. Savla JJ, Fisher BT, Faerber JA, Huang YV, Mercer-Rosa L: **Complete Versus Staged Repair for Neonates With Tetralogy of Fallot: Establishment and Validation of a Cohort of 2235 Patients Using Detailed Surgery Sequence Review of Health Care Administrative Data.** *Med Care* 2018, **56**(11):e76-e82.
7. Schneider C, McCrindle BW, Carvalho JS, Hornberger LK, McCarthy KP, Daubeney PE: **Development of Z-scores for fetal cardiac dimensions from echocardiography.** *Ultrasound Obstet Gynecol* 2005, **26**(6):599-605.
8. van Ark AE, Molenschot MC, Wesseling MH, de Vries WB, Strengers JLM, Adams A, Breur J: **Cardiac Valve Annulus Diameters in Extremely Preterm Infants: A Cross-Sectional Echocardiographic Study.** *Neonatology* 2018, **114**(3):198-204.
9. Bockeria LA, Podzolkov VP, Makhachev OA, Zelenikin MA, Alekian BG, Ilyin VN, Gadjiev AA, Shatalov KV, Kakuchaya TT, Khiriev T *et al*: **Surgical correction of tetralogy of Fallot with unilateral absence of pulmonary artery.** *Ann Thorac Surg* 2007, **83**(2):613-618.
10. Castleberry CD, Gudausky TM, Berger S, Tweddell JS, Pelech AN: **Stenting of the right ventricular outflow tract in the high-risk infant with cyanotic teratology of Fallot.** *Pediatr Cardiol* 2014, **35**(3):423-430.
11. Hirji A, Bernasconi A, McCrindle BW, Dunn E, Gurofsky R, Manlhiot C, Miner SE, Kingdom JC, Jaeggi ET, Van Arsdell G *et al*: **Outcomes of prenatally diagnosed tetralogy of Fallot: Implications for valve-sparing repair versus transannular patch.** *Can J Cardiol* 2010, **26**(1):e1-6.
12. Escribano D, Herraiz I, Granados M, Arbues J, Mendoza A, Galindo A: **Tetralogy of Fallot: prediction of outcome in the mid-second trimester of pregnancy.** *Prenat Diagn* 2011, **31**(12):1126-1133.
13. Arya B, Levasseur SM, Woldu K, Glickstein JS, Andrews HF, Williams IA: **Fetal echocardiographic measurements and the need for neonatal surgical intervention in Tetralogy of Fallot.** *Pediatr Cardiol* 2014, **35**(5):810-816.
14. Friedman K, Balasubramanian S, Tworetzky W: **Midgestation fetal pulmonary annulus size is predictive of outcome in tetralogy of fallot.** *Congenit Heart Dis* 2014, **9**(3):187-193.
15. Maeno Y, Himeno W, Fujino H, Sugahara Y, Furui J, Mizumoto Y, Kato H: **Progression of congenital heart disease in the prenatal period.** *Pediatr Int* 1999, **41**(6):709-715.
16. Jatavan P, Tongprasert F, Srisupundit K, Luewan S, Traisrisilp K, Tongsong T: **Quantitative Cardiac Assessment in Fetal Tetralogy of Fallot.** *J Ultrasound Med* 2016, **35**(7):1481-1488.
17. Quartermain MD, Glatz AC, Goldberg DJ, Cohen MS, Elias MD, Tian Z, Rychik J: **Pulmonary outflow tract obstruction in fetuses with complex congenital heart disease: predicting the need for neonatal**

Figures

Figure 1

Measurements of the fetal cardiac parameters. In the right ventricular outflow tract view, pulmonary valve annulus is measured from hinge point to hinge point at the location of maximal expansion during peak systole (A, arrow). In the three-vessel view, the right pulmonary artery is measured at the proximal part of its bifurcation (B, arrow). In the left ventricular outflow tract view, the aortic valve annulus is measured from hinge point to hinge point at the location of maximal expansion during peak systole (C, arrow). In the ductal arch view, pulmonary valve peak systolic velocity is measured at the level of the pulmonary valve using pulsed Doppler (D). In the aortic arch view, the descending aorta is measured at the level of the diaphragm (E, arrow).

Figure 2

Receiver-operating characteristic curves for the fetal cardiac parameters and postnatal operation type. The presented cardiac parameters of note include the pulmonary valve annulus z-score (A), right pulmonary artery (RPA) z-score (B), pulmonary valve peak systolic velocity (C), pulmonary valve annulus/aortic valve annulus ratio (D), and RPA/descending aorta ratio (E). AUC, area under the receiver-operating characteristic curve.