A situational analysis of child and adolescent mental health services and systems in the Western Cape Province of South Africa

Stella Mokitimi

University of Cape Town

Marguerite Schneider

University of Cape Town

Petrus J de Vries (✉ petrus.devries@uct.ac.za)

University of Cape Town <https://orcid.org/0000-0002-8915-1571>

Research Article

Keywords: Situational analysis, Child, adolescent, mental health, Services, Western Cape, South Africa, low- and middle-income countries, LMIC

Posted Date: June 30th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-524889/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background:

Child and Adolescent mental health services (CAMHS) and systems need strengthening globally. This requires knowledge of the hardware elements of the system (human resources, financing, medicines, technology, organisational structure, service infrastructure, and information systems). This study sought to examine these elements of CAMHS in the Western Cape Province of South Africa

Methods:

The World Health Organization Assessment Instrument of Mental Health Systems (WHO-AIMS) version 2.2 of 2005 was adapted to identify key variables of interest in CAMHS. Data were collected for the calendar year 2016 and focused on the public health sector. We outlined findings based on best available data across the six domains of the WHO-AIMS.

Results:

In domain 1, we found no provincial CAMHS policy or implementation plans to support the national CAMHS policy, and were unable to identify a CAMHS-specific budget. In domain 2, there was no dedicated provincial leadership structure for CAMHS, and no dedicated or 'child- and adolescent-friendly' mental health services at primary or secondary care levels. At tertiary level, there were only three specialist CAMH teams. The majority of CAMHS resources were based in the City of Cape Town, with limited resources in the rural districts. Essential medicines were available in all facilities, and the majority of children and adolescents had access to free services. In domain 3, data were limited about the extent of training offered to primary healthcare staff, and little or no psychosocial interventions were available in primary care. Domain 4 identified a small and variable CAMHS workforce across all levels of care. In domain 5 few public health campaigns focused on CAMH, and little evidence of formal intersectorial collaboration on CAMHS was identified. Domain 6 identified significant limitations in health information systems for CAMHS, including lack of child- and adolescent-specific and disaggregated data to establish baselines for policy development, monitoring, evaluation and CAMHS research.

Conclusions:

This study identified significant structural weaknesses in CAMHS and present a clear call for action to strengthen services and systems in the province and in South Africa. it would be important to expand research also to include provider and user perspectives for service-strengthening.

Background

The mental health of children is a global health priority, yet it is generally known that services are limited, particularly in low- and middle-income countries (LMIC) [1, 2]. Health system strengthening requires an understanding of multiple landscapes: the policy and resource landscape (representing mostly hardware elements of health systems), and the landscape as perceived by stakeholders (which would describe both hardware and software elements of the healthcare system) [3, 4]. Hardware typically refers to human resources, financing, medicines, technology, organisational structure, service infrastructure, and information systems, while the software typically refers to ideas and interests, relationships and power, values and norms, and the interactions between all factors and actors [3, 4].

As stated by the World Health Organization (WHO) [4] and summarised by Gilson [3], all the components of a health system should be balanced in order for the system to be responsive to the needs of the community it serves. A good enough health system should have sufficient resources that are equitably distributed, and human resources that have sufficient competencies to respond to the needs of the population [4].

The geographical focus of our work is South Africa, an upper-middle-income country that has some of the greatest economic and health disparities in the world [5, 6]. A situational analysis conducted by Kleintjes and colleagues in 2005 [2] evaluated key aspects of child and adolescent mental health services (CAMHS) in South Africa, Uganda, Zambia and Ghana. In comparison to the other three sub-Saharan African countries, South Africa had relatively more CAMH resources [2]. However, only 1.4% of all mental health outpatient services in the country, 3.8% of mental health beds in general hospitals, 1% of specialist mental health hospital beds, and 1% of day patient facilities in the country were dedicated to CAMHS. There were no specialist CAMH hospitals. The number of psychiatrists (general, and child & adolescent psychiatrists combined) was estimated at 0.28 per 100,000. No data were available on the number of mental health professionals in schools [7]. Qualitative data from semi-structured interviews with key stakeholders in the Kleintjes study [2] proposed three main themes as the reasons for the very low resources for CAMHS: first, the impact of stigma associated with mental health disorders; second, the low priority of all aspects of mental health in LMIC; and third, the lack of attention to the link between poverty and poor mental health [2]. While these proposed reasons are understandable when comparing mental health with physical health services, Kleintjes and colleagues did not attempt an explanation for the under-representation of child and adolescent *versus* adult mental health services.

No situational analysis has been performed since the study by Kleintjes and colleagues [2] and, to our knowledge, no study has conducted a finer-grained situational analysis of CAMHS in South Africa or any other LMIC.

The Western Cape Province of South Africa

In order to perform a more detailed situational analysis of CAMHS, we selected the Western Cape as the focus for our work. The Western Cape was selected for two reasons. Firstly, it is one of the provinces with better resources in terms of health services [8, 9] and therefore more accessible for research scrutiny in terms of data sources, documents and dedicated staff for mental health. Secondly, it is the base of our own clinical activities in the Division of Child and Adolescent Psychiatry (DCAP) at the University of Cape Town. It therefore represents the immediate and direct health system and services in which we are actors, and gives us a strong knowledge and experience base from which to look at the data.

In 2016, the year selected for data collection, South Africa had an overall estimated population of 55.9 million [10]. The Western Cape, one of the nine provinces of South Africa, was the fourth largest both in land size and population. The estimated population was 6.3 million of which 2.1 million (33.9%) were children and adolescents under the age of 19 years. The province has rural and urban areas and is served by one metropolitan municipality (City of Cape Town) and five district councils (West Coast, Cape Winelands, Overberg, Eden and Central Karoo). The metropolitan municipality is divided into four (4) main substructures: The Southern/Western substructure, the Klipfontein/Mitchell's Plain substructure, the Northern/Tygerberg substructure, and the Khayelitsha/Eastern substructure. Figure 1 shows the province and its urban and rural structures, substructures and districts. In addition, the figure shows the location of specialist CAMH services, and of the regional hospitals in the province.

In 2016, 63.8% of the population lived in the City of Cape Town, followed by the Cape Winelands (13.8%), Eden (9.7%), West Coast (6.9%), Overberg (4.6%), and the Central Karoo (1.2%) [10]. Table 1 shows a more detailed population distribution in the Western Cape by district in 2016. Unfortunately, no disaggregated data were published in the 2016 Census that would allow separation of adolescents (under the age of 19) and younger adults in the 15–34-year age groups.

Table 1. Population distribution by Western Cape district in 2016 (data from Stats SA, 2016)

2016 Community Survey						
General population			Children, adolescents and young adults			
			0–14 yrs		15–34 yrs*	
District / municipality	N	%	N	%	N	%
West Coast	436,403	6.9	113,113	25.9	153,472	35.2
Cape Winelands	866,001	13.8	230,708	26.6	316,210	36.5
Overberg	286,786	4.6	74,764	26.1	94,453	32.9
Eden	611,278	9.7	155,008	25.4	207,010	33.9
Central Karoo	74,247	1.2	18,862	25.4	27,936	37.6
City of Cape Town	4,005,016	63.8	1,042,259	26.0	1,331,960	33.3
Total	6,279,730	100.0	1,521,601	24.23	1,977,569	39.49

***The Census data did not provide any disaggregation between adolescents and younger adults in the 15–34-year age groups.**

The overall prevalence of CAMH disorders, adjusted for local population factors, comorbidity and adjusted according to the risk factors in the Western Cape, were estimated to be 17% [11]. It should be noted that these estimates did not include infant mental health disorders, autism spectrum disorders or neurodevelopmental disorders other than intellectual disability, suggesting that these rates may be an underestimate of the true prevalence of CAMH disorders in the province. Nevertheless, the estimates certainly highlight the burden of CAMH problems in the province, and the need for service provision.

Previous reviews of services and resources suggested a clear neglect of provision for CAMHS in the Western Cape [2, 7, 8, 12]. However, as acknowledged by previous authors, due to inadequate information systems, little or no data were available on children and adolescents with mental health problems in primary healthcare settings, general hospital outpatient and inpatient settings, or in specialist CAMHS facilities [7, 12].

Situational analysis and the World Health Organization Assessment Instrument for Mental Health Systems

The World Health Organization Assessment Instrument of Mental Health Systems (WHO-AIMS) version 2.2 of 2005 [13] is the WHO framework for conducting situational analyses in LMIC. It was conceptualised and developed by the Mental Health Evidence and Research Team (MER) of the Department of Mental Health and Substance Abuse (MSD) at the WHO, in collaboration with colleagues inside and outside the World Health Organization (WHO). The tool is used to collect information on mental health systems with the goal of providing baseline data that can be used to improve mental health systems and to monitor change. The WHO-AIMS framework includes six domains, 28 facets and 156 items to cover key aspects of mental health systems. It includes a brief version that can be used to collect data on a reduced number of variables about mental health systems. Both versions of the WHO-AIMS were developed with general mental health services (as opposed to CAMH) in mind. In fact, the WHO-AIMS framework has only one item specifically on CAMHS (Domain 2, item 2.2.6/Brief version item B13) on the number of children 18 years or younger treated at mental health outpatient facilities, including the proportion of under-19-year-olds treated in comparison to all patients treated. In South Africa, the WHO-AIMS Brief version was previously used to assess CAMHS at national level [2], and has been used to assess general mental health services at a regional/provincial level [14].

Here, we therefore set out to perform a fine-grained situational analysis of CAMHS in the Western Cape province. Whilst we firmly acknowledge that child and adolescent mental health problems cut across all systems and sectors, for the purpose of this study, given the time and resources constraints, this situational

analysis limited its scope to focus on the health system and mainly the Department of Health (DoH) public health system, rather than on other or multiple sectors.

Methods

Study design

This was a descriptive situational analysis of CAMHS in the Western Cape Province of South Africa.

Data collection

The Brief version of items from the WHO-AIMS version 2.2 of 2005 was adapted to collect data on CAMHS. We selected 37 items of direct relevance and modified them to focus on CAMHS. Table 2 shows the WHO-AIMS items adapted for this situational analysis. Data were collected for the calendar year January to December 2016. The situational analysis focused mainly on the DoH and the public health sector, and not on the private health sector or other government departments. WHO-AIMS checklists/survey questionnaires were distributed to key stakeholders in the provincial DoH. Based on the WHO-AIMS table of data sources, relevant documents and records were requested from regional offices, and from individual hospital/clinical facilities in order to complete the questionnaires and fill in any data gaps. The research team established and maintained contact with key stakeholder groups to follow up on the questionnaires, to identify and assist with any difficulties pertaining to the information required, and to verify the information provided. In addition, electronic searches for relevant information through provincial websites and Web of Science (version 5.34) were performed. After collation of all data, further clarifications were sought between January and May 2020 as a final validation check for any potential additional data that may have become available before preparation of this manuscript.

Table 2
Variables of interest adapted from the WHO-AIMS version 2.2 (Brief version) for this situational analysis

WHO-AIMS Domains	Western Cape provincial data collected
1. Policy and legislative framework	1.1 CAMH policies, plans, and legislations (B1, B3, B4) 1.2 Human rights legislation relevant to children and adolescents (B5) 1.3 Financing: Expenditure on CAMHS by the provincial DoH (B6)
2. Clinical services for children and adolescents with mental health disorders	2.1 Existence and functions of a regional CAMH authority (B9) 2.2 Organisation of CAMH services in terms of catchment areas (B10) 2.3 Outpatient services: Availability of CAMH outpatient facilities, and number/proportion of children and adolescents treated for mental health problems through outpatient facilities at primary, secondary and tertiary levels of care (B11, B12, B13) 2.4 Inpatient services: Availability of CAMH inpatient facilities, and number/proportion of children and adolescents treated (B15, B16, B17) 2.5 Availability of CAMH day patient facilities, community residential facilities, forensic facilities, or CAMH hospitals (B14, B18, B19, B25) 2.6 Interventions (medications): Psychotropic medicines appropriate for children and adolescents included on the essential medicines list; free access to essential psychotropic medicines, and availability of medicines in outpatient and inpatient settings at secondary and tertiary levels of care (B2, B8, B28, B29) 2.7 Interventions (psychosocial): Access to psychosocial interventions in outpatient and inpatient settings at secondary and tertiary levels of care (B26, B27)
3. CAMH in Primary Healthcare	3.1 Refresher training in CAMH provided to PHC doctors, nurses or other staff and interaction of PHC with specialist CAMHS (B31–B35) 3.2 Availability of medicines and psychosocial interventions in PHC facilities (B27, B33)
4. Human resources	4.1 Human resources in CAMHS (B38–B41)
5. Public education and links with other sectors	5.1 Public education and awareness campaigns about CAMH (B47)
6. Monitoring and research	6.1 Monitoring CAMHS (B52, B53) 6.2 Research in CAMH (B54)
CAMH = child and adolescent mental health; B items listed in brackets e.g. (B1) refer to items as listed in the WHO-AIMS Brief Version.	

Data capturing and analysis

All data sources were collated and numbered as primary sources of findings. Data sources and data source numbers (DSN) are shown in Table 3. Data were captured on the WHO-AIMS Excel Data Entry program version 2.2 of 2005 for analysis. Analysis was performed through descriptive statistics of variables of interest as outlined in Table 2.

Due to technical limitations, Table 3 is only available as a download in the supplementary files section

Results

Below we outline the results for each of the variables of interest shown in Table 2. Each finding includes one or more references as links to the evidence for the finding.

WHO-AIMS Domain 1: Policy and legislative framework

Policies, plans, and legislations

There was no provincial CAMH policy document and no provincial plan to give effect to the national CAMH policy [15, 16]. There was a general healthcare policy [16] which did not focus on CAMHS. Interestingly, the policy acknowledged the need to separate CAMHS from adult psychiatric services, but did not present any specific plan to do so. There was no province-specific CAMH mental health legislation. The National Mental Health Act no. 17 of 2002 [17] and the Child Care Act 74 of 1983 [18] were used as legal frameworks in the province [19].

Human rights legislation relevant to children and adolescents

To ensure that the human rights of all people (including children and adolescents) were met, a human rights review body (Mental Health Review Board) existed. Its functions included acting as licensing directorate for non-profit organisations, mental health programme audits, and monitoring [19, 20]. All specialist CAMH units had at least one annual external review/inspection of human rights protection of patients. In addition, all specialist CAMH inpatient units had admission, discharge, and seclusion policies, complaints and appeals processes, and procedures to ensure the protection of human rights [19].

Expenditure on child and adolescent mental health services by the Provincial Department of Health

There was no separate budget for CAMHS, and no budgetary information was presented in a way that could allow disaggregation of adult versus CAMH budgets [21–23]. Primary healthcare budgets were integrated into the overall District Health Services (Community Health Centres and Community-based services) budget. Secondary care budgets were integrated into District Hospital budgets. Tertiary care budgets were integrated either into Provincial Hospital Services (specialised psychiatric hospitals where some tertiary CAMH specialist units were situated) or into Central Hospital Services (central and tertiary hospitals where two tertiary CAMH specialist services were situated). CAMHS would have been delivered to outpatients at primary care, to outpatients and emergency inpatients at secondary level of care, and to outpatients, emergency inpatients and longer-term inpatients at tertiary level of care.

As shown in Table 4 the overall health services budget in the province for 2016/2017 was just under R20 billion (just over US\$1 billion). Of this amount, ~R7.8 billion (US\$ 511.1 million) (39.2%) was allocated to District Health Services, ~R5.6 billion(US\$371.8 million) (28.5%) to Central Hospital Services, ~R3.2 billion (US\$ 208.9 million) (16%) to Provincial Hospital Services, of which ~ R589 million (US\$38.5 million) (2.9% of the total health budget) was allocated to mental hospitals [21–23]. Given that none of these groupings provided services exclusively to children and adolescents with mental health problems, it was not possible to identify any CAMH-specific expenditure.

Table 4
Western Cape provincial budget for 2016/17

Total health budget in	Primary and secondary level	Tertiary level	
South African Rand	District Health Services	Provincial Hospital Services	Central Hospital Services
R19.983 billion (US\$1.307 billion)	R7.826 billion (US \$511.140 million) (39.2%)	R3.199 billion (US\$ 208.849 million) (16%)	R5.697 billion (US\$371.82 million) (28.5%)
		Non-mental health services	Mental health hospitals
		R2.610 billion (US \$170.427 million) (13.1%)	R589 million (US\$38.505 million) (2.9%)
R = South African Rand; At the time of publication (Aug 2020), R1 was equivalent to US\$0.058.			

WHO-AIMS Domain 2: Child and adolescent mental health resources

Existence and functions of a regional child and adolescent mental health authority

There was no provincial authority exclusively for CAMHS and no provincial director for mental health [19]. There was a provincial deputy-director for mental health and substance abuse. The role of the post-holder was to coordinate all mental health services (including CAMHS) within the province to a) ensure development and implementation of the national policy and legislation, b) oversee monitoring of services, c) facilitate equitable budgets for mental health, d) work closely with district health managers, all stakeholders and sectors, and e) evaluate services and policy implementation. The deputy director reported to the provincial director of health programmes [19].

Organisation of child and adolescent mental health services in terms of catchment areas

CAMHS were provided based on where families lived in the province, referred to as geographical service areas (GSAs) or catchment areas, as shown in Fig. 1. Services were provided across three levels of care: primary (level 1), secondary (level 2) and tertiary (level 3). The basic patient flow for service organisation was for children and families to start their CAMH 'journey' at level 1 services (their catchment area primary healthcare clinic), where they would be seen by a mental health nurse and, if required, a medical officer (a generally-trained doctor). When primary healthcare teams did not feel able to diagnose or treat the child and family, they would be referred to level 2 (their catchment area district hospital) where they would be seen by a general mental health nurse and, if required, a general psychiatrist. At level 2 children and families may also be seen by paediatric services depending on the reason and pathway for referral. When level 2 teams did not feel able to diagnose or treat the child and family, they would refer them to level 3 services. These are services with multidisciplinary expertise in child and adolescent mental health, including child mental health nurses, psychologists with expertise in child mental health, and subspecialist child & adolescent psychiatrists. We refer to level 3 services as 'specialist CAMHS'. Each specialist CAMH team served a specific metropolitan catchment area, specific district hospitals in that catchment area, as well as a specific rural area and its regional hospital. Table 5 shows the catchment areas for each specialist CAMHS unit [24–26].

Table 5
Catchment areas for specialist child and adolescent mental health services

Specialist CAMH unit	Metropolitan (urban)		Rural	
	Metro catchment area	District hospitals	Rural catchment area	Regional hospital
Division of Child and Adolescent Psychiatry (DCAP)	Southern	New Somerset Hospital	Eden	George Hospital
	Western	Victoria Hospital		
	Portion of Klipfontein	False Bay Hospital		
		Groote Schuur Hospital		
		Red Cross District Service		
Tygerberg Child and Adolescent Psychiatry team	Northern	Karl Bremer Hospital	Portion of Cape Winelands	Paarl Hospital
	Tygerberg	Eerste River Hospital	West Coast	
	Portion of Eastern	Tygerberg Hospital district service		
Lentegeur Child and Family Unit (CFU)	Khayelitsha	Khayelitsha Hospital	Overberg	Worcester Hospital
	Mitchell's Plain	Mitchell's Plain Hospital	Central Karoo	
	Portion of Eastern	Helderberg Hospital	Portion of Cape Winelands	
	Portion of Klipfontein			

In 2016 there were three specialist CAMH service units, all based in the metropolitan municipality. The Division of Child and Adolescent Psychiatry (DCAP) (University of Cape Town) was based at Red Cross War Memorial Children's Hospital in the Southern substructure, the Tygerberg Child and Adolescent Psychiatry team (Stellenbosch University) was based at Tygerberg Hospital, a general tertiary hospital in the Northern substructure, and the Lentegeur Child and Family Unit (University of Cape Town and Stellenbosch University) was based at Lentegeur Hospital, a mental hospital in the Mitchell's Plain substructure [24–26].

The specialist CAMH service units offered outpatient services to children and families in their catchment areas. Children and adolescents with any mental health problem could be referred by a specialist (psychiatrist or paediatrician) from a relevant catchment area [20, 24–26].

In the time period of this study (January–December 2016), the three units had areas of particular expertise: autism spectrum disorder and infant mental health at DCAP, neuropsychiatric conditions at Tygerberg, and substance abuse and rehabilitation at Lentegeur. Specialist inpatient services were available at each of the three specialist CAMH units but were not based on catchment areas. Instead, they were based on age and/or clinical profile of the child or adolescent. For instance, DCAP had an inpatient unit for children under 12, while Tygerberg and Lentegeur Hospitals had adolescent inpatient units [24–26].

There were also two units that could admit adolescents with intellectual disabilities, one based at Alexandra Hospital, a mental hospital for people with intellectual disabilities, and the other at Lentegeur Hospital. Both these hospitals were in the metropolitan municipality. The units offered inpatient and outpatient services for older adolescents with dual diagnosis (intellectual disabilities and a mental health problem) alongside adults with intellectual disabilities, and served all the districts of the Western Cape. The exclusion criteria for these units were social problems in the absence of intellectual disabilities and comorbid mental health problems, and patients with forensic histories [20, 24–26].

Even though catchment areas and GSAs were clear for general outpatient referrals at primary and secondary levels of care, this was not the case for particular areas of expertise at tertiary level of care, or for inpatient referrals at tertiary level. In addition, there were no specialist CAMHS in any of the rural districts [24–29].

Availability of child and adolescent mental health outpatient facilities, and number/proportion of children and adolescents treated for mental health problems through outpatient facilities at primary, secondary and tertiary levels of care

Outpatient services for children and adolescents with mental health problems were available at 317 health facilities spanning primary to tertiary levels of care. Services included limited mobile services and satellite clinics. Only three of all outpatient services (0.9%) were providing dedicated mental health services to children and adolescents. All other facilities (99.1%) were open to all ages [28].

Primary care (level 1)

There were 273 primary healthcare facilities (206 clinics, 58 community daycentres and 9 community health centres) that provided general mental health services, where children and adolescents were seen alongside adults [28]. Out of a total of 188,369 patients seen for mental health problems at primary level of care, 8,300 (4.4%) were children and adolescents. Table 6 shows the number of children and adolescents treated in 2016 in primary healthcare facilities per geographic service area. The majority of children and adolescents were treated in the City of Cape Town (6,300/8,300; 75.9%) followed by the Eden District and the Cape Winelands. The Central Karoo had the lowest number and proportion of children and adolescents seen for mental health disorders [28].

Table 6. The number and proportion of children and adolescents seen in 2016 in primary healthcare (level 1) outpatient settings for mental health problems in the Western Cape

Geographic service areas	Age distribution		Total (% children)
	>18 years	< 18 years	
City of Cape Town	131,836	6,330	138,166 (4.58%)
Cape Winelands District	16,986	609	17,595 (3.46%)
Central Karoo District	3,066	49	3,115 (1.57%)
Eden District	16,347	850	17,197 (4.94%)
West Coast District	11,834	462	12,296 (3.76%)
Total	180,069	8,300	188,369 (4.4%)

Secondary care (level 2)

Mental health services were provided in 34 district hospitals in the province in 2016 [28]. There were no separate outpatient CAMH facilities at secondary level of care [20, 28] children and adolescents were therefore seen alongside adult psychiatric patients. Out of 15,755 patients seen for mental health problems at secondary level in 2016, 1,145 (7.27%) were children or adolescents. Table 7 shows the numbers and proportion of children and adolescents seen at outpatient departments per GSAs in 2016 [28]. The largest number of children and adolescents were seen in the City of Cape Town (561 of 1,145; 48.9%), followed by the Eden District and the Cape Winelands. However, as a proportion of all mental health cases, all of the rural districts saw a greater proportion of under 19's, all in excess of 10% of patients seen (see Table 7).

Table 7
The number and proportion of children and adolescents seen in 2016 in secondary care (level 2) outpatient settings for mental health problems in the Western Cape

Geographic service area	Age distribution		Total (% children)
	> 18 years	< 18 years	
City of Cape Town	10,242	561	10,803 (5.19%)
Cape Winelands District	1,083	132	1,215 (12.19%)
Central Karoo District	162	29	191 (15.18%)
Eden District	2,064	248	2,312 (10.73%)
West Coast District	1,059	175	1,234 (14.18%)
Total	14,610	1,145	15,755 (7.27%)

Tertiary care (level 3)

Three specialist CAMHS in the province provided dedicated multidisciplinary outpatient services in 2016. Adolescents with intellectual disabilities and co-occurring psychiatric disorders were seen in the two specialist units for people with intellectual disabilities [24–26].

No formal summary data were available for Tygerberg and Lentegeur outpatient units in 2016 [28]. Patient numbers at DCAP are shown in Fig. 2, and self-declared ethnic/racial distribution in Fig. 3 [28]. A total of 3,639 children and adolescents were seen at DCAP in 2016. The majority of patients were male (63.2%) and the self-declared ethnicity (where provided) was coloured (1,943/2,945; 65.98%), followed by white (578/2,945; 19.63%) and black (424/2,945; 14.29%).

The Alexandra and Lentegeur units for intellectual disabilities treated three and 39 children and adolescents respectively with intellectual disabilities and co-occurring mental health disorders as outpatients in 2016 [27].

Child and adolescent inpatient services in 2016 were provided at secondary (level 2) and tertiary (level 3) levels. Emergency cases requiring 72-hour observations were admitted at secondary level and medium to long-term admissions were provided at tertiary level [28].

Secondary care (level 2)

Secondary level services were available in 42 health facilities [28]. There were no dedicated beds for children and adolescents with mental health problems in these hospitals. Children and adolescents who required inpatient emergency admission were admitted to paediatric units (up to age 12) or adult psychiatric wards (13–18 years). There were no disaggregated data for children and adolescents who were admitted to paediatric units or adult psychiatric wards [28]. Table 8 shows the numbers of children and adolescents per geographic service area who were admitted in level 2 inpatient emergency units in 2016 [28]. A total of 795 children and adolescents, representing 4.35% of all secondary care mental health admissions, were admitted. The majority of children and adolescents were treated in the City of Cape Town (473/795; 59.5%), followed by the Cape Winelands and Eden Districts (see Table 8). Interestingly, the rural districts had more than double the proportion of under-18-year-old admissions relative to total population numbers, compared to the City of Cape Town. In the City of Cape Town 3.31% of level 2 admissions were under the age of 18, while in rural districts the proportions were 7.27–9.66% [28].

Table 8
Children and adolescents with mental health problems admitted to secondary care (level 2) inpatient facilities per geographic service area in the Western Cape

Geographic service area	Age distribution		Total (% children)
	> 18 years	< 18 years	
City of Cape Town	13,824	473	14,297 (3.31%)
Cape Winelands District	1,339	105	1,444 (7.27%)
Central Karoo District	131	14	145 (9.66%)
Eden District	889	82	971 (8.44%)
Overberg District	487	42	529 (7.93%)
West Coast District	795	79	874 (9.03%)
Total	17,465	795	18,260 (4.35%)

Tertiary care (level 3)

Children and adolescents were admitted to one of three specialist CAMH inpatient units or to one of two intellectual disabilities inpatient units. Where this was not possible, admission to adult mental health units were made. All tertiary inpatient beds were in the City of Cape Town. A total of 346 children and adolescents were admitted to specialist inpatient units in 2016 [28]. The breakdown of admissions is outlined below.

The inpatient unit at DCAP (known as the Therapeutic Learning Centre) was a 6-bedded unit for 6–12-year-olds with complex or severe mental health problems, where children are typically admitted for 3–6 months. The unit admitted a total of 26 patients in 2016. The Tygerberg Child and Adolescent Psychiatry unit had a 16-bedded adolescent inpatient unit. Adolescents were admitted either for short-term diagnostic assessment or for longer-term intervention. The unit admitted 157 patients in 2016. The Lentegur Child and Family Unit had an 8-bedded adolescent unit, and also admitted adolescents for short-term diagnostic work-up, or short to medium-term therapeutic work. The unit admitted 152 patients in 2016 [28].

Eleven children or adolescents were admitted to adult mental health hospitals: Valkenberg = 8; Alexandra = 2; Stikland = 1 [28].

Availability of child and adolescent mental health day facilities, community residential facilities, forensic facilities or hospitals

Community or other residential facilities

There were no Department of Health day facilities and residential facilities exclusively for children and adolescents with mental health problems [28]. The Department of Social Development provided community residential facilities for children and adolescents with a range of psychosocial challenges including abuse, neglect or other social difficulties. Children and adolescents with mild mental health problems who needed care, were placed in these units. There were 69 of these facilities listed in the Western Cape in 2016 [28], but no data were available on the number of children in these facilities during the year. There were also special care centres and licensed homes for children and adults with severe and profound intellectual and physical disabilities, but no child and adolescent data were available [20].

Child and adolescent forensic and other residential facilities

Child and adolescent forensic mental health services were provided at one of the adult mental hospitals (Valkenberg Hospital) in the City of Cape Town [27]. A consultant psychiatrist, two child & adolescent psychiatrists, and five clinical psychologists provided sessions to the service. CAMH forensic services were aimed at providing the Department of Justice with criminal capacity/forensic psychiatric assessments for children in conflict with the law [29]. Court diversion programmes were also available to provide support for 10–18-year-olds who were both in conflict with the law and had co-occurring mental health needs. These therapeutic programmes were facilitated by a probation officer or social worker and were offered for a period of 3–12 months. Programmes were provided by the Department of Social Development and a range of non-governmental/non-profit organisations [30].

Child and adolescent mental hospitals

There were no dedicated CAMH hospitals in the province in 2016 [20, 24].

Interventions (medications): Psychotropic medicines appropriate for children and adolescents included on the essential medicines list, free access to essential psychotropic medicines, and availability of medicines in outpatient and inpatient settings at secondary and tertiary levels of care

All mental health facilities (outpatient and inpatient) at secondary and tertiary levels of care had at least one psychotropic medicine of each therapeutic class (antipsychotic, antidepressant, mood stabiliser, anxiolytic, stimulant and anti-epileptic medicines) available [27]. These drugs were fluoxetine, citalopram, haloperidol, chlorpromazine, imipramine, risperidone, methylphenidate, lorazepam, lithium carbonate, sodium valproate, carbamazepine, citalopram and biperiden [31]. About 4.8 million people including children and adolescents (76%) in the Western Cape qualified for free access to healthcare services and for those with mental health disorders, to essential psychotropic medicines if required [23].

Interventions (psychosocial): Access to psychosocial interventions in outpatient and inpatient settings at secondary and tertiary levels of care

Secondary level outpatient and inpatient services provided very limited access to psychosocial interventions for children and adolescents [27]. All specialist CAMH units provided multidisciplinary care at outpatient and inpatient level. This involved access to a range of psychosocial interventions appropriate for children and adolescents including psychoeducation, family interventions, counselling, individual psychological therapies, play-based and other evidence-based psychosocial interventions [15, 28].

WHO-AIMS Domain 3: Child and adolescent mental health in primary healthcare

Refresher training in child and adolescent mental healthcare provided to primary healthcare doctors, nurses or other staff and interactions of primary healthcare with specialist child and adolescent mental health services

The WHO-AIMS specifically seeks to quantify the proportion of mental health training to primary care doctors, nurses and other primary health care (PHC) staff in relation to other training provided in the year. The document also specifically seeks to quantify the number of primary healthcare staff that have received at least two days of refresher training in mental health in the last year. No data were available to answer these specific questions.

However, as actors in CAMHS, we know that specialist CAMHS provided outreach and support to colleagues at primary care level, including to medical doctors and nurses. Training was provided through workshops and seminars, and clinical consultation through a range of modalities including face-to-face, telephonic consultations, informal meetings, review of individual cases or through discussion of referral issues. However, there was no formal documentation of informal interactions with primary care staff in any health information systems, and the majority of these opportunities were with primary care colleagues in the metropolitan area [25, 29].

Availability of medicines and psychosocial interventions in primary healthcare facilities

All primary healthcare facilities had at least one psychotropic medicine of each therapeutic class (antipsychotic, antidepressant, mood stabiliser, anxiolytic, stimulant and anti-epileptic medicines) available [19]. The majority of the population had free access to medications. There was little if any access to psychosocial interventions for children or adolescents available at primary care level [23].

WHO-AIMS Domain 4: Human resources

Human resources in child and adolescent mental health services

No formal data were available on human resources specifically for CAMH. The majority of mental health human resources data represented staffing for all specialties across all ages. There were also no differentiation of data between staffing for outpatient versus inpatient care. Figure 4 shows the overall mental health human resources in the Western Cape in 2016. The majority of staff were mental health nurses (n = 1,079; 85.4% of all mental health staff), and there were few specialist psychiatrists (n = 51; 4.03% of all mental health staff). Eight of the 51 psychiatrists (15.7%) were qualified as child & adolescent psychiatrists. Trainees (psychiatric registrars and psychology interns) represented 5.38% of the workforce) [28].

In the absence of provincial human resources data for specialist CAMH, data were collected directly from the three tertiary CAMH units [28]. Figure 5 provides best estimates of human resources in the specialist CAMH facilities. As shown in Fig. 5 all these units had multidisciplinary teams which included child & adolescent psychiatry, psychology, nursing and other disciplines, but there was significant variability in the workforce profile between the specialist units. Psychiatrists and psychologists were appointed on contracts with 'joint conditions' between the Western Cape Government and respective universities, and were allocated 70% of their time to deliver clinical care and 30% of their time for academic activities (including teaching, supervision and research). Staff in other categories were appointed by the Western Cape Government with little to no time dedicated for teaching/ training, supervision and research.

In 2016, there were three posts in the province for psychiatrists to train as child & adolescent psychiatrists. However, there were no vacant positions in the government sector for child & adolescent psychiatrists once qualified. There was therefore no 'pipeline' that could support new child & adolescent psychiatrists to join the state sector. Newly qualified child & adolescent psychiatrists had to move into private practice, work in general psychiatry posts, or leave the country. There were no funded training programmes for psychologists, nurses, occupational therapists, or speech and language therapists to specialise in child and adolescent mental health.

WHO-AIMS Domain 5: Public education and links with other sectors

Public education and awareness campaigns about child and adolescent mental health

There were no formalised intersectoral collaborations between the DoH and other sectors specifically related to child and adolescent mental health and mental illness in children or adolescents [19].

One initiative, led by the Department of Social Development (DSD) and formally adopted by all provincial government departments, was related to infant mental health. The 'First 1,000 Days of Life' project was a collaborative cross-agency programme involving the DoH, DSD and the Western Cape Education Department (WCED), and was set up to identify and reduce risk factors for maternal and infant mental health problems in the first 1,000 days of life [32].

A number of other projects and campaigns were aimed at promoting child and adolescent well-being. These included a competition launched by the Salesian Life Choices (a youth organisation) to encourage young people to start discussions around health issues and depression [33], an anti-bullying campaign in schools run by the WCED in May 2016 [34] and the International 16 Days of Activism, a campaign aimed at increasing awareness of the negative impact of violence and abuse against women and children [35].

WHO-AIMS Domain 6: Monitoring and research

Monitoring of child and adolescent mental health services

There was a formally defined list of individual data items that ought to be collected by all child and adolescent mental health facilities, using CLINICOM and SENJANI information systems [19,28]. CLINICOM is a hospital information system that is used in various countries. It records the activities of a range of clinicians in outpatient and inpatient facilities and the data is saved comprehensively in one place. It keeps records of patients treated in medical facilities, medical documentation, laboratory test results, daily course of diseases, patient medication records, examination scheduling by specialists, rehabilitation and rehabilitation procedures. In South Africa and in the Western Cape, CLINICOM is a computer-based data capture programme used in secondary and tertiary hospitals in the province. All new patients and patient-related activities are expected to be captured on the CLINICOM system by clinic administrators. A sub-component of CLINICOM is used in primary healthcare facilities, to capture data on children and adolescents who present with mental health problems including personal information, inpatient and outpatient attendance, admissions, discharges, clinic, and diagnosis [19]. Despite the apparent availability of these information systems, it was not clear to what extent different healthcare facilities completed or utilised them. It was not possible for us to have direct access to the CLINICOM system to extract CAMH-related data. It was equally surprising that summative data, as presented in annual reports, were never disaggregated to present findings on children and adolescents separately from information on adults. There were no DoH or mental health reports in 2016 that included references to CAMH [19]. To our knowledge, there has never been a dedicated report from the provincial Department of Health on child and adolescent mental health.

Research in child and adolescent mental health

The three specialist CAMH units were linked to two universities (University of Cape Town and Stellenbosch University) and provided infrastructure and supervision for research in CAMH. In 2016 there were five research programmes, all led from the University of Cape Town, but with collaborations at Stellenbosch University. The programmes included the Adolescent Health Research Unit, the Centre for Autism Research in Africa, a Tuberous Sclerosis Complex Research Programme, a staff research development programme, and an Infant Mental Health research programme.

We performed a brief literature search for all publications relevant to CAMH published in 2016 by authors from the Western Cape or on participants from the Western Cape. We used Web of Science (version 5.34). Core Collection and the following search terms: "child AND adolescent AND mental AND health AND South Africa"; "autism AND South Africa"; "ADHD AND South Africa"; "adolescence AND South Africa"; "infant AND mental AND health AND South Africa" and "tuberous sclerosis". We then refined searches by selecting affiliations with institutions in the Western Cape. All identified abstracts were reviewed for relevance to CAMH. A total of 44 abstracts were identified. After removal of duplicates and abstracts not directly relevant to CAMH, a total of 26 articles were included [36–63]. A brief thematic grouping identified two articles on infant mental health, four on autism or ADHD, seven on adolescent mental health and five on tuberous sclerosis complex (TSC). It was of interest that a further six articles related to mental health aspects of HIV/AIDS, with the remaining three focusing on other themes of particular relevance in a South African setting: child abuse, foetal alcohol spectrum disorder (FAS), and cultural adaptation of an instrument to identify psychopathology. Table 9 provides a summary of the research relevant to CAMH and published in 2016 as identified by our search.

Table 9

Publications of research relevant to child and adolescent mental health that included authors from the Western Cape or a focus on the Western Cape in 2016

Research theme	Topic of research	Reference (First author, journal, volume: pages)
Infant Mental Health	Infant mental health and early childhood[39]	Worthman, <i>Social Science and Medicine</i> , 154: 62–69
	Reflective practice in infant mental health[40]	Berg, <i>Infant Mental Health Journal</i> , 37: 684–691
Autism and ADHD	Autism in Africa[41]	de Vries, <i>Current Opinion in Neurology</i> , 29: 130–136
	Performance of South African children on the CSBS, a tool for autism[42]	Chambers, <i>International Journal of Language and Communication Disorders</i> , 51: 265–275
	Theory of mind in autism[43]	Hamilton, <i>Journal of Child and Adolescent Mental Health</i> , 28: 233–241
	Management of ADHD in children and adolescents: clinical audit of ADHD assessment and treatment[44]	Vrba, <i>Journal of Child and Adolescent Mental Health</i> , 28: 1–19
HIV/AIDS	HIV-associated neurocognitive disorders in 6–16-year olds[45]	Hoare, <i>Neurology</i> , 87: 86–93
	The impact of household HIV on child development[46]	Sherr, <i>Child Care Health and Development</i> , 42: 890–899
	Mental health resilience in children who lost parents to HIV/AIDS[47]	Collishaw, <i>Journal of Abnormal Child Psychology</i> , 44: 719–730
	Social support for children affected by HIV/AIDS[48]	Sharer, <i>Aids Care – Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 28: 110–117
	Resilience in HIV-affected adolescents in South Africa[49]	Bhana, <i>Aids Care – Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 28: 49–59
	Correlates of emotional and behavioural problems in children with perinatally-acquired HIV[50]	Louw, <i>Aids Care – Psychological and Socio-Medical Aspects of AIDS/HIV</i> , 28: 842–850
Adolescent Mental Health	Mental health inequalities in adolescents[51]	Das-Munshi, <i>PloS One</i> , 11: 5
	Adolescent substance abuse[52]	Weybright, <i>Journal of Adolescence</i> , 49: 158–169
	Impact of family structure on adolescent psychological profile[53]	Daivids, <i>Journal of Psychology in Africa</i> , 26: 351–356
	Social protection and adolescent health[54]	Cluver, <i>PloS One</i> , 11: 10
	Parenting programme to prevent abuse of adolescents[55]	Cluver, <i>Trials</i> , 17: 328
	Reducing adolescent abuse in LMIC-[56]	Cluver, <i>BMC Public Health</i> , 16: 567
	Factors associated with readmission of adolescents discharged from inpatient units[57]	Pieterse, <i>Journal of Child and Adolescent Mental Health</i> , 28: [incomplete]
Tuberous Sclerosis Complex (TSC)	Clinical trial of everolimus for epilepsy in TSC[58]	French, <i>Lancet</i> , 388: 2153–2163
	Everolimus for neurocognitive problems in TSC[59]	Randell, <i>Trials</i> , 17: 398
	Long-Term use of everolimus for SEGA in TSC[60]	Franz, <i>PloS One</i> , 11: 6
	Towards an improved understanding of TSC-Associated Neuropsychiatric Disorders[61]	Leclezio, <i>Advances in Autism</i> , 2: 1–8
	Everolimus for renal angiomyolipomas in TSC[62]	Bissler, <i>Nephrology, dialysis and transplantation</i> , 31: 111–119
Other themes	Fatal child abuse [63]	Mathews, <i>South African Medical Journal</i> , 106: 22–25
	Theory of mind in children with FASD[64]	Lindinger, <i>Alcoholism – Clinical and Experimental Research</i> , 40: 367–376
	Cultural adaptation of the DISC-IV for Sotho-speaking South Africans [65]	Skinner, <i>Journal of Ethic & Cultural Diversity in Social Work</i> , 25: 1–19

Discussion

The aim of this study was to perform a fine-grained situational analysis of CAMHS in the Western Cape as a ‘case study’ for similar services in the rest country. Items from the WHO-AIMS 2.2 (Brief version) were adapted to capture information about the six fundamental domains of the CAMH health system in the province.

In terms of policy and legislative frameworks for CAMHS (WHO-AIMS Domain 1) There was an absence of any provincial CAMH policies or implementation plans [19]. This finding was in line with an earlier Policy analysis conducted by us [36]. We identified reasonable human rights policies and procedures for children. Of note was the consistent lack of disaggregated budgets for CAMHS. It was therefore not possible to comment on financing of CAMHS in the province. The budget for mental hospitals represented only 3% of the total health budget in the 2016/17 financial year. The CAMH budget would therefore have been a fraction of that 3%.

Policy development reflects commitment from the (provincial) government and relevant authorities towards the services, and provides a mandate to support funding mechanisms [64]. Lack of this important hardware element for CAMHS in the Western Cape therefore raises questions about the commitment of the provincial government and authorities towards strengthening of CAMHS. The lack of dedicated funding for CAMH was a second related concern. Our observations are underlined by the findings of Mokitimi and colleagues [65] and Docrat and colleagues [7] who confirmed disproportionately low expenditure on CAMHS in comparison to adult mental health services for the same period (January to December 2016). Adequate provision of resources in a health system requires adequate financing to ensure that a good enough service is delivered [4].

In terms of clinical services for CAMH (Domain 2), there was no dedicated CAMH authority. This signals a lack of dedicated leadership and governance of CAMH. Leadership and governance in a health system is important to ensure coordination of the services and equitable allocation of available resources [4]. Outpatient, but not inpatient, services were broadly organised in terms of catchment areas and across three levels of care (primary, secondary and tertiary). It was with great difficulty that we were able to disaggregate clinical data to identify that 8,300 under-18-year-olds were seen for mental health problems at primary, and 1,145 at secondary level. This represented 4.4% and 7.27% respectively of all people seen for mental health problems at primary and secondary levels of care. We were only able to find outpatient tertiary data for one of the three specialist CAMH units, where 3,639 children and adolescents were seen as outpatients in 2016. A total of 795 under-18-year-olds were admitted to inpatient units in secondary care (representing 4.35% of all secondary care mental health admissions) and a total of 346 in specialist CAMH inpatient units.

It was of concern to observe that all mental health services for children and adolescents at primary and secondary levels of care were mixed with adults. We were particularly concerned by the admission of eleven under-19-year-olds to adult mental hospitals. Again this finding concurs with Mokitimi and colleagues [65]. At all levels of care separate CAMHS should exist and adaptations are required to meet the needs of children and adolescents [4, 9]. Adult psychiatric environments are not conducive to the mental health and well-being of children and adolescents. It can jeopardise the safety of young people and can be frightening and traumatising, having a direct negative impact on the quality of care for children and adolescents [66]. International and national guidelines have previously recommended dedicated CAMHS [4, 9] and even the Western Cape General Health Policy [16] set out a plan to separate children and adolescents from adult psychiatric services. However, the findings presented here suggest that no implementation of these guidelines had taken place by the end of 2016.

A total of 63.8% of the Western Cape population lived in the City of Cape Town in 2016 [10]. Of all under-19-year-olds seen in primary care, 76.3% lived in the City of Cape Town. In contrast, at secondary care level, only 48.9% of the same age group outpatients and 59.5% of inpatients were in the metropolitan area. In the rural districts, by contrast, a greater proportion of under-18-year-olds were seen at secondary care facilities. All specialist CAMH units were based in the City of Cape Town. These results suggest that the lack of easy and close access to specialist CAMHS in the rural districts of the province may have added additional service pressure at the secondary level of care. This would be a very important empirical question to explore in future health systems research.

In terms of access to interventions, levels 1 and 2 primarily provided access to medication treatment, with little or no psychosocial support. Only at specialist CAMH level did children and adolescents have access to a range of medication, psychosocial treatment and the support of multidisciplinary teams.

Domain 3 of the WHO-AIMS focuses on CAMH in primary care (level 1). Even though 8,300 children and adolescents were seen in primary care settings, and despite the fact that there was anecdotal evidence of refresher training on CAMH for primary care staff, there were no sources documenting any access to refresher training on CAMH to PHC staff. As alluded to earlier, primary care staff had access to psychotropic medications on the essential drugs list, but very little if any access to psychosocial support. It is therefore of concern that we were not able to find clear data or evidence of consistent and sustained support to service providers, particularly at primary levels of care, where the majority of children and adolescents will first present with potential mental health problems. This observation is of particular importance in the context of South African prioritisation of public health and primary healthcare services.

It was difficult to access data on human resources for CAMH (Domain 4 of the WHO-AIMS) given that all CAMH staffing data were combined with adult mental health staff data in formal provincial datasets. Direct collection of data from specialist CAMH units allowed the basic description of the multidisciplinary staffing in these units. Data indicated high clinical and teaching activity in specialist CAMH units, but very limited dedicated resources for staff training and capacity-building, for instance, in training posts in CAMH.

In the 2016 data we could only identify one clear example of a joint initiative across Provincial Government Agencies (Health, Education, Social Development, Justice), focused on CAMH. The 'First 1,000 Days of Life' was cited as a positive initiative around maternal mental and physical health, and infant mental health. There was little evidence of other joint initiatives between sectors (WHO-AIMS, Domain 5). Again this lack of intersectoral collaboration and inequitable distribution of CAMH resources mirrors findings from a senior stakeholder group [65].

Domain 6 captured information on monitoring and research. Even though all levels of care apparently had information systems to collect data on patient care, we were not able to access any of the primary data collected through the CLINICOM system, and were not able to get access to any data where under-18-year-old mental health activity data were clearly disaggregated from other mental health data. This raised the question about how 'fit for purpose' the health information systems in the Western Cape may be for CAMH. Based on our observations, the existing information systems which lacks categories that would allow for disaggregation, may not be able to provide adequate and sufficiently nuanced information to help identify the problem areas in CAMH services, in order to inform the planning for responsive services and programmes [67], or for research on CAMH.

We were encouraged to see active research in CAMH in 2016 across a number of broad thematic domains, and that a significant proportion of CAMH research was on topics of particular resonance in the African context (e.g. HIV/AIDS, FASD, resilience, and child abuse).

Taking together the findings in the situational analysis, we identified a number of the gaps that will require urgent action. Table 10 shows an overview of our findings. Gaps were identified in all health systems domains, and at all levels of the healthcare system. These included the previously identified lack of policy

development and implementation for CAMHS, lack of resources (financial, governance, infrastructural and human), lack of public education, and lack of access to information services that could support monitoring and research in CAMHS.

Table 10
Health system gaps in child and adolescent mental health in the Western Cape as identified in this situational analysis

WHO-AIMS Domain	Gaps in CAMHS
Domain 1 Policy and legislative framework	<ul style="list-style-type: none"> • No provincial CAMH policy or implementation plans • No dedicated financing for CAMHS
Domain 2 Clinical services for children and adolescents with mental health disorders	<ul style="list-style-type: none"> • No dedicated leadership and governance structure for CAMH • No dedicated CAMHS lack of psychosocial interventions at secondary level • No specialist CAMHS in rural districts
Domain 3 CAMH in primary healthcare	<ul style="list-style-type: none"> • Inadequate documentation on training of professionals on CAMHS at primary care level • Lack of dedicated resources at secondary and tertiary care levels to support and train colleagues at primary level, particularly in rural districts • Lack of psychosocial interventions at primary care level
Domain 4 Human resources	<ul style="list-style-type: none"> • Limited information systems to access human resources data on CAMH • Limited human resources for CAMH at secondary and tertiary care levels
Domain 5 Public education and links with other sectors	<ul style="list-style-type: none"> • Limited public health campaigns on CAMH • Limited intersectoral collaboration about CAMH
Domain 6 Monitoring and research	<ul style="list-style-type: none"> • Lack of disaggregated and accessible information systems for CAMH

In recent work by Docrat and colleagues [7], a national survey of mental health costs was performed, using data for the same period as used in our study (2016/17). Apart from finding that the public mental health budget represented only 5% of the overall public health budget, they estimated that ~ 7.5% of the total uninsured population required some form of outpatient mental health care, and ~ 0.89% required inpatient mental health services [7]. Applying the same estimates to CAMH (as a rough calculation), we would have expected 157,500 children in the Western Cape to have required outpatient and 18,690 inpatient care. The data identified in this situational analysis identified a fraction of such expected numbers across all levels of care: 6.3% (9,922.5/157,500) of expected outpatient visits and 7.2% (1,346/18,690) of expected inpatient admissions. It is therefore very sobering to conclude that existing mental health services across all levels of care reached fewer than 10% of children and adolescents who may have needed mental healthcare in 2016. Of those reached, only those who accessed specialist CAMHS were likely to have received comprehensive evidence-based care.

Limitations Of The Study

The main limitation of this situational analysis was challenges of accessing accurate and disaggregated data through existing health information systems. We therefore acknowledge that our findings may not have been representative of the actual CAMH clinical activity and resources in the Western Cape in 2016. However, we tried to be rigorous and systematic in data collection and presented all data sources in a systematic way to optimise the accuracy of data collected. We recognise the lack of clear and accessible information systems as a limitation of our study, but also note this as a major gap in CAMHS in the province. We also acknowledge that, given the period of data identified and discussed (January–December 2016), these represent an historical analysis and that some of the findings may have changed since 2016. Finally, we acknowledge that we focused our analysis on the Department of Health to the exclusion of all the other sectors (Education, Social Development, Non-Profit etc.) that are also of fundamental importance in CAMH. Broadening out the focus to other sectors and to inter-sectorality would be a very important area for future CAMH research in the South African context.

Conclusions

In comparison to the majority of provinces in South Africa, the Western Cape is seen as relatively well-resourced in terms of health services. Our findings of very limited hardware for CAMHS (policies, dedicated funding, staffing, training, information systems), and clinical care provision to potentially fewer than 10% of the expected population of children and adolescents in need of mental healthcare, raises significant concerns about CAMHS, not only in the Western Cape, but also elsewhere in the country and in similar low-resource settings. Sadly, our findings are in keeping with previous literature that showed neglect of CAMH in South Africa and other LMIC despite the well-recognised burden of CAMH disorders [2, 9, 12, 14, 36, 65, 68]. While we focused on the hardware elements of CAMHS, our findings concurred with our study that explored the hardware and software elements of CAMHS [66]. Even though it will be important to expand our research to further explore software elements of the health system (e.g. through the experience of grassroots level service providers and users), the data presented here are already a clear call for action to find strategies and initiatives that will strengthen CAMHS in the province, the country and in other LMIC.

The WHO-AIMS was developed as a tool for situational analysis of adult mental health services and, in its original form, only have one item about CAMH services. As it stands, it is therefore not of particular use to generate a detailed situational analysis of CAMH in any country. We made a range of adaptations to the BRIEF version of the WHO-AIMS in order to collect much more breadth and depth about CAMH resources in our study. To our knowledge no other country has used a similar approach to generate a situational analysis specifically on CAMH. Given the vital importance of CAMH services throughout the world, but particularly in LMIC settings, development of a bespoke framework for CAMH situational analysis may be a very helpful action by the WHO. The version as adapted by us in this study may provide a useful starting place for such a global process. Taking together all service delivery elements from the

thesis, we therefore have to conclude that the health system in the province (Western Cape) was unable to meet the mental healthcare needs of children and adolescents in the province with a disproportionate impact in rural districts.

Abbreviations

CAMH
Child and Adolescent Mental Health
CAMHS
Child and Adolescent Mental Health Services
DCAP
Division of Child and Adolescent Psychiatry
DoH
Department of Health
DSD
Departement of Social Development
DSN
Data Source Numbers
GSAs
Geographical Service Areas
LMIC
Low- and Middle-Income Countries
PHC
Primary Health Care
WCED
Western Cape Education Departement
WHO-AIMS
World Health Organization Assessment Instrument of Mental Health Systems
WHO
World Health Organization

Declarations

Ethics approval and consent to participate

This study was approved by the UCT Human Research Ethics Committee (HREC 188/2016) and permission to conduct the study was received from the Western Cape Department of Health. The study adhered to the principles as set out in the Helsinki Declaration (2013).

Consent for publication

Not applicable

Availability of data material

The majority of data included are publically available. All other data are available from the authors.

Competing interests

The authors declare that they have no competing interests

Funding

To SM:

Research Funding: Department of Psychiatry, University of Cape Town

Spirit of 68 Scholarships: University of Cape Town

The Western Cape Department of Health

South African Medical Research Council PhD studentship

To PJdV:

National Research Foundation

Struengmann Fund

Authors' contributions

All three authors participated in the conception and design of the study, as well as reviewing the content for submission.

SM performed data collection, analysis, and interpretation of the data and prepared the first draft of the manuscript.

PJdV & MS contributed to analysis, interpretation of results and writing of the manuscript.

All authors approved the final version of the manuscript.

Acknowledgements

Prof Sharon Kleintjes for helpful discussions and advice, Marinda Roelofse for invaluable support around the Western Cape mental health system. Also to Drs Wendy Vogel, Rene Nassen and Sue Hawkridge, all heads of health facilities, and provincial administration staff for their support.

References

1. Juengsiragulwit D. Opportunities and obstacles in child and adolescent mental health services in low- and middle-income countries: a review of the literature. *Journal of Public Health*. 2015;4(2):110-22. doi:10.4103/2224-3151.206680.
2. Kleintjes S, Lund C, Flisher AJ, MHaPP Research Programme Consortium. A situational analysis of child and adolescent mental health services in Ghana, Uganda, South Africa and Zambia. *African Journal of Psychiatry*. 2010;13(2):132-9. doi:10.4314/ajpsy.v13i2.54360.
3. Gilson L. Health Policy and Systems Research: A Methodology Reader. 2012. https://www.who.int/alliance-hpsr/alliancehpsr_reader.pdf. Accessed 10 March 2020
4. World Health Organisation. Strengthening health systems to improve health outcomes. 2007. https://www.who.int/healthsystems/strategy/everybodys_business.pdf. Accessed 10 November 2017.
5. Mayosi BM, Benatar SR. Health and Health Care in South Africa — 20 Years after Mandela. *The New England Journal of Medicine*. 2014;371(14):1344-53. doi:10.1056/NEJMSr1405012.
6. The World Bank. World Bank Country and Lending Groups. 2018. <https://datahelpdesk.worldbank.org/knowledgebase/articles/906519-world-bank-country-and-lending-groups>. Accessed 18 May 2020.
7. Docrat S, Besada D, Cleary S, Daviaud E, Lund C. Mental health system costs, resources and constraints in South Africa: a national survey. *Health Policy Plan*. 2019;34:706-19. doi: 10.1093/heapol/czz085.
8. Lund C, Kleintjes S, Kakuma R, Flisher AJ, MHaPP Research Programme Consortium. Public sector mental health systems in South Africa: inter-provincial comparisons and policy implications. *Social Psychiatry and Psychiatric Epidemiology*. 2010;45(3):393–404. doi:10.1007/s00127-009-078-5.
9. Flisher A, Andrew D, Zuhayr K, Lund C, Sorsdahl K, Bronwyn M, et al. Child and adolescent mental health in South Africa. *Journal of Child and Adolescent Mental Health*. 2012, 24:149-161. .doi: 10.2989/17280583.2012.735505.
10. Statistics South Africa. Mid-year population estimates. 2016. <http://cs2016.statssa.gov.za/>. Accessed 10 November 2016 210.
11. Kleintjes S, Flisher AJ, Flick M, Railoun A, Lund C, Molteno C, et al. The prevalence of mental disorders among children, adolescents and adults in the Western Cape, South Africa. *South African Psychiatry Review*. 2006;9(3):157-60. doi: 10.4314/ajpsy.v9i3.30217.
12. Lund C, Flisher AJ. Norms for Mental Health Services in South Africa. *Social Psychiatry and Psychiatric Epidemiology*. 2006; 41(7):587-94. doi:10.1007/s00127-006-0057-z.
13. World Health Organisation. World Health Organisation Assessment Instrument for Mental Health Systems version 2.2 of 2005. 2005. http://www.who.int/mental_health/evidence/AIMS_WHO_2_2.pdf. Accessed 10 May 2016.
14. Petersen I, Bhana A, Campbell-Hall V, Mjadu S, Lund C, Kleintjes S, et al. Planning for district mental health services in South Africa: a situational analysis of a rural district site. *Health Policy Plan*. 2009;24(2):140-50. doi:10.1093/heapol/czn049.
15. Department of Health. Republic of South Africa. Policy Guidelines. Child and Adolescent Mental Health. 2003. <https://www.gov.za/documents/policy-guidelines-child-and-adolescent-mental-health#> Accessed 15 March 2021.
16. Department of Health. Western Cape Government. Republic of South Africa. Healthcare 2030: The Road to Wellness. 2014. <https://www.westerncape.gov.za/assets/departments/health/healthcare2030.pdf> . Accessed 10 June 2016.
17. Mental Health Care Act no. 17 of 2002. 2004. http://www.justice.gov.za/legislation/acts/2002-017_mentalhealthcare.pdf. Accessed 6 June 2016.
18. Department of Social Development. Republic of South Africa. (1 February 1987). Child Care Act 74 of 1983. https://www.westerncape.gov.za/assets/departments/social-development/child_care_act_74_of_1983.pdf. Accessed 16 May 2020.

19. Mokitimi S. Indepth interview with the Western Cape Provincial Deputy-Director for Mental Health and Substance Abuse, February 9, 2017.
http://www.cara.uct.ac.za/sites/default/files/image_tool/images/444/A.%20Interview%20with%20the%20Western%20Cape%20Provincial%20Deputy%20
Accessed 12 May 2021.
20. Western Cape Government. Provincial Mental Health Directory 2015.2015. <https://pmhp.za.org/wp-content/uploads/DoH-Mental-Health-Resource-Directory-2015.pdf>. Accessed 10 May 2016.
21. Western Cape Government. Budget 2016 Summary. 2016.
https://www.westerncape.gov.za/assets/departments/treasury/Documents/Budget/2016/2016_budget_summary_budget_day_3_march_2016.pdf.
Accessed 10 November 2017.
22. Provincial Treasury. Western Cape Government. Budget Overview of Provincial Revenue and Expenditure 2016.
https://www.westerncape.gov.za/assets/departments/treasury/Documents/Budget/2016/2016_overview_of_prov_rev_exp_march_web.pdf. Accessed 17 March 2019.
23. Provincial Treasury. Western Cape Government. Budget Estimates of Provincial Revenue and Expenditure 2016.
https://www.westerncape.gov.za/assets/departments/treasury/Documents/Budget/2016/2016_estimates_prov_rev_exp_march_2016_incl_addendums.pdf
Accessed 17 March 2019.
24. Western Cape Government. Mental Health Services in the Western Cape. 2016. <https://www.westerncape.gov.za/general-publication/mental-health-services-western-cape>. Accessed 10 November 2017.
25. Western Cape Government. Division of Child and Adolescent Psychiatry (DCAP). 2016.
https://www.westerncape.gov.za/dept/health/documents/public_info/D/39513?toc_page=7. Accessed 10 November 2017.
26. Western Cape Government. (n.d). Mental Health Hospital Services. <https://www.westerncape.gov.za/service/mental-health-hospital-services>. Accessed 10 November 2017.
27. Mokitimi S. Indepth interview with the Head of Clinical Unit, Division of Child and Adolescent Psychiatry, March 12, 2020.
http://www.cara.uct.ac.za/sites/default/files/image_tool/images/444/B.%20Catchment%20areas%20for%20specialist%20child%20and%20adolescent%20
Accessed 12 May 2021.
28. Mokitimi S. Provincial Mental Health Information on CAMH (Mental Health Data). 2016.
http://www.cara.uct.ac.za/sites/default/files/image_tool/images/444/C.%20Additional%20data%20tables%20-%20Mokitimi%20et%20al.pdf. Accessed 12 May 2021.
29. Western Cape Government. Tygerberg Hospital Annual Report 2016. 2016.
https://www.westerncape.gov.za/sites/www.westerncape.gov.za/files/tygerberg_hospital_annual_report_2016_web.pdf. Accessed 24 November 2017.
30. Department of Social Development (Western Cape Government). (n.d). What is Diversion?. <https://www.westerncape.gov.za/general-publication/what-diversion>. Accessed 16 May 2020.
31. Department of Health. Republic of South Africa. Standard Treatment Guidelines and Essential Medicines List for South Africa Hospital Level Paediatrics 2017 Edition. 2017. The National Department of Health, Pretoria, South Africa . <http://www.health.gov.za/index.php/standard-treatment-guidelines-and-essential-medicines-list/category/456-hospital-level-paediatrics>. Accessed 10 December 2019.
32. Western Cape Government. First 1 000 Days Campaign. 2017. <https://www.westerncape.gov.za/general-publication/first-1-000-days-campaign>. Accessed 13 August 2019.
33. 2016 Annual Report, Salesian Life Choices. https://www.lifechoices.co.za/sites/default/files/2017-10/lc_annual_report_2016-final_update_2.pdf.
Accessed 15 March 2021
34. Western Cape Government. How to handle bullying. 2016. <https://www.westerncape.gov.za/general-publication/how-handle-bullying>. Accessed 10 August 2017.
35. Western Cape Government. 16 Days of Activism For no violence against Women and Children. 2016. <https://www.westerncape.gov.za/general-publication/what-16-days-activism>. Accessed 17 November 2017.
36. Mokitimi S, Schneider M, de Vries PJ. Child and adolescent mental health policy in South Africa: History, current policy development and implementation, and policy analysis. *International Journal of Mental Health Systems*. 2018;12(36). doi:10.1186/s13033-018-0213-3
37. Shatkin, J. P., & Belfer, M. L. (2004). The Global Absence of Child and Adolescent Mental Health Policy. *Child and Adolescent Mental Health*, 9(3), 104–108. doi:10.1111/j.1475-3588.2004.00090.x
38. Worthman CM, Tomlinson M, Rotheram-Borus MJ. When can parents most influence their child's development? Expert knowledge and perceived local realities. *Soc Sci Med*. 2016 (154): 62–69.doi: 10.1016/j.socscimed.2016.02.040.
39. Berg, A. Reflective practice in infant mental health—A South African perspective. *Infant Mental Health Journal*. 2016, 37(6), 684–691. doi.org/10.1002/imhj.21605.
40. de Vries PJ. Thinking globally to meet local needs: autism spectrum disorders in Africa and other low-resource environments. *Current Opinion in Neurology*. 2016;29(2):130-6. doi: 10.1097/WCO.0000000000000297.
41. Chambers N, Stronach ST, Wetherby AM. Performance of South African children on the Communication and Symbolic Behavior Scales—Developmental Profile (CSBS DP). *International Journal of Language and Communication Disorders*. 2016. 51(3): 265–275 .doi.org/10.1111/1460-6984.12203.
42. Hamilton K, Hoogenhout M, Malcolm-Smith S. Neurocognitive considerations when assessing Theory of Mind in Autism Spectrum Disorder. *Journal of Child and Adolescent Mental Health*. 2016 28(3):233-241.doi: 10.2989/17280583.2016.1268141.

43. Vrba K, Vogel W, de Vries PJ. Management of ADHD in children and adolescents: clinical audit in a South African setting. *Journal of Child and Adolescent Mental Health*. 2016;28(1):1-19. doi: 10.2989/17280583.2015.1128437.
44. Hoare J, Phillips N, Joska JA, Paul R, Donald KA, Stein DJ, et al. Applying the HIV-associated neurocognitive disorder diagnostic criteria to HIV-infected youth. *Neurology*. 2016;87(1):86-93. doi: 10.1212/WNL.0000000000002669.
45. Sherr L, Skeen S, Hensels IS, Tomlinson M, Macedo A. The effects of caregiver and household HIV on child development: a community-based longitudinal study of young children. *Child: Care, Health and Development*. 2016. 42(6): 890–899. doi.org/10.1111/cch.12387.
46. Collishaw B, Gardner F, Aber JL, Cluver L. Predictors of Mental Health Resilience in Children who Have Been Parentally Bereaved by AIDS in Urban South Africa. *Journal of Abnormal Child Psychology*. 2016 ;44(4):719-30. doi: 10.1007/s10802-015-0068-x.44.
47. Sharer M, Cluver L, Shields JJ & Ahearn F. The power of siblings and caregivers: under-explored types of social support among children affected by HIV and AIDS. *Psychological and Socio-medical Aspects of AIDS/HIV*. 2016; 28:110-117. doi.org/10.1080/09540121.2016.1178942.
48. Bhana A, Mellins CA, Small L, Nestadt DF, Leu C, Petersen I, et al. Resilience in perinatal HIV+ adolescents in South Africa. *AIDS Care*. 2016;28(2): 49–59. doi: 10.1080/09540121.2016.1176676.
49. Louw K N, Ipser J, Phillips N & Hoare J. Correlates of emotional and behavioural problems in children with perinatally acquired HIV in Cape Town, South Africa. *AIDS Care*. 2016: 28(7), 842-850. doi: 10.1080/09540121.2016.1140892.
50. Das-Munshi J, Lund C, Mathews M, Clark C, Rothon C, Stansfeld S. Mental Health Inequalities in Adolescents Growing Up in Post-Apartheid South Africa: Cross-Sectional Survey, SHaW Study. *PloS One*, 2016, 11:5. doi.org/10.1371/journal.pone.0154478.
51. Weybright EH, Caldwell LL, Ram N, Smith EA, Wegner L. Trajectories of adolescent substance use development and the influence of healthy leisure: A growth mixture modeling approach. *Journal of Adolescence*. 2016, 49:158–169. doi: 10.1016/j.adolescence.2016.03.012.
52. Davids, EL, Ryan J, Yassin Z, Hendrickse S, Roman NV. Family structure and functioning: Influences on adolescents psychological needs, goals and aspirations in a South African setting. *Journal of Psychology in Africa*. 2016, 26(4): 351 – 356. doi.org/10.1080/14330237.2016.1208929.
53. Cluver LD, Orkin FM, Meinck F, Boyes ME, Yakubovich AR, Sherr L. Can Social Protection Improve Sustainable Development Goals for Adolescent Health? *PLoS ONE*. 2016. 11:10. doi.org/10.1371/journal.pone.0164808
54. Cluver L, Meinck F, Shenderovich Y, Ward CL, Romero RH, Redfern A, et al. A parenting programme to prevent abuse of adolescents in South Africa: study protocol for a randomised controlled trial. *Trials*. 2016, 17:328. doi.org/10.1186/s13063-016-1452-8.
55. Cluver L, Meinck F, Yakubovich A, Doubt J, Redfern A, Ward C, et al. Reducing child abuse amongst adolescents in low- and middle-income countries: A pre-post trial in South Africa. *BMC Public Health*. 2016, 16:567. doi.org/10.1186/s12889-016-3262-z.
56. Pieterse D, Temmingh H, Vogel W. Factors associated with readmission in South African adolescents discharged from two inpatient psychosocial rehabilitation units. *Journal of Child and Adolescent Mental Health*. 2016;28(3):199-212. doi: 10.2989/17280583.2016.1259165. PMID: 27998264.
57. French JA, Lawson JA, Yapici Z, Ikeda H, Polster T, Nabbout R, et al. Adjunctive everolimus therapy for treatment-resistant focal-onset seizures associated with tuberous sclerosis (EXIST-3): a phase 3, randomised, double-blind, placebo-controlled study. *Lancet*. 2016. 388(10056), 2153–2163. doi.org/10.1016/S0140-6736(16)31419-2.
58. Randell E, McNamara R, Davies DM, Jones EO, Kirby N, Angel L, et al. *Trials*. The use of everolimus in the treatment of neurocognitive problems in tuberous sclerosis (TRON): study protocol for a randomised controlled trial. 2016. 17:398. doi 10.1186/s13063-016-1446-6.
59. Franz DN, Belousova E, Sparagana S, Bebin EM, Frost MD, Kuperman R, et al. Long-Term Use of Everolimus in Patients with Tuberous Sclerosis Complex: Final Results from the EXIST-1 Study. *PloS One*. 2016, 11: 6. doi.org/10.1371/journal.pone.0158476.
60. Leclezio L, de Vries PJ. Towards an improved understanding of TSC-Associated Neuropsychiatric Disorders (TAND). *Advances in Autism*. 2016, 2(2):76-83. doi: 10.1108/AIA-12-2015-0025.
61. Bissler JJ, Kingswood JC, Radzikowska E, Zonnenberg BA, Frost M, Belousova E, et al. Everolimus for renal angiomyolipoma in patients with tuberous sclerosis complex or sporadic lymphangioleiomyomatosis: extension of a randomized controlled trial. *Nephrol Dial Transplant*. 2016; 31(1):111-9. doi: 10.1093/ndt/gfv249.
62. Mathews S, Martin LJ. Developing an understanding of fatal child abuse and neglect: Results from the South African child death review pilot study. *South African Medical Journal*. 2016;106(12):1160-1163. doi:10.7196/SAMJ.2017.v106i12.12130.
63. Lindinger NM, Malcolm-Smith S, Dodge NC, Molteno CD, Thomas KGF, Meintjes EM, et al. Theory of Mind in Children with Fetal Alcohol Spectrum Disorders. *Alcoholism Clinical and Experimental Research*. 2016. 40(2):367-376. doi: 10.1111/acer.12961.
64. Skinner D, Sharp CS, Serekoane M & Ross MR. The Cultural Adaptation of the DISC-IV: Appropriateness for Sotho-Speaking South Africans. *Journal of Ethnic & Cultural Diversity in Social Work*. 2016, 25 (1). 1-19. doi.org/10.1080/15313204.2015.1129521.
65. Mokitimi S, Jonas K, Schneider M, de Vries PJ. Child and Adolescent Mental Health Services in South Africa—Senior Stakeholder Perceptions of Strengths, Weaknesses, Opportunities and Threats in the Western Cape Province. *Frontiers in Psychiatry*. 2019; 10. doi: 10.3389/fpsy.2019.00841
66. Shatkin JP, Belfer ML. The Global Absence of Child and Adolescent Mental Health Policy. *Child and Adolescent Mental Health*. 2004; 9(3):104-8. doi:10.1111/j.1475-3588.2004.00090.x
67. Gerson R, Havens J. The Child and Adolescent Psychiatric Emergency: A Public Challenge. *Psychiatric Times*. 2015;32(11). <https://www.psychiatristimes.com/view/child-and-adolescent-psychiatric-emergency-public-health-challenge>. Accessed 16 July 2020.
68. World Health Organisation. Monitoring the building blocks of health systems: a handbook of indicators and their measurement strategies 2010. https://www.who.int/healthinfo/systems/WHO_MBHSS_2010_full_web.pdf?ua=1World. Accessed 12 July 2019.
69. Kapp SA, Petr CG, Robbins ML, Choi JJ. Collaboration Between Community Mental Health and Juvenile Justice Systems: Barriers and Facilitators. *Child Adolescent Social Work Journal*. 2013;30(6):505-517. doi:10.1007/s10560-013-0300x.

Figures

Figure 1

Map of the Western Cape Province showing the metropolitan area (the City of Cape Town and its substructures) and rural health districts. Urban areas are indicated in green; rural areas are indicated in blue. The figure also shows the location of regional hospitals and specialist CAMH units.

Figure 2

Number and sex of children and adolescents treated as outpatients at the Division of Child and Adolescent Psychiatry, University of Cape Town in 2016.

Figure 3

Self-declared ethnicity of children and adolescents treated as outpatients at the Division of Child and Adolescent Psychiatry in 2016.

Figure 4

Staff proportions for mental health services in the Western Cape in 2016

Figure 5

Human resources in specialist child and adolescent mental health units in 2016

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [MokitimiTable3120521.docx](#)