

Healthcare Utilization and Out-of-Pocket Expenditures Associated with Depression in Adults: A Cross-Sectional Analysis in Nepal

Selina Rajan

London School of Hygiene and Tropical Medicine Population Studies Group

Sujit D Rathod

London School of Hygiene and Tropical Medicine Department of Population Health

Nagendra P Luitel (✉ luiteln@gmail.com)

Transcultural Psychosocial Organization (TPO) Nepal <https://orcid.org/0000-0002-8291-0205>

Adrianna Murphy

London School of Hygiene and Tropical Medicine Department of Population Health

Tessa Roberts

London School of Hygiene and Tropical Medicine Department of Population Health

Mark JD Jordans

King's College London

Research article

Keywords: healthcare, Nepal, Adults, depression, cost

Posted Date: December 31st, 2019

DOI: <https://doi.org/10.21203/rs.2.14587/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Health Services Research on March 25th, 2020. See the published version at <https://doi.org/10.1186/s12913-020-05094-9>.

Abstract

Background: Despite attempts to improve universal healthcare coverage (UHC) in low income countries like Nepal, most healthcare utilization is still financed by out-of-pocket (OOP) payments, with detrimental effects on the poorest and most in need. Evidence from high income countries shows that depression is associated with increased healthcare utilization, which may lead to increased OOP expenditures, placing greater stress on families. To inform policies for integrating mental healthcare into UHC in LMIC, we must understand healthcare utilization and OOP expenditure patterns in people with depression.

Aims: We examined associations between symptoms of depression and frequency and type of healthcare utilization and OOP expenditure among adults in Chitwan District, Nepal.

Methods: We analysed data from a population-based survey of 2040 adults in 2013, who completed the PHQ-9 screening tool for depression and answered questions about healthcare utilization. We modelled associations between increasing PHQ-9 score and healthcare utilization frequency and OOP expenditure using negative binomial regression. We also compared sector-specific utilization of outpatient healthcare and their related costs among adults with and without probable depression, determined by a PHQ-9 score of 10 or more.

Results: We classified 80 (3.6%) participants with probable depression, 70.9% of whom used some form of healthcare in the past year compared to 43.9% of people without probable depression. Mean annual OOP healthcare expenditures were \$118 USD in people with probable depression, compared to \$110 USD in people without. With each unit increase in PHQ-9 score, there was a 14% increase in total healthcare visits (95% CI 7%-22%, $p < 0.0001$) and \$9 USD increase in OOP expenditures (95% CI \$2-\$17; $p < 0.0001$). People with depression sought most healthcare from pharmacists (30.1%) but reported the greatest expenditure on specialist doctors (\$36 USD).

Conclusions: In this population-based sample from Central Nepal, we identified dose-dependent increases in healthcare utilization and OOP expenditure with increasing PHQ-9 scores. Strengthening UHC to include early detection and treatment for people affected by depression as an integrated component of healthcare should lead to a reduction in financial pressures on families, which is likely to reduce the incidence of depression in Nepal.

Introduction

In an international pursuit to reduce global poverty and improve population health, the world's governments have committed, in the Sustainable Development Goals, to achieve universal healthcare coverage (UHC), including financial risk protection by 2030.¹ A critical decision in the progressive realisation of UHC is which health services to include in essential packages of care. There is evidence that mental health interventions can be cost-effective^{2,3}, and several prominent mental health advocates have recommended their inclusion in UHC.⁴⁻⁷ Besides being a major contributor to the burden of

disability worldwide⁸, much of which falls on low- and middle-income countries (LMIC)⁹, there are also strong economic arguments for the inclusion of basic mental health care in UHC. The global economic burden of mental disorders comes to \$2.5 trillion USD per year, projected to double by 2030, of which two thirds arises from lost earnings and one third directly from healthcare expenditure².

To inform national policies for achieving UHC, and decisions around which interventions to prioritise in low-resource settings, we need to understand patterns of healthcare utilization and out-of-pocket payments (OOP). There is evidence from North America and Europe that healthcare expenditure is more than twice as high in people with depression than those without¹⁰⁻¹⁴ suggesting that this group places a disproportionate burden on health systems, whose needs are important to consider when planning UHC. However, it is unknown whether this finding is generalisable to other countries with vastly different health systems. In LMIC, where healthcare is predominantly financed by OOP payments¹⁵, any increased utilization can lead to impoverishment, itself a major risk factor for depression¹⁶, and can result in the poorest individuals foregoing necessary care¹⁷.

Nepal is a low-income fragile state, where prolonged conflict, trauma and natural disasters have contributed to a weakened health system in which 60% of total healthcare expenditure is privately funded, of which 80% is OOP.¹⁸ The Nepalese government has made significant efforts to improve UHC in the last 5 years, with social health insurance now available in 42 out of 77 of its districts.¹⁹ However, uptake remains low, with only 6% of the population signing up to the scheme¹⁹, and specialist mental health services remain unavailable for the majority of individuals with mental disorders.²⁰ At present there is no data from Nepal on healthcare utilization by people with depression, in order to inform policies around including depression treatment care in UHC to address this group's needs and reduce OOP spending. While there is some evidence that OOP expenditures increase among older people with depression in southern India,²¹ the available evidence from elsewhere in LMIC shows that community healthcare utilization by people with depression varies considerably between countries.²²

Recent evidence from the Programme for Improving Mental Health Care (PRIME)²³ demonstrates the feasibility^{24,25} and effectiveness of integrating mental health services into primary healthcare settings in Nepal, based on the WHO Mental Health Gap Action Programme (mhGAP).²⁶ Modelled estimates suggest that this kind of integration could be scaled-up nationally for an investment of \$5.55 US dollars per capita, spread over 7 years.²⁷ The PRIME approach provides a potential model upon which to base integrated services if mental health care were included in UHC. Understanding patterns of healthcare utilization and OOP expenditure by people with depression can help us to predict the likely financial and health service impacts of implementing this policy in Nepal.

In this secondary analysis of data from the PRIME study, we set out to examine associations between increasing symptoms of depression and healthcare utilization and related OOP expenditure among adults in one district in Central Nepal. In our definition of healthcare utilization, we included hospitalizations and any use of any outpatient healthcare services. These included both generalist and specialist doctors,

mental health specialists, pharmacists and providers of traditional and complementary medicine. Our secondary objective was to compare sector-specific utilization of outpatient healthcare among adults with and without probable depression and to estimate the mean expenditure associated with each provider type.

Methods

Setting

This study was carried out in Chitwan district in central Nepal, where some mental health services were already in existence at the time of study, as described previously.^{28,29} In previous studies, approximately 41% of people reported symptoms of depression in the north-western mountains,³⁰ compared to 3% in this sample from the central plains³¹, of whom only 8.1% accessed treatment.³¹ Despite Chitwan having adopted the National Mental Health Policy in 1996, there were only 2 psychiatrists serving its population of over half a million³² at the time of the study.

Sampling and participants

The rationale, study design and data collection procedures for the PRIME study have been detailed previously.³¹ Briefly, we conducted a population-based survey of adults in Chitwan District. Using household and population data from Village Development Committees (VDCs; similar to municipalities, with community participation in their administration) in Chitwan, the lead investigator randomly selected houses from each VDC, and field workers enumerated the adults in the household and a family member randomly selected 1 adult from a series of concealed papers for recruitment into the study. Eligibility criteria included age of 18 years or over, residency in Chitwan, fluency in the Nepali language and willingness and ability to provide informed consent. Between May and August 2013, 99% of eligible adults provided informed consent and 2040 adults participated in the study.

Each participant completed a two-part structured questionnaire. All participants completed part 1, which consisted of sections including demographic characteristics and screening for probable depression (described below) and alcohol use disorder. A total of 479 participants who either screened positive for depression or reported having chronic symptoms of depression (213) or alcohol-use disorder (78) or both (18) and 10% of randomly selected screen negative participants (170) completed part 2 as shown in Additional File 1. Part 2 consisted of sections including household economic status healthcare utilization and OOP expenditure for both inpatient (12-month recall) and outpatient (3-month recall) healthcare, which are described in more detail below. We used this design feature to facilitate the analysis of secondary research questions whilst maintaining sufficient power without overburdening research staff and participants. We used a threshold of 10% to enable collection of health services data on equal numbers of screen positive and screen negative participants, (based on an assumed prevalence of alcohol use disorder or depression of 10%) and used sampling weights to account for this in the analysis.

The questionnaire was orally administered by a trained fieldworker in the Nepali language and responses were collected on a questionnaire application, which was programmed onto an Android mobile device.

Measures

Depression status

We screened participants for depression with the 9-item Patient Health Questionnaire (PHQ-9)³³. The PHQ-9 grades self-reported symptoms of depression in the previous 2 weeks and severity is determined by increasing scores. We interpreted a score of 10 or more to reflect probable depression, which in a Nepali validation study had a sensitivity of 0.94 and specificity of 0.80.³⁴ In this sample, the PHQ-9 had a Cronbach's alpha of 0.79.³¹

Household Economic Status

Household economic status was evaluated on the basis of Household economic status was evaluated on the basis of household assets (such as water supply, sanitation facilities, power supply, radios, televisions, mobile phones, cooking fuel, a separate kitchen and type of flooring) using a household asset score, which is described further in Additional File 2 and was developed specifically for use in Nepal for the purposes of this study. Employing the methods described by Vyas and Kumaranayake³⁵, we used principal component analysis to generate a relative wealth index, which we subsequently categorized into thirds to reflect low, average and high wealth categories.

Healthcare utilization

We assessed healthcare utilization and OOP healthcare expenditure using a version of the Client Socio-demographic and Service Receipt Inventory³⁶, which has been adapted for use in LMIC.³⁷ To measure healthcare utilization, we asked participants to report the number of times they had been admitted to hospital in the previous 12 months, if at all. We then asked participants to report the number of times, if any, they had visited outpatient services for any health problem (including but not exclusively for depression) over the previous 3 months including seven types of healthcare providers: traditional healers, community workers, nurse/midwives, pharmacists, general doctors, specialist doctors, and psychiatrists and other mental health workers. In order to make comparisons between inpatient and outpatient healthcare utilization and to calculate an estimate of total utilization, we standardized the number of visits in 3 months to reflect annual outpatient healthcare utilization as reported in similar analyses.^{10,17,38} We also recorded data on the presenting health complaint.

Healthcare expenditure

For each inpatient hospital admission reported in the past 12 months, we asked participants to report all payments for hospital fees, medicines, biomedical investigations and transportation incurred both personally and by friends and family but for the interviewee's benefit only. We used the sum of all these payments to estimate the annual inpatient OOP expenditure. The inclusion of payments for transportation to and from health facilities in the definition of OOP expenditures is consistent with other OOP cost studies from LMIC.³⁹

For each episode of outpatient healthcare utilization over 3 months, we also asked participants to report all individual OOP payments for biomedical or traditional practitioners as well as return transportation. We also standardised these payments to reflect annual outpatient OOP expenditure. Finally, we summed the annual inpatient and outpatient OOP expenditures to estimate the total annual OOP expenditure. We did not include opportunity costs or indirect costs such as lost productivity and all costs were defined from the user's perspective. All expenditures were reported in Nepali Rupees and converted to US dollars according to the exchange rate at the end of data collection (1 USD:96.997 Nepali rupee on 02 Aug 2013). We observed one implausible outlier for outpatient OOP expenditure and replaced it with the sample's mean outpatient OOP cost.

Statistical analysis

First, we report participants' demographic⁴⁰ characteristics, PHQ-9 scores, frequency of healthcare utilization and mean OOP expenditures, stratified by depression screening status. Categorical measures are summarised as proportions, age as means and standard deviations and PHQ-9 scores and healthcare utilization (whose distribution was skewed), as medians and interquartile ranges. We present mean annual OOP expenditures, overall and by depression screening status, employing bootstrapping methods with 1000 replications to estimate standard errors. This method enables estimation of mean expenditures required for health budgeting⁴¹, whilst accounting for the skewed distribution typical of cost data^{42,43} and has been used with survey data of this nature previously.⁴⁴ One participant who had missing age was imputed to 39.8 years, the mean value for the overall sample. Second, we assessed the relative changes in annual inpatient, outpatient and total healthcare utilization for each unit increase in PHQ-9 score in all participants. Given the skew in the distribution of healthcare utilization, which precluded use of linear regression, we used negative binomial regression. Third, we estimated the relative changes in inpatient, outpatient and total OOP healthcare expenditure for each unit increase in PHQ-9 score using bootstrapped linear regression models. We adjusted for age, sex and relative wealth index in all models. We used the relative wealth index as a proxy for household economic status and specifically for access to basic necessities, which has been shown in Nepal to predict healthcare utilization.¹⁵ In exploratory analyses, we also adjusted for education and occupation as potential confounders of the association between depression and healthcare utilization but omitted these from the primary model to avoid collinearity. We also checked for interactions between PHQ-9 score and relative wealth group. Finally, we report utilization and mean OOP expenditures for each outpatient provider type, stratified by probable depression status.

For each step, survey-adjusted methods were used to account for the complex sampling design and sampling probability weights.⁴⁵ Data were analysed using Stata version 14.2.⁴⁰ Ethical approval was obtained from the Nepal Health Research Council (Kathmandu, Nepal), the World Health Organization (Geneva, Switzerland) and the London School of Hygiene and Tropical Medicine.

Results

Demographic Characteristics, PHQ–9 Scores, Healthcare Utilization, and OOP Expenditures

Table 1 displays the demographic characteristics, PHQ–9 scores, health care utilization and expenditure of community survey participants. The mean age of all 2040 participants was 39.8 years (SD 15.2), of whom 59.8% were female and 54.5% had secondary level education or higher. The median PHQ–9 score was 2 with a range 0 to 24 and the point prevalence of probable depression (PHQ–9 \geq 10) was 3.6%. Of all 479 adults who answered questions about healthcare utilization, 233 had not used any healthcare, whereas 18 had been admitted to hospital, 197 had utilised outpatient services and 31 had used both. People with probable depression were twice as likely to be admitted to hospital (15.8%) as people without (7.8%) and were also more likely to visit outpatient services (69.7%) than people without (39.8%). Nearly all participants who utilised healthcare incurred some OOP expenditure and the mean total OOP expenditure was \$118 USD/year (SD 265) in adults with probable depression and \$110 USD (SD 879) in people without depression.

INSERT TABLE 1

Healthcare Utilization and Expenditures and Depression Severity

The associations between PHQ–9 score and healthcare utilization and expenditure are displayed in Table 2. After adjustment for age, sex and relative wealth, each unit increase in PHQ–9 score was associated with a 14% increase in total healthcare consultations (95% CI 7%–22%, $P < 0.0001$). Outpatient utilization increased by 14% with each unit increase in PHQ–9 score (95% CI 6%–23%, $P < 0.0001$), whereas the association between PHQ–9 scores and inpatient admissions was weaker at 7% (95% CI –6%–22%, $P = 0.314$).

For each increment in PHQ–9 score, mean OOP expenditure on outpatient and total healthcare increased by \$5.36 USD (95% CI 2.36–8.47) and \$9.37 USD (95% CI 2.15–16.58) per year respectively after adjustment for age, sex and relative wealth (Table 2). There was only weak evidence for differences in inpatient expenditures. Although we present estimates from multivariable models for both utilization and OOP expenditure, the estimated associations were similar in univariate models and were not modified by relative wealth index during initial inspection of the data. Exploratory analyses adjusting additionally for

educational attainment and occupation yielded similar associations between depression and total healthcare utilization (17% (95% CI 9%–25%) (including outpatient 17% (95% CI 9%–27%); and inpatient 12% (–3%–29%)) and total OOP healthcare expenditures (\$9.84 (1.39–18.30) (including \$5.35 (95% CI 2.30–8.40) for outpatient and \$4.49 (–2.55–11.54) for inpatient).

INSERT TABLE 2

Healthcare Utilization by Provider Type

Figure 1 displays the proportion of people with and without probable depression who used each type of outpatient healthcare. For each provider, the proportion of people who reported any outpatient consultations in the past 3 months was greater among those with probable depression than those without. Of note, people with depression most commonly sought treatment from pharmacists (30.1%), who they were 3 times more likely to visit than people without probable depression. General doctors were the second most common source of treatment seeking in the depressed group (21.2%). In comparison, people without depression consulted (non-mental health) specialist doctors (17.7%) most frequently, whilst few people overall visited traditional healers (6.7%) or psychiatrists (0.2%).

INSERT FIGURE 1

Provider-Specific Healthcare Expenditure

Participants with probable depression incurred higher annual OOP expenditures for each provider type on average than participants without, as shown in [Figure 2](#). In contrast to utilization patterns, these expenditures were highest for visits to specialist doctors (\$35.99 USD (SD 15.12)), followed by general doctors (\$12.44 USD (SD 6.11)) and pharmacists (\$7.97 USD (SD 3.28)), compared to \$3.48 USD (SD 14.21) for traditional healers and \$3.03 USD (SD 24.26) for psychiatrists.

INSERT FIGURE 2

Discussion

Principal findings

Our results demonstrate that people with probable depression in Central Nepal used more healthcare and incurred greater OOP expenditures than people without depression, and that both utilization and OOP expenditure increased significantly with increasing depression screening scores. People with probable depression incurred total mean expenditures of \$118 (SD 29) USD/year OOP on healthcare, which is likely to represent a substantial proportion of household budgets considering that median annual income for adults with depression in this area has been estimated to be \$501.¹⁷ The association between symptom

severity and total healthcare utilization and OOP costs was attributable to outpatient rather than inpatient service use, and was not modified by relative wealth.

Our findings also show that when seeking healthcare, individuals with probable depression most often sought care from pharmacists, with very few consulting mental health specialists for healthcare. Despite their more frequent use, annual pharmacist costs were much lower than for consultations with specialist doctors (\$7.97 USD/year compared to \$36.99 USD/year) and general doctors (\$12.04 USD/year).

Strengths and limitations

This is the only population-based study that we are aware of from Nepal or any other low-income country to simultaneously evaluate dose-dependent associations between depression and healthcare utilization and OOP expenditure and to examine sector-specific utilization. We used a large, population-based sample, in which secondary level educational attainment was fairly representative of national estimates, suggesting that this was not an obvious source of selection bias.⁴⁶ We also minimised the risk of misclassification bias because we validated the PHQ-9 in this setting and modelled symptom score as a continuous variable for most analyses.

However, our findings should be interpreted in light of some limitations. Firstly, due to the cross-sectional design, (and because the recall periods for the PHQ-9 and healthcare utilization were 2 months and 3 to 12 months respectively), it is difficult to infer causality or determine the true temporal relationship between depression and healthcare utilization. Multiple testing may also have led to spurious results, although the pattern of results was consistent across outcomes and similar to other studies from other resource-poor settings.^{22,37,47} However, depression typically adopts a chronic relapsing and remitting course and it is likely that these symptoms preceded the 2 week recall period." . Secondly, in the absence of reliable health service records, we measured utilization and expenditure by self-report, which may be subject to recall bias, although we restricted the recall period for outpatient utilization to three months to minimise this. Conversely, by extrapolating outpatient healthcare utilization from 3 months to 1 year, we may have introduced infrequency bias due to seasonal fluctuations, which could have led us to over or underestimate these associations. Thirdly, we cannot rule out unmeasured confounding from urban or rural residence or physical comorbidities, which are known to influence healthcare access in Nepal^{48,49}, as neither were recorded in this study. Consequently, we are unable to quantify the relative contributions of other serious health conditions to healthcare utilization and OOP expenditure. However, previous studies that have adjusted for comorbidities have nevertheless found residual associations²², while studies restricted to populations with chronic conditions still find increased healthcare expenditures among people with mental illness⁵⁰. Finally, the OOP expenditures that we report are derived from healthcare utilization by the interviewee alone but may impose a wider burden to the household and to friends, whose associated costs were also included. However, we did not collect income or expenditure data at the household level and were therefore unable to model either expenditures as a proportion of

available household budget, or catastrophic healthcare expenditures, which are more likely to have a direct impact on family members.

Comparison with previous literature

The pattern of increased healthcare consumption by people with probable depression in Nepal reported here is consistent with observations from high-income countries, where the greatest direct healthcare costs are also attributable to general healthcare rather than specialist mental healthcare utilization.¹² Our finding that excess expenditure by people with depression was largely accounted for by outpatient service use is consistent with findings from India^{21,51}, Brazil⁵² and other middle-income countries²², which demonstrate 14 to 36% increases in community healthcare utilization among people with depression, and suggest that depression is a risk factor for catastrophic health expenditure.³⁷ This is the first study that we know of to independently report hospital admission frequency and OOP healthcare expenditures in people with symptoms of depression in LMIC, although a study from Canada also found no significant increases in health service related costs of admissions.¹²

Whether people with depression seek more help from pharmacists in LMIC has not been studied before, but in Brazil it has been reported that depressed individuals use more medications than the general population.⁵² It is important to note that the role of pharmacists in Nepal is generally limited to the provision of over the counter drugs rather than diagnoses or prescriptions. As reported elsewhere in Nepal⁴⁸, we detected minimal utilization of traditional healers for overall healthcare needs. This may be due to increased stigmatization of the use of traditional healers⁵³ and a general preference for 'pill-based' treatments in South Asia⁵⁴. Conversely, only 2% of people with depression who utilized outpatient services were seen by a psychiatrist, which is consistent with previous literature and the shortage of mental health specialists in the area³².

Mechanisms and implications

Given the limitations of the data described above, there are multiple potential explanations of the association between depression symptoms and healthcare utilization and expenditure, which are not necessarily mutually exclusive: (1) The first explanation is that depression leads to increased utilization of services through somatization—that is, the manifestation of psychological distress as physical symptoms—which has been demonstrated previously⁵⁵ (figure 3a). (2) Alternatively, depression may lead to increased healthcare utilization because of its association with physical comorbidities, which are both a risk factor for and a consequence of depression^{56–58} (figure 3b). (3) Finally, OOP healthcare expenditure can result in poverty³⁸, which in turn is a major risk factor for depression¹⁶, raising the possibility that the causal pathway acts in the opposite direction (figure 3c).

There is evidence to support the plausibility of each of these pathways. In support of the first and second hypotheses, which posit a causal role of depression, associations have been shown prospectively between depression and health service utilization in Canada¹² and the United States.⁵⁹ Further, somatization is a common presentation of depression in South Asia^{60–64} and particularly in Nepal⁶⁵, which could explain the increased demand for physical rather than mental healthcare. Previous analyses have also shown that only 8.1% of people with probable depression in this sample reported actively seeking treatment specifically for depression symptoms⁶⁶, which is far lower than the 71% who sought any healthcare at all. This may relate to the lack of effective detection and treatment of depression in primary care^{67,68}, and possibly to low perceived need for intervention for psychological symptoms.⁶⁹

At the same time, a growing evidence base also demonstrates that depression increases the risk of a variety of physical health problems^{70–73}, which could mediate the relationship between depression and healthcare utilization, although we were unable to explore this. Cross-sectional evidence from Nepal supports an association between depression and chronic comorbidities^{74,75}. Nonetheless, studies that have controlled for these comorbidities or restricted to populations with chronic conditions have still found a residual association between depression and increased healthcare utilization, suggesting that depression also exerts an independent effect on treatment-seeking.²²

Recent evidence shows that primary care workers can be trained²⁴ to improve detection of depression and deliver mhGAP-based interventions²⁶ (including psychoeducation and medication) to effectively treat depression and reduce disability in Nepal²⁵. If depression is the primary driver of increased healthcare utilization among this group (either directly, by somatization, or indirectly, by impacting on general health) then we would expect the roll-out of such services to reduce excess healthcare use and associated OOP expenditure. Elsewhere, the integration of mental health services into primary care in Andhra Pradesh in India³⁹ has been shown to substantially reduce OOP healthcare expenditures. In Nepal, the estimated combined costs of scaling-up services for psychosis, depression and epilepsy are less than \$1.30 USD/capita/year.²⁷ In contrast to the additional \$9 USD/year OOP incurred by people in this study for each increment in PHQ–9 score, this is a relatively small investment, which might minimise financial impoverishment, whilst reducing symptoms of depression and hopefully break the link between the two.

However, the notion that OOP healthcare expenditures lead to depression (figure 3c) also finds some empirical support. Healthcare expenditure represents an exceptionally large share of total household expenditure in Nepal¹⁵ and therefore poses a risk of catastrophic expenditure, impoverishment and widened inequalities¹⁵, which are significant determinants of depression.¹⁶ In support of this, we found that the poorest individuals were more likely to suffer from depression but did not appear to forego care, which we would expect to cause further impoverishment. Thus, the lack of financial protection for people with depression is likely to reinforce a vicious cycle of further healthcare utilization and greater impoverishment. According to this model, introducing financial protections would reduce depression prevalence. Supporting the uptake of social health insurance schemes in Nepal could therefore break the cycle between the impoverishing effects of healthcare utilization and the associated risks of depression.

In support of this hypothesis, studies from the United States show that introduction of Medicaid led to 9% reductions in depression prevalence and near elimination of catastrophic expenditure.⁷⁶

Given previous evidence of the inter-relationships between poverty and financial shocks, depression, and physical health, it seems probable that the observed association between depression and healthcare utilization and expenditure results from a combination of these pathways (figure 3d). Indeed, the idea that physical health, mental health and poverty interact synergistically through multiple complex pathways underlies syndemics theory.⁷⁷⁻⁷⁹ As we did not collect data on all of these factors, further research would be necessary to empirically confirm the relative contribution of each of these mechanisms in the Nepali context. However, a combination of increased financial protection for healthcare and the provision of effective depression treatment might reasonably be recommended to break this cycle.

In terms of how to provide depression treatment as part of UHC, the low frequency of hospital admissions support proposals to integrate mental health services into primary rather than secondary care in LMIC. The low cost and frequent consultation of pharmacists, relative to general and specialist doctors, is also relevant to policy as it indicates an additional cadre of healthcare providers who are in frequent contact with the target population. Evidence from PRIME demonstrates that task-sharing by training and supervising non-specialist health workers can improve depression detection rates⁶⁸ and mhGAP-based interventions for depression in Nepal have been found to be most effective when they were supplemented with psychosocial treatments delivered in the community.²⁴ Our findings suggest that enlisting pharmacists to identify potential depression patients and refer them to community counsellors to receive appropriate treatments such as the Healthy Activity Program⁸⁰, might be a fruitful strategy to both reduce the provision of unnecessary medications (and therefore OOP costs) and ultimately provide both mental and physical health benefits.²⁵

Unanswered questions and future research

This study contributes to the evidence base on healthcare utilization and expenditure by demonstrating that, in Nepal, adults with probable depression use a disproportionate amount of healthcare and incur increased healthcare expenditures. However, as this is a secondary analysis, our findings are more exploratory than confirmatory and it is necessary to design adequately powered prospective studies that also record physical health outcomes to confirm these findings, establish the direction of causation and understand the mechanisms that underlie these associations. Future studies should also investigate whether providing mhGAP-based interventions for depression mitigates excess utilization of health services by people with depression, to inform their inclusion in social health insurance packages of care and to prevent impoverishment in this group. Modelling studies have estimated that strengthening primary health care in LMIC to diagnose and treat depression could result in productivity gains as high as \$4 USD for every \$1 USD invested⁸¹ and the real-world effect of such an approach should therefore be evaluated. Equally, the impact of expanding financial protection for depression as part of UHC should

also be evaluated in LMIC. Finally, the feasibility of training pharmacists in Nepal to recognise depression and signpost individuals with depression to appropriate care should be investigated.

Conclusion

Our findings show that people with symptoms of depression in central Nepal utilise more healthcare and incur greater OOP costs compared to people without depression. This economic burden is accounted for by frequent visits to pharmacists and costly visits to general and specialist doctors, with minimal use of specialist mental health care. These findings reinforce the emerging consensus that policy-makers in low-income countries must improve universal access to cost-effective treatments for depression through integration of mental healthcare services into primary healthcare, as well as expanding financial protection as part of universal health care initiatives, to reduce the financial burden of depression on patients. Engaging pharmacists and primary care doctors as key stakeholders may help to improve case detection and access to appropriate evidence-based services, which can be delivered in the community. Future research should investigate the impact of financial protection on depression, and of depression treatment on healthcare utilization and expenditure.

Declarations

Ethics Approval and Consent to Participate

Ethical approval was obtained from the Nepal Health Research Council (Kathmandu, Nepal) and the World Health Organization (Geneva, Switzerland) and informed consent was sought from each participant.

Consent for Publication

Not applicable

Availability of Data and Material

The datasets analysed during the current study are not publicly available immediately due to organisational policy and ethical considerations. In the consent forms, we did not ask for consent from participants for their data to be made publicly available. However, we have set up a system outlined in our publication policy where interested researchers can apply for access to the data via the PRIME consortium Expression of Interest form which is available here: <http://www.prime.uct.ac.za/contact-us>. The data access committee is made of the members of the PRIME management group, led by the PRIME CEO, Prof Crick Lund at the University of Cape Town. All requests for data will go through the PRIME Expression of Interest form which is available in PRIME website <http://www.prime.uct.ac.za/contact-us>. Anyone who is interested in collaboration or using PRIME data can fill in the form and submit it. As

described in our publication policy, the request is vetted by the PRIME Management Team and the PRIME consortium. Additional details can be obtained from Erica Breuer, PRIME project manager (Erica.breuer@uct.ac.za).

Competing Interests

The authors declare that they have no competing interests.

Funding

This document is an output from the Programme for Improving Mental Health Care Research

Programme Consortium, funded by the UK Department of International Development for the benefit of LMICs. No funding bodies had any role in study design, data collection and analysis, decision to publish or preparation of the manuscript. The views expressed in this publication are not necessarily those of the funders. AM is funded by a Wellcome Trust Research Fellowship (number 104349/Z/14/Z).

Author's Contributions

SR was the lead author, and was ultimately responsible for the conceptualisation, study design, statistical analysis, interpretation of results and writing and re-drafting of the manuscript;

SDR contributed to funding acquisition, data curation, study design, statistical analysis and writing and gave detailed comments and feedback on earlier drafts.

NPL was involved in funding acquisition, project administration, data curation and the study design and approved the final drafts.

AM contributed significantly to drafting the manuscript, interpretation of results and writing and re-drafting of the manuscript.

TR contributed significantly to drafting the manuscript, interpretation of results and writing and re-drafting of the manuscript.

MJDJ was the lead supervisor and contributed to the conceptualisation, funding acquisition, study design and reviewed and provided detailed feedback and comments on earlier drafts.

All authors read and approved the final manuscript.

Acknowledgements

We thank the Transcultural Psychosocial Organization Nepal research teams in Chitwan for their work on this study. We would also like to thank Mary de Silva for her supervision early on in the conceptualisation of this study and to James Shearer at King's College London for his advice on elements of the economic analysis.

Author Correspondence

Mark J. D. Jordans, King's College London, Institute of Psychiatry, Psychology and Neuroscience, 16 De Crespigny Park, Camberwell, London SE5 8AF, UK. Email: mark.jordans@kcl.ac.uk

Author's Information

Selina Rajan, MSc, MFPH, Specialist Public Health Registrar, Department of Health Services, Policy and Planning, London School of Hygiene and Tropical Medicine; Sujit D. Rathod, PhD, Assistant Professor, Department of Population Health, London School of Hygiene and Tropical Medicine, UK; Nagendra P. Luitel, MA, Researcher, Transcultural Psychosocial Organization; Adrianna Murphy, SM, PhD., Centre for Global Chronic Conditions; Department of Health Services Research and Policy; Faculty of Public Health and Policy; London School of Hygiene and Tropical Medicine; Tessa Roberts MSc, PhD, Health Service & Population Research Department, Institute of Psychiatry, Psychology & Neuroscience, King's College London, UK; Mark J. D. Jordans, PhD, Reader, Centre for Global Mental Health, Institute of Psychiatry, Psychology and Neuroscience, King's College London, UK; and Transcultural Psychosocial Organization, Nepal.

References

1. World Health Organization. Tracking Universal Health Coverage: 2017 Global Monitoring Report.; 2017. doi:Licence: CC BY-NC-SA 3.0 IGO
2. Chisholm D, Sanderson K, Ayuso-Mateos JL, Saxena S. Reducing the global burden of depression: Population-level analysis of intervention cost-effectiveness in 14 world regions. *Br J Psychiatry*. 2004;184(MAY):393–403. doi:10.1192/bjp.184.5.393
3. Patel V, Chisholm D, Rabe-Hesketh S, Dias-Saxena F, Andrew G, Mann A. Efficacy and cost-effectiveness of drug and psychological treatments for common mental disorders in general health care in Goa, India: a randomised, controlled trial. *Lancet*. 2003;361(9351):33–39. doi:10.1016/S0140–6736(03)12119–8
4. Kleinman A, Estrin GL, Usmani S, et al. Time for mental health to come out of the shadows. *Lancet*. 2016;387(10035):2274–2275. doi:10.1016/S0140–6736(16)30655–9

5. Patel V, Saxena S, Lund C, et al. The Lancet Commission on global mental health and sustainable development. *Lancet*. October 2018. doi:10.1016/S0140-6736(18)31612-X
6. Chisholm D, Johansson KA, Raykar N, et al. Universal Health Coverage for Mental, Neurological, and Substance Use Disorders: An Extended Cost-Effectiveness Analysis. The International Bank for Reconstruction and Development / The World Bank; 2016. doi:10.1596/978-1-4648-0426-7_CH13
7. Thornicroft G, Patel V. Including mental health among the new sustainable development goals. *BMJ*. 2014;349:g5189. doi:10.1136/bmj.g5189
8. Global Burden of Disease study. GBD Results Tool | GHDx. <http://ghdx.healthdata.org/gbd-results-tool>. Accessed February 3, 2019.
9. Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJL. Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet*. 2006;367(9524):1747-1757. doi:10.1016/S0140-6736(06)68770-9
10. König H, König H-H, Konnopka A. The excess costs of depression: a systematic review and meta-analysis. *Epidemiol Psychiatr Sci*. April 2019;1-16. doi:10.1017/S2045796019000180
11. Simon GE, VonKorff M, Barlow W. Health care costs of primary care patients with recognized depression. *Arch Gen Psychiatry*. 1995;52(10):850-856. doi:10.1001/archpsyc.1995.03950220060012
12. Chiu M, Lebenbaum M, Cheng J, De Oliveira C, Kurdyak P. The direct healthcare costs associated with psychological distress and major depression: A population-based cohort study in Ontario, Canada. *PLoS One*. 2017;12(9). doi:10.1371/journal.pone.0184268
13. Bosmans JE, de Bruijne MC, de Boer MR, van Hout H, van Steenwijk P, van Tulder MW. Health care costs of depression in primary care patients in The Netherlands. *Fam Pract*. 2010;27(5):542-548. doi:10.1093/fampra/cmq033
14. Brilleman SL, Purdy S, Salisbury C, Windmeijer F, Gravelle H, Hollinghurst S. Implications of comorbidity for primary care costs in the UK: a retrospective observational study. *Br J Gen Pract*. 2013;63(609):e274-e282. doi:10.3399/bjgp13X665242
15. Van Doorslaer E, O'Donnell O, Rannan-Eliya RP, et al. Catastrophic payments for health care in Asia. *Health Econ*. 2007;16(11):1159-1184. doi:10.1002/hec.1209
16. Lund C, Brooke-Sumner C, Baingana F, et al. Social determinants of mental disorders and the Sustainable Development Goals: a systematic review of reviews. *The Lancet Psychiatry*. 2018;5(4):357-369. doi:10.1016/S2215-0366(18)30060-9
17. Lund C, Docrat S, Abdulmalik J, et al. Household economic costs associated with mental, neurological and substance use disorders: a cross-sectional survey in six low-and middle-income countries. 2019. doi:10.1192/bjo.2019.20
18. World Health Organization (WHO). GHO | By category | Health expenditure ratios, by country, 1995-2014 - Nepal. WHO. <http://apps.who.int/gho/data/view.main.HEALTHEXPRATIONPL?lang=en>. Published 2014. Accessed November 23, 2017.

19. Government Of Nepal Health Insurance Board. District wise Insurees. <http://dashboard.hib.gov.np:81/>. Accessed July 19, 2019.
20. World Health Organization. Mental Health ATLAS 2017 Member State Profile: Nepal. [https://www.who.int/mental_health/evidence/atlas/profiles-2017/NPL.pdf?ua = 1](https://www.who.int/mental_health/evidence/atlas/profiles-2017/NPL.pdf?ua=1). Published 2017. Accessed July 19, 2019.
21. Brinda EM, Rajkumar AP, Enemark U, Prince M, Jacob KS. Nature and determinants of out-of-pocket health expenditure among older people in a rural Indian community. *Int Psychogeriatrics*. 2012;24(10):1664–1673. doi:10.1017/S104161021200083X
22. Albanese E, Liu Z, Acosta D, et al. Equity in the delivery of community healthcare to older people: findings from 10/66 Dementia Research Group cross-sectional surveys in Latin America, China, India and Nigeria. *BMC Health Serv Res*. 2011. doi:10.1186/1472-6963-11-153
23. Lund C, Tomlinson M, De Silva M, et al. PRIME: A Programme to Reduce the Treatment Gap for Mental Disorders in Five Low- and Middle-Income Countries. *PLoS Med*. 2012;9(12):e1001359. doi:10.1371/journal.pmed.1001359
24. Jordans MJD, Luitel NP, Kohrt BA, et al. Community-, facility-, and individual-level outcomes of a district mental healthcare plan in a low-resource setting in Nepal: A population-based evaluation. Geng EH, ed. *PLOS Med*. 2019;16(2):e1002748. doi:10.1371/journal.pmed.1002748
25. Jordans MJD, Luitel NP, Garman E, et al. Effectiveness of psychological treatments for depression and alcohol use disorder delivered by community-based counsellors: two pragmatic randomised controlled trials within primary healthcare in Nepal. *Br J Psychiatry*. January 2019:1–9. doi:10.1192/bjp.2018.300
26. World Health Organization (WHO). *Scaling up Care for Mental, Neurological, and Substance Use Disorders*.; 2008. doi:ISBN: 9789241596
27. Chisholm D, Heslin M, Docrat S, et al. Scaling-up services for psychosis, depression and epilepsy in sub-Saharan Africa and South Asia: Development and application of a mental health systems planning tool (OneHealth). *Epidemiol Psychiatr Sci*. 2017. doi:10.1017/S2045796016000408
28. Jordans MJDD, Luitel NP, Pokhrel P, Patel V. Development and pilot testing of a mental healthcare plan in Nepal. *Br J Psychiatry*. 2015;208:1–33. doi:10.1192/bjp.bp.114.153718
29. Jordans MJ, Luitel NP, Tomlinson M, Komproe IH. Setting priorities for mental health care in Nepal: a formative study. *BMC Psychiatry*. 2013;13(1):332. doi:10.1186/1471-244X-13-332
30. Kohrt B a, Hruschka DJ, Worthman CM, et al. Political violence and mental health in Nepal: prospective study. *Br J Psychiatry*. 2012;201(4):268–275. doi:10.1192/bjp.bp.111.096222
31. Rathod SD, De Silva MJ, Ssebunnya J, et al. Treatment Contact Coverage for Probable Depressive and Probable Alcohol Use Disorders in Four Low- and Middle-Income Country Districts: The PRIME Cross-Sectional Community Surveys. Fischer G, ed. *PLoS One*. 2016;11(9):1–15. doi:10.1371/journal.pone.0162038
32. Hanlon C, Luitel NP, Kathree T, et al. Challenges and Opportunities for Implementing Integrated Mental Health Care: A District Level Situation Analysis from Five Low- and Middle-Income Countries.

- Montazeri A, ed. PLoS One. 2014;9(2):e88437. doi:10.1371/journal.pone.0088437
33. Kroenke K. Depression screening is not enough. *Ann Intern Med.* 2001;134(5):418–420.
 34. Kohrt BA, Luitel NP, Acharya P, Jordans MJD. Detection of depression in low resource settings: validation of the Patient Health Questionnaire (PHQ–9) and cultural concepts of distress. *BMC Psychiatry.* 2016;16:58. doi:10.1186/s12888–016–0768-y
 35. Vyas S, Kumaranayake L. Constructing socio-economic status indices: How to use principal components analysis. *Health Policy Plan.* 2006;21(6):459–468. doi:10.1093/heapol/czl029
 36. Chisholm D, Sekar K, Kumar KK, et al. Integration of mental health care into primary care. Demonstration cost-outcome study in India and Pakistan. *Br J Psychiatry.* 2000;176(JUN.):581–588. doi:10.1192/bjp.176.6.581
 37. Patel V, Chisholm D, Kirkwood BR, Mabey D. Prioritizing health problems in women in developing countries: comparing the financial burden of reproductive tract infections, anaemia and depressive disorders in a community survey in India. *Trop Med Int Health.* 2007;12(1):130–139. doi:10.1111/j.1365–3156.2006.01756.x
 38. van Doorslaer E, O'Donnell O, Rannan-Eliya RP, et al. Effect of payments for health care on poverty estimates in 11 countries in Asia: an analysis of household survey data. *Lancet.* 2006;368(9544):1357–1364. doi:10.1016/S0140–6736(06)69560–3
 39. Raja S, Kippen S, Janardhana N, et al. Evaluating economic outcomes of the mental health and development model in Andhra Pradesh, India. *Basic Needs Rep.* 2008:1–29.
 40. StataCorp. *Stata Statistical Software: Release 14.* College Station, TX: StataCorp LP. 2015. 2015. doi:10.2307/2234838
 41. Thompson SG, Barber JA. How should cost data in pragmatic randomised trials be analysed? *BMJ.* 2000;320(7243):1197–1200. doi:10.1136/BMJ.320.7243.1197
 42. Efron B, Tibshirani R. *An Introduction to the Bootstrap.* Chapman & Hall; 1994.
 43. Mihaylova B, Briggs A, O'Hagan A, Thompson SG. Review of statistical methods for analysing healthcare resources and costs. *Health Econ.* 2011;20(8):897–916. doi:10.1002/hec.1653
 44. The UK Small Aneurysm Trial Participants. Health service costs and quality of life for early elective surgery or ultrasonographic surveillance for small abdominal aortic aneurysms. UK Small Aneurysm Trial Participants. *Lancet (London, England).* 1998;352(9141):1656–1660. doi:10.1016/S0140–6736(98)10284–2
 45. Groves RM, Fowler FJ, Couper M, Lepkowski JM, Singer E, Tourangeau R. *Survey Methodology.* Wiley; 2009.
 46. World Bank. *Education Statistics - All Indicators | DataBank.* <https://databank.worldbank.org/data/reports.aspx?source = Education Statistics>. Accessed May 11, 2019.
 47. Hailemichael Y, Hanlon C, Tirfessa K, et al. Catastrophic health expenditure and impoverishment in households of persons with depression: A cross-sectional, comparative study in rural Ethiopia. *BMC*

- Public Health. 2019;19(1). doi:10.1186/s12889-019-7239-6
48. Saito E, Gilmour S, Yoneoka D, et al. Inequality and inequity in healthcare utilization in urban Nepal: a cross-sectional observational study. *Health Policy Plan.* 2016;31(7):817–824. doi:10.1093/heapol/czv137
49. Saito E, Gilmour S, Rahman MM, Gautam GS, Shrestha PK, Shibuya K. Catastrophic household expenditure on health in Nepal: a cross-sectional survey. *Bull World Health Organ.* 2014;92(10):760–767. doi:10.2471/BLT.13.126615
50. Sporinova B, Manns B, Tonelli M, et al. Association of Mental Health Disorders With Health Care Utilization and Costs Among Adults With Chronic Disease. *JAMA Netw Open.* 2019;2(8):e199910-e199910. doi:10.1001/JAMANETWORKOPEN.2019.9910
51. Soni A, Fahey N, Byatt N, et al. Association of common mental disorder symptoms with health and healthcare factors among women in rural western India: results of a cross-sectional survey. *BMJ Open.* 2016;6(7):e010834. doi:10.1136/BMJOPEN-2015-010834
52. Huang H, Menezes PR, Da Silva SA, Tabb K, Barkil-Oteo A, Scazufca M. The association between depressive disorders and health care utilization: Results from the São Paulo Ageing and Health Study (SPAH). *Gen Hosp Psychiatry.* 2014. doi:10.1016/j.genhosppsy.2013.11.003
53. Kohrt BA, Harper I. Navigating diagnoses: Understanding mind-body relations, mental health, and stigma in Nepal. *Cult Med Psychiatry.* 2008;32(4):462–491. doi:10.1007/s11013-008-9110-6
54. Jain S, Jadhav S. Pills that Swallow Policy: Clinical Ethnography of a Community Mental Health Program in Northern India. *Transcult Psychiatry.* 2009;46(1):60–85. doi:10.1177/1363461509102287
55. Barsky AJ, Orav EJ, Bates DW. Somatization increases medical utilization and costs independent of psychiatric and medical comorbidity. *Arch Gen Psychiatry.* 2005;62(8):903–910. doi:10.1001/archpsyc.62.8.903
56. Druss BG, Walker ER. Mental disorders and medical comorbidity. *Synth Proj Res Synth Rep.* 2011; (21):1–26.
57. Firth J, Siddiqi N, Koyanagi A, et al. The Lancet Psychiatry Commission: a blueprint for protecting physical health in people with mental illness. *The Lancet Psychiatry.* 2019;6(8):675–712. doi:10.1016/S2215-0366(19)30132-4
58. Prince M, Patel V, Saxena S, et al. No health without mental health. *Lancet.* 2007;370(9590):859–877. doi:10.1016/S0140-6736(07)61238-0
59. Unutzer J, Patrick DL, Simon G, et al. Depressive symptoms and the cost of health services in HMO patients aged 65 years and older. A 4-year prospective study. *JAMA.* 1997;277(20):1618–1623.
60. Simon GE, VonKorff M, Piccinelli M, Fullerton C, Ormel J. An international study of the relation between somatic symptoms and depression. *N Engl J Med.* 1999;341(18):1329–1335. doi:10.1056/NEJM199910283411801
61. Gururaj, Girish I, Gururaj G, Girish N, Isaac M K. NCMH Background Papers·Burden of Disease in India Mental, neurological and substance abuse disorders: Strategies towards a systems approach. Burd

- Dis India. 2005;226.
62. Raguram R, Weiss MG, Keval H, Channabasavanna SM. Cultural dimensions of clinical depression in Bangalore, India. *Anthropol Med*. 2001;8(1):31–46. doi:10.1080/13648470120063889
 63. Shidhaye R, Mendenhall E, Sumathipala K, Sumathipala A, Patel V. Association of somatoform disorders with anxiety and depression in women in low and middle income countries: a systematic review. *Int Rev Psychiatry*. 2013;25(1):65–76. doi:10.3109/09540261.2012.748651
 64. Rao D, Young M, Raguram R. Culture, Somatization, and Psychological Distress: Symptom Presentation in South Indian Patients from a Public Psychiatric Hospital. *Psychopathology*. 2007;40(5):349–355. doi:10.1159/000106312
 65. Kohrt BA, Hruschka DJ. Nepali concepts of psychological trauma: The role of idioms of distress, ethnopsychology and ethnophysiology in alleviating suffering and preventing stigma. *Cult Med Psychiatry*. 2010;34(2):322–352. doi:10.1007/s11013-010-9170-2
 66. Luitel NP, Jordans MJD, Kohrt BA, Rathod SD, Komproe IH. Treatment gap and barriers for mental health care: A cross-sectional community survey in Nepal. *PLoS One*. 2017. doi:10.1371/journal.pone.0183223
 67. Patel V, Ramasundarahettige C, Vijayakumar L, et al. Suicide mortality in India: A nationally representative survey. *Lancet*. 2012;379(9834):2343–2351. doi:10.1016/S0140-6736(12)60606-0
 68. Rathod SD, Roberts T, Medhin G, et al. Detection and treatment initiation for depression and alcohol use disorders: Facility-based cross-sectional studies in five low-income and middle-income country districts. *BMJ Open*. 2018;8:23421. doi:10.1136/bmjopen-2018-023421
 69. Andrade LH, Alonso J, Mneimneh Z, et al. Barriers to mental health treatment: Results from the WHO World Mental Health surveys. *Psychol Med*. 2014. doi:10.1017/S0033291713001943
 70. Gan Y, Gong Y, Tong X, et al. Depression and the risk of coronary heart disease: A meta-analysis of prospective cohort studies. *BMC Psychiatry*. 2014;14(1). doi:10.1186/s12888-014-0371-z
 71. Pan A, Sun Q, Okereke OI, Rexrode KM, Hu FB. Depression and risk of stroke morbidity and mortality: a meta-analysis and systematic review. *JAMA*. 2011;306(11):1241–1249. doi:10.1001/jama.2011.1282
 72. Chida Y, Hamer M, Wardle J, Steptoe A. Do stress-related psychosocial factors contribute to cancer incidence and survival? *Nat Clin Pract Oncol*. 2008;5(8):466–475. doi:10.1038/ncponc1134
 73. Knol MJ, Twisk JWR, Beekman ATF, Heine RJ, Snoek FJ, Pouwer F. Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. *Diabetologia*. 2006;49(5):837–845. doi:10.1007/s00125-006-0159-x
 74. Neupane D, Panthi B, McLachlan CS, Mishra SR, Kohrt BA, Kallestrup P. Prevalence of undiagnosed depression among persons with hypertension and associated risk factors: a cross-sectional study in urban Nepal. *PLoS One*. 2015;10(2):e0117329. doi:10.1371/journal.pone.0117329
 75. Niraula K, Kohrt BA, Flora M, et al. Prevalence of depression and associated risk factors among persons with type-2 diabetes mellitus without a prior psychiatric history: a cross-sectional study in clinical settings in urban Nepal. *BMC Psychiatry*. 2013;13(1):309. doi:10.1186/1471-244X-13-309

76. Bernstein M, Gruber JH, Newhouse JP, Schneider EC, Wright BJ. The Oregon Experiment—Effects of Medicaid on Clinical Outcomes. 2013;1713–1722. doi:10.1056/NEJMsa1212321
77. Mendenhall E, Kohrt BA, Norris SA, Ndeti D, Prabhakaran D. Non-communicable disease syndemics: poverty, depression, and diabetes among low-income populations. *Lancet*. 2017;389(10072):951–963. doi:10.1016/S0140–6736(17)30402–6
78. Mendenhall E, Omondi GB, Bosire E, et al. Stress, diabetes, and infection: Syndemic suffering at an urban Kenyan hospital. *Soc Sci Med*. 2015;146:11–20. doi:10.1016/j.socscimed.2015.10.015
79. Sweetland AC, Kritski A, Oquendo MA, et al. Addressing the tuberculosis–depression syndemic to end the tuberculosis epidemic. *Int J Tuberc Lung Dis*. 2017;21(8):852–861. doi:10.5588/ijtld.16.0584
80. Chowdhary N, Anand A, Dimidjian S, et al. The Healthy Activity Program lay counsellor delivered treatment for severe depression in India: systematic development and randomised evaluation. *Br J Psychiatry*. 2016;208(4):381–388. doi:10.1192/bjp.bp.114.161075
81. Chisholm D, Sweeny K, Sheehan P, et al. Scaling-up treatment of depression and anxiety: A global return on investment analysis. *The Lancet Psychiatry*. 2016;3(5):415–424. doi:10.1016/S2215–0366(16)30024–4

Tables

Table 1: PHQ-9 scores, sociodemographic and healthcare-related characteristics of adults in Chitwan District, Nepal, 2013.

	N	Overall (%)	PHQ-9 Negative N=1960 (96.41%)	PHQ-9 positive N=80 (3.56%)
PHQ-9 Scores	2040			
Range		0-24	0-9	10-24
Median (IQR)		2 (0-4)	2 (0-4)	11 (10-14)
Demographic Characteristics	2040			
Mean age (SD)		39.77 (15.23)	39.54 (15.06)	45.87 (18.49)
Female	1336	59.75	59.96	59.74
Educational Status				
Sub primary / illiterate	275	12.83	12.38	25.00
Primary / literate	686	32.68	32.26	44.08
Secondary or higher	1079	54.49	55.36	31.02
Occupation				
Agriculture, Labour or Service	1409	66.40	65.93	79.11
Business and Service	302	15.46	15.74	8.01
Unemployed	94	4.88	4.92	4.01
Student	161	9.99	10.12	6.33
Other	74	3.26	3.29	2.54
Relative wealth index*				
Low	155	22.27	21.75	35.33
Average	162	32.57	32.75	28.04
High	161	45.16	45.49	36.63
Healthcare Utilization	479			
Inpatient	49			
Percentage with any admissions		8.02	7.82	15.83
Median annual number of visits (IQR)		0 (0-0)	0 (0-0)	0 (0-0)
Outpatient	228			

Percentage with any outpatient visits	39.99	38.83	69.70
Median annual number of visits (IQR)	0 (0-8)	0 (0-4)	4 (0-8)
Total	246		
Percentage with any healthcare utilization	44.93	43.91	70.85
Median annual number of visits (IQR)	1 (0-8)	0 (0-5)	4 (0-8)
Healthcare Expenditure	479		
Inpatient	47		
Percentage with any expenditure	99.06	99.19	97.46
Mean annual expenditure in USD (SD)**	65.94 (766.01)	68.12 (820.59)	53.95 (194.63)
Outpatient	191		
Percentage with any expenditure	85.58	85.60	88.65
Mean annual expenditure in USD (SD)	44.85 (130.81)	41.44 (124.81)	63.71 (161.17)
Total	212		
Percentage with any expenditure	87.86	87.44	97.43
Mean annual expenditure in USD (SD)	110.80 (822.78)	109.55 (878.97)	117.66 (265.46)

Unless otherwise stated, numbers represent percentages.

*Relative wealth index, healthcare utilization and healthcare expenditure were assessed in 479 of the total 2040 participants who answered part 2 of the survey and are presented as tertiles reflecting low, average and high wealth.

The percentage with any expenditure is calculated for those who consumed that type of healthcare only

All healthcare expenditures reflect OOP expenditures in US Dollars.

**SD calculated using bootstrapping techniques with 1000 bootstrap samples.

Table 2. Associations between PHQ-9 scores and annual healthcare utilization and OOP expenditure among adults in Chitwan district,

Nepal, 2013.

	Utilization (Number of visits)		OOP Expenditure (US dollars)	
	IRR (95% CI)	P value	β (95% CI)	P value
Inpatient Admissions	1.07 (0.94, 1.22)	0.314	4.00 (-1.84, 9.84)	0.179
Outpatient Consultations	1.14 (1.06, 1.23)	<0.0001	5.36 (2.26, 8.47)	0.001
Total Healthcare	1.14 (1.07, 1.22)	<0.0001	9.37 (2.15, 16.58)	0.011

Incidence rate ratios (IRR) 95% CI and p-values estimated using survey-adjusted negative binomial regression.

β coefficients, 95% CI and P-values estimated using adjusted bootstrapped linear regression.

Estimates reflect changes associated with a one unit increase in PHQ-9 score after adjustment for age, gender and relative wealth index.

Supplementary Files Legend

Additional File 1.doc

Title: Sampling Procedure

Description: Sampling Procedure for the Community Surveys for the Programme for Improving Mental Health CarE (PRIME) In Nepal


Additional File 2.doc

Title: Relative Wealth Asset Index

Description: Relative Wealth Asset Index to Determine Household Economic Status in the Programme for Improving Mental Health CarE (PRIME) in Nepal

Figures

Figure 1. Outpatient Provider-Specific Healthcare utilization by probable depression status among adults in Chitwan District, Nepal, 2013




Survey-adjusted percentages are reported with respect to all 479 participants who answered questions about healthcare utilization as the denominator. Many participants visited more than one provider. The sum of all provider-specific proportions therefore exceeded the 70% and 39% of participants with and without depression who used any outpatient services.

Figure 1

Outpatient Provider-Specific Healthcare utilization by probable depression status among adults in Chitwan District, Nepal, 2013.

Figure 2. Outpatient sector-specific mean OOP healthcare expenditures (US Dollars/yr) by probable depression status



Mean costs were calculated according to responses from all 479 participants who answered questions about healthcare utilization. Expenditures for community workers, nurses and midwives were negligible

Figure 2

Outpatient sector-specific mean OOP healthcare expenditures (US Dollars/yr) by probable depression status.

Figure 3. Schematic Representation of Potential Causal Pathways Between Depression and Healthcare Utilization and OOP Expenditure




Figure 3

Schematic Representation of Potential Causal Pathways Between Depression and Healthcare Utilization and OOP Expenditure.

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Additionalfile1.docx](#)
- [AdditionalFile2.docx](#)