Efficiency of Non-invasive Prenatal Screening in Pregnant Women at Advanced Maternal Age

Hui Zhu

Jiaxing University Affiliated Women and Children Hospital; Women's Hospital, School of Medicine, Zhejiang University

Xiaoxiao Jin

Women's Hospital, School of Medicine, Zhejiang University

Yuqing Xu

Women's Hospital, School of Medicine, Zhejiang University

Weihua Zhang

Jiaxing University Affiliated Women and Children Hospital

Xiaodan Liu

Jiaxing University Affiliated Women and Children Hospital

Jinglei Jin

Women's Hospital, School of Medicine, Zhejiang University

Yeqing Qian

Women's Hospital, School of Medicine, Zhejiang University

Min-Yue Dong (✉ dongmy@zju.edu.cn)

Women's hospital, School of Medicine, Zhejiang University <https://orcid.org/0000-0002-4344-7924>

Research

Keywords: Advanced maternal age, Trisomy, Noninvasive prenatal screening, Fetal aneuploidies, Prenatal screening method

Posted Date: August 6th, 2020

DOI: <https://doi.org/10.21203/rs.3.rs-52792/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at BMC Pregnancy and Childbirth on January 26th, 2021. See the published version at <https://doi.org/10.1186/s12884-021-03570-6>.

Efficiency of Non-invasive Prenatal Screening in Pregnant Women at Advanced Maternal Age

Hui Zhu^{1,2}, Xiaoxiao Jin², Yuqing Xu², Weihua Zhang¹, Xiaodan Liu¹, Jinglei Jin²,
Yeqing Qian², Minyue Dong^{2*}

¹ Jiaxing University Affiliated Women and Children Hospital, Jiaxing, 314051, Zhejiang, China.

² Women's Hospital, School of Medicine, Zhejiang University, Hangzhou, 310006, Zhejiang, China.

*Correspondence: dongmy@zju.edu.cn

Abstract

Background: To evaluate the efficiency of noninvasive prenatal screening (NIPS) in the prenatal screening for fetal aneuploidies in pregnant women at advanced maternal age (AMA).

Results: From February 1, 2015, to December 31, 2018, 29,443 pregnant women at AMA underwent NIPS and followed-up were recruited. Their detailed clinical data and follow-up outcomes were collected and analyzed. The sensitivity, specificity, positive predictive value (PPV), negative predictive value (NPV) of NIPS for detecting fetal trisomy 21 were 99.11%, 99.96%, 90.98%, and 100%, respectively. These same parameters for detecting fetal trisomy 18 were 100%, 99.94%, 67.92%, and 100%, respectively. Finally, these parameters for detecting trisomy 13 were 100%, 99.96%, 27.78%,

25 and 100%, respectively. The prevalence of fetal trisomy 21 increased
26 with maternal age. There were significant differences in the
27 prevalence of fetal trisomy 21 among the groups I (35-37-year-old),
28 II (38-40-year-old) and III (41-year-old and older) ($P<0.05$).

29 **Conclusion:** It is indicated that NIPS is an effective prenatal
30 screening method in pregnant women at AMA.

31 **Keywords:** Advanced maternal age, Trisomy, Noninvasive prenatal
32 screening, Fetal aneuploidies, Prenatal screening method

1. Background

With the implementation of the two-child policy in China, the number of pregnant women at advanced maternal age (AMA) has increased dramatically^[1]. It has been shown that the pregnant women at AMA account for 33.4%-46% in prenatal diagnosis centers^[2-5].

The risk for fetal aneuploidies increases with maternal age. Invasive prenatal diagnosis should be recommended for women at AMA in some countries including China. However, the invasive prenatal diagnosis is not accepted by some pregnant women. Non-invasive prenatal screening (NIPS) is the alternative choice for these women.

Cell-free fetal DNA-based NIPS has been proven to be of high sensitivity and specificity for detecting common chromosomal aneuploidies (trisomy 21, trisomy 18 and trisomy 13), with low false positive and false negative rates. Moreover, clinical observations have indicated that NIPS has excellent performance in either high-risk or low-risk population of serological screening, and the detection efficiency is much higher than that of serological screening^[6-9]. However, the application of NIPS in the prenatal screening of women at AMA is still rare, especially in Chinese

52 population.

53 In this study, 29,343 AMA pregnant women (35 years of age or
54 over) were recruited. All of them underwent NIPS and were
55 followed-up. We intended to explore the clinical significance of
56 NIPS to detect trisomies 21, 18 and 13 in AMA pregnant women and
57 to provide an appropriate prenatal screening program for AMA
58 pregnant women.

59

60 **2. Results**

61 **2.1 NIPS results**

62 29,443 women at AMA who underwent NIPS and completed
63 their pregnancy outcome follow-up were enrolled (Table 1). A total
64 of 145 pregnant women were identified as being high-risk for fetal
65 trisomy 21. Among them, 111 cases were confirmed to carry fetuses
66 of trisomy 21. Eleven were false positive. Twenty-three cases did not
67 receive diagnosis. Among them, included was one case of
68 spontaneous abortion due to premature rupture of membranes, three
69 of stillbirth, five termination of pregnancies (TOP) due to fetal

70 abnormalities (two cases of cardiac abnormalities, three cases of
71 multiple malformations indicated), 13 of TOP without diagnosis, and
72 one case who refused prenatal diagnosis but delivered a baby of
73 leukemia without karyotyping. Moreover, one case was determined
74 to be a false negative. In this 39-year-old pregnant woman, NIPS
75 was conducted at gestational age of 13 weeks and a negative result
76 was obtained. Ultrasound revealed fetal edema and complete
77 endocardial cushion defect. The pregnancy was terminated after fetal
78 trisomy 21 was confirmed.

79 A total of 70 pregnant women were identified to be at high risk
80 for trisomy 18 (Table 1). Among them, 53 cases received diagnosis
81 and 36 cases were confirmed to carry fetuses of trisomy 18. In
82 addition, 17 cases did not receive the prenatal diagnosis, including
83 three cases of stillbirth, 11 TOP due to fetal malformations (eight of
84 multiple malformations, three of cardiac abnormalities), and three
85 TOP without diagnosis.

86 Moreover, a total of 22 pregnant women were identified to be at
87 high risk for trisomy 13 (Table 1). Among them, 18 cases received
88 diagnosis and five were determined to carry fetuses of trisomy 13.
89 Thirteen were false positive. Four cases did not receive the prenatal

diagnosis and terminated their pregnancies, including three cases of multiple malformations, and one who refused diagnosis.

Table 1 Screening results of common fetal chromosomal aneuploidies

Trisomy	No. of cases with high-risk results	TP	FP	No. of cases without prenatal diagnosis	FN
T21	145	111	11	23	1
T18	70	36	17	17	0
T13	22	5	13	4	0

Data are presented as n (%), unless otherwise indicated. TP (true positive), trisomy is verified; FP (false positive), trisomy is incorrectly classified as trisomy; FN (false negative), trisomy is incorrectly classified as normal.

2.2 Detection efficiency in AMA pregnant women

In order to estimate the detection efficiency of NIPS, we focused on the sensitivity, specificity, PPV, and NPV of trisomies 21, 18 and 13. As shown in Table 2, the sensitivity, specificity and NPV for NIPS for fetal trisomy 21, trisomy 18 and trisomy 13 were all over 99%. And, the PPV for fetal trisomy 21, trisomy 18 and trisomy 13 were 90.98%, 67.92%, and 27.78%, respectively.

Table 2 Detection efficiency of common chromosomal aneuploidies by NIPS

Trisomy	Sensitivity	Specificity	PPV(95% CI)	NPV(95% CI)	Rate of TP
T21	99.11 (94.62-99.99)	99.96 (99.93-99.98)	90.98 (84.08-95.19)	100 (99.98-100)	0.382
T18	100 (88.53-100)	99.94 (99.91-99.97)	67.92 (53.55-79.70)	100 (99.98-100)	0.123
T13	100 (51.09-100)	99.96 (99.92-99.98)	27.78 (12.17-51.20)	100 (99.98-100)	0.017

Data are presented as n (%), unless otherwise indicated. PPV, positive perspective value; NPV, negative perspective value; CI, confidence intervals.

2.3 The correlation between the incidence of fetal trisomy 21 and maternal age

As shown in Table 3, the rates of high-risk and true positive for fetal trisomy 21 were positively correlated with maternal age ($P<0.01$). Then, we classified the subjects into three groups (I: 35-37 years old; II: 38-40 years old; III: 40 years old and over). Both the high-risk rate and true positive rate for trisomy 21 were significantly different ($P<0.05$) between groups I and II. The high-risk rate differed significantly ($P<0.05$) between groups II and III.

Table 3 High-risk rate and true positive percentage of fetal trisomy 21 in pregnant women in different age groups

Age(y)	No. of cases	No. of cases with high-risk	Rate of cases with high-risk	TP	Rate of TP
35	7044	19	0.27	16	0.23
36	6557	26	0.40	16	0.24
37	4921	16	0.33	13	0.26
38	3824	28	0.73	21	0.55
39	2669	8	0.30	6	0.22
40	1863	12	0.64	10	0.54
41	1044	17	1.63	14	1.34
42	672	6	0.89	6	0.89
43	389	6	1.54	4	1.03
≥44	330	7	2.12	5	1.52

Data are presented as n (%), unless otherwise indicated. TP (true positive), trisomy is verified.

3. Discussion

In this study, we confirmed that NIPS had a high sensitivity, specificity and PPV for detecting trisomies 21, 18 and 13 in AMA pregnant women. In addition, the high-risk rates and incidence of trisomy 21 increased with maternal age. These findings point to the clinical significance of NIPS to detect trisomies 21, 18 and 13 in AMA pregnant women and may help doctors and pregnant women to choose a suitable prenatal screening and diagnosis way.

Serological screening is widely used for fetal aneuploidies. At present, as for the high-risk pregnant women, fetal karyotyping with

amniotic fluid cells or cord blood cells is used as the diagnosis for fetal chromosomal abnormalities. In China, all AMA pregnant women are advised to undergo prenatal diagnosis. However, with the implementation of the two-child policy, the number of AMA pregnant women has increased significantly^[1], which has greatly increased the demand for prenatal diagnosis^[10]. However, amniotic fluid sampling or umbilical cord blood collection are invasive procedures, with the risk of miscarriage which was estimated at 0.5 to 1.0%^[11,12]. There is also a risk of infection in such procedures^[13]. As a result, the overall utilization rate of both methods is low. Moreover, some pregnant women may have contraindications for invasive prenatal diagnosis, such as the high risk of inducing abortion, fever, increased tendency for bleeding, and infection^[14]. Therefore, it is needed to find prenatal screening methods that better meet the clinical needs.

NIPS is a noninvasive prenatal screening technique for fetal aneuploidies. NIPS is based on high-throughput sequencing to detect cell-free fetal DNA (cffDNA) in maternal peripheral blood. In 1997,

Lo et al.^[15] found cffDNA in maternal blood and revealed that cffDNA was suitable for prenatal examination. However, it was not widely applied in clinic until the emergence of high-throughput sequencing ^[16]. Bianchi et al ^[17] compared NIPS and serological screening in general population, which recruited 1,914 women with singleton pregnancies from 21 centers in USA. Each sample was tested by both methods. The positive predictive values for NIPS and standard screening were 45.5% and 4.2% for trisomy 21, and 40.0% and 8.3% for trisomy 18, respectively. NIPS showed significantly better performances than serological screening. Meanwhile, Bianchi et al ^[17] also found that the false negative rates were 0.3% and 0.2% for trisomies 21 and 18 as detected by NIPS, respectively, which were much lower than those of serological screening (3.6% and 0.6%, respectively). Similarly, in a study of 146,958 women ^[18], it revealed that the sensitivity was 99.17%, 98.24% and 100%, that the specificity was 99.95% ,99.95% and 99.96%, that the PPV was

92.19%, 76.61% and 32.84%, and that the NPV was 99.99%, 100% and 100%, for trisomies 21,18 and 13, respectively. Using expanded noninvasive prenatal screening (“NIPS-Plus”), which is considered a better screening method than usual NIPS, Liang ^[19] demonstrated that the PPVs were 95%, 82% and 46%, for trisomies 21, 18 and 13, respectively. Those findings obtained from large size of general populations were consistent with the results of ours, indicating NIPS is suitable for pregnant women at AMA.

Lots of investigations demonstrated that NIPS is superior to serological screening and suitable for the detection of trisomy 21, trisomy 18 and trisomy 13 in all high risk or low risk populations, AMA or not^[17-19]. Thus, the International Society for Prenatal Diagnosis (ISPD), the American College of Obstetricians and Gynecologists (ACOG), the Royal College of Obstetricians and Gynecologists (RCOG), and the American College of Medical Genetics and Genomics (ACMG), have recommended NIPS as the preferred screening method for all pregnant women. Additionally, NIPS has been included in a national policy or national program in 14 European countries ^[20]. Considering the excellent efficiency of

189 NIPS, NIPS could be promoted as the preferred screening method
190 for AMA pregnant women. However, invasive screening methods
191 such as amniotic fluid analysis and cord blood collection are still
192 needed to carry out karyotype analysis for high-risk women
193 identified by NIPS.

194 Since maternal age is closely associated with the incidence of
195 fetal chromosomal abnormalities ^[21], we also studied the correlation
196 between maternal age and the incidence of trisomy 21. Generally,
197 the incidence increased with maternal age. This is consistent with
198 previous report^[22].

199 We noted some shortcomings of this study. On the one hand,
200 there were some cases without diagnosis in the high-risk population
201 detected by NIPS. Many of cases might have had fetal aneuploidy,
202 especially those with imaging abnormalities or fetal death. Therefore,
203 the positive predictive values of fetal trisomy 21, trisomy 18, and
204 trisomy 13 detected by NIPS were likely to be higher than what were
205 described here. The low incidence of trisomy 18 and trisomy 13

made it impossible to carry out an age stratification study as was done for the trisomy 21. Multicenter studies with larger sample sizes are expected in the future and that should provide additional data in support of optimizing prenatal screening and diagnosis strategies for AMA pregnant women.

4. Conclusions

In summary, by analyzing the data from 29,343 AMA pregnant women, we have found that NIPS has good detection efficiency and it is suitable for screening for fetal aneuploidy in AMA pregnant women.

5. Subjects and methods

5.1 Subjects

From February 1, 2015, to December 31, 2018, 29,443 AMA pregnant women (35 years old or older) who underwent NIPS and completed their pregnancy outcome follow-up at Jiaying Maternal and Child Health Hospital and Women's Hospital, Zhejiang University School of Medicine were recruited. The maternal age ranged from 35 to 55 years with a median of 37, and 151 cases were

45 years of age and over. The gestational age ranged from 12 to 30 weeks, and the median was 16 weeks. The gestational age was calculated according to their last menstruation period and verified by ultrasound in early pregnancy. The pregnancy outcomes were followed-up by checking the prenatal diagnosis database, delivery and infant record of hospital or registration system, or followed-up approximately three months after deliveries. The use of the data was approved with the Institutional Review Board and all participants gave their informed consent.

5.2 NIPS

The venous blood of pregnant women was collected and stored in blood collection tubes supplemented with EDTA at 4°C. The plasma DNA was extracted and used for library construction. The fetal cell-free DNA fragments in the plasma were analyzed using the BGISEQ-100 sequencing platform. The rates of risk for fetal trisomy 21, trisomy18, and trisomy13 were obtained through bioinformatics analysis^[23,24]. The normal range of the Z score was between -3.0 and

3.0.

5.3 Fetal karyotyping

Fetal karyotyping was recommended for women who were identified to be at high risk using NIPS. The exact process of fetal karyotyping was described previously^[25].

5.4 Statistics

SPSS (SPSS Company, Chicago, IL, USA) version 19.0 software was used. Data were compared using the chi-square test, and $P < 0.05$ was used to indicate significant differences. The sensitivity, specificity, PPV, and NPV of NIPS were also calculated.

Abbreviations

NIPS: Noninvasive prenatal screening; AMA: Advanced maternal age; PPV: Positive predictive value; NPV: Negative predictive value; TOP: Termination of pregnancies; TP: True positive; FP: False positive; FN: False negative; cffDNA: Cell-free fetal DNA; CI: Confidence intervals; ISPD: International Society for Prenatal Diagnosis; ACOG: American College of Obstetricians and Gynecologists; RCOG: Royal College of Obstetricians and Gynecologists, ACMG: American College of Medical

Genetics and Genomics

Acknowledgements

We honestly appreciate our pregnant women at AMA for their participation in this study and all the colleagues in two hospitals (Jiaxing Maternal and Child Health Hospital and Women's Hospital, Zhejiang University School of Medicine). We would like to thank Dr. Jiong Gao (BGI Genomics, BGI-shenzhen 518083, China) for modifying the article.

Authors' contributions

ZH and JXX collected clinical data and analyzed the data, ZWH and LXD performed the NIPS and karyotype analysis, XYQ was a major contributor in drafting and revising the manuscript, JJL followed up the patients, QYQ took a plan for the research, DMY designed the work and revised the manuscript. All authors read and approved the final manuscript.

Funding

This work is supported by the Medical Scientific Research Foundation of Zhejiang Province (2018KY809) and the Technology Bureau Program of

Jiaxing, Zhejiang Province (2020AD300035).

Availability of data and materials

The data used or analyzed during the current study are included within the article.

Ethics approval and consent to participate

The study was approved by the Ethics Committee of Jiaxing Maternal and Child Health Hospital. All participants gave their informed consent in this study.

Consent for publication

Not applicable.

Competing interests

The authors have no conflicts of interest to declare.

Author details

¹Jiaxing University Affiliated Women and Children Hospital, Jiaxing, 314051, Zhejiang, China.

²Women's Hospital, School of Medicine, Zhejiang University, Hangzhou,

310006, Zhejiang, China.

References

1. Zhang H X, Zhao Y Y, Wang Y Q. Analysis of the Characteristics of Pregnancy and Delivery before and after Implementation of the Two-child Policy. *Chin Med J (Engl)*. 2018; 131(1): 37-42.<https://doi.org/10.4103/0366-6999.221268>
2. Chen TF, Mao QQ, Lu LP, et al. Relationship between fetal karyotype and age of pregnancy. *Chinese Journal of Laboratory Medicine*. 2016; 39(6): 423-426
3. Lin XJ, Sun QM, He XC, et al. Clinical analysis of fetal chromosomes karyotype abnormalities. *Chinese Journal of Obstetrics & Gynecology and Pediatrics*. 2016; 12(2): 173-178
4. Chen YP, Zheng FX, Zhou Q, et al. Application of high-throughput sequencing in the diagnosis of fetal aneuploidies in women with advanced maternal age. *Reproduction & Contraception*. 2016; 36(9): 709
5. Zhu YN. Study on fetal chromosome abnormality and appropriate prenatal diagnostic technique. Zhengjiang: Zhejiang University. 2015
6. Iwarsson E, Jacobsson B, Dagerhamn J, et al. Analysis of cell-free fetal DNA in maternal blood for detection of trisomy 21, 18 and 13 in a general pregnant population and in a high risk population - a systematic review and meta-analysis. *Acta Obstet Gynecol Scand*. 2017; 96(1): 7-18.<https://doi.org/10.1111/aogs.13047>
7. Gil M M, Quezada M S, Bregant B, et al. Implementation of maternal blood cell-free DNA testing in early screening for aneuploidies. *Ultrasound Obstet Gynecol*.

2013; 42(1): 34-40.<https://doi.org/10.1002/uog.12504>

8. Mackie F L, Hemming K, Allen S, et al. The accuracy of cell-free fetal DNA-based non-invasive prenatal testing in singleton pregnancies: a systematic review and bivariate meta-analysis. *Bjog*. 2017; 124(1): 32-46.<https://doi.org/10.1111/1471-0528.14050>

9. Drury S, Hill M, Chitty L S. Cell-Free Fetal DNA Testing for Prenatal Diagnosis. *Adv Clin Chem*. 2016; 76: 1-35.<https://doi.org/10.1016/bs.acc.2016.05.004>

10. Gong W, Xu D R, Caine E D. Challenges arising from China's two-child policy. *Lancet*. 2016; 387(10025): 1274.[https://doi.org/10.1016/s0140-6736\(16\)30020-4](https://doi.org/10.1016/s0140-6736(16)30020-4)

11. Tabor A, Alfirevic Z. Update on procedure-related risks for prenatal diagnosis techniques. *Fetal Diagn Ther*. 2010; 27(1): 1-7.<https://doi.org/10.1159/000271995>

12. Beta J, Lesmes-Heredia C, Bedetti C, et al. Risk of miscarriage following amniocentesis and chorionic villus sampling: a systematic review of the literature. *Minerva Ginecol*. 2018; 70(2): 215-219.<https://doi.org/10.23736/s0026-4784.17.04178-8>

13. Odibo A O, Gray D L, Dicke J M, et al. Revisiting the fetal loss rate after second-trimester genetic amniocentesis: a single center's 16-year experience. *Obstet Gynecol*. 2008; 111(3): 589-95.<https://doi.org/10.1097/AOG.0b013e318162eb53>

14. Stranc L C, Evans J A, Hamerton J L. Chorionic villus sampling and amniocentesis for prenatal diagnosis. *Lancet*. 1997; 349(9053): 711-4.[https://doi.org/10.1016/s0140-6736\(96\)08169-x](https://doi.org/10.1016/s0140-6736(96)08169-x)

15. Lo Y M, Corbetta N, Chamberlain P F, et al. Presence of fetal DNA in maternal plasma and serum. *Lancet*. 1997; 350(9076): 485-7.[https://doi.org/10.1016/s0140-6736\(97\)02174-0](https://doi.org/10.1016/s0140-6736(97)02174-0)

16. Chiu R W, Akolekar R, Zheng Y W, et al. Non-invasive prenatal assessment of

352 trisomy 21 by multiplexed maternal plasma DNA sequencing: large scale validity
 353 study. *Bmj*. 2011; 342: c7401.<https://doi.org/10.1136/bmj.c7401>

354 17. Bianchi D W, Rava R P, Sehnert A J. DNA sequencing versus standard prenatal
 355 aneuploidy screening. *N Engl J Med*. 2014; 371(6):
 356 578.<https://doi.org/10.1056/NEJMc1405486>

357 18. Zhang H, Gao Y, Jiang F, et al. Non-invasive prenatal testing for trisomies 21, 18
 358 and 13: clinical experience from 146,958 pregnancies. *Ultrasound Obstet Gynecol*.
 359 2015; 45(5): 530-8.<https://doi.org/10.1002/uog.14792>

360 19. Liang D, Cram D S. Clinical utility of noninvasive prenatal screening for
 361 expanded chromosome disease syndromes. 2019; 21(9): 1998-
 362 2006.<https://doi.org/10.1038/s41436-019-0467-4>

363 20. Gadsboll K, Petersen O B, Gatinois V. Current use of noninvasive prenatal testing
 364 in Europe, Australia and the USA: A graphical presentation.
 365 2020.<https://doi.org/10.1111/aogs.13841>

366 21. Heffner L J. Advanced maternal age--how old is too old? *N Engl J Med*. 2004;
 367 351(19): 1927-9.<https://doi.org/10.1056/NEJMp048087>

368 22. Yamada T, Sekizawa A, Fujii Y, et al. Maternal age-specific risk for trisomy 21
 369 based on the clinical performance of NIPT and empirically derived NIPT age-specific
 370 positive and negative predictive values in Japan. 2018; 63(10): 1035-
 371 1040.<https://doi.org/10.1038/s10038-018-0453-8>

372 23. Jiang F, Ren J, Chen F, et al. Noninvasive Fetal Trisomy (NIFTY) test: an
 373 advanced noninvasive prenatal diagnosis methodology for fetal autosomal and sex
 374 chromosomal aneuploidies. *BMC Med Genomics*. 2012; 5:

375 57.<https://doi.org/10.1186/1755-8794-5-57>

376 24. Chen S, Lau T K, Zhang C, et al. A method for noninvasive detection of fetal
377 large deletions/duplications by low coverage massively parallel sequencing. *Prenat*
378 *Diagn.* 2013; 33(6): 584-90.<https://doi.org/10.1002/pd.4110>

379 25. Chen M, Fu X Y, Luo Y Q, et al. Detection of fetal duplication 16p11.2q12.1 by
380 next-generation sequencing of maternal plasma and invasive diagnosis. *J Matern Fetal*
381 *Neonatal Med.* 2019; 32(1): 38-45.<https://doi.org/10.1080/14767058.2017.1369947>

382