Comparison of the effects of empagliflozin and glimepiride on endothelial function in patients with type 2 diabetes: A randomized controlled study

Haruka Tamura (✉ faraway02february@gmail.com)

Chigasaki Shiritsu Byoin <https://orcid.org/0000-0001-5997-8431>

Yoshinobu Kondo

Chigasaki Shiritsu Byoin

Kohei Ito

Chigasaki Shiritsu Byoin

Masanori Hasebe

Yokohama Shiritsu Daigaku Igakubu Daigakuin Igaku Kenkyuka

Shinobu Satoh

Chigasaki Shiritsu Byoin

Yasuo Terauchi

Yokohama Shiritsu Daigaku Igakubu Daigakuin Igaku Kenkyuka

Research article

Keywords: Empagliflozin, Endothelial function, Glimepiride, Diabetes, Flow-mediated dilation

Posted Date: February 26th, 2020

DOI: <https://doi.org/10.21203/rs.2.14604/v4>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Version of Record: A version of this preprint was published at Diabetes on June 22nd, 2018. See the published version at <https://doi.org/10.2337/db18-1138-P>.

Abstract

Abstract Background : Among patients with type 2 diabetes and established cardiovascular disease, those receiving empagliflozin have a lower rate of primary composite cardiovascular outcome and death from any cause. While treatment with sulfonylurea reduces microvascular complications in diabetes, it increases cardiovascular hospitalization or mortality in combination with metformin. In the present study, we assessed the effects of empagliflozin and glimepiride, a commonly prescribed sulfonylurea, on endothelial function using flow-mediated dilation (FMD) to estimate arteriosclerosis and cardiovascular events in patients with type 2 diabetes. **Methods :** In this prospective, open-label, randomized, parallel-group study, 58 patients with type 2 diabetes received metformin and glargine before bedtime for 12 weeks. This was followed by additional treatment with either 10 mg empagliflozin or 0.5 mg glimepiride for 12 weeks. The primary outcome was the change in the FMD measurement (DFMDs), measured prior to and after 12 weeks of additional treatment. Secondary outcomes comprised changes in metabolic markers and body composition. **Results :** Analysis of the empagliflozin group (n = 30) and glimepiride group (n = 28) showed no significant differences in DFMDs (empagliflozin, $-0.19 \pm 2.34\%$; glimepiride, $-0.37 \pm 2.77\%$; $P = 0.79$). Similarly, changes in glycated hemoglobin (HbA1c) were similar between the two groups. However, significant difference in body weight changes were observed between the two groups (empagliflozin, -0.59 ± 2.5 kg; glimepiride, 1.2 ± 3.0 kg; $P = 0.02$). In addition, an analysis of the body composition revealed that body fluid volume significantly decreased after empagliflozin treatment (baseline, 35.8 ± 6.8 L; after 12 weeks, -0.33 ± 0.72 L; $P = 0.03$). **Conclusions :** Empagliflozin did not improve endothelial function when compared to glimepiride in patients with type 2 diabetes but decreased their body fluid volume. This suggested that the coronary protective effect of empagliflozin is not derived by protection of the endothelial function but rather from reduction in the risk of heart failure. **Trial Registration :** This trial was registered on September 13, 2016; UMIN000024001 . **Keywords:** Empagliflozin, Endothelial function, Glimepiride, Diabetes, Flow-mediated dilation

Background

In the treatment of coronary artery disease, the assessment of coronary endothelial vasoreactivity has important diagnostic and prognostic implications. Specifically, an estimate of the arteriosclerosis and cardiovascular events can be made by measuring endothelial dysfunction [1]. Thus, this parameter may be useful for assessing cardiovascular risk in diabetes as well. The flow-mediated dilation (FMD) method evaluates endothelial function in a noninvasive manner, based on the intrinsic ability of blood vessels to respond to blood flow [2]. FMD involves high frequency ultrasonographic imaging of the brachial artery by reflecting nitric oxide production within the artery. Since FMD can predict cardiovascular events, it has been utilized in numerous investigations of arteriosclerosis [3-5]. However, variables such as age, systolic blood pressure, body mass index (BMI), sex, presence of diabetes mellitus, lipid-lowering medication, smoking, and a decrease in visceral adipose tissue mass can impact FMD measurements [6-8].

Blood glucose levels in diabetes can be decreased by preventing proximal tubular glucose reabsorption and increasing urinary glucose excretion, using an inhibitor of sodium glucose cotransporter 2 (SGLT2). Notably, such inhibitors can also reduce blood pressure [9] and body weight [9, 10]. The EMPA-REG OUTCOME trial revealed that treatment with empagliflozin, a SGLT2 inhibitor, reduced the risk of cardiovascular outcomes and death from all causes in patients with pre-existing atherosclerotic cardiovascular disease. However, the incidence of myocardial infarction or stroke did not differ significantly between the empagliflozin and placebo groups [11].

Furthermore, a meta-analysis showed that SGLT2 inhibitors improved the composite outcome of myocardial infarction, stroke, or cardiovascular death [12]. However, the effects varied according to the presence or absence of atherosclerotic cardiovascular disease. These inhibitors could reduce the composite of major adverse cardiovascular events, albeit only in patients with existing atherosclerotic cardiovascular disease. In addition, hospitalization following heart failure was reduced with SGLT2 inhibitor treatment, regardless of the status of atherosclerotic cardiovascular disease or heart failure at baseline. However, the anti-atherosclerotic mechanism of SGLT2 inhibitors remains unclear [12].

In the UK Prospective Diabetes Study (UKPDS) 33 Group, microvascular complications were reduced with sulfonylurea therapy, but no effect on macrovascular disease was seen. The control of blood glucose by sulfonylureas decreased the risk of microvascular complications. [13]. However, the observation that a combination therapy of sulfonylurea and metformin increased cardiovascular hospitalization or mortality was concerning [14]. The lack of effect of sulfonylurea on macrovascular disease could be a result of non-action on endothelial function. The effect of reduced microvascular complication was derived from the strict control of blood glucose and decreased glycated hemoglobin (HbA1c). We thus hypothesized that sulfonylureas may not improve endothelial function. Consequently, in this study we used FMD to compare endothelial function in patients with type 2 diabetes treated with either an SGLT2 inhibitor (empagliflozin) or a sulfonylurea (glimepiride).

Methods

Study design

This study was registered with the University Hospital Medical Information Network Clinical Trial Registry (UMIN000024001) as a prospective, open-label, randomized, parallel-group comparison study. Approval for this study was obtained from the ethics committee of Chigasaki Municipal Hospital (approval No. 2016-04). The study protocol conforms to the provisions of the revised Declaration of Helsinki guidelines. Written, informed consent was obtained from all the participating patients.

Inclusion and exclusion criteria

Patients with type 2 diabetes (aged: 20–80 years old) who were hospitalized at the Chigasaki Municipal Hospital for diabetes education and blood glucose control were. These patients received metformin and basal insulin therapy prior to discharge and had a BMI ≤ 45 kg/m².

Patients were excluded if they showed severe renal (estimated glomerular filtration rate [eGFR] < 45 ml min⁻¹ 1.73 m⁻²) or liver dysfunction; were on steroid therapy; experienced cardiovascular disease and a cerebral infarction within 24 weeks of the study; had cancer, severe infection, or were traumatized; were or could become pregnant; were allergic to empagliflozin, glargine, glimepiride, or metformin; or if the supervising doctor decided that the patient did not qualify for this study.

Treatment and interventions

Patients who were treated with metformin and glargine for 12 weeks from discharge were randomized to receive either 10 mg empagliflozin or 0.5 mg glimepiride daily. According to the Japanese guidelines, a 10 mg dose of empagliflozin is recommended as the initial starting dose. In case of glimepiride, we selected a dose of 0.5 mg since this dose had similar efficacy in lowering blood glucose as 10 mg empagliflozin. Randomization was stratified by age, glycated hemoglobin (HbA1c), and FMD. Blood samples were obtained in a fasting state. Postprandial plasma glucose (PPG) was measured by self-monitoring blood glucose system (ONE TOUCH Verio IQ; Johnson and Johnson Co., New Brunswick, NJ, USA). Initial measurements of FMD were made prior to additional treatment and randomization. Subsequent measurements were made following additional treatment for 12 weeks. In general, treatment was not changed after randomization. However, if the fasting plasma glucose was maintained under 90 mg/dL, glargine dose was decreased by one unit, once weekly. Hypoglycemia (i.e., blood glucose < 70 mg/dL) was ascertained using the values recorded by the patients. We instructed the patients to self-monitor blood glucose and hypoglycemia symptoms twice a day.

Endpoints and assessments

The primary outcome was the change in FMD and was measured prior to and following 12 weeks of additional treatment (Fig. 1). Secondary outcomes included changes in the levels of metabolic markers in plasma during fasting state and were measured prior to and following 12 weeks of treatment. Body composition components, such as skeletal muscle and total fat mass, and body fluid volume were assessed using a multifrequency bioelectrical impedance analyzer (BIA) (InBody720; InBody Co., Ltd., Seoul, South Korea).

Sample size and statistical analysis

The effect of SGLT2 inhibitors on endothelial function, as measured by FMD, was unknown at the time this protocol was developed. We estimated that the Δ FMD with empagliflozin treatment for 12 weeks would be $1.5 \pm 2.0\%$ based on the results from a previous study [15]. Based on a two-sided P -value of 5% and a power of 80%, a sample size of 58 patients was required to detect a significant difference amongst the two treatment groups. We used Students' t -test to compare the averages of Δ FMD between the two groups, before and after additional treatment.

Results

Patients

Between June 2016 and 2018, 69 patients with type 2 diabetes agreed to participate in this study. A total of 63 patients were randomized at a one-to-one ratio to either empagliflozin or glimepiride group. Ultimately, 30 patients in the empagliflozin group and 28 patients in the glimepiride group were analyzed (Fig. 2). The baseline clinical and biochemical characteristics in the two treatment groups were statistically similar (Table 1). Patients received the following drugs in the empagliflozin vs. glimepiride groups prior to enrollment in the study: metformin (13.3% vs. 0%), DPP-4 inhibitor (16.7% vs. 17.9%), SGLT2-inhibitor (10.0% vs. 0%), sulfonylurea (3.3% vs. 3.6%), α -glucosidase inhibitor (0% vs. 3.6%), and insulin (40.0% vs. 46.4%).

Endothelial function

The average baseline FMD values before and after 12-week treatment with empagliflozin and glimepiride are presented in Table 2. No significant difference was observed in the changes in FMD in empagliflozin vs. glimepiride groups.

Metabolic markers

The fasting plasma glucose remained unchanged in both the groups, with no significant difference evident between the two groups following treatment ($P = 0.69$) (Table 3). However, HbA1c and glycated albumin (GA) levels significantly decreased in both the groups ($P \leq 0.05$). However, the changes in these metabolic markers were not significantly different among the two groups (Δ HbA1c, $P = 0.75$ and Δ GA, $P = 0.42$). Uric acid (UA) significantly decreased in the empagliflozin group ($P < 0.001$), and significantly increased in the glimepiride group ($P = 0.01$). A significant difference between the two groups was also noted in the Δ UA ($P < 0.001$). Systolic ($P = 0.59$) and diastolic ($P = 0.74$) blood pressures along with the heart rate ($P = 0.81$) did not change significantly between the two groups. Empagliflozin treatment significantly increased low-density lipoprotein- C (LDL-C) ($P < 0.001$), whereas triglycerides (TG) ($P =$

0.31) and high-density lipoprotein- C (HDL-C) ($P = 0.37$) did not change significantly between the two groups. Glimepiride treatment led to a significant increase in body weight ($P < 0.05$) after treatment. In addition, a significant difference was observed in the Δ body weight between the two groups ($P = 0.02$). However, in the subgroup of patients in whom decreased body weight was observed, Δ FMD was not significantly different between the two groups ($P = 0.62$) (Table 4). Glimepiride also led to a significant increase of waist circumference after treatment ($P = 0.004$). Moreover, a significant difference was observed in the Δ waist circumference between the two groups ($P = 0.008$). With regard to body composition, glimepiride led to a significant increase in total fat mass ($P = 0.02$). In comparison, empagliflozin significantly decreased body fluid volume ($P = 0.03$).

The fasting C-peptide immunoreactivity (CPR) ($P = 0.55$), homeostasis model assessment 2 steady state beta cell (%B) function ($P = 0.2$), homeostasis model assessment 2 insulin sensitivity (%S) ($P = 0.09$), and homeostasis model assessment 2 insulin resistance ($P = 0.22$) were not significantly different between the two groups (Table S1). In this study, empagliflozin and glimepiride did not affect insulin secretion, insulin sensitivity, pancreatic β -cell function, or insulin resistance. The associations between arterial sclerosis markers and Δ FMD, Δ HbA1c, Δ body weight, Δ HDL-C, and Δ TG were related to Δ FMD. Additionally, the Δ FMD may have been greater if the baseline FMD was lower (Table S2).

Discussion

Based on the results from this study, we found that although fasting plasma glucose (FPG), HbA1c, and GA were equally improved in both empagliflozin and glimepiride groups, Δ FMD was not significant. A previous study has reported that glimepiride does not demonstrate an improvement in the endothelial function [15]. In this study, empagliflozin had no effect towards the improvement of endothelial function, irrespective of the improvement in glucose levels. To observe the changes in endothelial function induced by drugs that lack the ability to improve the glucose levels, this study was conducted in patients with a steady glucose-controlled state. Although a baseline HbA1c was targeted in this study, severe hypoglycemia did not occur after reducing glargine dose, as the fasting plasma glucose was maintained under 90 mg/dL. Additionally, GA improved significantly in both the groups. It is possible that glucose variation was improved by both the additional treatment. The insulin dose was also decreased significantly in both the groups, and this may have resulted in the increase in body weight. The improvements in Δ HbA1c, body weight, HDL-C, and TG and the reduction in HbA1c in the empagliflozin group did not affect the FMD, although the correlation between the change in the levels of a decrease in visceral adipose tissue mass, waist circumference, and Δ FMD has been shown in previous studies [8, 16]. These findings suggest that empagliflozin had no effect in improving FMD although empagliflozin is effective in improving HbA1c.

Although long-term administration of empagliflozin has been shown to maintain a stable eGFR [17], we observed that the renal function worsened in the empagliflozin group during the relatively short observation period. It was reported that empagliflozin decreased hyperfiltration by diuresis in the initial term [18]. This may be attributed to the changes in renal dynamics induced by empagliflozin. In a recent

study, canagliflozin reduced the risk of kidney failure at a median follow up time of 2.62 years [19]. Losartan showed a rapid initial decline in the renal function which led to a slower decrease in long-term renal function [20]. However, it has been reported that an initial rapid decline of renal function leads to kidney failure [21]. Therefore, empagliflozin would be expected to prevent the progression of kidney disease over an extended period of time.

In Body720, an eight-polar BIA was shown to accurately estimate total and appendicular body composition, independent of age and sex [20]. The body fluid volume was significantly decreased following 12 weeks of empagliflozin treatment. However, it was unclear whether the effect of empagliflozin on the body fluid volume was greater than that of glimepiride. Empagliflozin has been shown to improve hospitalization rates after heart failure [21] and is thought to ameliorate the effects of this disease by decreasing body fluid volume. During the observational period of our study, heart failure was not observed in either treatment group. This suggests that empagliflozin might have a coronary protective effect that is not derived from the impact on the endothelial function. This diuretic medication is known to decrease congested heart failure, although its effect in reduction of the cardiovascular event risk is unclear. It was reported that SGLT2 inhibitors reduce more interstitial fluid volume than blood volume compared to the loop diuretics [22]. It was hypothesized that improvement of systemic congestion and renal function, derived from decreased hyperfiltration, resulted in prevention of cardiovascular event without reducing arterial filling and perfusion.

The FMD may be improved when the baseline FMD is low [23]. However, in both the subgroups that showed a lower than median baseline FMD, the Δ FMD did not significantly differ between the two groups ($P = 0.83$) (Table 6). Notably, although SGLT2 inhibitors have a secondary preventive role in adverse cardiovascular events, they lack a primary preventive role [12]. However, results from the DECLARE-TIMI 58 study demonstrated that dapagliflozin had both primary as well as secondary roles in preventing adverse cardiovascular events. Dapagliflozin decreased cardiovascular death and hospitalization due to heart failure in patients without previous cardiovascular disease [24]. However, in this study, no patient had a similar background, and hence it was necessary to undertake a secondary intervention.

The effects of other oral hypoglycemic agents, like pioglitazone and glucagon-like peptide 1 (GLP-1) analogs, in improving endothelial function have been reported previously [15, 25]. A dipeptidyl peptidase 4 (DPP-4) inhibitor either improved [23], had no effect [26], or worsened [27] endothelial function but did not affect cardiovascular events [28, 29]. In another study, GLP-1 analog treatment enhanced [30, 31] or had no effect on [32] endothelial function. In patients with type 2 diabetes, liraglutide, a GLP-1 analog, was successful in preventing nonfatal myocardial infarction or stroke, along with death from cardiovascular causes [33]. Similar to these findings, results from a meta-analysis showed that significant heterogeneity existed between DPP-4 inhibitor and GLP-1 [34]. Factors such as the size of the study, duration of intervention, and age or sex of the participants did not affect the mean difference in FMD [34]. Rather, a change in FMD was found to be dependent on the baseline FMD. The low baseline FMD in our study compared to other reported studies [23, 25, 35] might have an effect on the results.

Several limitations were evident in this study. First, as the study participants were outpatients, it was not possible to completely exclude patients who smoked or had a meal before the FMD examination. Moreover, non-compliance of dietary requirements in some patients might have affected the observed change in body weight. Additionally, even though we had included wash out periods of anti-diabetic therapy for 12 weeks before the study, the long-term effects of anti-diabetic therapy before admission cannot be completely ruled out. Finally, although the number of patients to detect a significant difference in comparisons of the two groups was adequate, the observation period was relatively short. Although, an evidence of observation period assessed by FMD has not been established yet, a longer study period to monitor any adverse events is required for future studies.

Conclusions

Overall, we found that compared to glimepiride, empagliflozin did not improve endothelial function, at least in a 12-week treatment period, in patients with type 2 diabetes without previous cardiovascular disease. However, empagliflozin significantly reduced body fluid volume. Thus, the coronary protective effect of empagliflozin may be derived not from the prevention of endothelial dysfunction, but rather from reducing the chances of heart failure.

Abbreviations

ARB/ACE-I, angiotensin type II receptor blocker/angiotensin converting enzyme I; BMI, body mass index; Cr, serum creatinine; CRP, C-reactive protein; CysC, cystatin C; dBp, diastolic blood pressure; DPP, dipeptidyl peptidase; eGFR, estimated glomerular filtration rate; eGFR_{cys}, estimated glomerular filtration rate by cystatin C; F-CPR, fasting C-peptide immunoreactivity; FMD, flow-mediated dilation; FPG, fasting plasma glucose; GA, glycated albumin; GLP, glucagon-like peptide; HbA1c, glycated hemoglobin; HDL, high-density lipoprotein; HOMA2%B, Homeostasis Model Assessment 2 steady state beta cell (%B) function; HOMA2%S, Homeostasis Model Assessment 2 insulin sensitivity (%S); HOMA2IR, Homeostasis Model Assessment 2 insulin resistance (IR); HR, heart rate; LDL, low-density lipoprotein; L-FABP, liver-type fatty acid binding protein; sBP, systolic blood pressure; TG, triglycerides; UA, uric acid; U-Alb, urine albumin

Declarations

Ethics approval and consent to participate

Approval for this study was obtained from the ethics committee of Chigasaki Municipal Hospital, approval No. 2016-04. Written, informed consent was obtained from all patients.

Consent for publication

Not applicable

Availability of data and materials

The datasets used and analyzed during the current study are available from the corresponding author on reasonable request.

Competing interests

The authors declare that they have no competing interests.

Funding

We have not accepted any financial support for this study.

Authors' contributions

YK provided advice on study design; HT and YK designed the study. HT collected and analyzed the data. HT performed statistical analysis. HT, KI, and MH interpreted the data. HT wrote the manuscript. YK, SS, and YTDrafted the work or substantively revised it. All authors read and approved the final manuscript.

Acknowledgements

Not applicable.

References

1. Schachinger V, Britten MB, Zeiher AM. Prognostic impact of coronary vasodilator dysfunction on adverse long-term outcome of coronary heart disease. *Circulation* 2000, 101(16):1899-1906.
2. Corretti MC, Anderson TJ, Benjamin EJ, Celermajer D, Charbonneau F, Creager MA, et al. Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J Am Coll Cardiol* 2002, 39(2):257-265.
3. Yeboah J, Folsom AR, Burke GL, Johnson C, Polak JF, Post W, et al. Predictive value of brachial flow-mediated dilation for incident cardiovascular events in a population-based study: the multi-ethnic study of atherosclerosis. *Circulation* 2009, 120(6):502-509.
4. Ter Avest E, Stalenhoef AF, de Graaf J. What is the role of non-invasive measurements of atherosclerosis in individual cardiovascular risk prediction? *Clin Sci (Lond)* 2007, 112(10):507-516.
5. Deanfield JE, Halcox JP, Rabelink TJ. Endothelial function and dysfunction: testing and clinical relevance. *Circulation* 2007, 115(10):1285-1295.
6. Maruhashi T, Soga J, Fujimura N, Idei N, Mikami S, Iwamoto Y, et al. Relationship between flow-mediated vasodilation and cardiovascular risk factors in a large community-based study. *Heart* 2013, 99(24):1837-1842.

7. Benjamin EJ, Larson MG, Keyes MJ, Mitchell GF, Vasan RS, Keaney JF, Jr., et al. Clinical correlates and heritability of flow-mediated dilation in the community: the Framingham Heart Study. *Circulation* 2004, 109(5):613-619.
8. Rittig K, Hieronimus A, Thamer C, Machann J, Peter A, Stock J, et al. Reducing visceral adipose tissue mass is essential for improving endothelial function in type 2 diabetes prone individuals. *Atherosclerosis* 2010, 212(2):575-579.
9. Kadowaki T, Haneda M, Inagaki N, Terauchi Y, Taniguchi A, Koiwai K, et al. Efficacy and safety of empagliflozin monotherapy for 52 weeks in Japanese patients with type 2 diabetes: a randomized, double-blind, parallel-group study. *Advances in therapy* 2015, 32(4):306-318.
10. Roden M, Merker L, Christiansen AV, Roux F, Salsali A, Kim G, et al. Safety, tolerability and effects on cardiometabolic risk factors of empagliflozin monotherapy in drug-naïve patients with type 2 diabetes: a double-blind extension of a Phase III randomized controlled trial. *Cardiovasc Diabetol* 2015, 14:154.
11. Zinman B, Wanner C, Lachin JM, Fitchett D, Bluhmki E, Hantel S, et al. Empagliflozin, Cardiovascular Outcomes, and Mortality in Type 2 Diabetes. *N Engl J Med* 2015, 373(22):2117-2128.
12. Zelniker TA, Wiviott SD, Raz I, Im K, Goodrich EL, Bonaca MP, et al. SGLT2 inhibitors for primary and secondary prevention of cardiovascular and renal outcomes in type 2 diabetes: a systematic review and meta-analysis of cardiovascular outcome trials. *Lancet* 2019, 393(10166):31-39.
13. Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 1998, 352(9131):837-853.
14. Rao AD, Kuhadiya N, Reynolds K, Fonseca VA. Is the combination of sulfonylureas and metformin associated with an increased risk of cardiovascular disease or all-cause mortality?: a meta-analysis of observational studies. *Diabetes Care* 2008, 31(8):1672-1678.
15. Papathanassiou K, Naka KK, Kazakos N, Kanioglou C, Makriyiannis D, Pappas K, et al. Pioglitazone vs glimepiride: Differential effects on vascular endothelial function in patients with type 2 diabetes. *Atherosclerosis* 2009, 205(1):221-226.
16. Miyazaki S, Hiasa Y, Takahashi T, Tobetto Y, Chen H, Mahara K, et al. Waist circumference reduction is more strongly correlated with the improvement in endothelial function after acute coronary syndrome than body mass index reduction. *J Cardiol* 2010, 55(2):266-273.
17. Wanner C, Inzucchi SE, Lachin JM, Fitchett D, von Eynatten M, Mattheus M, et al. Empagliflozin and Progression of Kidney Disease in Type 2 Diabetes. *N Engl J Med* 2016, 375(4):323-334.
18. Vallon V, Gerasimova M, Rose MA, Masuda T, Satriano J, Mayoux E, et al. SGLT2 inhibitor empagliflozin reduces renal growth and albuminuria in proportion to hyperglycemia and prevents glomerular hyperfiltration in diabetic Akita mice. *Am J Physiol Renal Physiol* 2014, 306(2):F194-204.
19. Perkovic V, Jardine MJ, Neal B, Bompoint S, Heerspink HJL, Charytan DM, et al. Canagliflozin and Renal Outcomes in Type 2 Diabetes and Nephropathy. *N Engl J Med* 2019, 380(24):2295-2306.

20. Malavolti M, Mussi C, Poli M, Fantuzzi AL, Salvioli G, Battistini N, et al. Cross-calibration of eight-polar bioelectrical impedance analysis versus dual-energy X-ray absorptiometry for the assessment of total and appendicular body composition in healthy subjects aged 21-82 years. *Annals of human biology* 2003, 30(4):380-391.
21. Fitchett D, Zinman B, Wanner C, Lachin JM, Hantel S, Salsali A, et al. Heart failure outcomes with empagliflozin in patients with type 2 diabetes at high cardiovascular risk: results of the EMPA-REG OUTCOME(R) trial. *Eur Heart J* 2016, 37(19):1526-1534.
22. Hallow KM, Helmlinger G, Greasley PJ, McMurray JJV, Boulton DW. Why do SGLT2 inhibitors reduce heart failure hospitalization? A differential volume regulation hypothesis. *Diabetes Obes Metab* 2018, 20(3):479-487.
23. Shigiyama F, Kumashiro N, Miyagi M, Iga R, Kobayashi Y, Kanda E, et al. Linagliptin improves endothelial function in patients with type 2 diabetes: A randomized study of linagliptin effectiveness on endothelial function. *J Diabetes Investig* 2017, 8(3):330-340.
24. Wiviott SD, Raz I, Bonaca MP, Mosenzon O, Kato ET, Cahn A, et al. Dapagliflozin and Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2019, 380(4):347-357.
25. Tsuchiya K, Akaza I, Yoshimoto T, Hirata Y. Pioglitazone improves endothelial function with increased adiponectin and high-density lipoprotein cholesterol levels in type 2 diabetes. *Endocr J* 2009, 56(5):691-698.
26. Nomoto H, Miyoshi H, Furumoto T, Oba K, Tsutsui H, Inoue A, et al. A Randomized Controlled Trial Comparing the Effects of Sitagliptin and Glimepiride on Endothelial Function and Metabolic Parameters: Sapporo Athero-Incretin Study 1 (SAIS1). *PLoS One* 2016, 11(10):e0164255.
27. Ayaori M, Iwakami N, Uto-Kondo H, Sato H, Sasaki M, Komatsu T, et al. Dipeptidyl peptidase-4 inhibitors attenuate endothelial function as evaluated by flow-mediated vasodilatation in type 2 diabetic patients. *J Am Heart Assoc* 2013, 2(1):e003277.
28. Green JB, Bethel MA, Armstrong PW, Buse JB, Engel SS, Garg J, et al. Effect of Sitagliptin on Cardiovascular Outcomes in Type 2 Diabetes. *N Engl J Med* 2015, 373(3):232-242.
29. Scirica BM, Bhatt DL, Braunwald E, Steg PG, Davidson J, Hirshberg B, et al. Saxagliptin and cardiovascular outcomes in patients with type 2 diabetes mellitus. *N Engl J Med* 2013, 369(14):1317-1326.
30. Lambadiari V, Pavlidis G, Kousathana F, Varoudi M, Vlastos D, Maratou E, et al. Effects of 6-month treatment with the glucagon like peptide-1 analogue liraglutide on arterial stiffness, left ventricular myocardial deformation and oxidative stress in subjects with newly diagnosed type 2 diabetes. *Cardiovasc Diabetol* 2018, 17(1):8.
31. Ceriello A, Esposito K, Testa R, Bonfigli AR, Marra M, Giugliano D. The possible protective role of glucagon-like peptide 1 on endothelium during the meal and evidence for an "endothelial resistance" to glucagon-like peptide 1 in diabetes. *Diabetes Care* 2011, 34(3):697-702.
32. Nomoto H, Miyoshi H, Furumoto T, Oba K, Tsutsui H, Miyoshi A, et al. A Comparison of the Effects of the GLP-1 Analogue Liraglutide and Insulin Glargine on Endothelial Function and Metabolic

Parameters: A Randomized, Controlled Trial Sapporo Athero-Incretin Study 2 (SAIS2). PLoS One 2015, 10(8):e0135854.

33. Marso SP, Daniels GH, Brown-Frandsen K, Kristensen P, Mann JF, Nauck MA, et al. Liraglutide and Cardiovascular Outcomes in Type 2 Diabetes. N Engl J Med 2016, 375(4):311-322.

34. Batzias K, Antonopoulos AS, Oikonomou E, Siasos G, Bletsas E, Stampouloglou PK, et al. Effects of Newer Antidiabetic Drugs on Endothelial Function and Arterial Stiffness: A Systematic Review and Meta-Analysis. J Diabetes Res 2018, 2018:1232583.

35. Nakamura K, Oe H, Kihara H, Shimada K, Fukuda S, Watanabe K, et al. DPP-4 inhibitor and alpha-glucosidase inhibitor equally improve endothelial function in patients with type 2 diabetes: EDGE study. Cardiovasc Diabetol 2014, 13:110.

Tables

Table 1 Baseline characteristics of patients

	Empagliflozin group ($\pm$ SD)	Glimepiride group ($\pm$ SD)	<i>P</i> value
Age [years]	58.6 $\pm$ 8.5	54.3 $\pm$ 12.2	0.13
Gender (male/female)	18 / 12	18 / 10	0.74
Estimated duration of diabetes [years]	6.1 $\pm$ 7.2	4.9 $\pm$ 4.9	0.44
Current smoker [%]	20.0	39.3	0.11
FMD [%]	5.49 $\pm$ 2.05	5.46 $\pm$ 2.2	0.96
HbA1c [%]	6.9 $\pm$ 1.1	6.6 $\pm$ 0.7	1.0
FPG [mg/dL]	136.9 $\pm$ 65.2	127.1 $\pm$ 52.6	0.37
GA [%]	17.0 $\pm$ 3.7	16.3 $\pm$ 3.4	0.48
Body weight [kg]	70.0 $\pm$ 11.3	69.6 $\pm$ 17.1	0.92
BMI [kg/m ²]	26.1 $\pm$ 3.7	25.9 $\pm$ 5.4	0.9
Cr [mg/dL]	0.76 $\pm$ 0.16	0.73 $\pm$ 0.16	0.54
eGFR [mL min ⁻¹ 1.73 m ⁻²]	74.9 $\pm$ 12.9	81.9 $\pm$ 19.7	0.11
UA [mg/dL]	5.5 $\pm$ 1.2	5.4 $\pm$ 1.5	0.96
LDL-C [mg/dL]	94.0 $\pm$ 26.7	89.2 $\pm$ 28.1	0.51
HDL-C [mg/dL]	54.7 $\pm$ 16.0	57.4 $\pm$ 16.4	0.52
TG [mg/dL]	197.7 $\pm$ 101.1	176.0 $\pm$ 131.5	0.2
sBP [mmHg]	129.6 $\pm$ 14.9	130.1 $\pm$ 20.5	0.9
dBp [mmHg]	80.8 $\pm$ 10.0	78.0 $\pm$ 9.3	0.28
HR [bpm]	79.4 $\pm$ 14.8	77.5 $\pm$ 14.3	0.64

Values are shown as the means $\pm$ standard deviation (SD). Paired Student's *t*-tests were used to compare values between different groups.

FMD, flow-mediated dilation; HbA1c, glycated hemoglobin; FPG, fasting plasma glucose; GA, glycated albumin; BMI, body mass index; Cr, serum creatinine; eGFR, estimated glomerular filtration rate

Table 2 FMD (% $\pm$ SD) with treatment

	Empagliflozin	Glimepiride	
FMD (0)	5.49 ± 2.05	5.46 ± 2.2	<i>P</i> = 0.96
FMD (12)	5.3 ± 2.28	5.09 ± 1.86	<i>P</i> = 0.71
	<i>P</i> = 0.66	<i>P</i> = 0.49	
ΔFMD (12) - (0)	-0.19 ± 2.34	-0.37 ± 2.77	<i>P</i> = 0.79

Values are shown as the means ± standard deviation (SD).

FMD, flow-mediated dilation

FMD (0) means the baseline FMD value at 0 week. FMD (12) means the FMD value at 12 weeks after additional treatment.

Δ indicates the change in the FMD value between 0 and 12 week.

P values in each row show the comparison of FMD values at the baseline and at week 12. P value in each line shows the comparison of changes in FMD values for both the groups.

Table 3 Changes in metabolic markers

	Empagliflozin	Glimepiride	
Fasting plasma glucose (mg/dL) [mean $\pm$ SD]			
Week 12	124.5 $\pm$ 52.1	119.9 $\pm$ 49.1	
	$P = 0.11$	$P = 0.47$	
Δ FPG	-12.4 $\pm$ 44.4	-7.3 $\pm$ 52.0	$P = 0.69$
Postprandial plasma glucose (mg/dL) [mean $\pm$ SD]			
Week 12	169.6 $\pm$ 87.0	165.1 $\pm$ 61.1	
	$P = 0.07$	$P = 0.26$	
Δ PPG	-16.5 $\pm$ 43.9	-9.5 $\pm$ 27.6	$P = 0.63$
HbA1c (%) [mean $\pm$ SD]			
Week 12	6.7 $\pm$ 1.1	6.4 $\pm$ 0.78	
	$P = 0.001$	$P = 0.01$	
Δ HbA1c	-0.22 $\pm$ 0.36	-0.26 $\pm$ 0.5	$P = 0.75$
GA (%) [mean $\pm$ SD]			
Week 12	16.0 $\pm$ 3.2	15.7 $\pm$ 3.1	
	$P < 0.001$	$P = 0.05$	
Δ GA	-0.97 $\pm$ 1.3	-0.65 $\pm$ 1.6	$P = 0.42$
Renal function			
Cr (mg/dL) [mean $\pm$ SD]			
Week 12	0.79 $\pm$ 0.15	0.74 $\pm$ 0.16	
	$P = 0.03$	$P = 0.8$	
eGFR (mL min ⁻¹ 1.73 m ⁻²) [mean $\pm$ SD]			
Week 12	72.1 $\pm$ 11.7	81.5 $\pm$ 20.2	
	$P = 0.02$	$P = 0.8$	
UA (mg/dL) [mean $\pm$ SD]			
Week 12	4.8 $\pm$ 1.2	5.7 $\pm$ 1.4	
	$P < 0.001$	$P = 0.01$	
Δ UA	-0.64 $\pm$ 0.9	0.26 $\pm$ 0.5	$P < 0.001$

Body weight (kg) [mean $\pm$ SD]			
Week 12	69.4 $\pm$ 12.0	70.8 $\pm$ 18.2	
	$P = 0.22$	$P < 0.05$	
Δ Body weight	-0.59 $\pm$ 2.5	1.2 $\pm$ 3.0	$P = 0.02$
Waist circumference (cm) [mean $\pm$ SD]			
baseline	91.6 $\pm$ 9.2	91.2 $\pm$ 15.7	0.27
Week 12	90.9 $\pm$ 8.9	92.3 $\pm$ 15.3	
	$P = 0.07$	$P = 0.004$	
Δ Waist circumference	-0.64 $\pm$ 1.8	1.1 $\pm$ 2.8	$P = 0.008$
Blood pressure (mmHg) [mean $\pm$ SD]			
sBP			
Week 12	130.2 $\pm$ 14.2	128.9 $\pm$ 19.3	
	$P = 0.79$	$P = 0.62$	
Δ sBP	0.69 $\pm$ 13.7	-1.3 $\pm$ 13.5	$P = 0.59$
dBP			
Week 12	80.6 $\pm$ 7.6	78.6 $\pm$ 12.0	
	$P = 0.91$	$P = 0.72$	
Δ dBP	-0.24 $\pm$ 10.8	0.64 $\pm$ 9.3	$P = 0.74$
Heart rate			
Week 12	80.1 $\pm$ 15.4	79.0 $\pm$ 12.8	
	$P = 0.76$	$P = 0.55$	
Δ Heart rate	0.68 $\pm$ 11.2	1.5 $\pm$ 11.6	$P = 0.81$
Lipids (TG, HDL-C, LDL-C) (mg/dL) [mean $\pm$ SD]			
LDL-C			
Week 12	107.5 $\pm$ 33.0	93.9 $\pm$ 29.6	
	$P < 0.001$	$P = 0.33$	
Δ LDL-C	13.5 $\pm$ 19.3	4.7 $\pm$ 25.2	$P = 0.14$
HDL-C			
Week 12	56.0 $\pm$ 12.5	56.4 $\pm$ 13.1	

	$P = 0.43$	$P = 0.61$	
Δ HDL-C	1.4 ± 9.4	-1.1 ± 10.9	$P = 0.37$
TG			
Week 12	177.1 ± 86.9	185.7 ± 123.7	
	$P = 0.29$	$P = 0.8$	
Δ TG	-20.6 ± 103.8	9.6 ± 121.5	$P = 0.31$
Body fluid volume (L) [mean $\pm$ SD]			
Baseline	35.8 ± 6.8	36.6 ± 8.6	0.69
Week 12	35.4 ± 6.9	36.3 ± 8.7	
	$P = 0.03$	$P = 0.32$	
Δ Body fluid volume	-0.33 ± 0.72	-0.35 ± 1.8	$P = 0.94$
Total fat mass (kg) [mean $\pm$ SD]			
Baseline	21.1 ± 6.8	20.5 ± 11.0	0.8
Week 12	20.6 ± 6.8	21.7 ± 11.6	
	$P = 0.16$	$P = 0.02$	
Δ Total fat mass	-0.58 ± 2.1	1.2 ± 2.3	$P = 0.006$
Adverse events (%)			
	rash on both arms (3.3%)	hypoglycemia (3.6%)	
Glargine (U) [mean $\pm$ SD]			
Baseline	9.4 ± 4.6	11.7 ± 9.0	0.6
Week 12	8.4 ± 5.1	9.6 ± 8.9	
	$P = 0.02$	$P < 0.001$	
Δ Glargine	-1.0 ± 2.4	-2.1 ± 3.5	$P = 0.16$

Values are shown as the means $\pm$ standard deviation (SD).

Δ indicates the changes in the FMD values between 0 and 12 week.

P values in each row show the comparison of FMD values at the baseline and at week 12. P value in each line shows the comparison of changes in FMD values for both the groups.

FMD, flow-mediated dilation; HbA1c, glycated hemoglobin; GA, glycated albumin; Cr, serum creatinine; eGFR, estimated glomerular filtration rate; UA, uric acid; sBP, systolic blood pressure; dBP, diastolic blood pressure; LDL-C, low-density lipoprotein cholesterol; HDL-C, high-density lipoprotein cholesterol; TG, triglycerides

Table 4 Change of FMD (% ± SD) with treatment in the subgroup of patients with decreased body weight during the observation period

	Empagliflozin (<i>n</i> = 17)	Glimepiride (<i>n</i> = 8)	
ΔFMD (12) - (0)	0.41 ± 2.55	0.94 ± 2.16	<i>P</i> = 0.62

FMD, flow-mediated dilation
Δ indicates the change in FMD value between 0 and 12 week.
P value in each line shows the comparison of changes in FMD values of both the groups.

Table 6 Change of FMD (% ± SD) after additional treatment in subgroups with lower than median baseline FMD

	Empagliflozin (<i>n</i> = 14)	Glimepiride (<i>n</i> = 15)	
ΔFMD(12)-(0)	0.66 ± 1.96	0.82 ± 1.92	<i>P</i> = 0.83

FMD, flow-mediated dilation
Δ indicates the change in the FMD value between 0 and 12 week.
P value in each line shows the comparison of changes in FMD values for both the groups.

Figures

Figure 1

Study flow chart. Patients who took metformin and glargine were randomized to empagliflozin or glimepiride groups. Flow-mediated dilation (FMD), blood examination, body weight, and blood pressure were checked at the study baseline and endpoints.

Figure 2

Consort 2010 flow diagram of patient selection. Ultimately, 30 patients in the empagliflozin group and 28 patients in the glimepiride group were analyzed

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [AdditionalFile1.pdf](#)