

# Does Cystic Fibrosis Constitute an Advantage in COVID-19 Infection?

**Valentino Bezzeri**

Azienda Ospedaliera Universitaria Integrata Verona <https://orcid.org/0000-0002-6849-4487>

**Francesca Lucca**

Azienda Ospedaliera Universitaria Integrata Verona <https://orcid.org/0000-0002-5294-9301>

**Sonia Volpi**

Azienda Ospedaliera Universitaria Integrata Verona

**Marco Cipolli** (✉ [marco.cipolli@aovr.veneto.it](mailto:marco.cipolli@aovr.veneto.it))

Azienda Ospedaliera Universitaria Integrata Verona

---

## Short Report

**Keywords:** COVID-19, SARS-CoV-2, Cystic Fibrosis

**Posted Date:** August 6th, 2020

**DOI:** <https://doi.org/10.21203/rs.3.rs-53181/v1>

**License:** © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

---

**Version of Record:** A version of this preprint was published at Italian Journal of Pediatrics on October 6th, 2020. See the published version at <https://doi.org/10.1186/s13052-020-00909-1>.

# Abstract

The Veneto region is one of the most affected Italian regions by COVID-19. Chronic lung diseases, such as chronic obstructive pulmonary disease (COPD), may constitute a risk factor in COVID-19. Moreover, respiratory viruses were generally associated with severe pulmonary impairment in cystic fibrosis (CF). We would have therefore expected numerous cases of severe COVID-19 among the CF population. Surprisingly, we found that CF patients were significantly protected against infection by SARS-CoV-2. We discussed this aspect formulating some reasonable theories.

## Introduction

The comorbidities of obesity, hypertension, diabetes, heart failure, and chronic lung disease have been associated with poor outcome in coronavirus disease 2019 (COVID-19).<sup>1</sup> Once Severe Acute Respiratory Syndrome (SARS)– Coronavirus (CoV)-2 has infected host cells, excessive inflammatory and thrombotic processes take place. A cytokine storm release with markedly elevated IL-6 levels are associated with increased lethality.<sup>2</sup> Increased neutrophilia is part of that inflammatory response and has been associated with a poorer outcome. Increased neutrophil extracellular traps (NET) have been hypothesized to be predictive of poor outcome in COVID-19 patients.<sup>3</sup>

Cystic fibrosis (CF) lung disease, characterized by the release of a many proinflammatory cytokines and chemokines, is the major cause of morbidity and mortality in CF. Excessive NET accumulation has been observed in the CF airway.<sup>4</sup> Moreover, influenza and other respiratory viral infections are known to cause severe pulmonary impairment in CF.<sup>5</sup> Because of the risk factors of chronic lung disease and host hyper-immune status, one would surmise that CF patients would be at an increased risk of developing severe COVID-19 illness.

## Methods

We conducted a retrospective study of 532 CF patients followed at the Cystic Fibrosis Center of Verona, Italy. SARS-CoV-2 positivity was tested by collecting combined nose-throat swabs and subsequent Real-Time PCR using the Nimbus MuDT tm (Seegene, Seoul, South Korea), and Simplexa COVID-19 Direct kits (DiaSorin, Stillwater, MN, USA), following the manufacturer guidelines.

## Results

The Veneto region is one of the Italian regions most strongly affected by COVID-19, with 19,729 cases and 2,062 related fatalities (as of 23 July 2020). The Veneto region health system has carried out 1,149,084 multiple COVID-19 tests on 465,433 subjects so far (data from the Italian Ministry of Health).

We contacted all the 532 patients from Veneto, of whom 118 subjects (22.2%) undergone further combined nose and throat swab test for SARS-CoV-2. Surprisingly, we found that CF patients were

significantly protected against infection by SARS-CoV-2. We identified only one patient, a 35 year-old man, with COVID-19 infection. Furthermore, this individual reported only mild symptoms including dry cough and body temperature, and he did not require intensive care unit (ICU) admission. These data indicate that the COVID-19 infection rate in the Veneto regional CF population is 0.19%, in contrast to 0.40% in the general population (Table 1).

**Table 1. Demographic and Epidemiologic Data from the Veneto Region**

Condition	Population	Tested subjects (% of regional population)	COVID-19 cases	Rate of infection (% of regional population)	COVID-19 fatalities	CFR
General population	4,905,854*	465,433 (9.5%)	19,729	0.40%	2,062	10.4%
CF population	532	118 (22.2%)	1	0.19%	0	-

CFR, case fatality rate.

\* Data from the Istituto Nazionale di Statistica (ISTAT) of Italy, updated to 1 January 2019

## Discussion

CF condition is supposed to represent a comorbidity for COVID-19. Our data show instead that the prevalence of COVID-19 in CF population of one of the Italian region most affected by the SARS-CoV-2 outbreak, namely Veneto region, is reduced compared with general population. Interestingly, our data are in line with emerging evidence observed by the “COVID-CF project in Europe” of the European Cystic Fibrosis Society (<https://www.ecfs.eu/covid-cf-project-europe>). According to preliminary data from this project, among approximately 48,000 European CF patients, eight patients needed ICU admission and only three died for COVID-19 (CFR 3.2%) (as of 15 July 2020).

Several factors might play a role in reducing the prevalence and, perhaps, the lethality of COVID-19: typical behavior in patients with CF, who might wear protective masks and avoid contact with sick people, use of antibiotics, or host factors. A more mild form of COVID-19 might be expected because disruption of IL-6 signaling in CF lungs occurs through elevated serine protease release and subsequent cleavage of both membrane-bound and soluble IL-6 receptors.<sup>6</sup> Chronic anti-inflammatory therapies, including azithromycin and DNase, administered to 56% and 51% of our patients, respectively, might also contribute to the protection.

Altogether, these data strengthen the hypothesis that CF may constitute an advantage in COVID-19, encouraging the scientific community to look for the possible reasons behind this unexpected scenario.

# Abbreviations

CF: Cystic Fibrosis

ICU: intensive care unit

SARS-CoV-2: Severe Acute Respiratory Syndrome-Coronavirus-2

COVID-19: Coronavirus disease 2019

NET: Neutrophil extracellular trap

# Declarations

## Ethics approval and consent to participate

This preliminary study reported only a retrospective analysis of data available through the Institutional Database. Personal data were restricted to essential information, and were treated in order to guarantee the respect of privacy of the involved patients, as specifically stated by Italian Law D.Lgs n.196 of 2003 about personal data protection. Therefore, the study did not require preliminary evaluation by the local Ethical Committee.

## Consent for publication

Not applicable

## Availability of data and material

Data were obtained from the Institutional Database at Azienda Ospedaliera Universitaria Integrata, Verona.

## Competing interests

The authors declare that they have no competing interests.

## Funding

Not applicable

## Authors' contributions

VB interpreted the data and drafted the manuscript; FL, SV and MC collected and reviewed clinical data; MC conceived the idea and critically reviewed the manuscript. All the authors read and approved the final manuscript.

## Acknowledgements

The authors thank Dr Seth J Corey (Cleveland Clinic, OH, USA) for the useful suggestions and comments on this work and for his linguistic assistance.

## References

1. Remuzzi A, Remuzzi G. COVID-19 and Italy: what next? *Lancet*. 2020;395(10231):1225-1228.
2. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. *Lancet*. 2020;395(10229):1054-1062.
3. Barnes BJ, Adrover JM, Baxter-Stoltzfus A, et al. Targeting potential drivers of COVID-19: neutrophil extracellular traps. *J Exp Med*. 2020;217(6):e20200652.
4. Dwyer M, Shan Q, D'Ortona S, et al. Cystic fibrosis sputum DNA has NETosis characteristics and neutrophil extracellular trap release is regulated by macrophage migration-inhibitory factor. *J Innate Immun*. 2014;6(6):765-779.
5. Viviani L, Assael BM, Kerem E, groupEAHNS. Impact of the A (H1N1) pandemic influenza (season 2009-2010) on patients with cystic fibrosis. *J Cyst Fibros*. 2011;10(5):370-376.
6. McGreal EP, Davies PL, Powell W, et al. Inactivation of IL-6 and soluble IL-6 receptor by neutrophil derived serine proteases in cystic fibrosis. *Biochim Biophys Acta*. 2010;1802(7-8):649-658.