Renal Replacement Therapy and Concurrent Fluconazole Therapy Increase Linezolid-Related Thrombocytopenia Among Adult Patients

Yueh-chun Hsu

E-Da Hospital <https://orcid.org/0000-0002-1061-8109>

Szu-ying Chen

E-Da Hospital

Yung-jun Hung

E-Da Hospital

Yu-wei Huang (✉ yw.huang0211@gmail.com)

E-Da Hospital <https://orcid.org/0000-0002-5858-5369>

Research

Keywords: Linezolid, Thrombocytopenia, Renal replacement therapy, Fluconazole

Posted Date: May 24th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-532929/v1>

License: This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Abstract

Background: Linezolid, an antibiotic used against gram-positive pathogens, has been reported to be associated with thrombocytopenia. However, limited information is available on susceptibility to thrombocytopenia after linezolid usage. We aimed to investigate the risk factors for linezolid-associated thrombocytopenia.

Methods: We conducted a retrospective cohort study of patients aged ≥ 18 years who received linezolid for ≥ 5 d during hospitalization in 2019. Demographic information, hospitalized information and laboratory data were extracted from electronic medical records. Thrombocytopenia was defined as a platelet count of $<100,000$ cells/mm³ or a reduction from baseline $\geq 25\%$. Binary logistic regression and survival analyses were used to evaluate the risk factors for linezolid-associated thrombocytopenia.

Results: A total of 98 patients were enrolled. Thrombocytopenia occurred in 52 patients (53.1%), with a median first presentation 9 d after initiation of linezolid. There was no significant difference in the mortality or proportion of platelet transfusions between patients with and without thrombocytopenia. A higher risk of linezolid-associated thrombocytopenia was found in patients who received renal replacement therapy (RRT) (OR 4.8 [1.4-16.4]), concurrent fluconazole (OR 3.5 [1.2-9.8]), or a longer duration of linezolid treatment (OR 1.1 [1.0-1.1]). Patients who received RRT or concurrent fluconazole therapy had a shorter median time to develop thrombocytopenia (with vs. without RRT: 8 vs. 15 d; with vs. without fluconazole: 11 vs. 15 d). Those who simultaneously received RRT and fluconazole had the shortest median of time (6.5 d) and the highest risk of developing thrombocytopenia (87.5%).

Conclusions: Patients who received RRT, concurrent fluconazole treatment, or a longer-duration linezolid had a higher risk of developing linezolid-associated thrombocytopenia.

Background

Linezolid, a synthetic oxazolidinone antibiotic, has a fair activity against gram-positive pathogens. Because of its good tissue penetration, linezolid is a key therapeutic choice for nosocomial lung infections and skin or skin structure infections [1]. An oral bioavailability of 100% makes it easy to adjust for administration. According to the manufacturer's recommendation [2], there is no need to adjust the dose for patients with renal impairment or mild-to-moderate hepatic impairment.

Regarding the safety of linezolid, a variety of adverse effects, such as myelosuppression, peripheral and optic neuropathy, serotonin syndrome, and lactic acidosis have been reported [2]. In comparator-controlled phase III studies [3], diarrhea, nausea, and headaches were frequent adverse effects. Although under longer treatment durations, linezolid-treated patients had a higher potential for lower platelet counts compared to the comparator, the difference was not statistically significant. However, emerging observational studies have demonstrated the thrombocytopenic effect of linezolid [4–13]. Depending on different definitions of thrombocytopenia, the incidence of thrombocytopenia ranged from 16.7–48.4% in adults treated with intravenous or oral linezolid [4–13].

Although several studies have highlighted the susceptible population of linezolid-associated thrombocytopenia, the results are inconclusive. Possible risk factors including high daily dose per kg [4, 6], long duration [5, 8, 11, 13], renal impairment [5–8, 10–13], and low baseline platelet counts [9, 11, 12] have been reported. One study [9] showed that carbapenem combination therapy and low baseline platelet count were significant predictors, and the duration of linezolid administration or renal function had little effect. However, a study conducted by Choi *et al.* [11] revealed that low-dose aspirin was a possible risk factor, and the effect of carbapenem combination therapy was not significant.

More than 60% of patients admitted to the intensive care unit may suffer thrombocytopenia [14–16]. Multiple factors predispose an individual to platelet deficiency, including sepsis, major trauma, and drug exposure [14–16].

Medications that may induce thrombocytopenia are as follows: abciximab [15], amiodarone [11], aspirin [11, 15], clopidogrel [15], digoxin [11], dipyridamole [15], eptifibatide [15], fluconazole [17], haloperidol [11], heparin [11, 14–16], oxaliplatin [16], piperacillin [11, 16], rifampin [11, 16], ticlopidine [15], trimethoprim-sulfamethoxazole [11, 15, 16], valproic acid [11, 15], and vancomycin [15, 16].

In addition to linezolid, a variety of predisposing factors and drugs may contribute to thrombocytopenia. For clinicians, it is urgent to define a susceptible population to avoid harmful outcomes. Therefore, we conducted an observational study to evaluate the risk factors of linezolid-associated thrombocytopenia. However, whether a patient's predisposing factors or combination treatment with certain drugs aggregates linezolid-related thrombocytopenia is still inconclusive.

Methods

Study design and population

We designed a retrospective cohort study to investigate the possible risk factors for linezolid-associated thrombocytopenia. Patients ≥ 18 years and hospitalized at the E-Da hospital from January 1, 2019 to December 31, 2019, were recruited into this study. We reviewed the electronic medical records of patients who received linezolid orally or intravenously for ≥ 5 d. All linezolid prescriptions were approved by the infection specialist. Patients with any of the following were excluded: A linezolid treatment duration < 5 d, baseline platelet count $< 100,000$ cells/mm³, received platelet transfusions within 10 d before the linezolid treatment, platelet count not monitored after starting therapy, had undergone chemotherapy or radiation for malignancy, presence of hematological diseases. Our study adhered to ethical restrictions and was approved by the Institutional Review Board of E-Da Hospital (EMRP-109-142).

Individual characteristics

Individual characteristics and information from hospitalization, including age, sex, body weight, height, comorbidity, history of renal replacement therapy (RRT), date of admission, and date of discharge were collected from the medical information system. Diseases that had been documented in medical records more than three times in six months prior to the initiation until the end of therapy were defined as comorbidities. In addition, admission to the intensive care unit during hospitalization, and history of receiving chemotherapy or radiotherapy in the six months prior to initiation until discontinuation of linezolid were also collected.

Information about linezolid usage such as type of infection, route of administration, dosage, and dosing interval were collected from electronic medical records. Previously prescribed glycopeptides, concurrent antibiotics, and combination medications that might affect platelet count and the date of platelet transfusions were reviewed. Combination medications were defined as medications that were co-administrated with linezolid for ≥ 3 d.

Laboratory data

Laboratory data, such as microbiological isolates, platelet count, and serum creatinine, were collected from electronic medical records.

The baseline platelet count was defined as the latest platelet count prior to linezolid treatment. The percentage change in platelet count was calculated by baseline and the lowest platelet count after initiating therapy. According to the latest version of the Common Terminology Criteria for Adverse Events (CTCAE), version 5 [18], thrombocytopenia was

defined as a platelet count $< 100,000$ cells/mm³ [15] or a reduction from baseline $\geq 25\%$. Platelet counts that recovered to $\geq 150,000$ cells/mm³ or baseline, and platelet counts after linezolid discontinuation ≥ 14 d were also collected.

The latest data on serum creatinine levels before initiation of linezolid were collected. Baseline renal function was estimated using the Cockcroft-Gault equation [19]. Enrolled patients were classified into four groups: creatinine clearance (CrCL) < 30 , $30 \leq \text{CrCL} < 60$, $\text{CrCL} \geq 60$ mL/min, and RRT.

Statistical analysis and sample size calculation

Continuous and categorical variables were analyzed using the t-test and chi-square test, respectively. For nonparametric variables, the Mann–Whitney U test or Fisher’s exact test was used instead. The association between risk factors and thrombocytopenia was evaluated using binary logistic regression and adjustment for potential confounders. Variables with a P -value < 0.1 , in the univariable model were further evaluated in the multivariable model. Analyses were presented as estimates and 95% confidence intervals, and P -values < 0.05 , were considered significant. The time between initiating the treatment and the occurrence of thrombocytopenia was analyzed using Kaplan-Meier curves and was tested using the log-rank test. Statistical analyses were performed using SAS version 9.4.

Results

A total of 216 patients were prescribed intravenous or oral linezolid in 2019. After excluding 118 patients with, 98 patients were finally included in this study (Fig. 1). Table 1 shows the individual characteristics of the 98 study subjects stratified by thrombocytopenia. The mean age of all study subjects was 69.0 ± 15.7 years and was equal in both sexes. Sixty-five patients (66.3%) received intravenous linezolid. The average treatment duration of linezolid was 13.9 ± 7.2 d. Urinary tract infection was observed in 48.0% of patients, and 52.0% of patients had a history of intensive care unit admission. Vancomycin-resistant *Enterococcus* isolates and oxacillin-resistant *Staphylococcus* isolates were present in 71.7% and 13.0% of patients, respectively.

Thrombocytopenia occurred in 52 patients (53.1%), with a median of 9 d after initiation of linezolid, and 8 of them (15.4%) received platelet transfusions. After discontinuation, the median duration of platelet counts returned to normal range was 10 d.

There was no significant difference in mortality (26.9% and 23.9%, $P = 0.7330$) and proportion of platelet transfusions (15.4% and 4.4%, $P = 0.0716$) between those with and without thrombocytopenia. Patients who developed thrombocytopenia had a significantly higher proportion of patients receiving RRT (36.5% vs. 10.9%), intravenous administration of linezolid (75.0% vs. 56.5%), male (59.6% vs. 39.1%), and longer median duration (14.0 vs. 10.5 d). In addition, a higher ratio of diabetes mellitus, chronic kidney disease, and concurrent use of fluconazole were found in thrombocytopenia (Table 1).

Table 2 shows the odds ratios of the risk factors associated with linezolid-associated thrombocytopenia. In the crude model, patients who were male, diabetic, receiving RRT, intravenous linezolid, or concurrent fluconazole therapy were found to have increased odds ratios of linezolid-associated thrombocytopenia compared with their counterparts. In the main model, we found increased odds ratios of linezolid-associated thrombocytopenia in patients who received longer duration of linezolid, RRT, and concurrent fluconazole therapy, with an OR of 1.071 (95% CI 1.000–1.146), OR 4.773 (95% CI 1.390–16.389), and OR 3.474 (95% CI 1.230–9.810), respectively.

Figure 2 illustrates the Kaplan-Meier survival curves of linezolid-associated thrombocytopenia events in patients who received RRT or fluconazole treatment. We found that patients who underwent RRT had a shorter median time (with vs. without RRT: 8 vs. 15 d, log-rank test $P = 0.0043$; Fig. 2A) and higher risk (with vs. without RRT: 79.2% vs. 44.6%, $P = 0.0032$) of developing linezolid-associated thrombocytopenia. However, there was no significant difference in the incidence of thrombocytopenia between $\text{CrCL} < 30$, $30 \leq \text{CrCL} < 60$ and $\text{CrCL} \geq 60$ mL/min (results detailed in Additional file 1: Fig. 1S). Among the 24 patients who received RRT, 16 (67%) had end-stage renal disease (ESRD) who underwent RRT, while the others underwent emergent RRT for acute kidney injury (AKI). A slightly higher percentage of thrombocytopenia was noted among ESRD patients than among AKI patients, but the difference was not statistically significant (ESRD vs. AKI: 81.25% vs. 75.00%, $P = 0.7223$).

Patients who received fluconazole were also observed to have a shorter median time (with vs. without fluconazole: 11 vs. 15 d, log-rank test $P = 0.0332$; Fig. 2B) and higher risk (with vs. without fluconazole: 66.7% vs. 45.2%, $P = 0.0397$) of developing linezolid-associated thrombocytopenia.

Figure 3 illustrates the subgroup analyses of linezolid-associated thrombocytopenia stratified by RRT and fluconazole therapy. The incidence of linezolid-associated thrombocytopenia was 34.8%, 60.7%, 70.5%, and 84.5%, respectively, showing an incremental trend in subgroups without RRT and fluconazole, with fluconazole, with RRT, and with both RRT and fluconazole. Multiple logistic regression showed that the odds ratios were 13.125 (95% CI 1.482–1116.267), 5.625 (95% CI 1.558–20.312), and 2.898 (95% CI 1.097–7.654), respectively, among patients who received concurrent RRT and fluconazole, RRT alone, and fluconazole alone compared with patients who did not receive RRT and fluconazole therapy.

Table 1
Individual and characteristics data of enrolled patients

Characteristics	With thrombocytopenia (n = 52)	Without thrombocytopenia (n = 46)	P-value
Age (year), mean $\pm$ SD	71.0 $\pm$ 16.1	66.7 $\pm$ 15.2	0.1824
Male	31 (59.62)	18 (39.13)	0.0430
BMI (kg/m ²), mean $\pm$ SD	24.9 $\pm$ 4.4	25.0 $\pm$ 6.3	0.9529
Renal function, mL/min			
CrCL $\geq$ 60	12 (23.08)	18 (39.13)	0.0277
30 $\leq$ CrCL < 60	11 (21.15)	12 (26.09)	
CrCL < 30	10 (19.23)	11 (23.91)	
Renal replacement therapy	19 (36.54)	5 (10.87)	
ICU admission, no (%)	30 (57.69)	21 (45.65)	0.2338
Comorbidity, no (%)			
Hypertension	36 (69.23)	30 (65.22)	0.6724
Diabetes mellitus	29 (55.77)	14 (30.43)	0.0117
CKD	20 (38.46)	7 (15.22)	0.0102
Liver diseases	9 (17.31)	6 (13.04)	0.5585
Platelet (10 ³ cells/mm ³), mean $\pm$ SD			
Baseline	283.8 $\pm$ 136.1	270.7 $\pm$ 101.3	0.5870
Nadir during therapy	131.3 $\pm$ 82.9	290.7 $\pm$ 111.7	< .0001
After discontinuation ($\geq$ 14 d)	280.5 $\pm$ 148.9	300.0 $\pm$ 122.6	0.5491
Previous glycopeptide, no (%)			
None	24 (46.15)	24 (52.17)	0.4908
Vancomycin	5 (9.62)	7 (15.22)	
Teicoplanin	22 (42.31)	13 (28.26)	
Both vancomycin and teicoplanin	1 (1.92)	2 (4.35)	
Median duration of linezolid (day)	14.0	10.5	0.0310
Route of linezolid, no (%)			

Characteristics	With thrombocytopenia (n = 52)	Without thrombocytopenia (n = 46)	P value
Oral	7 (13.46)	16 (34.78)	0.0456
Intravenous	39 (75.00)	26 (56.52)	
IV and oral	6 (11.54)	4 (8.70)	
Types of infections, no (%)			
Urinary tract infection	25 (48.08)	22 (47.83)	0.4087
Skin and soft tissue infection	9 (17.31)	5 (10.87)	
Bacteremia	8 (15.38)	7 (15.22)	
Intra-abdominal infection	6 (11.54)	4 (8.70)	
Respiratory infection	1 (1.92)	6 (13.04)	
Others	3 (5.77)	2 (4.35)	
Microbiological isolate, no (%)			0.3857
Enterococcus sp.	6 (11.54)	3 (6.52)	
VRE strain	37 (71.15)	29 (60.00)	
Oxacillin-resistant Staphylococcus	4 (7.69)	8 (17.39)	
Others	5 (9.62)	6 (13.04)	
Concurrent medications, no (%)			0.7799
Aspirin or clopidogrel	9 (17.31)	7 (15.22)	
Amiodarone	2 (3.85)	5 (10.87)	0.2478
Piperacillin/tazobactam	5 (9.62)	4 (8.70)	0.2695
Cephalosporin	15 (28.85)	10 (21.74)	0.4205
Carbapenem	27 (51.92)	16 (34.78)	0.0879
Colistin	10 (19.23)	9 (19.57)	0.9667
Fluconazole	24 (46.15)	12 (26.09)	0.0397
Fluoroquinolone	16 (30.77)	13 (28.26)	0.7860

Characteristics	With thrombocytopenia (n = 52)	Without thrombocytopenia (n = 46)	P-value
Data are presented as mean ± standard deviation; BMI, body mass index; CrCL, creatinine clearance; ICU, intensive care unit;			
IV, intravenous; CKD, Chronic kidney disease; VRE, vancomycin-resistant enterococcus.			

Table 2
Association between risk factors and incidence of linezolid-associated thrombocytopenia

	Crude model			Main model		
	OR	95% CI	P-value	OR	95% CI	P-value
Age	1.018	0.992–1.045	0.1851	-	-	-
Male	2.296	1.021–5.166	0.0445	2.210	0.829–5.890	0.1127
BMI	0.998	0.926–1.075	0.9512	-	-	-
Linezolid treatment duration	1.057	0.992–1.125	0.0861	1.071	1.000–1.146	0.0506
Routes of linezolid						
IV	3.427	1.239–9.480	0.0176	2.950	0.884–9.847	0.0786
Both IV and oral	3.427	0.731–16.080	0.1183	1.314	0.187–9.236	0.7840
Baseline platelet count	1.001	0.998–1.004	0.5900	-	-	-
ICU admission	1.623	0.730–3.611	0.2349	-	-	-
Renal replacement therapy	4.721	1.593–13.993	0.0051	4.773	1.390–16.389	0.0130
Hypertension	1.200	0.515–2.795	0.6726	-	-	-
Diabetes mellitus	2.882	1.253–6.628	0.0128	1.698	0.635–4.544	0.2917
Liver diseases	1.395	0.456–4.273	0.5596	-	-	-
Cephalosporin	1.459	0.580–3.671	0.4218	-	-	-
Carbapenem	2.025	0.896–4.574	0.0897	1.056	0.397–2.813	0.9125
Fluconazole	2.428	1.033–5.708	0.0419	3.474	1.230–9.810	0.0187
Fluoroquinolone	1.128	0.472–2.695	0.7865	-	-	-
Main model was adjusted for gender, treatment duration, routes of administration, renal replacement therapy, diabetes mellitus, concurrent carbapenem, and concurrent fluconazole therapy.						
BMI, body mass index; ICU, intensive care unit; IV, intravenous; CI, confidence interval						

Discussion

To the best of our knowledge, this study is the first to demonstrate that patients who received RRT or concurrent fluconazole treatment may have a higher incidence of linezolid-associated thrombocytopenia. In addition, we revealed

the real-world situation and the association between clinical risk factors and linezolid-associated thrombocytopenia.

By performing binary logistic regression, RRT, concurrent fluconazole therapy, and longer duration of linezolid were identified as risk factors for linezolid-associated thrombocytopenia.

We discovered that patients who received RRT had a significantly higher risk (with vs. without RRT: 79.2% vs. 44.6%, $P = 0.0032$) and shorter median time (with vs. without RRT: 8 vs. 15 days, $P = 0.0043$) to have thrombocytopenia. Our finding was compatible with Hanai *et al.*'s results, which showed an onset time of 8.5 d in patients undergoing hemodialysis [8]. Hirano *et al.* found that patients with CrCL < 30 mL/min (60.0%) had a significantly higher incidence of thrombocytopenia than those with CrCL > 60 mL/min (26.4%) [5]. Moraza *et al.* and Hanai *et al.* [8] reported similar findings [7]. Several studies have revealed that renal function plays a critical role in thrombocytopenia associated with linezolid [5–8, 10, 12, 20]. The manufacturer recommends that the pharmacokinetic parameters of the parent drug are not transformed [2], and that no dosage adjustment is needed for renal impairment. However, a higher incidence of thrombocytopenia was noted among our patients who received RRT. Brier *et al.* concluded that the levels of two primary metabolites (PNU-142586 and PNU-142300) were higher in patients requiring hemodialysis [21]. Matsumoto *et al.* suggested that renal impairment elevated linezolid trough concentration, and higher drug exposure might be related to thrombocytopenia [22]. Considering the higher possibility of thrombocytopenia in renal dysfunction, physicians should be aware of the harmful outcomes, although the mechanism of linezolid-associated remains unclear.

The literature reports fluconazole-associated thrombocytopenia limited to several case reports. Pasikhova *et al.* reported a case of fluconazole-associated agranulocytosis with thrombocytopenia [17]. One population-based study enrolled 54,803 users of either fluconazole or itraconazole, and only 1 of 34,220 fluconazole users had thrombocytopenia [23]. So far, no interaction between fluconazole and linezolid has been found [24]. Little information is available regarding the underlying fluconazole and linezolid-associated thrombocytopenia. We suppose that the use of fluconazole may indicate more severe sepsis, and sepsis may be the direct result of thrombocytopenia [25]. Therefore, further studies are necessary.

We found that patients treated with linezolid for ≥ 14 d had a relatively higher frequency of thrombocytopenia (duration ≥ 14 d vs. $14 >$ duration ≥ 5 d: 63.83% vs. 43.14%, $P = 0.0403$) (results detailed in Additional file 1: Fig. 2S). Our results are compatible with those of other studies. Hirano *et al.* [5] and Takahashi *et al.* [13] reported a higher incidence of thrombocytopenia among patients with linezolid duration ≥ 14 d. Choi *et al.* [11] even suggested an increased odds ratio for linezolid duration ≥ 7 d. Therefore, we suggest monitoring platelet counts for prolonged durations of linezolid administration, especially for those who received RRT.

According to recent studies, the incidence of thrombocytopenia ranged from 16.7–48.4% in adults treated with linezolid [4–13]. Kim *et al.* [10] revealed a 48.3% risk of thrombocytopenia in a cohort of 60 ICU patients, and thrombocytopenia was defined as a platelet count of < 150,000 cells/mm³ or a decrease of at least 50% from the baseline. The relatively higher incidence (53.1%) in our study could have resulted from different disease severities and definitions of thrombocytopenia.

Our data showed that patients with diabetes mellitus had twice the risk of thrombocytopenia. However, this was not significant after adjusting for confounding factors. Diabetic kidney disease, a microvascular complication, frequently leads patients to undergo dialysis or renal transplantation [26]. Consequently, a higher proportion of diabetes mellitus in the thrombocytopenic group is anticipated. Regarding the administration route, oral-form linezolid was mostly prescribed for discharge in our study. This could be a possible explanation for the higher ratio of intravenous linezolid

in patients with thrombocytopenia. Our results were compatible with the findings of Takahashi *et al.* [13], which showed that oral linezolid decreased the risk of thrombocytopenia.

However, the mechanism underlying linezolid-associated thrombocytopenia remains unclear. Several mechanisms have been proposed. For instance, immune-mediated thrombocytopenia has been proposed by Bernstein *et al.* [27]. Other theories, including the suppression of platelets release from mature megakaryocytes [28], and Tsuji *et al.* assumed that the inhibition of the platelet formation was the most common mechanism [29]. Many studies reported that the linezolid-associated adverse reactions were related to linezolid trough concentration ($C_{\min}$) [30, 31]. Cojutti *et al.* [32] monitored patients' $C_{\min}$ for dose adjustment at day 3–5 of therapy, and 2–8mg/L was taken as desired range for $C_{\min}$. Peripheral venous blood samples were collected 5 min prior to the subsequent dose. Levels of linezolid were analyzed by a validated HPLC analysis method. However, more studies are necessary to clarify the approach of dose adjustment.

Several clinical factors might also affect platelets, such as a history of surgery and recent transfusions [33]. Using radionuclides, the mean life span of transfused platelets in humans was 9.9 ± 0.6 d [34–37]. Therefore, we excluded patients who received platelet transfusions ≤ 10 d before initiation of linezolid. Moreover, the material of the dialyzer may cause thrombocytopenia; polysulfone membranes and amounts of polyvinylpyrrolidone are thought to be related to its occurrence, but the exact mechanism remains to be elucidated [38].

Our observational cohort study has several limitations. First, due to the retrospective nature of the study design, we could not make causal inferences. Second, means of monitoring concentrations of linezolid and its metabolites were not available, and it was difficult to evaluate the pharmacokinetic parameters of linezolid in thrombocytopenic patients. Monitoring linezolid $C_{\min}$ at steady-state condition may be a feasible approach. However, therapeutic drug monitoring of linezolid is in experimental stage, and more studies are warranted.

Conclusions

Renal adjustment is not required according to the official recommendation. However, in our study, we found that the duration of treatment, RRT, and concurrent fluconazole were significantly associated with the risk of thrombocytopenia. Patients who received RRT had a significantly shorter median time to thrombocytopenia after initiation of linezolid. Careful monitoring of platelets is warranted for patients with renal impairment, especially for those already receiving RRT. However, due to the retrospective nature of our study, large randomized control trials are warranted to verify the association between possible clinical factors and the risk of linezolid-associated thrombocytopenia.

Abbreviations

AKI: acute kidney injury; BMI: body mass index; CI: Confidence interval; CKD: Chronic kidney disease; CrCL: creatinine clearance; CTCAE: Common Terminology Criteria for Adverse Events; ESRD: end stage renal disease; ICU: intensive care unit; IV: intravenous; OR: odds ratio; RRT: renal replacement therapy; SD: standard deviation; VRE: vancomycin-resistant enterococcus.

Declarations

Competing interests

The authors declare that they have no competing interests

Funding

This study was not funded by any party.

Authors' contributions

YC Hsu designed the study, reviewed the medical records, collected data, performed statistical analysis and wrote the manuscript. YW Huang designed the study, provided medical opinions and revised the manuscript. SY Chen provided statistical assistance and revised the manuscript. YJ Hung provided hospital resources and related assistance and helped seeking statistical advice.

Acknowledgments

We would like to thank Lei-Yu Chang for offering help to submit ethics approval and Wan-Yi Chang-Chien for providing assistance on performing statistical analysis.

Availability of data and materials

No applicable.

Ethics approval and consent to participate

This study adhered to ethical restriction and was approved by institutional review board of E-Da hospital. Approval number was EMRP-109-142.

Consent for publication

No applicable.

Author details

¹Department of Pharmacy, E-Da hospital, Kaohsiung 824, Taiwan. ²Department of Anesthesiology, Emergency and Critical Care Center, E-Da hospital, I-Shou University, Kaohsiung 824, Taiwan; Department of Nursing, Fooyin University, Kaohsiung 831, Taiwan. ³Division of Occupational Medicine, E-Da Hospital, I-Shou University, Kaohsiung 824, Taiwan. ⁴Division of Surgical Intensive Care, Department of Critical Care Medicine, E-Da Hospital, I-Shou University, Kaohsiung 824, Taiwan; Department of Nursing, Fooyin University, Kaohsiung 831, Taiwan.

References

1. Dryden MS. Linezolid pharmacokinetics and pharmacodynamics in clinical treatment. *J Antimicrob Chemother.* 2011;66(Suppl 4):iv7–15.
2. Product Information. ZYVOX(R) intravenous injection, oral tablets, oral suspension, linezolid intravenous injection, oral tablets, oral suspension. <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>. Accessed 4 November 2020.
3. Rubinstein E, Isturiz R, Standiford HC, Smith LG, Oliphant TH, Cammarata S, Hafkin B, Le V, Remington J. Worldwide assessment of linezolid's clinical safety and tolerability: comparator-controlled phase III studies. *Antimicrob Agents Chemother.* 2003;47:1824–31.
4. Niwa T, Suzuki A, Sakakibara S, Kasahara S, Yasuda M, Fukao A, Matsuura K, Goto C, Murakami N, Itoh Y. Retrospective cohort chart review study of factors associated with the development of thrombocytopenia in adult

- Japanese patients who received intravenous linezolid therapy. *Clin Ther*. 2009;31:2126–33.
5. Hirano R, Sakamoto Y, Tachibana N, Ohnishi M. Retrospective analysis of the risk factors for linezolid-induced thrombocytopenia in adult Japanese patients. *Int J Clin Pharm*. 2014;36:795–9.
 6. Natsumoto B, Yokota K, Omata F, Furukawa K. Risk factors for linezolid-associated thrombocytopenia in adult patients. *Infection*. 2014;42:1007–12.
 7. Moraza L, Leache L, Aquerreta I, Ortega A. Linezolid-induced haematological toxicity. *Farm Hosp*. 2015;39:320–6.
 8. Hanai Y, Matsuo K, Ogawa M, Higashi A, Kimura I, Hirayama S, Kosugi T, Nishizawa K, Yoshio T. A retrospective study of the risk factors for linezolid-induced thrombocytopenia and anemia. *J Infect Chemother*. 2016;22:536–42.
 9. Kaya Kilic E, Bulut C, Sonmezer MC, Ozel O, Ataman Hatipoglu C, Tuncer Ertem G, Tulek N, Kinikli S. Risk factors for linezolid-associated thrombocytopenia and negative effect of carbapenem combination. *J Infect Dev Ctries*. 2019;13:886–91.
 10. Kim HS, Lee E, Cho YJ, Lee YJ, Rhie SJ. Linezolid-induced thrombocytopenia increases mortality risk in intensive care unit patients, a 10 year retrospective study. *J Clin Pharm Ther*. 2019;44:84–90.
 11. Choi GW, Lee JY, Chang MJ, Kim YK, Cho Y, Yu YM, Lee E. Risk factors for linezolid-induced thrombocytopenia in patients without haemato-oncologic diseases. *Basic Clin Pharmacol Toxicol*. 2019;124:228–34.
 12. Lima LS, Brito E, Mattos K, Parisotto EB, Perdomo RT, Weber SS. A retrospective cohort study to screen linezolid-induced thrombocytopenia in adult patients hospitalized in the Midwestern Region of Brazil. *Hematol Transfus Cell Ther*. 2020;42:230–7.
 13. Takahashi Y, Takesue Y, Nakajima K, Ichiki K, Tsuchida T, Tatsumi S, Ishihara M, Ikeuchi H, Uchino M. Risk factors associated with the development of thrombocytopenia in patients who received linezolid therapy. *J Infect Chemother*. 2011;17:382–7.
 14. Zarychanski R, Houston DS: Assessing thrombocytopenia in the intensive care unit: the past, present, and future. *Hematology Am Soc Hematol Educ Program* 2017; 2017:660–666.
 15. Ostadi Z, Shadvar K, Sanaie S, Mahmoodpoor A, Saghaleini SH. Thrombocytopenia in the intensive care unit. *Pak J Med Sci*. 2019;35:282–7.
 16. Aluru N, Samavedam S. Thrombocytopenia in Intensive Care Unit. *Indian J Crit Care Med*. 2019;23:185-s188.
 17. Pasikhova Y, Ludlow SP. Fluconazole associated agranulocytosis and thrombocytopenia. *Int J Clin Pharm*. 2014;36:268–70.
 18. Common Terminology Criteria for Adverse Events (CTCAE). Version 5.0
https://ctep.cancer.gov/protocolDevelopment/electronic_applications/docs/CTCAE_v5_Quick_Reference_5x7.pdf. Accessed November 1 2020.
 19. Cockcroft DW, Gault MH. Prediction of creatinine clearance from serum creatinine. *Nephron*. 1976;16:31–41.
 20. Kato H, Hamada Y, Hagihara M, Hirai J, Yamagishi Y, Matsuura K, Mikamo H. Bicytopenia, especially thrombocytopenia in hemodialysis and non-hemodialysis patients treated with linezolid therapy. *J Infect Chemother*. 2015;21:707–12.
 21. Brier ME, Stalker DJ, Aronoff GR, Batts DH, Ryan KK, O'Grady M, Hopkins NK, Jungbluth GL. Pharmacokinetics of linezolid in subjects with renal dysfunction. *Antimicrob Agents Chemother*. 2003;47:2775–80.
 22. Matsumoto K, Takeshita A, Ikawa K, Shigemi A, Yaji K, Shimodozono Y, Morikawa N, Takeda Y, Yamada K. Higher linezolid exposure and higher frequency of thrombocytopenia in patients with renal dysfunction. *Int J Antimicrob Agents*. 2010;36:179–81.

23. Bradbury BD, Jick SS. Itraconazole and fluconazole and certain rare, serious adverse events. *Pharmacotherapy*. 2002;22:697–700.
24. Hashemian SMR, Farhadi T, Ganjparvar M. Linezolid: a review of its properties, function, and use in critical care. *Drug Des Devel Ther*. 2018;12:1759–67.
25. Larkin CM, Santos-Martinez MJ, Ryan T, Radomski MW. Sepsis-associated thrombocytopenia. *Thromb Res*. 2016;141:11–6.
26. Braunwald E. Diabetes, heart failure, and renal dysfunction: The vicious circles. *Prog Cardiovasc Dis*. 2019;62:298–302.
27. Bernstein WB, Trotta RF, Rector JT, Tjaden JA, Barile AJ. Mechanisms for linezolid-induced anemia and thrombocytopenia. *Ann Pharmacother*. 2003;37:517–20.
28. Tajima M, Kato Y, Matsumoto J, Hirosawa I, Suzuki M, Takashio Y, Yamamoto M, Nishi Y, Yamada H. Linezolid-Induced Thrombocytopenia Is Caused by Suppression of Platelet Production via Phosphorylation of Myosin Light Chain 2. *Biol Pharm Bull*. 2016;39:1846–51.
29. Tsuji Y, Holford NHG, Kasai H, Ogami C, Heo YA, Higashi Y, Mizoguchi A, To H, Yamamoto Y. Population pharmacokinetics and pharmacodynamics of linezolid-induced thrombocytopenia in hospitalized patients. *Br J Clin Pharmacol*. 2017;83:1758–72.
30. Nukui Y, Hatakeyama S, Okamoto K, Yamamoto T, Hisaka A, Suzuki H, Yata N, Yotsuyanagi H, Moriya K. High plasma linezolid concentration and impaired renal function affect development of linezolid-induced thrombocytopenia. *J Antimicrob Chemother*. 2013;68:2128–33.
31. Kawasuji H, Tsuji Y, Ogami C, Kimoto K, Ueno A, Miyajima Y, Kawago K, Sakamaki I, Yamamoto Y. Proposal of initial and maintenance dosing regimens with linezolid for renal impairment patients. *BMC Pharmacol Toxicol*. 2021;22:13.
32. Cojutti PG, Merelli M, Bassetti M, Pea F. Proactive therapeutic drug monitoring (TDM) may be helpful in managing long-term treatment with linezolid safely: findings from a monocentric, prospective, open-label, interventional study. *J Antimicrob Chemother*. 2019;74:3588–95.
33. Gauer RL, Braun MM. Thrombocytopenia. *Am Fam Physician*. 2012;85:612–22.
34. Harker LA, Finch CA. Thrombokinetis in man. *J Clin Invest*. 1969;48:963–74.
35. Richard J. Davey JPA: Post-Transfusion Red Blood Cell and Platelet Survival and Kinetics: Basic Principles and Practical Aspects. 2007:455–466.
36. H. AR: Blood platelet kinetics and platelet transfusion. *The Journal of clinical investigation* 2013; 123:4564–4565.
37. Richard JDavey. JPA: Blood Banking and Transfusion Medicine. Second edn; 2007.
38. Daugirdas JT, Bernardo AA. Hemodialysis effect on platelet count and function and hemodialysis-associated thrombocytopenia. *Kidney Int*. 2012;82:147–57.

Figures

Figure 1

Flow chart showing enrollment into the study and clinical outcomes.

(*significant difference at $p \leq 0.05$)

Figure 2

Kaplan-Meier survival curves to reveal the impact on the incidence of thrombocytopenia. (A) Renal replacement therapy (RRT). (B) Fluconazole. Difference between groups were tested by log-rank test.

(*significant difference at $p \leq 0.05$)

Figure 3

Incidence of linezolid-associated thrombocytopenia among patients who received both RRT and fluconazole, RRT alone, fluconazole alone, and denied RRT and fluconazole

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementaryinformation.docx](#)