Comparison of Prevalence of Degenerative Findings on Lumbar MRI among Sciatica Patients classified using the McKenzie Method.

Sinikka Kilpikoski (✉ sinikka.kilpikoski@kolumbus.fi)

Department of Physical and Rehabilitation Medicine, Central Finland Health Care District Hospital, Jyväskylä

Eetu Suominen

Faculty of Medicine, University of Turku, Turku

Jussi Repo

Department of Orthopaedics and Traumatology, Tampere University Hospital Tampere

Arja H Häkkinen

Health Sciences, University of Jyväskylä, Jyväskylä

Kati Kyrölä

Department of Orthopaedics and Traumatology, Central Finland Health Care District Hospital, Jyväskylä

Hannu Kautiainen

Primary Health Care Unit, Kuopio University Hospital, Finland; Folkhälsan Research Center, Helsinki

Jari Ylinen

Department of Physical and Rehabilitation Medicine, Central Finland Health Care District Hospital, Jyväskylä

Keywords:

Posted Date: May 26th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-536100/v1>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

The authors have requested that this preprint be removed from Research Square.