

Prevalence and factors associated with skin-to-skin contact (SSC) practices: findings from a population-based cross-sectional survey in 10 selected districts of Bangladesh

Nazia Binte Ali (✉ nazia.ali@icddr.org)

International Centre for Diarrhoeal Disease Research Bangladesh

Sabrina Sharmin Priyanka

International Centre for Diarrhoeal Disease Research Bangladesh

Bal Ram Bhui

Save the Children

Samantha Herrera

PATH

Md. Rashidul Azad

International Centre for Diarrhoeal Disease Research Bangladesh

Afsana Karim

Save the Children

Shams Zubair

Save the Children

Mohammad Mahmoodur Rahman

International Centre for Diarrhoeal Disease Research Bangladesh

SM Rokonzaman

International Centre for Diarrhoeal Disease Research Bangladesh

Umme Salma Jahan Meena

Save the Children

Shams El Arifeen

International Centre for Diarrhoeal Disease Research Bangladesh

Sk Masum Billah

International Centre for Diarrhoeal Disease Research Bangladesh

Research Article

Keywords: Skin to skin contact, Essential Newborn Care (ENC), Facility delivery, Caeserean Section, Antenatal Care (ANC), Sociodemographic Characteristics, Low and Middle Income Countries, Bangladesh

Posted Date: May 27th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-537359/v1>

License:  This work is licensed under a Creative Commons Attribution 4.0 International License.

[Read Full License](#)

Abstract

Background: Skin-to-skin contact (SSC) is one of the eight proven Essential Newborn Care (ENC) practices that improve newborn survival. Despite having myriads of benefits, it is one of the least used ENC interventions especially in the low and middle-income countries (1% to 74%). In Bangladesh, the prevalence of SSC practices was 26% in 2014. To advance the use of this intervention, it is important to assess its prevalence within the population and to identify factors that facilitate or inhibit SSC practices in Bangladesh.

Methods: We used baseline household survey data of USAID's MaMoni MNCSP project conducted in 10 districts of Bangladesh in 2019. Our analysis included 13,695 recently delivered women (RDW) with a live birth outcome. Our primary outcome was the mother's reported SSC practice. We examined various antepartum, intrapartum, newborn, and sociodemographic factors associated with SSC using a multivariable generalized linear model. Our findings were reported using adjusted Prevalence Risk Ratios (aPRRs) and 95% Confidence Intervals (CIs).

Results: Overall, 28% of RDW reported practicing SSC across the 10 surveyed districts. Our multivariable analysis showed that public facility delivery (aPRR 2.01; 95%CI: 1.80, 2.26), private facility delivery (aPRR 1.23; 95%CI: 1.06, 1.42) and 4 antenatal care (ANC) visits at least one from a medically trained provider (MTP) (aPRR 1.17; 95%CI: 1.03, 1.26) had a significant positive association with SSC practices. Caesarean section (aPRR 0.64; 95%CI: 0.56, 0.73) had a significant negative association with SSC practices. We also found a significant positive association of SSC practices with mothers' who perceived the birth size of their baby to be small, mothers with a higher education level (≥ 10 years), and mothers from households in the highest wealth quintile.

Conclusions:

The prevalence of SSC is very low in the surveyed districts of Bangladesh. Considering the factors associated with SSC, relevant stakeholders need to increase their efforts on improving ANC and facility delivery coverages as well as improving SSC practices in the facilities especially after caesarean deliveries. Countries with a high burden of home deliveries also need to emphasize community-based interventions and increasing coverage of skilled birth attendance for improving this life-saving intervention.

Background

The first 28 days of life, known as the neonatal period, is the most crucial time for newborn survival. Globally, around 2.5 million babies died in the neonatal period in 2017(1). Despite making significant improvements in the reduction of mortality among under-five children, the reduction of mortality in the neonatal period remains an unfinished Millennium Development Goal (MDG) agenda for many low and middle-income countries(2–5). In Bangladesh, the neonatal mortality rate remains high at 30 per 1000 live births, constituting around 67 % of all under-five deaths in 2017(6). To reduce newborn death, the

World Health Organization (WHO) recommends a set of simple life-saving interventions, known as Essential Newborn Care (ENC) practices (7, 8). The core interventions included in ENC are early initiation of breastfeeding, thermal regulation (drying/wrapping the baby, skin-to-skin contact (SSC) and delayed bathing), and appropriate cord care. (7, 8). A significant positive association of SSC has been shown with thermal regulation and early initiation of breastfeeding practices (9).

WHO defines SSC as “*placing the baby naked on the mother’s bare chest, in a prone position covered by a cloth/blanket*” (10). SSC is recommended to start immediately or within 10 minutes of birth after vaginal deliveries and as soon as the mother regains consciousness after cesarean Sect. (11). It is suggested to continue for an hour or longer if well tolerated by both the mother and the newborn (12). The benefits of SSC on newborn and mother’s health are myriad. A recent Cochrane review of 46 randomized control trials including 3850 women and their infants suggested that immediate and early SSC (immediately after birth up till one hour) significantly improved the initiation of breastfeeding rates, duration of breastfeeding, and exclusive breastfeeding rates (13–15). Newborns who received SSC had better stability in physiological parameters; cardiorespiratory parameters and blood glucose level (13). Moreover, this critical period of contact provides an opportunity for neurobehavioral development of the newborn through the establishment of mother-infant attachment (13–15). SSC benefits extend to the mother also, with improvements in the shortening of the third stage of labor, placental separation, uterine contraction, and milk let down (7).

Despite the numerous benefits, globally the more common practice is to separate the newborn from the mother after birth in both hospital and community settings (15). A recent systematic review of literature from 28 countries found that the prevalence of SSC varied between 1–74% in the low and middle-income countries (16). The review also showed a low prevalence of immediate SSC among countries in the South-East Asian region; Nepal and India at 15% (17). Similar to other neighboring countries, the prevalence of SSC is low in Bangladesh at 26% in 2014 (18).

For Bangladesh to achieve the Sustainable Development Goal (SDG) target for neonatal mortality, at 12 deaths per 1000 live births by 2030, the country will need to substantially improve essential newborn care coverage, especially SSC practices (19). The first step in improving SSC practices is to identify the evidence on factors influencing the practice to develop context-specific solutions. The challenges in using existing evidence are that very few studies document the population-level determinants of SSC practice in Bangladesh, and the most recent data from Bangladesh on SSC practices are outdated (latest reflects data from 2012) and do not reflect the substantial changes in maternal care-seeking practices that have occurred over the past years (18). For example, between 2014 and 2017, the coverage of four or more antenatal care (ANC) visits increased from 31–47% while facility delivery increased from 37–50%. Both are important predictors of SSC (20). Considering these evidence gaps, it is important to undertake an updated analysis using the most recent data to understand the current context of SSC practices in the local context.

In this study, we estimated the prevalence and determined the factors associated with SSC practices in 10 surveyed districts in Bangladesh. Findings from this study will generate updated evidence for program managers, researchers, and policymakers to develop and test context-specific solutions for improving coverage of this potentially life-saving intervention.

Methods

Data and settings:

We used baseline household survey data of the MaMoni Maternal Newborn Care Strengthening Project (MNCSP) conducted in 10 districts in Bangladesh in 2019: Noakhali, Feni, Chandpur, Lakshmipur, Brahmanbaria, Habiganj, Manikganj, Faridpur, Madaripur, Khushtia. MaMoni MNCSP is a USAID-funded five-year project (2018–2022) focusing on strengthening maternal and newborn care interventions through the public health system in Bangladesh. This project aims to ensure equitable and high-quality maternal and newborn care service delivery, addressing health system challenges. icddr,b, a member of the MaMoni MNCSP consortium, conducted the baseline population-based survey to measure coverage of key maternal and newborn health interventions.

The survey collected information from 14,982 recently delivered women (RDW) who had a birth outcome (abortion/stillbirth/live births) in the preceding 15 months using a structured questionnaire adapted from BDHS 2017. For this paper, we included 13,695 women who had a live birth outcome (1287 women having a stillbirth or abortion were excluded). We used data on women's sociodemographic characteristics, newborn characteristics, ANC, place, and mode of delivery, and SSC practices, to estimate the prevalence and factors associated with SSC practices in the study areas.

Sample size and sampling:

The survey sample size was calculated to provide district-level estimates at a 5% level of precision for selected maternal, newborn, and family planning indicators. Considering the $\pm 5\%$ error margin and adjusting for clustering, an average of 120 clusters from each district with a cluster size of 10 was required. Our cluster size was 100 households. In this study, we included RDW with a birth outcome in the preceding 15 months to better evaluate changes over the life of the project. The sampling of clusters was done using the probability proportional to size (PPS) technique. The sampling frame for the PPS was prepared using census data from the Bangladesh Bureau of Statistics community series 2011 and population size for the study areas was adjusted using growth rates.

Data collection and quality assurance:

A household listing exercise was first conducted to identify eligible RDW for the interviews. To identify the cluster boundaries, data collectors (listers) first identified an index household using a random number table from the eligible couple register of the Family Welfare Assistants (FWA). One hundred households were then listed starting from the index household in the clockwise direction to define the cluster boundary. During the listing exercise, listers also identified all RDW having any birth outcome (live birth,

stillbirth, or abortion) in the three years preceding the survey and collected household-level data. Later, a list of survey respondents (RDW with birth outcomes in the past 15 months of the survey) was generated using an automated system. Interviewers visited the sampled women and conducted full interviews in Bangla. Data collection was carried out between March and September 2019.

To ensure the quality of data collection, our data collectors (listers/ interviewers) and field supervisors who had extensive experience in survey data collection, were extensively trained on the data collection tools between February and March 2019. Training included an in-depth review of the questionnaires, data collection methods, mock interviews, and field practices. Refresher training was provided as required. We used an Andriod-based system with an in-built consistency check system(range and skips) for minimizing data entry error. A real-time dashboard was used for data quality monitoring. Supervisors and investigators conducted occasional field visits for observing data collection activities and provided post-visit feedback to the data collectors and supervisors in review meetings.

Variable Definitions:

Outcome variables:

Our primary outcome variable was SSC practices. We used mothers' reported data for defining SSC (18). Mothers were asked for their last live birth in the previous 15 months if "*the baby was placed naked directly on the bare skin of her chest immediately after birth*". The responses were recorded as either yes or no.

Explanatory variables:

We identified a list of explanatory variables based on an extensive literature review. Four domains of variables were tested; antepartum, intrapartum, newborn, and sociodemographic characteristics of the mother.

The frequency of ANC visits at least one from a medically trained provider (MTP) was considered as an antepartum factor. Categories include no ANC from an MTP, 1–3 ANC visits (at least one from an MTP), and ≥ 4 ANC visits (at least one from an MTP). Any ANC from a "*medical doctor, nurse, midwife, paramedic, family welfare visitor (FWV), community skilled birth attendant (CSBA), or medical assistant/sub-assistant community medical officer (MA/SACMO)*" was defined as ANC from an MTP.

Intrapartum factors included the places of delivery (home, public facility, private facility, and NGO facilities), modes of delivery (vaginal delivery and cesarean sections), and whether the mother reported any complications during delivery. "*Severe headache with blurry vision, convulsions, high blood pressure, excessive vaginal bleeding, prolonged labor > 12 hours, retained placenta, and mal-presentation during delivery*" were included as complications during delivery.

We also examined newborn factors; including the mother's perceived birth size of the baby (average, small or large), birth order (categories are 1–2, 3–4, and 5+), and sex of the baby.

The mother's age categories, the highest level of educational attainment, current occupation (homemaker, unskilled worker, and skilled worker), religion, and household wealth quintiles were the sociodemographic factors analyzed in this study. We used principal component analysis (PCA) for generating the household wealth index. Variables included in the PCA were housing characteristics, ownership of land, household assets, and household possession of domestic animals and transport vehicles. The PCA score was then divided into quintiles to assign household wealth quintile.

Data analysis:

We first examined the frequency and distribution of all explanatory and outcome variables. Sample weight was calculated to adjust for the survey design. We reported numbers and weighted proportions for presenting descriptive statistics. In our bivariate analysis, we used generalized linear models (GLM) to examine the association of explanatory variables with the outcome variable. Bivariate models were adjusted for weighting. Variables having a significant association in bivariate analysis were further examined using a multivariable model. GLM was also used for multivariable analysis. We specified binomial family, log link function (as our outcome is binary), robust estimator (to adjust for nonparametric distribution), and adjusted for sampling weight in our multivariable model. Findings from the bivariate and multivariable models were reported using Prevalence Rate Ratios (PRRs) and 95% confidence intervals. We consider P-value < 0.05 as significant. Stata 14 statistical software was used for analysis.

Ethics:

The survey received ethical approval from the Institutional Review Board of icddr,b and Save the Children US Ethical Review Committee. The survey was conducted following ethical principles of research with human participants. Data collectors read out the informed consent forms and explained the purpose of the survey, risks, benefits, use of data, procedures for ensuring privacy and confidentiality, and the voluntary nature of participation to the study participants. Data collection commenced upon receiving informed written consent.

Results

Out of 13,695 mothers included in the study, more than half (60%) were between the age of 20 to 29 years (Table 1). Almost two-thirds of mothers had some secondary or higher education while only 7% of mothers had no formal education. The majority of mothers surveyed were homemakers (97%).

Overall, 29% of the study participants did not report any ANC from an MTP whereas 45% reported 1–3 ANC (at least one from an MTP) and 27% reported receiving ≥ 4 ANC (at least one from an MTP) (Table 1). Forty-seven percent of the mothers delivered at home, 37% in a private facility, and 15% in a public facility. Among all surveyed mothers, the prevalence of cesarean section was high at 36%. The majority of the mothers who delivered in the private facilities reported cesarean deliveries (85%) while less than one-third of mothers who delivered in the public facilities reported having a cesarean delivery (Fig. 1). One in five mothers reported experiencing complications during delivery. More than two-thirds of

mothers reported that the perceived size of their child at birth was average while one in 10 mothers reported that their child was small at birth. Around 64% of the babies were first or second born.

Table 1

Sociodemographic, antepartum, intrapartum, and newborn characteristics of RDWs in surveyed districts of Bangladesh, 2019

Variables	Categories	Frequency (n) N= 13,695	Percentage* %(95%CI)
Sociodemographic characteristics			
Mother's age	15–19	2028	14.8(14.2, 15.4)
	20–29	8166	59.6(58.8, 60.4)
	30–39	3318	24.2(23.5, 24.9)
	40–49	183	1.3(1.1, 1.5)
Mother's educational level	No formal education	986	7.2(6.8, 7.6)
	Primary incomplete(1-4years)	1570	11.5(10.9, 12.0)
	Primary complete (5 years)	2024	14.8(14.2, 15.4)
	Secondary incomplete (6–9)	6205	45.3(44.5, 46.1)
	Secondary and above (≥ 10)	2910	21.3(20.6, 21.9)
Mother's current occupation	Homemaker	13268	96.9(96.6, 97.2)
	Skilled worker	375	2.7(2.5, 3.0)
	Unskilled worker	52	0.4(0.3, 0.5)
Mother's religion	Muslim	12912	94.3(93.9, 94.7)
	Hindu /others	783	5.7(5.3, 6.1)
Wealth Quintiles	Lowest	2657	19.4(18.7, 20.1)
	Second	2671	19.5(18.8, 20.2)
*weighted			

Variables	Categories	Frequency (n) N = 13,695	Percentage* %(95%CI)
	Middle	2777	20.3(19.6, 20.9)
	Fourth	2758	20.1(19.5, 20.8)
	Highest	2832	20.7(20.0, 21.4)
Antepartum characteristics			
Frequency of ANC visits (at least one from a medically trained provider (MTP))	No ANC from MTP	3948	28.8(28.0, 29.6)
	1–3 ANC (at least one from an MTP)	6091	44.5(43.6, 45.3)
	≥4 (at least one from an MTP)	3656	26.7(25.9, 27.4)
Intrapartum characteristics			
Places of delivery	Home delivery	6417	46.9(46.0, 47.7)
	Public facility	2040	14.9(14.3, 15.5)
	Private facility	5025	36.7(35.9, 37.5)
	NGO facility	133	0.9(0.8, 1.1)
	Missing	80	0.6(0.4, 0.7)
Modes of delivery	Normal delivery	8774	64.1(63.2, 64.9)
	Caeserean section	4914	35.9(35.1,36.7)
	Missing	9	0.1(0.0,0.1)
Any reported complication during delivery	No	10743	78.4(77.7, 79.1)
	Yes	2952	21.5(20.9, 22.2)
New-born characteristics			
*weighted			

Variables	Categories	Frequency (n) N = 13,695	Percentage* %(95%CI)
Birth size	Small	1363	9.9(9.5, 10.5)
	Average	9393	68.6(67.8, 69.4)
	Large	2893	21.1(20.4, 21.8)
	Missing	46	0.3(0.2, 0.4)
Birth order	1–2	8787	64.2(63.4, 64.9)
	3–4	4149	30.3(29.5, 31.1)
	5+	759	5.5(5.2, 5.9)
Sex of the child	Male	7017	48.3(47.4, 49.1)
	Female	6609	51.2(50.4, 52.1)
	Missing	69	0.5(0.3, 0.6)
*weighted			

*Created by Authors

Prevalence of reported SSC practices:

Overall, 28% of mothers reported SSC practices (Table 2). The reported practice of SSC was higher among mothers who had ≥ 4 ANC visits (at least one from an MTP)(30%; 95%CI: 28.8, 31.8) compared to mothers who had no ANC from an MTP (25%; 95%CI: 23.9, 26.7). More mothers who delivered in the public facilities reported practicing SSC (42%; 95%CI: 39.5, 43.8) with their newborn than mothers who delivered at home (26%; 95%CI: 25.3, 27.5) or in a private facility (26%; 95%CI: 24.3, 26.7). The practice of SSC was lower among mothers who delivered by cesarean section (25%; 95%CI: 23.8, 26.2) compared to those who had a vaginal delivery (30%; 95%CI: 29.2, 31.2). Figure 2 showed that reported SSC practices were higher for normal deliveries in both public facilities and private facilities compared to home deliveries, however, the opposite was observed for cesarean deliveries.

Among the newborn characteristics, the reported practice of SSC decreased with increasing the perceived birth size of the baby. It was reported among 34% (95%CI: 31.5, 36.6) of babies perceived to be small,

28% (95%CI: 27.2, 28.9) of babies perceived to be average, and 27% (95%CI: 24.9, 28.2) of babies perceived to be large. The practice of reported SSC declined with increasing birth order; 29% (95%CI: 28.1, 29.9) among first or second birth order, 28% (95%CI: 26.4, 29.1) among third or fourth birth order, and 24% (95%CI: 20.8, 27.0) among fifth or higher birth order. Mothers with a higher level of education and those who belong to higher wealth quintiles reported a higher prevalence of SSC.

Table 2

SSC prevalences by antepartum, intrapartum, newborn, and sociodemographic factors in surveyed districts of Bangladesh, 2019

Variables	Categories	Overall (N = 13695)	Frequency (n = 3878)	Percentages* %(95%CI)
The overall prevalence of SSC		13695	3878	28.3(27.6, 29.1)
Antepartum characteristics				
Frequency of antenatal care(ANC) visits with at least one ANC from a medically trained provider (MTP)	No ANC from MTP	3948	1000	25.3(23.9, 26.7)
	1–3 ANC (at least 1 from an MTP)	6091	1772	29.1(27.9, 30.2)
	≥ 4 (at least 1 from an MTP)	3656	1107	30.3(28.8, 31.8)
Intrapartum characteristics				
Place of delivery	Home delivery	6417	1694	26.4(25.3, 27.5)
	Public facility	2040	850	41.7(39.5, 43.8)
	Private facility	5025	1282	25.5(24.3, 26.7)
	NGO facility	133	30	22.6(15.7, 30.0)
Mode of deliveries	Normal delivery	8774	2650	30.2(29.2, 31.2)
	Caeserean section	4914	1227	25.0(23.8,26.2)
Any complication during delivery	No	10743	2962	27.6(26.7, 28.4)
	Yes	2952	917	31.1(29.4, 32.8)
Newborn characteristics				
Birth size	Small	1363	464	34.0(31.5, 36.6)
* Weighted				

Variables	Categories	Overall (N = 13695)	Frequency (n = 3878)	Percentages* %(95%CI)
	Average	9393	2637	28.1(27.2, 28.9)
	Large	2893	770	26.6(25.0, 28.3)
Birth order	1–2	8787	2548	29.0(28.1, 29.9)
	3–4	4149	1150	27.7(26.4, 29.1)
	5+	759	181	23.8(20.8, 27.0)
Sex of the child	Male	7017	2019	28.8(27.7, 29.8)
	Female	6609	1838	27.8(26.7, 28.9)
Sociodemographic characteristics				
Mother's age	15–19	2028	591	29.1(28.3, 32.5)
	20–29	8166	2324	28.5(28.6, 30.6)
	30–39	3318	926	27.9(28.6,30.6)
	40–49	183	38	20.9(15.8, 28.6)
Mother's educational level	No formal education	986	243	24.6(21.9, 27.5)
	Primary incomplete(1-4years)	1570	379	24.2(22.1, 26.4)
	Primary complete (5 years)	2024	504	24.9(23.0, 26.8)
	Secondary incomplete (6–9)	6205	1829	29.5 (28.3, 30.6)
	Secondary and above (≥ 10)	2910	923	31.7 (30.0, 33.4)
* Weighted				

Variables	Categories	Overall (N = 13695)	Frequency (n = 3878)	Percentages* %(95%CI)
Mother's current occupation	Homemaker	13268	3773	28.4(27.7, 29.2)
	Skilled worker	375	92	24.7(20.3, 29.2)
	Unskilled worker	52	14	26.6(15.6, 41.0)
Mother's religion	Muslim	12912	3642	28.2(27.4, 28.9)
	Hindu /others	783	237	30.2(27.1, 33.6)
Wealth quintiles	Lowest	2657	664	25.0(23.3, 26.7)
	Second	2671	706	26.4(24.8, 28.2)
	Middle	2777	813	29.3(27.6, 31.0)
	Fourth	2758	827	29.9(28.3, 31.7)
	Highest	2832	868	30.7(30.0, 32.4)
* Weighted				

*Created by Authors

Factors associated with SSC practices

Table 3 illustrates the adjusted and unadjusted association of SSC with antepartum, intrapartum, newborn, and sociodemographic factors. Our bivariate analysis shows a significant positive association of SSC with 1–3 ANC visits at least one from an MTP, four or more ANC visits at least one from MTP during pregnancy, delivery in a public facility, the experience of complications during delivery, perceived smaller newborn birth size, mother's secondary level of educational attainment, and household wealth quintiles. On the other hand, having a cesarean section delivery, increasing birth order and mother's age between 40–49 years, were inversely associated with SSC.

Our multivariable model found a significant positive association of SSC with receiving one to three ANC visits (at least one from an MTP) (aPRR 1.14; 95%CI: 1.03, 1.26) and ≥ 4 ANC visits (at least one from an

MTP) (aPRR 1.17; 95%CI: 1.03, 1.26) compared to mothers who reported no ANC from an MTP. Compared to the women who delivered at home, delivery at a health facility increased SSC practices by two-fold for public facilities (aPRR 2.01; 95%CI: 1.80, 2.26) and 23% for private facilities (aPRR 1.23; 95%CI: 1.06, 1.42). Women who had a caesarean section were significantly less likely to practice SSC; (aPRR 0.64; 95%CI: 0.56, 0.73) compared to women who had a vaginal delivery. Newborns with mothers having complications during delivery were more likely to receive SSC than their counterparts. Babies who were perceived to be small at birth had a higher likelihood of receiving SSC compared to babies perceived to be the average size; (aPRR 1.28; 95%CI: 1.12, 1.45). Mothers having higher educational attainment were more likely to report SSC compared to women having no formal education. Newborns of mothers from the highest wealth quintile had a significantly higher likelihood of receiving SSC (aPRR 1.17; 95%CI: 1.02, 1.35) than the newborns of mothers from the lowest wealth quintile.

Table 3
Factors associated with SSC in surveyed districts of Bangladesh, 2019

Variables	Categories	Bivariate analysis		Multivariable analysis	
		uPRR ¹	95%CI	aPRR ²	95%CI
Antepartum characteristics					
Frequency of antenatal care(ANC) visits with at least one ANC from a medically trained provider (MTP)	No ANC from an MTP	ref		ref	
	1–3 ANC (at least 1 from an MTP)	1.16***	1.09, 1.24	1.14**	1.03, 1.26
	≥4 (at least 1 from an MTP)	1.23***	1.14, 1.32	1.17*	1.04, 1.30
Intrapartum characteristics					
Place of delivery	Home delivery	ref		ref	
	Public facility	1.60***	1.50, 1.71	2.01***	1.80, 2.26
	Private facility	1.02	0.95, 1.09	1.23**	1.06, 1.42
	NGO facility	0.85	0.62, 1.17	0.88	0.58, 1.33
Mode of delivery	Vaginal delivery	ref		ref	
	Caesarean section	0.88***	0.83, 0.93	0.64***	0.56, 0.73
Any complication during delivery	No	ref		ref	
	Yes	1.13***	1.06, 1.20	1.13**	1.04, 1.25
Newborn characteristics					
Birth size	Average	ref		ref	

* Pvalue < 0.05

* *Pvalue < 0.001

* **Pvalue < 0.000

¹Unadjusted Prevalence Rate Ratios

²Adjusted Prevalence Rate Ratios

		Bivariate analysis		Multivariable analysis	
	Small	1.21*	1.13, 1.32	1.28***	1.12, 1.45
	Large	0.95	0.89, 1.02	0.95	0.86, 1.04
Birth order	1–2	ref		ref	
	3–4	0.95	0.90, 1.01	0.97	0.88, 1.07
	5+	0.80*	0.70, 0.92	0.86	0.70, 1.07
Sex of the child	Female	ref			
	Male	1.04	0.98, 1.10		
Sociodemographic characteristics					
Mother's age	15–19	ref		ref	
	20–29	0.97	0.90, 1.05	1.00	0.90, 1.13
	30–39	0.96	0.88, 1.05	1.08	0.92, 1.25
	40–49	0.71*	0.53, 0.96	0.83	0.55, 1.23
Mother's educational level	No formal education	ref		ref	
	Primary incomplete(1-4years)	0.98	0.84, 1.12	0.95	0.78, 1.15
	Primary complete (5 years)	1.01	0.88, 1.16	0.97	0.80, 1.16

* Pvalue < 0.05

* *Pvalue < 0.001

* **Pvalue < 0.000

¹Unadjusted Prevalence Rate Ratios

²Adjusted Prevalence Rate Ratios

		Bivariate analysis		Multivariable analysis	
	Secondary incomplete (6–9)	1.20*	1.07, 1.35	1.18*	1.00, 1.41
	Secondary and above (≥ 10)	1.31***	1.16, 1.48	1.31**	1.09, 1.60
Mother's current occupation	Homemaker	ref			
	Skilled worker	0.89	0.74, 1.06		
	Unskilled worker	0.91	0.57, 1.47		
Mother's religion	Muslim	ref			
	Hindu /others	1.09	1.00, 1.21		
Wealth Quintiles	Lowest	ref		ref	
	Second	1.06	1.01, 1.17	1.01	0.89, 1.15
	Middle	1.18***	1.08, 1.29	1.14*	1.00, 1.29
	Fourth	1.21***	1.10, 1.32	1.14	1.00, 1.30
	Highest	1.25***	1.14, 1.36	1.17*	1.02, 1.35
* Pvalue < 0.05					
* *Pvalue < 0.001					
* **Pvalue < 0.000					
¹ Unadjusted Prevalence Rate Ratios					
² Adjusted Prevalence Rate Ratios					

Discussion

Skin-to-skin contact is one of the eight proven effective ENC interventions that improve newborn survival(9, 21). The prevalence of reported SSC practices was low at 28% in ten surveyed districts in Bangladesh. Results from our multivariable model identified two critical positive factors for improving SSC practices; i) facility delivery and ii) ANC from MTPs. In contrast, cesarean delivery significantly

reduces the practice of SSC in facilities. Researchers, program implementers, and policymakers need to consider these determinants when identifying and designing essential interventions to improve SSC practices at the population level.

Despite being an easy and no costs intervention with enormous benefits, SSC is one of the least used ENC practices in developing countries(15, 17, 22, 23). Our findings reiterate this evidence for 10 surveyed districts in Bangladesh. A similarly low prevalence of SSC practices was reported in previous studies in Bangladesh, which showed a prevalence between 26% and 30%(18, 24). Our reported prevalence of SSC is higher than the reported prevalence in India (14.5%), Nepal (16.5%), Tanzania (9.7%), and Ethiopia (24.3%). It is lower than the reported prevalence in Sri Lanka at 50% and the Gambia at 35.7% (24–28). The low prevalence of SSC practices in these developing countries could be attributed to various factors such as the high burden of home deliveries, low ANC coverage, availability of community-based interventions, and sociodemographic and health system factors(24, 25, 27–29).

One striking finding of our study is that mothers who delivered in the facilities have a higher likelihood of SSC practices compared to the mothers who delivered at home. Previous studies from Bangladesh, Gambia, and Ethiopia also found facility delivery to be a significant positive predictor of SSC practices (7, 27, 28). Although Bangladesh is observing an increasing trend in coverage of facility deliveries, 50% of women are still delivering at home(6). This finding highlights the need for increasing coverage of facility delivery as well as improving SSC practices after home deliveries in Bangladesh. Lack of awareness, social norms, distance to the facilities, lack of transport facilities, cost of care-seeking, and poor quality of care are some of the key barriers preventing women from delivering in facilities in rural areas of Bangladesh(30–35). Alliance for Maternal and Newborn Health Improvement (AMANHI) project implementing 24/7 obstetric services, training of the health care providers, community advocacy, comprehensive birth planning counseling, financial incentives for covering delivery costs, and emergency referral transport in Sylhet Division of Bangladesh, significantly improved coverage of facility deliveries from 25–78% in the intervention areas (36). Policymakers and health programmers need to consider a similar integrated package of interventions addressing both supply-side and demand-side barriers to improve coverage of facility deliveries in this setting.

On the other hand, to improve SSC practices after home deliveries, community-based interventions such as community skilled birth attendants (CSBA) and community demand generation activities may play a significant role. Community-based maternal and newborn health package piloting in Bangladesh, Malawi, Nepal, and India showed significant improvement in all four ENC practices including SSC (25, 29). The study deployed trained community health workers/ volunteers for regular home visits during and soon after delivery. Community health workers/ volunteers provided routine maternal and newborn care, promoted routine health service utilization, supported birth preparedness, conducted counseling on danger signs and essential newborn care, and identified and referred women and newborns having maternal and newborn danger signs to the nearest health facilities during regular home visits. Although community-based interventions showed some positive impact on ENC practices, considering very low coverage of SBA among home deliveries (3%) in Bangladesh and resource-intensive nature of the

community based maternal and newborn health programming, the question remains “*which investment will bring maximum impact on SSC practices in these resource-limited settings; facility delivery or home delivery or both?*” (20, 24)

Another important finding of our study is that despite having a significant positive association of facility delivery on SSC practices, less than half of the women who delivered in the facilities reported SSC. This signifies that increasing coverage of facility delivery alone will not be able to bring the change in SSC practices without improving its practices in the facilities. Shortage of skilled workforce, lack of time, difficulty in assessing eligibility for SSC, interference with routine procedures after birth, lack of motivation among the mothers and family members, and cultural practices were identified as some of the major barriers in implementing SSC after facility deliveries (37–39). WHO and UNICEF recommend implementation of the baby-friendly hospital initiatives to improve SSC and breastfeeding practices through training of the health care providers and creating an enabling hospital environment such as a standard guideline on SSC, availability of skilled workforce, and availability of sufficient postnatal beds with the provision of privacy (12, 40). The Government of Bangladesh has already started the implementation of baby-friendly hospital initiatives in public health facilities. Experience from implementation showed a significant positive impact on improving SSC and early initiation of breastfeeding practices in several other settings (41, 42). Additionally, strengthening monitoring and adoption of essential measures to address the barrier and challenges of implementing SSC in the facilities using the Plan-Do-Act-Check (PDCA) model may improve SSC practice in facilities (43–45). Save the Children’s MaMoni MNCSP project has recently started implementing the PDCA model for improving ENC practices including SSC in the public health facilities of Manikganj and Madaripur district of Bangladesh. Evaluation of this PDCA model will generate evidence on the effectiveness of this intervention in improving SSC practices in the local contexts.

Our result showed that having a cesarean delivery reduced the likelihood of receiving SSC by 36% compared to the mothers having a vaginal birth. Findings from other studies also suggested a negative impact of cesarean section on SSC practices and early initiation of breastfeeding (11, 46–50). WHO recommends starting SSC immediately (if possible in the operation theater) after cesarean delivery as soon as the mother regains consciousness in the absence of any precarious complications (12). As uncomplicated cesarean deliveries are performed using spinal anesthesia where mothers remain conscious at the time of operation, SSC can be initiated in the operating theater. Two different intervention modalities such as the Plan, Do Study, Act (PDSA) model and Practice Reflection, Education, and training, Combined with the Ethnography for Sustainable Success (PRECESS) model showed a significant improvement in SSC practices after cesarean births (51–54). The PDSA model piloted SSC in the operation theater using a flow chart and developed a monitoring and feedback mechanism to address the barriers and challenges of implementation. The PRECESS project educated the staff on standard steps of SSC implementation created a staff monitoring mechanism on SSC after cesarean births and shared videos with mothers on SSC (51, 52). One limitation of these studies is that the sample sizes are small. Contextual adaptation of these evidence-based interventions must be tested using a rigorous

implementation research design, with a sufficient sample size for generating strong evidence on the efficacy of these interventions in the local context.

Another challenge in improving SSC practices in Bangladesh is the low coverage of SSC in private and NGO facilities. High rates of cesarean deliveries in private facilities (85%) and NGO facilities (45 %) could be the reason for low SSC practices. Our national-level data also showed that 84% of women who delivered in the private facilities had a cesarean Sect. (20). The WHO threshold of optimal cesarean section rates for any country is 15% which signifies that unnecessary cesarean deliveries are happening mostly in the profit-driven private sectors(55). It's crucial to implement proven interventions including accreditation of private facilities, implementation of Robson's classification, and the second review to regulate cesarean sections in the private facilities (56–58). In essence, implementing programs to initiate SSC after cesarean sections in the operating theater along with optimization of cesarean deliveries are necessary to bring about the potential improvement in SSC practices.

Our study also showed that mothers having one to three ANC visits and four or more ANC visits (at least one from an MTP) had better reported SSC practices. Our finding is consistent with previous studies in Bangladesh, Gambia, and Ethiopia. Our finding is consistent with previous studies in Bangladesh, Gambia, and Ethiopia(7, 27, 28). ANC has the potential to improve SSC practices in two ways. First, ANC is an important predictor of facility delivery and it is evident from our findings that facility delivery improves SSC practices (59–62). Second, lack of motivation among the mother and the family members is a potential barrier to SSC practices(37, 39). Recent evidence identifies ANC counseling as a potential opportunity for motivating mothers for uptake of SSC(37).

This study has the following limitations. First, we used mothers' self-reported data for defining SSC practices. Previous studies reported mixed evidence on the validity of using mothers' reported data for defining SSC compared to observation-based data (63, 64). However, observation-based data collection is expensive and not feasible during population-based surveys. Additionally, we found that major national-level population-based surveys such as DHS in Bangladesh and Nigeria collect mothers' reported data for defining SSC and we adopted questions from BDHS for defining SSC in this study [7]. We recommend further exploration to identify appropriate wording for defining SSC during population-level surveys and to collect more in-depth information regarding the actual timing of the start and duration of SSC to improve validity. Second, recall bias may also be an issue which we tried to address by interviewing women with shorter recall periods in this survey(20).

Despite the limitations, our study is unique in terms of generating evidence on the prevalence and factors associated with SSC practices, which are not often studied at a population level (18). It is crucial to know the population-level estimates of SSC and to understand the factors influencing its practices, to better understand the barriers to its uptake, and what can be done to improve coverage of the practice in the local context.

Conclusions

There is no doubt that SSC has numerous benefits for both the mother and the baby(65). It is high time to act to improve SSC practices in the facilities and coverage of facility deliveries. Implementation of the baby-friendly hospital initiative may improve practices among the health care providers and create an enabling environment to increase SSC practices after facility deliveries. Emphasis is needed on the initiation of SSC after the cesarean section in the absence of maternal and newborn complications. It is also important to regulate unnecessary cesarean deliveries. To increase facility deliveries, it is important to address both the supply and demand-side barriers through an integrated package of interventions. Community-based interventions are important for improving coverage of SSC practices, especially in settings with a high burden of home births. ANC from MTPs could be the potential platform for not only encouraging mothers for facility deliveries but also counseling mothers on this potentially lifesaving intervention. Further research is needed to test evidence-based solutions for improving SSC practices in the local context.

Abbreviations

ANC
Antenatal Care
GLM
Generalized Linear Model
MaMoni MNCSP
MaMoni Maternal Newborn Care Strengthening Project
MTP
Medically Trained Provider
RDW
Recently Delivered Women
SSC
Skin-to-Skin Contact
UNICEF
United Nations Children’s Fund
USAID
United States Agency for International Development
WHO
World Health Organization

Declarations

Ethics approval and consent to participate:

The study received ethical approval from the Institutional Review Board of icddr,b and Save the Children US Ethical Review Committee. Our data collectors took informed written consent from the study participants (recently delivered women with a birth outcome in the past 15 months preceding survey) for

data collection. They read out the consent forms in Bangla and explained every section such as the purpose of the survey, risks, benefits, use of data, procedures for ensuring privacy and confidentiality, and the voluntary nature of participation to the study participants. Data collection began upon receiving informed written consent.

Consent for publication

Not applicable

Availability of data and materials

Due to ethical restrictions related to protecting study participants privacy and confidentiality, data access is restricted by the Ethical Review Committee of icddr,b. According to the icddr,b data policy (<http://www.icddrb.org/policies>), interested parties may contact Ms. Armana Ahmed (aahmed@icddrb.org) with further inquiries related to data access.

Competing interests

The authors declare no competing interests.

Funding

United States Agency for International Development (USAID). USAID played no role in design of the study, data collection, analysis, interpretation of data and writing of this manuscript.

Authors' contributions

This paper was designed by NBA. SKMB, SEA, BRB, and USJM conceptualized the MaMoni MNCSP project, did fund acquisition, guided protocol development and seeking ethical approval, supervised data collection and data analysis. SSP, NBA, SZ, AK, MMR, and SMR contributed to protocol development, project implementations, and data collection. BRB, SH, and MRA assisted NBA in developing the first draft. All authors reviewed and finalized the manuscript.

Acknowledgments

Save the Children's MaMoni MNCSP project was funded by the United States Agency for International Development (USAID). We acknowledge with gratitude the commitment of USAID to their research efforts in Bangladesh. We are also grateful to the Government of the People's Republic of Bangladesh; Global

Affairs Canada (GAC); Swedish International Development Cooperation Agency (Sida) and the Department for International Development, (UKAid) for providing core/unrestricted support to icddr,b. We are thankful to our study participants and our data collection and data management team for their contribution in this study.

Authors' information

¹Maternal and Child Health Division, International Centre for Diarrhoeal Disease Research, Bangladesh (icddr,b), Dhaka, Bangladesh

²Save the Children, Dhaka, Bangladesh

³PATH, Washington, DC., USA (formerly Save the Children)

⁴The University of Sydney School of Public Health, Sydney, NSW 2006, Australia

References

1. Hug L, Alexander M, You D, Alkema L, Estimation UNI-aGfCM. National, regional, and global levels and trends in neonatal mortality between 1990 and 2017, with scenario-based projections to 2030: a systematic analysis. *Lancet Glob Health*. 2019;7(6):e710-e20.
2. WHO WHO. Global Health Observatory (GHO) data 1990, 2017 [Available from: https://www.who.int/gho/child_health/mortality/neonatal_infant_text/en/].
3. WHO WHO. Success factors for women's and children's health: Rwanda. 2015.
4. Unicef. Millenium Development Goals. 2014.
5. Victora CG, Requejo JH, Barros AJ, Berman P, Bhutta Z, Boerma T, et al. Countdown to 2015: a decade of tracking progress for maternal, newborn, and child survival. *Lancet* (London, England). 2016;387(10032):2049-59.
6. National Institute of Population Research and Training (NIPORT), Mitra and Associates (MA), International I. Bangladesh Demographic and Health Survey 2017. Dhaka, Bangladesh, and Rockville, Maryland, USA: NIPORT, Mitra and Associates, and ICF International.; 2019.
7. Singh K, Khan SM, Carvajal-Aguirre L, Brodish P, Amouzou A, Moran A. The importance of skin-to-skin contact for early initiation of breastfeeding in Nigeria and Bangladesh. *Journal of global health*. 2017;7(2).
8. Darmstadt GL, Kinney MV, Chopra M, Cousens S, Kak L, Paul VK, et al. Who has been caring for the baby? *The Lancet*. 2014;384(9938):174-88.
9. Khadivzadeh T, Karimi A. The effects of post-birth mother-infant skin to skin contact on first breastfeeding. 2009.

10. Cadwell K, Brimdyr K, Phillips R. Mapping, Measuring, and Analyzing the Process of Skin-to-Skin Contact and Early Breastfeeding in the First Hour After Birth. *Breastfeed Med.* 2018;13(7):485-92.
11. Stevens J, Schmied V, Burns E, Dahlen H. Immediate or early skin-to-skin contact after a Caesarean section: a review of the literature. *Matern Child Nutr.* 2014;10(4):456-73.
12. WHO. Protecting, Promoting and Supporting Breastfeeding in Facilities Providing Maternity and Newborn Services 2017 WHO2017 [Available from: <http://apps.who.int/iris/bitstream/>].
13. Moore ER, Anderson GC, Bergman N, Dowswell T. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev.* 2012;5(5):CD003519-CD.
14. Moore ER, Anderson GC, Bergman N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev.* 2007(3):Cd003519.
15. Moore ER, Bergman N, Anderson GC, Medley N. Early skin-to-skin contact for mothers and their healthy newborn infants. *Cochrane Database Syst Rev.* 2016;11(11):CD003519-CD.
16. Abdulghani N, Edvardsson K, Amir LH. Worldwide prevalence of mother-infant skin-to-skin contact after vaginal birth: A systematic review. *PLoS One.* 2018;13(10):e0205696-e.
17. Abdulghani N, Edvardsson K, Amir LH. Worldwide prevalence of mother-infant skin-to-skin contact after vaginal birth: A systematic review. *PloS one.* 2018;13(10).
18. Singh K, Khan SM, Carvajal-Aguirre L, Brodish P, Amouzou A, Moran A. The importance of skin-to-skin contact for early initiation of breastfeeding in Nigeria and Bangladesh. *Journal of global health.* 2017;7(2):020505-.
19. UN. Sustainable Development Goals: United Nations; 2015 [Available from: <http://www.un.org/sustainabledevelopment/health/>].
20. National Institute of Population Research and Training (Bangladesh), Mitra and Associates (Firm), MEASURE DHS (Program). Bangladesh demographic and health survey, 2017 : key indicators. Dhaka: National Institute of Population Research and Training, Ministry of Health and Family Welfare; 2019. vii, 56 pages p.
21. Barros FC, Bhutta ZA, Batra M, Hansen TN, Victora CG, Rubens CE. Global report on preterm birth and stillbirth (3 of 7): evidence for effectiveness of interventions. *BMC Pregnancy Childbirth.* 2010;10 Suppl 1(Suppl 1):S3.
22. de Graft-Johnson J, Vesel L, Rosen HE, Rawlins B, Abwao S, Mazia G, et al. Cross-sectional observational assessment of quality of newborn care immediately after birth in health facilities across six sub-Saharan African countries. *BMJ open.* 2017;7(3):e014680-e.
23. de Jonge E, Azad K, Hossen M, Kuddus A, Manandhar DS, van de Poel E, et al. Socioeconomic inequalities in newborn care during facility and home deliveries: a cross sectional analysis of data from demographic surveillance sites in rural Bangladesh, India and Nepal. *Int J Equity Health.* 2018;17(1):119-.
24. Crowe S, Prost A, Hossen M, Azad K, Kuddus A, Roy S, et al. Generating insights from trends in newborn care practices from Prospective Population-Based Studies: examples from India, Bangladesh and Nepal. 2015;10(7):e0127893.

25. Upadhyay RP, Rai SK, Anand KJAP. Community neonatal practices and its association with skilled birth attendance in rural Haryana, India. 2012;101(12):e535-e9.
26. Cederfeldt J, Carlsson J, Begley C, Berg MJS, Healthcare R. Quality of intra-partum care at a university hospital in Nepal: A prospective cross-sectional survey. 2016;7:52-7.
27. Nigatu D, Abeje G, Mekonnen AG, Azage M, Bogale D. Maternal Health Service Uptake Is Associated with a Higher Skin-to-Skin Care Practice in Ethiopia: Result from a National Survey. *Biomed Res Int*. 2020;2020:8841349-.
28. Ekholuenetale M, Onikan A, Ekholuenetale CEJJotEPHA. Prevalence and determinants of mother and newborn skin-to-skin contact in The Gambia: a secondary data analysis. 2020;95(1):1-9.
29. Sitrin D, Guenther T, Waiswa P, Namutamba S, Namazzi G, Sharma S, et al. Improving newborn care practices through home visits: lessons from Malawi, Nepal, Bangladesh, and Uganda. *Global health action*. 2015;8:23963-.
30. Bohren MA, Hunter EC, Munthe-Kaas HM, Souza JP, Vogel JP, Gülmezoglu AM. Facilitators and barriers to facility-based delivery in low- and middle-income countries: a qualitative evidence synthesis. *Reprod Health*. 2014;11(1):71.
31. Borghi J, Ensor T, Somanathan A, Lissner C, Mills A. Mobilising financial resources for maternal health. *Lancet (London, England)*. 2006;368(9545):1457-65.
32. Nahar S, Costello A. The hidden cost of 'free' maternity care in Dhaka, Bangladesh. *Health Policy Plan*. 1998;13(4):417-22.
33. Nguyen HT, Hatt L, Islam M, Sloan NL, Chowdhury J, Schmidt JO, et al. Encouraging maternal health service utilization: an evaluation of the Bangladesh voucher program. *Social science & medicine (1982)*. 2012;74(7):989-96.
34. Saksena P, Xu K, Elovainio R, Perrot J. Utilization and expenditure at public and private facilities in 39 low-income countries. *Tropical medicine & international health : TM & IH*. 2012;17(1):23-35.
35. Sarker BK, Rahman M, Rahman T, Hossain J, Reichenbach L, Mitra DK. Reasons for Preference of Home Delivery with Traditional Birth Attendants (TBAs) in Rural Bangladesh: A Qualitative Exploration. *PloS one*. 2016;11(1):e0146161.
36. Rahman S, Choudhury AA, Khanam R, Moin SMI, Ahmed S, Begum N, et al. Effect of a package of integrated demand- and supply-side interventions on facility delivery rates in rural Bangladesh: Implications for large-scale programs. *PLoS One*. 2017;12(10):e0186182-e.
37. Alenchery AJ, Thoppil J, Britto CD, de Onis JV, Fernandez L, Suman Rao PN. Barriers and enablers to skin-to-skin contact at birth in healthy neonates - a qualitative study. *BMC pediatrics*. 2018;18(1):48-.
38. Chan G, Bergelson I, Smith ER, Skotnes T, Wall S. Barriers and enablers of kangaroo mother care implementation from a health systems perspective: a systematic review. *Health Policy Plan*. 2017;32(10):1466-75.
39. Abdulghani N, Edvardsson K, Amir LH. Health care providers' perception of facilitators and barriers for the practice of skin-to-skin contact in Saudi Arabia: A qualitative study. *Midwifery*. 2020;81:102577.

40. WHO. Baby-friendly hospital initiative: revised, updated and expanded for integrated care. 2009.
41. Philipp BL. The Importance of the Baby-Friendly Hospital Initiative. *JAMA pediatrics*. 2017;171(3):305-.
42. Rowe-Murray HJ, Fisher JR. Baby friendly hospital practices: cesarean section is a persistent barrier to early initiation of breastfeeding. *Birth*. 2002;29(2):124-31.
43. Saxena S, Ramer L, Shulman IA. A comprehensive assessment program to improve blood-administering practices using the FOCUS–PDCA model. *Transfusion*. 2004;44(9):1350-6.
44. Redick EL. Applying FOCUS-PDCA to solve clinical problems. *Dimensions of Critical Care Nursing*. 1999;18(6):30.
45. Farahbakhsh M, Tabrizi J, NIKNIAZ A. The use of Focus-PDCA in primary health care performance improvement: case study of East Azerbaijan health centers. 2010.
46. Brady K, Bulpitt D, Chiarelli C. An interprofessional quality improvement project to implement maternal/infant skin-to-skin contact during cesarean delivery. *Journal of Obstetric, Gynecologic, & Neonatal Nursing*. 2014;43(4):488-96.
47. Wu Y, Wang Y, Huang J, Zhang Z, Wang J, Zhou L, et al. The association between caesarean delivery and the initiation and duration of breastfeeding: a prospective cohort study in China. *Eur J Clin Nutr*. 2018;72(12):1644-54.
48. Karim F, Billah SM, Chowdhury MAK, Zaka N, Manu A, El Arifeen S, et al. Initiation of breastfeeding within one hour of birth and its determinants among normal vaginal deliveries at primary and secondary health facilities in Bangladesh: A case-observation study. *PLoS One*. 2018;13(8).
49. Islam MA, Mamun A, Hossain MM, Bharati P, Saw A, Lestrel PE, et al. Prevalence and factors associated with early initiation of breastfeeding among Bangladeshi mothers: A nationwide cross-sectional study. *PLoS One*. 2019;14(4):e0215733.
50. Hobbs AJ, Mannion CA, McDonald SW, Brockway M, Tough SC. The impact of caesarean section on breastfeeding initiation, duration and difficulties in the first four months postpartum. *BMC Pregnancy Childbirth*. 2016;16:90.
51. Hung KJ, Berg OJMTAJoMCN. Early skin-to-skin after cesarean to improve breastfeeding. 2011;36(5):318-24.
52. Crenshaw JT, Cadwell K, Brimdyr K, Widström A-M, Svensson K, Champion JD, et al. Use of a video-ethnographic intervention (PRECESS Immersion Method) to improve skin-to-skin care and breastfeeding rates. 2012;7(2):69-78.
53. Nolan A, Lawrence CJJoO, Gynecologic, Nursing N. A pilot study of a nursing intervention protocol to minimize maternal-infant separation after cesarean birth. 2009;38(4):430-42.
54. Gouchon S, Gregori D, Picotto A, Patrucco G, Nangeroni M, Di Giulio PJNr. Skin-to-skin contact after cesarean delivery: an experimental study. 2010;59(2):78-84.
55. Haider MR, Rahman MM, Moinuddin M, Rahman AE, Ahmed S, Khan MM. Ever-increasing Caesarean section and its economic burden in Bangladesh. *PLoS One*. 2018;13(12):e0208623.

56. Betrán AP, Temmerman M, Kingdon C, Mohiddin A, Opiyo N, Torloni MR, et al. Interventions to reduce unnecessary caesarean sections in healthy women and babies. *The Lancet*. 2018;392(10155):1358-68.
57. Begum T, Nababan H, Rahman A, Islam MR, Adams A, Anwar I. Monitoring caesarean births using the Robson ten group classification system: A cross-sectional survey of private for-profit facilities in urban Bangladesh. *PLoS One*. 2019;14(8):e0220693.
58. Begum T, Rahman A, Nababan H, Hoque DME, Khan AF, Ali T, et al. Indications and determinants of caesarean section delivery: Evidence from a population-based study in Matlab, Bangladesh. *PLoS One*. 2017;12(11):e0188074.
59. Shahabuddin ASM, Delvaux T, Utz B, Bardají A, De Brouwere V. Determinants and trends in health facility-based deliveries and caesarean sections among married adolescent girls in Bangladesh. *BMJ open*. 2016;6(9):e012424-e.
60. Kamal SM. Preference for institutional delivery and caesarean sections in Bangladesh. *J Health Popul Nutr*. 2013;31(1):96-109.
61. Rahman M. Deliveries among adolescent mothers in rural Bangladesh: who provides assistance? *World health & population*. 2009;11(2):5-14.
62. Anwar I, Sami M, Akhtar N, Chowdhury ME, Salma U, Rahman M, et al. Inequity in maternal health-care services: evidence from home-based skilled-birth-attendant programmes in Bangladesh. *Bull World Health Organ*. 2008;86(4):252-9.
63. Blanc AK, Warren C, McCarthy KJ, Kimani J, Ndwiga C, RamaRao S. Assessing the validity of indicators of the quality of maternal and newborn health care in Kenya. *Journal of global health*. 2016;6(1):010405-.
64. Stanton CK, Rawlins B, Drake M, Dos Anjos M, Cantor D, Chongo L, et al. Measuring coverage in MNCH: testing the validity of women's self-report of key maternal and newborn health interventions during the peripartum period in Mozambique. *PLoS One*. 2013;8(5):e60694-e.
65. Safari K, Saeed AA, Hasan SS, Moghaddam-Banaem L. The effect of mother and newborn early skin-to-skin contact on initiation of breastfeeding, newborn temperature and duration of third stage of labor. *Int Breastfeed J*. 2018;13:32-.

Figures

Modes of Delivery by Places of Delivery




Figure 1

Distribution of modes of delivery by places of delivery in surveyed districts of Bangladesh, 2019*.

Reported Skin-to-Skin Practices by Modes and Places of Delivery




Figure 2

Distribution of SSC by modes and places of delivery in surveyed districts of Bangladesh, 2019*