Validation of a QTL for grain size and weight using an introgression line from a cross between *Oryza sativa* and *Oryza minuta*

Yue Feng

China National Rice Research Institute

Xiaoping Yuan

China National Rice Research Institute

Yiping Wang

China National Rice Research Institute

Yaolong Yang

China National Rice Research Institute

Mengchen Zhang

China National Rice Research Institute

Hanyong Yu

China National Rice Research Institute

Qun Xu

China National Rice Research Institute

Shan Wang

China National Rice Research Institute

Xiaojun Niu

China National Rice Research Institute

Xinghua Wei (✉ weixinghua@caas.cn)

<https://orcid.org/0000-0002-0883-3694>

Original article

Keywords: *Oryza sativa*, *Oryza minuta*, Introgressive line, Grain size and weight, Quantitative trait loci

Posted Date: January 4th, 2021

DOI: <https://doi.org/10.21203/rs.3.rs-54218/v2>

License: © ⓘ This work is licensed under a Creative Commons Attribution 4.0 International License. [Read Full License](#)

Version of Record: A version of this preprint was published at Rice on May 20th, 2021. See the published version at <https://doi.org/10.1186/s12284-021-00472-1>.

Abstract

Background: Grain size and weight are important target traits determining grain yield and quality in rice. Wild rice species possess substantial elite genes that can be served as an important resource for genetic improvement of rice. In this study, we identify and validate a novel QTL on chromosome 7 affecting the grain size and weight using introgression lines from cross of *Oryza sativa* and *Oryza minuta*.

Results: An introgression line 'IL188' has been achieved from a wild species *Oryza minuta* (2n = 48, BBCC, W303) into *O. sativa japonica* Nipponbare. The F₂ and F₂:3 populations derived from a cross between IL188 and Nipponbare were used to map QTLs for five grain size traits, including grain length (GL), grain width (GW), grain length to width ratio (LWR), grain thickness (GT) and thousand grain weight (TGW). A total of 12 QTLs for the five grain traits were identified on chromosomes 1, 2, 3, 6, 7, and 8. The QTL-*qGL7* controlling GL on chromosome 7 was detected stably in the F₂ and F₂:3 populations, and explained 15.09-16.30% of the phenotypic variance. To validate the effect of *qGL7*, eight residual heterozygous line (RHL) populations were developed through selfing four F₂:3 and four F₂:4 plants with different heterozygous segments for the target region. By further developing SSR and Indel markers in the target interval, *qGL7* was delimited to a ~261 kb region between Indel marker Y7-12 and SSR marker Y7-38, which also showed significant effects on grain width and thousand grain weight. Comparing with the reference genome of Nipponbare, stop or frameshift mutations in the exon of the three putative genes *LOC_Os07g36830*, *LOC_Os07g36900* and *LOC_Os07g36910* encoding F-box domain-containing proteins may be the candidate genes for *qGL7*. Scanning electron microscopy analysis of the glume's epidermal cells showed that the cell length and width of NIL-*qGL7*IL188 was higher than NIL-*qGL7*Nip, indicating that *qGL7* increases grain size and weight by regulating cell expansion.

Conclusions: In this study, we detected 12 QTLs regulating grain size and weight using an introgression line from a cross between an *Oryza sativa* and *Oryza minuta*. Of these loci, we confirmed and delimited the *qGL7* to a ~261 kb region. Three putative genes, *LOC_Os07g36830*, *LOC_Os07g36900* and *LOC_Os07g36910* encoding F-box domain-containing proteins may be the candidate genes for *qGL7*. These results provide a basis for map-based cloning of the *qGL7* gene and useful information for marker assisted selection in rice grain quality improvement.

Background

Rice (*Oryza sativa* L.) is one of the most important cereal crops in Asia and is the main staple food for the majority of peoples in the world. Breeding of high-yielding rice is crucial for meeting the food demand of the increasing world population (Ikeda et al. 2013). Grain yield in rice is determined by three major components: the number of panicles, the number of grains per panicle, and grain weight (Huang et al. 2013). Among these, the most reliable trait is grain weight, which is largely determined by grain size, which is specified by its three dimensions (length, width, and thickness) and the degree of filling (Xiong and Zhang 2010).

Grain size and weight are important components determining rice grain yield, and they are controlled by multiple quantitative trait loci (QTLs) (Zhang et al. 2012; Yu et al. 2018). To date, over 400 QTLs

modulating grain size and weight have been identified and are distributed on each of rice's 12 chromosomes (Huang et al. 2013; Zuo and Li 2014; Kashifet al. 2020). However, only a few major QTLs including *GS3*, *qSW5*, *GW2*, *qGL3/GL3.1*, *GW8*, *GL7/GW7*, *TGW6* and *GS9* have been isolated by map-based cloning methods (Mao et al. 2010; Shomura et al. 2008; Song et al. 2007; Qi et al. 2012; Zhang et al. 2012; Wang et al. 2012; Wang et al. 2015a; Wang et al. 2015b; Ishimaru et al. 2013; Zhao et al. 2018). The isolation of these genes has enhanced our knowledge of the molecular regulatory mechanisms responsible for grain size and weight (Song et al. 2008).

Oryza minuta (2n = 48, BBCC) is an allotetraploid wild species, which is endemic to Philippines and Papua New Guinea. This species belongs to the *Oryza officinalis* complex and harbors useful genes for resistance to blast blight, bacterial blight, brown planthopper and sheath blight (Amante et al. 1992; Brar and Khush 1997). However, low crossability and limited recombination between unrelated genomes limit the transfer of QTLs from *Oryza minuta* to cultivars. Following the availability of advanced backcross quantitative trait loci (AB-QTL) approach proposed by Tanksley and Nelson (1996), several studies have been reported to identify the QTLs controlling yield and quality-related traits and to simultaneously transfer them from wild to cultivated species (Xiao et al. 1998; Thomson et al. 2003; Yoon et al. 2006; Tian et al. 2006; Mallikarjuna Swamy et al. 2012; Yun et al. 2016). However, few attempts have been made to identify and capture yield-related QTLs from *Oryza minuta* into cultivars.

In the present study, we used an advanced across line IL188 from a cross between the japonica variety, Nipponbare, *Oryza sativa* and a wild accession, W303, *Oryza minuta*, as the donor parent, to map QTLs for rice grain size traits. The objectives of this study were: (1) to reliably identify novel genomic regions associated with grain size traits from W303 (*Oryza minuta*), (2) to evaluate the effects of introgressive segments on grain size traits, (3) to fine mapping the QTL-*qGL7* and validate the effects of *qGL7* for rice grain size and weight on chromosome 7.

Results

Genetic background of IL188

A total of 512 SSR markers were screened for polymorphism between W303 and Nipponbare. Among them, 185 markers produced polymorphic bands between the parents. These 185 polymorphic markers were further used to assay the genotype of IL188. Thirty of these markers (16.2%) showed W303 genotype, which covered 11 regions distributed on six chromosomes. The introgressed segments distributed on chromosomes 1, 2 (two), 3, 5, 6 (two), 7 (two), and 8 (two), respectively (Fig. 3). These 30 markers were further used to genotype the F₂ and F₂:3 populations derived from a cross between Nipponbare and IL188.

Trait performance of the parents, F₂ and F₂:3 populations

The phenotypic values of the two parents of five agronomic traits, including GL, GW, GT, LWR, and TGW were shown in Table 1. Compared with Nipponbare, IL188 had higher values for GL, GW, LWR, and TGW but lower values for GT. The frequency distributions of the five grain size traits in the F₂ and F₂:3 populations displayed a continuous variation (Fig. 4). All these traits except TGW showed two-way

transgressive segregation and followed a near normal distribution in the both populations. The results fulfill the requirement of QTL mapping.

Correlation analysis of five grain traits

The correlation coefficients among the five grain traits in the F2 and F2:3 populations were shown in Table 2. The correlation coefficients ranged from -0.677 to 0.614 and -0.662 to 0.598, respectively, in the F2 and F2:3 populations. Significant correlation was observed for each pair-wise combination except that between GL and GT, LWR and TGW. In the both populations, GL showed positive correlation with GW, LWR and TGW, GW displayed strong positive correlation with GT and TGW, while LWR showed negative correlation with GW and GT.

QTLs for grain size traits in the F2 and F2:3 populations

A total of 12 QTLs for five grain size traits were detected on chromosomes 1, 2, 3, 6, 7, and 8 in the F2 and F2:3 populations (Table 3). The phenotypic variance explained by each QTL ranged from 4.72% to 16.30%. Four of these regions were found to affect two traits. The RM7341–RM128 interval on the chromosome 1 and RM12924–RM5812 interval on chromosome 2 showed consistent effects on GL and LWR in both populations. In the RM7341–RM128 interval, *qGL1* and *qLWR1* explained phenotypic variances by 8.77% and 7.55% in the F2 population, and 9.13% and 7.40% in the F2:3 population, respectively. In the RM12924–RM5812 interval, *qGL2* and *qLWR2* explained phenotypic variances by 6.90% and 5.02% in the F2 population, and 7.09% and 5.68% in the F2:3 population, respectively. The enhancing alleles of these QTLs all derived from IL188. The RM500–RM429 interval on chromosome 7 showed consistent effects on GL and TGW in the both populations. The *qGL7* and *qTGW7* explained phenotypic variances by 16.30% and 9.97% in the F2 population, and 15.09% and 6.65% in the F2:3 population, respectively. The enhancing alleles of the two QTLs also derived from IL188. The RM3845–RM6948 on chromosome 8 exhibited significant effects on GT and LWR only in the F2:3 population. The *qGT8* and *qLWR8* explained 7.41% and 4.72% of phenotypic variances, with enhancing alleles derived from Nipponbare and IL188, respectively. The other four regions, which covered RM6307–RM5807, RM3199–RM3684, RM7158–RM276, RM408–RM3702 on chromosome 2, 3, 6 and 8, respectively, were each detected for a single trait, with *R*² ranging from 4.85% to 7.06%.

Among these regions, the RM500–RM429 interval on chromosome 7 showed the largest effect for GL and relatively stable QTLs for TGW. Therefore, the region was chosen for further validation. For ease of description, the *qGL7* and *qTGW7* detected in this region were integrated as *qGL7*.

Substitution mapping of *qGL7* and sequence analysis of candidate genes

Four NIL-F2 populations carrying heterozygous segments overlapped in the RM500–RM429 interval were constructed, including R1, R2, R3 and R4. Significant genotypic effects were detected for the three grain size traits in R2 and R3. In the two populations, the additive effects were 0.115 and 0.109 for GL, 0.065 and 0.050 for GW, 0.621 and 0.907 for TGW, explained phenotypic variances by 19.54% and 15.46%, 21.65% and 10.24%, and 16.09% and 16.86% (Table 4). The enhancing allele was derived from IL188, the same as what was found in the F2 and F2:3 populations. The additive effects and *R*² were similar between R2 and R3, indicated that *qGL7* located in the common segregating regions of the two populations. In

R1 and R4, no significant effect was detected for any traits, indicated that *qGL7* located outside of segregating regions of the two populations. As shown in Fig. 5, this is an interval flanked by markers Y7-3 and Y7-4, corresponding to a 725-kb region in the Nipponbare genome (Fig. 5).

Following the update target regions, other four NIL-F2 populations were developed, including R5, R6, R7 and R8. Significant genotypic effects were detected in R6 and R8, but not in R5 and R7. In R6 and R8, the additive effects were 0.129 and 0.074 for GL, 0.026 and 0.031 for GW, 0.595 and 0.494 for TGW, explained phenotypic variances by 48.52% and 18.12%, 16.44% and 12.77%, and 25.38% and 9.92% (Table 4). Again, the enhancing allele was derived from IL188. These results indicated that *qGL7* was located within the common segregating regions of R6 and R8 but outside the segregating region of R5 and R7. Consequently, *qGL7* was delimited into a 261-kb region flanked by Y7-12 and Y7-38 (Fig. 5).

In the 261-kb genomic region of the Nipponbare genome, a total of thirty-seven putative genes were predicted based on the Nipponbare sequence (Os-Nipponbare-Reference-IRGSP-1.0). Whole-genome resequencing was performed on the NIL-*qGL7*/IL188. Compared with the Nipponbare reference genome, seven stop code mutations and ten frameshift mutations were identified in the exonic region of the twelve genes (Table S3). One of them, *LOC_Os07g36850* encodes a putative transposon protein, four genes encode retrotransposon proteins and four genes encode expressed protein of unknown function. In addition, *LOC_Os07g36900* encodes a protein containing F-box and LRR motifs, both *LOC_Os07g36830* and *LOC_Os07g36910* encode [F-box proteins](#).

Histocytological Analysis

The homozygous plants were selected from R6 and R8. They were selfed to develop two NIL populations. The effect of *qGL7* was further validated using these two populations. Compared with NIL-*qGL7*/Nip, the GL and GW in the NIL-*qGL7*/IL188 were significantly larger (Fig. 6A-C), thus resulting in a larger TGW (Fig. 6D). These indicated that *qGL7* had stable effects on grain size traits.

In addition, we examined the cell length and width of epidermal cells of the outer and inner spikelet hulls of NIL-*qGL7*/Nip and NIL-*qGL7*/IL188 by scanning electron microscopy. Both the length and width of epidermal cells of the outer and inner spikelet hulls were increased in NIL-*qGL7*/IL188 compared with those in NIL-*qGL7*/Nip (Fig. 6E-L). Additionally, we investigated the cell number in the vertical and lateral direction of the outer spikelet hulls of the NIL-*qGL7*/Nip and NIL-*qGL7*/IL188 by scanning electron microscopy. No significant difference in total cell number in the longitudinal and lateral direction of spikelet hulls was observed between NIL-*qGL7*/Nip and NIL-*qGL7*/IL188 (Figure S1B-C). These results indicate that *qGL7* regulates grain size by promoting cell expansion.

Discussion

Common wild rice is the wild ancestor of cultivated rice (Second, 1982; Oka, 1988; Wang et al. 1992). As the ancestor of cultivated rice, wild rice has been well recognized as an extremely important resource for rice improvement, since it carries many beneficial agronomic traits which have been lost in the cultivated rice through natural and human selection (Sun et al. 2001; Sakai and Itoh 2010). In the present study, an advanced backcross line, IL188, was developed from a cross between Nipponbare and *O. minuta*. The F2 and

F2:3 populations derived from a cross between Nipponbare and IL188 was used to identify the QTLs controlling grain size and grain weight. A total of 12 putative QTLs for grain size and grain weight were detected in the F2 and F2:3 populations, and 9 of which were commonly detected in both populations.

A comparison of the QTL regions from this study with those seen in previous rice linkagemaps (<http://www.gramene.org>) revealed that six regions were shared across studies. For GL and LWR, one QTL was detected in the interval RM7341–RM128 on chromosome 1. Wan et al. (2005) also detected a stable QTL for the same traits in the similar regions on chromosome 1, and Qi et al. (2017) detected a QTL for LWR closely linked with the marker RM128. One locus associated with GL and LWR (*qGL2*, and *qLWR2*) were located in the interval RM12924–RM5812 on chromosome 2. Interestingly, Yoon et al. (2006) confirmed a locus associated with TGW, GW, GT, and LWR in the same region using an advanced backcross population between *O. Grandiglumis* and *O. sativa*. Two QTLs for GW and GT were respectively located in the vicinity of QTLs detected in previous studies (Yoon et al. 2006; MallikarjunaSwamy et al. 2012; Qi et al. 2017). In our study, one QTL was detected for TGW in the interval RM6307–RM5807 on chromosome 2, and Xue et al. (2019) also detected a QTL for TGW in the nearby region.

More importantly, one major QTL, *qGL7*, was detected for GL and TGW in the interval RM500–RM429 on chromosome 7 in the F2 and F2:3 populations. The *O. minuta* introgressive line alleles could increase GL and TGW. The *qGL7* could explain 15.09–16.30% and 6.65–9.97% of the phenotypic variation for GL and TGW, respectively. The *O. minuta* allele at locus *qGL7* increased GL and TGW by an average of 0.19 mm and 0.65 g, respectively. Interestingly, Rahman et al. (2007) also detected a QTL for GL and TGW in the same region using an F2:3 population between *O. minuta* introgression line and *O. sativa*. This result indicated that there really exist a stable QTL controlling grain size and grain weight, and the *O. minuta* allele could positively regulate grain size and grain weight. As we known, some major-effect QTLs for grain size and weight on chromosome 7 have been fine mapped and cloned in previous studies. Bai et al. (2010) detected a pleiotropic QTL for grain size and this QTL *qGL7* was narrowed down to within a 258-kb region. Shao et al. and Qiu et al. identified a major QTL *GS7/qSS7* on the long arm of chromosome 7 for grain size. Subsequently, This *GL7/GW7* (the same as *GS7/qSS7*) gene has been cloned by Wang et al. (2015a) and Wang et al. (2015b). They have found that copy number variations at *GL7/GW7* locus cause elevated expression of *GL7* and thus an increase in grain length. The grain size gene *GLW7*, encoding the plant-specific transcription factor OsSPL13, has been isolated and functionally characterized using GWAS approach (Si et al. 2016). Xu et al. (2015) identified a dominant big grain mutant BG2 that encoded a cytochrome P450, OsCYP78A13 on chromosome 7. Here, we have defined the locus *qGL7* to a 261 kb region on the long arm of chromosome 7. By comparing the physical location of *qGL7* with the reported grain size QTLs on chromosome 7, we found that *qGL7* is a novel QTL for regulating rice grain size.

Compared with the Nipponbare reference genome, seven stop code mutations and ten frameshift mutations were identified in the exonic region of the twelve genes of NIL-*qGL7* IL188 in the 261-kb region. Three putative genes of them, *LOC_Os07g36830*, *LOC_Os07g36900* and *LOC_Os07g36910* encoded F-box domain-containing proteins. F-box proteins are the substrate-recognition components of SCF (SKP1-Cul1-F-box) type E3 ubiquitin protein ligases (Skowyra et al. 1997; Feldman et al. 1997), which participate in the regulation of many physiological processes and play a key role in cell division, signal transduction, development and metabolism

(Patton et al. 1998). In rice, *GW2* encodes a RING-type protein with E3 ubiquitin ligase activity, which is known to function in the degradation by the ubiquitin-proteasome pathway (Song et al. 2007). In addition, Chen et al. 2013 reported that overexpression of *OsFBK12* (encoding an F-box protein) could increase grain size in rice. Therefore, we suppose that the three putative genes, *LOC_Os07g36830*, *LOC_Os07g36900* and *LOC_Os07g36910* encoded F-box domain-containing proteins may be the candidate genes for *qGL7*. In future, transgenic studies will be carried out for the three F-box domain-containing genes identified in the *qGL7* locus to further elucidate the molecular mechanism of *qGL7* involving in regulation of rice grain size.

Classical quantitative genetics assumes that trait correlations are the result of either pleiotropic effects or the tight linkage of genes (Wan et al. 2005). In this study, *qGL1/qLWR1* and *qGL2/qLWR2* were mapped in the same interval on chromosome 1 and chromosome 2, respectively, and the positive alleles were all derived from *O. minuta*. As well, *qGL7* and *qTGW7* shared the same confidence interval on chromosome 7 and their effect acted in the same direction. Co-localization of these QTLs, as the result of either pleiotropic effects or close linkage, could provide an explanation for the genetic basis of high trait correlations, which ranged from 0.488 between GL and LWR to 0.614 between GL and TGW.

Transfer and utilization of useful genes from wild rice into cultivated varieties are effective and aim to improve grain yield, quality, and crop genetic diversity (Brar and Khush 1997; Xie et al. 2008; Yun et al. 2016; Qi et al. 2017). However, efforts to improve rice grain traits of modern cultivars using *O. minuta* as donor parents are limited. In the present study, *O. minuta* alleles increase rice grain traits in the Nipponbare background at most QTLs, revealing the possibility that *O. minuta* alleles could improve grain traits. Although a number of genes/QTLs involved in the regulation of grain size have been cloned in rice, the molecular mechanisms of how grain size is regulated remain unknown. In this study, we found that *qGL7* could increase both grain length and grain weight, and the isolation of *qGL7* will be beneficial in better understanding of the regulation mechanism of grain size in rice. In addition, our continuous work will be helpful in improving rice yield and quality by molecular design breeding.

Conclusions

An introgression line IL188 was identified, which exhibited increased grain size and weight. A total of 12 QTLs for five grain traits were detected using F2 and F2:3 populations derived from crosses between IL188 and Nipponbare. One of the QTLs, *qGL7* was delimited to a ~261 kb region on the long arm of chromosome 7, and three putative genes, *LOC_Os07g36830*, *LOC_Os07g36900* and *LOC_Os07g36910* encoding F-box domain-containing proteins may be the candidate genes for *qGL7*. The *qGL7* increases grain size and weight by regulating cell expansion. These results will be helpful not only for understanding the genetic basis of grain size traits, but also simultaneously improving grain size and weight through marker-assisted selection (MAS) in rice breeding programs.

Materials And Methods

Plant materials

The introgression line, IL188, derived from an interspecific cross between *Oryza sativa* japonica Nipponbare and a wild species *Oryza minuta* W303 collected from the Germplasm Resource Center of IRRI, followed by three backcrosses with Nipponbare and aided by embryo rescue and subsequently self-pollinated for four generations. IL188 showed significantly longer grain length and higher grain weight than the recurrent parent Nipponbare (Fig. 1). To elucidate the genetic basis of the grain size and weight variation, an F₂ population consisting of 166 individuals was constructed by selfing the F₁ between the female parent IL188 and male parent Nipponbare, and the F₂:3 population was derived from the selfed seeds of the F₂ plants.

Following the initial outcome of QTL analysis, four residual heterozygous plants were selected from the F₂:3 populations, carrying sequential heterozygous segments covering the interval RM500–RM429. They were selfed, and four NIL-F₂ populations were constructed. They contained 180, 184, 184 and 195 plants and were named as R1, R2, R3 and R4, which carried overlapping heterozygous segments in the interval RM11–RM1135, RM11–Y7-2, RM11–Y7-2 and RM11–Y7-4, respectively (Fig. 2). The genomic background of the R1, R2, R3 and R4 populations was shown in Table S1. The R1, R2, R3 and R4 populations which were respectively only heterozygous in the target interval and basically homozygous in the genomic background were used for the substitution mapping of *qGL7*.

Four other plants were further selected from the R3 population carrying sequential heterozygous segments covering the interval Y7-3–Y7-4. They were selfed, and four NIL-F₂ populations were constructed. They contained 130, 144, 146 and 140 plants and were named as R5, R6, R7 and R8, which carried overlapping heterozygous segments in the interval Y7-4–RM21787, Y7-4–RM455, RM21787–Y7-12 and RM21787–Y7-13, respectively (Fig. 2). The R5, R6, R7 and R8 populations were used for further substitution mapping of *qGL7*. Non-recombinant homozygous plants were further identified in the R6 and R8 populations and selfed. Two sets of NILs were developed, each consisting of 20 IL188 homozygous lines and 20 Nipponbare homozygous lines.

The F₂ and F₂:3 populations were grown at the Hangzhou Experiment Station of China Rice Research Institute (CNRRI), Zhejiang (N 30°32', E 120°12'), China, and the Lingshui Experiment Station of CNRRI, Hainan (N 18°48', E 110°02'), China, in the summer and winter of 2014. The NIL-F₂ populations and two sets of NILs were planted at the Hangzhou Experiment Station of CNRRI in the summer of 2015, 2016 and 2017. The F₂ and NIL-F₂ populations were planted with 20 cm between plants and 30 cm between rows. The F₂:3 families and two sets of NILs were grown in a randomized complete block design with two replications, five rows per plot, 8 plants per row, 20 cm between plants within each row and 30 cm between rows. The field management followed the standard agronomic practices.

Grain size trait evaluation

For the F₂ and NIL-F₂ populations, the plants were individually harvested for trait evaluation. For the F₂:3 population and NILs-*qGL7*-Nip and NILs-*qGL7*-IL188, ten plants in each line were harvested in bulk for trait evaluation. Five grain size traits were evaluated in each population. For grain length (GL), grain width (GW) and grain thickness (GT), 20 full-filled rice grains were randomly selected and individually measured using an electronic digital display vernier caliper. The averaged values of the 20 grains were used for data analysis. The

grain length-width ratio (LWR) is equal to GL divided by its GW. Thousand grain weight (TGW) was evaluated by measuring the weight of 200 randomly selected full-filled grains per F2 plant. The phenotypic evaluations of F2:3 family, NIL-F2 population and NIL lines were the same as those for F2 plants described above.

Scanning Electron Microscopy

The spikelets of NIL-*qGL7*Nip and NIL-*qGL7*IL188 were collected at maturity stage. The samples were fixed in FAA solution (formalin:glacial acetic acid: ethanol in 1:1:18 ratio by volume) at 4 °C for 24h, then dehydrated by a graded ethanol series, and were dried by critical-point drying method. The samples were observed under the scanning electron microscope (HITACHI, S-3000N). The spikelet epidermal cell size was measured using image J software.

DNA extraction and molecular marker analysis

DNA was extracted from fresh leaves samples following the CTAB method (Murray and Thompson 1980) with minor modifications. A total of 512 SSR markers with good genome coverage were selected to detect the polymorphisms between parents W303 and Nipponbare, 185 of which distributed across all 12 chromosomes showed polymorphisms between the two parents. Furthermore, 30 polymorphic SSR markers between IL188 and Nipponbare were used to genotype the F2 and F2:3 populations. Sixteen markers were used for fine mapping (Table S2).

Linkage map construction and data analysis

A genetic linkage map was constructed using MAPMAKER/EXP version 3.0 (Lander et al. 1987). The Kosambi mapping function (Kosambi, 1944) was used to transform the recombination frequency into cM. Composite interval mapping (CIM) was carried out to scan the introgressive genomic regions for putative QTLs using Windows QTL Cartographer 2.5 (<http://statgen.ncsu.edu/qtlcart/WQTLCart.htm>). The LOD threshold of 2.5 was used for declaring the presence of a putative QTL in a given genomic region. Nomenclature of QTLs was conducted as described by McCouch et al. (1997).

Phenotypic differences between IL188 and Nipponbare and between two homozygous lines in the NIL populations were compared using the student's test. Correlation analysis of grain size traits were performed using SPSS software.

Additional Files

Supplementary information

Supplementary Table S1: The genomic background of the R1-R4 populations.

Supplementary Table S2: Primers used for fine mapping.

Supplementary Table S3: The variations of NIL-*qGL7*IL188 in the 261-kb region compared with the Nipponbare reference genome.

Supplementary Figure S1: Comparison of grain size and cell number in the outer spikelet hulls along the vertical and lateral direction between NIL-*qGL7*Nip and NIL-*qGL7*L188. Scale bar, 1 mm. A, Mature grains of NIL-*qGL7*L188 (left) and NIL-*qGL7*Nip (right). B-C, The cell number in the longitudinal and lateral direction of outer spikelet hulls.

Declarations

Acknowledgments

We wish to thank Dr. Zhang Zhenhua for careful corrections and valuable suggestions on the revision.

Authors' Contributions

YF designed the experiments; XY, YW, YY, MZ, HY, QX, SW, and XN performed experiments and analyzed the data; YF and XW wrote the manuscript. All authors read and approved the final version of the manuscript.

Funding

This study was supported by the Major Scientific and Technological Project for New Varieties Breeding of Zhejiang Province (2016C02050-6-1), Zhejiang Provincial Natural Science Foundation of China (LY15C130005).

Availability of Data and Materials

The datasets supporting the conclusions of this article are included within the article.

Consent for Publication

All authors are consent for publication.

Competing interests

The authors declare that they have no competing interests.

Abbreviations

QTL: Quantitative trait locus; IL: Introgression line; RHL: Residual heterozygous line; SSR: Simple sequence repeat; GWAS: Genome wide association study; MAS: Molecular assisted selection; NIL: Near isogenic line

References

Amante-Bordeos A, Sitch LA, Nelson R, Dalmacio RD, Oliva NP, Aswidinnoor H, Leung H (1992) Transfer of bacterial blight and blast resistance from the tetraploid wild rice *Oryza minutata* cultivated rice, *Oryza sativa*. Theor Appl Genet 84:345–354

- Bai XF, Luo LJ, Yan WH, Kovi MR, Zhan W, Xing YZ (2010) Genetic dissection of rice grain shape using arecombinant inbred line population derived from two contrasting parents and fine mapping a pleiotropic quantitative trait locus *qGL7*. *BMC genet* 11:16
- Brar DS, Khush GS (1997) Alien introgression in rice. *Plant Mol Biol* 35:35–47
- Chen Y, Xu YY, Luo W, Li WX, Chen N, Zhang DJ, Chong K (2013) The F-box protein OsFBK12 targets OsSAMS1 for degradation and affects pleiotropic phenotypes, including leaf senescence in rice. *Plant Physiol* 163:1673–1685
- Feldman RM, Correll CC, Kaplan KB, Deshaies RJ (1997) A complex of Cdc4p, Skp1p, and Cdc53p/cullin catalyzes ubiquitination of the phosphorylated CDK inhibitor Sic1p. *Cell* 91:221–230
- Huang R, Jiang L, Zheng J, Wang T, Wang H, Huang Y, Hong Z (2013) Genetic bases of rice grain shape: so many genes, so little known. *Trends Plant Sci* 18:218–226
- Ikeda M, Miura K, Aya K, Kitano H, Matsuoka M (2013) Genes offering the potential for designing yield-related traits in rice. *Curr Opin Plant Biol* 16:213–220
- Huang RY, Jiang LR, Zheng JS, Wang TS, Wang HC, Huang YM, Hong ZJ (2013) Genetic bases of rice grain shape: so many genes, so little known. *Trends Plant Sci* 18:218–226
- Ishimaru K, Hirotsu N, Madoka Y, Murakami N, Hara N, Onodera H, Kashiwagi T, Ujiie K, Shimizu B, Onishi A, Miyagawa H, Katoh E (2013) Loss of function of the IAA-glucose hydrolase gene *TGW6* enhances rice grain weight and increases yield. *Nat Genet* 45(6):707–711
- Kashif H, Zhang YX, Workie A, Aamir R, Adil A, Md. Hasanuzzaman R, Wang H, Shen XH, Cao LY, Cheng SH (2020) Association mapping of quantitative trait loci for grain size in introgression line derived from *Oryza rufipogon*. *Rice Sci* 27(3):246–254
- Kosambi, DD (1944) The estimation of map distance from recombination values. *Ann Eugen* 12:172–175.
- Lander ES, Green P, Abrahamson J, Barlow A, Daly MJ, Lincoln SE, Newburg L (1987) MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1:174–181
- Mallikarjuna Swamy BPM, Kaladhar K, Shobha Rain N, Prasad GSV, Viraktamath BC, Ashok Reddy G, Sarla N (2012) QTL analysis for grain quality traits in 2 BC₂F₂ populations derived from crosses between *Oryza sativa* cv Swarna and 2 accessions of *O. nivara*. *J Hered* 103(3):442–452
- Mao HL, Sun SY, Yao JL, Wang CR, Yu SB, Xu CG, Li XH, Zhang QF (2010) Linking differential domain functions of the GS3 protein to natural variation of grain size in rice. *Proc Natl Acad Sci USA* 107:19579–19584
- McCouch SR, Cho YG, Yano M, Paul E, Blinstrub M, Morishima H, Kinoshita T (1997) Report on QTL nomenclature. *Rice Genet Newsl* 14:11–13

Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res* 8:4321–4325

Oka HI (1988) Origin of cultivated rice. *Developments in Crop Science*. Vol. 14, Japan Scientific Society Press okyo and Elsevier. Amsterdam.

Patton EE, Willerns AR, Tyers M (1998) Combinatorial control in ubiquitin-dependent proteolysis: don't Skp the F-box hypothesis. *Trends Genet* 14:236–243

Qi L, Sun Y, Li J, Sun L, Zheng XM, Wang X, Li K, Yang Q, Qiao W (2017) Identify QTLs for grain size and weight in common wild rice using chromosome segment substitution lines across six environments. *Breeding Sci* 67:472–482

Qi P, Lin YS, Song XJ, Shen JB, Huang W, Shan JX, Zhu MZ, Jiang LW, Gao JP, Lin HX (2012) The novel quantitative trait locus *GL3.1* controls rice grain size and yield by regulating Cyclin-T1;3. *Cell Res* 22(12):1666–1680

Qiu XJ, Gong R, Tan YB, Yu SB (2012) Mapping and characterization of the major quantitative trait locus *qSS7* associated with increased length and decreased width of rice seeds. *Theor Appl Genet* 125(8):1717–1726

Rahman ML, Chu SH, Choi M, Qiao YL, Jiang WZ, Piao R, Khanam S, Cho Y, Jeung J, Jena KK, Koh H (2007) Identification of QTLs for some agronomic traits in rice using an introgression line from *Oryza minuta*. *Mol Cells* 24(1): 16–26

Sakai H, Itoh T (2010) Massive gene losses in Asian cultivated rice unveiled by comparative genome analysis. *BMC Genom* 11:121

Second G (1982) Origin of the genic diversity of cultivated rice (*Oryza spp.*): study of the polymorphism scored at 40 isozyme loci. (*Jpn J Genet*) 57:25–57

Shao GN, Wei XJ, Chen ML, Tang SQ, Luo J, Jiao GA, Xie LH, Hu PS (2012) Allelic variation for a candidate gene for *GS7*, responsible for grain shape in rice. *Theor Appl Genet* 125(6):1303–1312

Shomura A, Izawa T, Ebana K, Ebitani T, Kanegae H, Konishi S, Yano M (2008) Deletion in a gene associated with grain size increased yields during rice domestication. *Nat Genet* 40(8):1023–1028

Si LZ, Chen JY, Huang XH, Gong H, Luo JH, Hou QQ, Zhou TY, Lu TT, Zhu JJ, Shangguan YY, Chen EW, Gong CX, Zhao Q, Jing YF, Zhao Y, Li Y, Cui LL, Fan DL, Lu YQ, Weng QJ, Wang YC, Zhan QL, Liu KY, Wei XH, An K, An G, Han B (2016) *OsSPL13* controls grain size in cultivated rice. *Nat Genet* 48:447–456

Skowyra, D, Craig KL, Tyers M, Elledge SJ, Harper JW (1997) F-box proteins are receptors that recruit phosphorylated substrates to the SCF ubiquitin-ligase complex. *Cell* 91:209–219

Song XJ, Ashikari M (2008) Toward an optimum return from crop plants. *Rice* 1:135–143

- Song XJ, Huang W, Shi M, Zhu MZ, Lin HX (2007) A QTL for rice grain width and weight encodes a previously unknown RING-type E3 ubiquitin ligase. *Nat Genet* 39:623–630
- Sun CQ, Wang XK, Li ZC, Yoshimura A, Iwata N (2001) Comparison of the genetic diversity of common wild rice (*Oryza rufipogon* Griff.) and cultivated rice (*O. sativa* L.) using RFLP markers. *Theor Appl Genet* 102:157–162
- Swamy BPM, Kaladhar K, Rani NS, Prasad GSV, Viraktamath BC, Reddy GA, Sarla N (2012) QTL analysis for grain quality traits in 2 BC₂F₂ populations derived from crosses between *Oryza sativa* cv Swarna and 2 accessions of *O. nivara*. *J Hered* 103(3):442–452
- Tanksley SD, Nelson JC (1996) Advanced backcross QTL analysis: a method for the simultaneous discovery and transfer of valuable QTLs from unadapted germplasm into elite breeding lines. *Theor Appl Genet* 92:191–203
- Thomson MJ, Tai TH, McClung AM, Lai XH, Hinga ME, Lobos KB, Xu Y, Martinez CP, McCouch SR (2003) Mapping quantitative trait loci for yield, yield components and morphological traits in an advanced backcross population between *Oryza rufipogon* and the *Oryza sativa* cultivar Jefferson. *Theor Appl Genet* 107:479–493
- Tian F, Li DJ, Fu Q, Zhu ZF, Fu YC, Wang XK, Sun CQ (2006) Construction of introgression lines carrying wild rice (*Oryza rufipogon* Griff.) segments in cultivated rice (*Oryza sativa* L.) background and characterization of introgressed segments associated with yield-related traits. *Theor Appl Genet* 112:570–580
- Wan XY, Wan JM, Weng JF, Jiang L, Bi JC, Wang CM, Zhai HQ (2005) Stability of QTLs for rice grain dimension and endosperm chalkiness characteristics across eight environments. *Theor Appl Genet* 110:1334–1346
- Wang SK, Li S, Liu Q, Wu K, Zhang JQ, Wang SS, Wang Y, Chen XB, Zhang Y, Gao CX, Wang F, Huang HX, Fu XD (2015a) The *OsSPL16-GW7* regulatory module determines grain shape and simultaneously improves rice yield and grain quality. *Nat Genet* 47(8):949–954
- Wang SK, Wu K, Yuan QB, Liu XY, Liu ZB, Lin XY, Zeng RZ, Zhu HT, Dong GJ, Qian Q, Zhang GQ, Fu XD (2012) Control of grain size, shape and quality by *OsSPL16* in rice. *Nat Genet* 44(8):950–954
- Wang YX, Xiong GS, Hu J, Jiang L, Yu H, Xu J, Fang YX, Zeng LJ, Xu EB, Xu J, Ye WJ, Meng XB, Liu RF, Chen HQ, Jing YH, Wang YH, Zhu XD, Li JY, Qian Q (2015b) Copy number variation at the *GL7* locus contributes to grain size diversity in rice. *Nat Genet* 47(8):944–948
- Wang ZY, Second G, Tanksley SD (1992) Polymorphism and phylogenetic relationships among species in the genus *Oryza* as determined by analysis of nuclear RFLPs. *Theor Appl Genet* 83:565–581
- Xiao JH, Li JM, Grandillo S, Ahn SN, Yuan LP, Tanksley SD, McCouch SR (1998) Identification of trait-improving quantitative trait loci alleles from a wild rice relative, *Oryza rufipogon*. *Genetics* 150:899–909
- Xie X, Song MH, Jin F, Ahn SN, Suh JP, Hwang HG, Kim YG, McCouch SR (2008) Fine mapping of a yield-enhancing QTL cluster associated with transgressive variation in an *Oryza sativa* × *O. rufipogon* cross. *Theor Appl Genet* 116:613–622

Xie XB, Song MH, Jin FX, Ahn SN, Suh JP, Hwang HG, McCouch SR (2006) Fine mapping of a grain weight quantitative trait locus on rice chromosome 8 using near-isogenic lines derived from a cross between *Oryza sativa* and *Oryza rufipogon*. *Theor Appl Genet* 113:885–894

Xiong YZ, Zhang QF (2010) Genetic and molecular bases of rice yield. *Annu Rev Plant Biol* 61:421–442

Xu F, Fang J, Ou SJ, Gao SP, Zhang FX, Du L, Xiao YH, Wang HR, SunXH, Chu JF, Wang GD, Chu CC (2015) Variations in *CYP78A13* coding region influence grain size and yield in rice. *Plant Cell & Environ* 38(4):800–811

Xue P, Zhang YX, Lou XY, Zhu AK, Chen YY, Sun B, Yu P, Cheng SH, Cao LY, Zhan XD (2019) Mapping and genetic validation of a grain size QTL *qGS7.1* in rice (*Oryza sativa* L.). *J Integr Agric* 18(8):1838–1850

Yoon DB, Kang KH, Kim HJ, Ju HG, Kwon SJ, Suh JP, Jeong OY, Ahn SN (2006) Mapping quantitative trait loci for yield components and morphological traits in an advanced backcross population between *Oryza grandiglumis* and the *O. sativa japonica* cultivar Hwaseongbyeon. *Theor Appl Genet* 112:1052–1062

Yu JP, Miao JL, Zhang ZY, Xiong HY, Zhu XY, Sun XM, Pan YH, Liang YT, Zhang Q, Rehman RMA, Li JJ, Zhang HL, Li ZC (2018) Alternative splicing of *OsLG3b* controls grain length and yield in japonica rice. *Plant Biotechnol J* 16(9):1667–1678

Yun YT, Chung CT, Lee YJ, Na HJ, Lee JC, Lee SG, Lee KW, Yoon YH, Kang JW, Lee HS, Lee JY, Ahn SN (2016) QTL mapping of grain quality traits using introgression lines carrying *Oryza rufipogon* chromosome segments in japonica rice. *Rice* 9:62

Zhang Q, Yao GX, Hu GL, Chen C, Tang B, Zhang HL, Li ZC (2012) Fine Mapping of *qTGW3-1*, a QTL for 1000-Grain Weight on Chromosome 3 in Rice. *J Integr Agri* 11(6):879–887

Zhang XJ, Wang JF, Huang J, Lan HX, Wang CL, Yin CF, Wu YY, Tang HJ, Qian Q, Li JY, Zhang HS (2012) Rare allele of *OsPPKL1* associated with grain length causes extra-large grain and a significant yield increase in rice. *Proc Natl Acad Sci USA* 109:21534–21539

Zhao DS, Li QF, Zhang CQ, Zhang C, Yang QQ, Pan LX, Ren XY, Lu J, Gu MH, Liu QQ (2018) *GS9* acts as a transcriptional activator to regulate rice grain shape and appearance quality. *Nat Commun* 9(1):1240

Zuo JR, Li JY (2014) Molecular genetic dissection of quantitative trait loci regulating rice grain size. *Ann Rev Genet* 48:99–118

Figures

Figure 1

Comparison of whole-plant rice and grain performance between IL188 and Nipponbare. A, Plant type of IL188 (left) and Nipponbare (right). B, Paddy and brown rice grains of IL188 (left) and Nipponbare (right). Bars: 10 cm in A, 6 mm in B.

Figure 2

A scheme showing how plants materials were developed.

Figure 3

Genetic linkage map showing QTL positions detected in the F₂ and F_{2:3} populations. White and black shapes indicate F₂ and F_{2:3} population, respectively.

Figure 4

Frequency distribution of five grain shape traits in the F₂ and F_{2:3} populations. (P₁: Nipponbare, P₂: IL188). The vertical axis of each figure represents the number of F₂ and F_{2:3} plants, blue bars and red bars indicate F₂ and F_{2:3}, respectively.

Figure 5

Genotypes compositions of NIL populations in the target region. NIL, near-isogenic line.

Figure 6

Comparison of grain size and spikelets epidermal cells between NIL-qGL7Nip and NIL-qGL7IL188. A, Mature grains of NIL-qGL7Nip and NIL-qGL7IL188. Scale bar, 6 mm. B-D, Grain length (GL), grain width (GW), and 1000-grain weight (TGW) for NIL-qGL7Nip and NIL-qGL7IL188. Data are given as mean $\pm$ (n = 20). ** indicate significant difference at 0.01 level. E-F, Outer epidermal cells of grain hulls of NIL-qGL7Nip and NIL-qGL7IL188. Bars = 100 μ m. G-H, The average length and width of outer epidermal cells. (n = 10). I-J, Inner epidermal cells of grain hulls of NIL-qGL7Nip and NIL-qGL7IL188. Bars = 100 μ m. K-L, The average length and width of inner epidermal cells. (n = 10).

Supplementary Files

This is a list of supplementary files associated with this preprint. Click to download.

- [Supplementarytable.docx](#)
- [SupplementaryFig.1.tif](#)